



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

LANE

MEDICAL



LIBRARY

Gift

Dr. H. C. McClenahan



A
SYSTEM OF SURGERY.

VOL. IV.

**DISEASES OF THE ORGANS OF LOCOMOTION, OF
INNERVATION, OF DIGESTION, OF RESPIRATION,
AND OF THE URINARY ORGANS.**

LONDON: PRINTED BY
SPOTTISWOODE AND CO., NEW-STREET SQUARE
AND PARLIAMENT STREET

A
SYSTEM OF SURGERY

THEORETICAL AND PRACTICAL

IN

TREATISES BY VARIOUS AUTHORS.

EDITED BY

T. HOLMES, M.A. CANTAB.

SURGEON AND LECTURER ON SURGERY AT ST. GEORGE'S HOSPITAL
MEMB. CORRESP. DE LA SOCIÉTÉ DE CHIRURGIE DE PARIS.

SECOND EDITION, IN FIVE VOLUMES.

With Illustrations.

*VOL. IV.—DISEASES OF THE ORGANS OF LOCOMOTION, OF
INNERVATION, OF DIGESTION, OF RESPIRATION,
AND OF THE URINARY ORGANS.*

LANE LIBRARY

LONDON:
LONGMANS, GREEN, AND CO.
1870.

The right of translation is reserved.

YSAABU! 3BA!

74
1.4
870

CONTENTS

OF

THE FOURTH VOLUME.



DISEASES OF THE ORGANS OF LOCOMOTION (continued).

DISEASES OF THE JOINTS.

By ATHOL A. JOHNSTONE, Esq.

LATE SURGEON TO THE HOSPITAL FOR SICK CHILDREN.

PART I.

DISEASES COMMON TO ALL THE JOINTS.

DISEASES OF SYNOVIAL MEMBRANES :	PAGE
Inflammation of synovial membranes	2
Simple synovitis; acute	3
Acute articular abscess	6
Chronic synovitis	7
Treatment of synovitis	8
Dropsy of joints	13
Scrofulous synovitis (pulpy degeneration)	16
Rheumatic synovitis	22
Chronic osteo-arthritis (chronic rheumatic arthritis— rheumatic gout)	22
Gouty synovitis	29
Pyæmic "	33
Gonorrhœal synovitis (gonorrhœal rheumatism)	34
Syphilitic "	36
Loose cartilages in joints	36
DISEASES OF THE ARTICULAR EXTREMITIES OF THE BONES :	
Simple inflammation and its consequences	41
Scrofulous inflammation	42
Rheumatic and syphilitic inflammation	51

7788

Rachitic affections	
Tumours of the articular ends of the bones	
ABNORMAL CONDITIONS OF THE ARTICULAR CARTILAGES :	
Physiological anatomy of articular cartilage	
Hypertrophy	
Atrophy	
Gouty deposits	
Fatty degeneration	
Ossification	
Ulceration	
DISEASES OF OTHER TISSUES IN AND AROUND THE JOINTS :	
Of the ligamentous structures	
" muscles	
" areolar and adipose tissues	
" synovial bursæ	
" integuments	
ANCHYLOSIS	
ARTICULAR NEURALGIA	
WOUNDS OF JOINTS	

PART II.

DISEASES OF INDIVIDUAL JOINTS.

Of the hip	
Acute arthritis	
Common synovitis	
Morbus coxarius (strumous synovitis and ostitis)	
Morbus coxæ senilis	
Neuralgia	
Of the pubic, sacro-iliac, and sacro-coccygeal joints	
" knee	
Of bursæ near the knee	
Of the ankle	
" joints of the foot	
" sterno-clavicular, acromio-clavicular, and sternal joints	
" shoulder	
" elbow	
" wrist and other articulations of the hand	
" temporo-maxillary joint	

DISEASE OF THE SPINE.

BY A. SHAW, ESQ.

CONSULTING SURGEON TO THE MIDDLESEX HOSPITAL.

Caries of the spine; nature of the morbid process	
Symptoms of the incipient stage of the disease	

	PAGE
Angular projection	104
Rigidity of the affected portion	105
Heat and swelling	106
Pain	106
Diagnosis from neuralgia of the spine	107
Description of angular deformity	109
Changes in chest and abdomen	112
Abscesses from disease of the spine	113
Anchylosis	115
Spurious anchylosis	116
Psoas and lumbar abscess	118
Psoas abscess	119
'Neck' of the abscess	121
Lumbar abscess	122
Spinal abscess in the neck	122
Varieties in spinal abscess	123
Bifurcation of the sac	123
Presenting above Poupart's ligament	123
" in gluteal region	123
" at external abdominal ring	123
Contents discharged by lung	124
Opening into intestine	125
" urinary bladder	125
Diagnosis of spinal abscess	126
Treatment of external spinal abscess	127
Affection of the spinal cord from extension of the disease	129
Disorganisation of the cord ; paraplegia	131
Spasmodic actions of the paralysed limbs	133
Diagnosis	134
Prognosis in paraplegia	134
Treatment of spinal disease	135
The prone-couch	137
Counter-irritants	138
Disease of the atlas and axis	138
Symptoms	143
Prognosis	145
Treatment	146
Necrosis of atlas	146

DISEASES AND INJURIES OF NERVES.

PART I.

NERVE-LESIONS AND THEIR MORE IMMEDIATE EFFECTS.

BY J. LOCKHART CLARKE, M.D. F.R.S.

Diseases of nerves	148
Inflammation ; acute	148

	PAGE
Inflammation ; chronic	150
" idiopathic	150
" traumatic	151
Treatment of neuritis	151
Ulceration of nerves	152
Tumours of nerves	153
Neuroma	154
Tubercle	158
Cancer	158
Hypertrophy of nerves	158
Atrophy "	159
Injuries of nerves	159
Contusions	159
Compression and distension	159
Laceration	161
Ligature	162
Complete division	163
Incomplete division	164
Puncture	164
Incised wounds	166
Foreign bodies in nerves	168
Treatment of injuries of nerves	169
Neuralgia	172
Nutritive and other changes resulting from injuries and diseases of nerves	173
Diminution of temperature in parts supplied	174
Effects on skin and its appendages	174
Alteration of secretions	177
Periodical swelling of joints and other parts	177
Atrophy and contraction of muscles	177
Treatment	181
Changes due to idiopathic affections of nerves	181

PART II.

REMOTER CONSEQUENCES OF NERVE-LESIONS.

BY C. E. BROWN-SÉQUARD, M.D., F.R.S.

INTRODUCTION—Kinds of symptoms dependent on lesions of nerves	184
SECTION I. Diseases of the nervous centres and other organs, caused by an injury to, or a disease of, a nerve	186
Epilepsy	186
Tetanus	188
Hysteria	190
Cataplexy	191
Chorea	191

FOURTH VOLUME.

ix

	PAGE
Hydrophobia	191
Tremulous movements	192
Rotatory convulsions	192
Local convulsions	192
Contracture of blood-vessels	193
Paralysis	193
Anæsthesia	196
Amaurosis	196
Deafness	196
Collapse	197
Loss of smell, taste, and hearing	197
Neuralgia	197
Delirium	198
Aphasia	199
Inflammation	199
Coma	200
Apoplexy	201
Muscular atrophy	201
Atrophy of the cellular tissue	201
Hypertrophy	202
Various alterations of nutrition and secretion	202
SECTION II. General features and rules of treatment of the various affections caused by an irritation of a nerve	203
Local treatment	206
General "	207

LOCOMOTOR ATAXY.

BY J. LOCKHART CLARKE, M.D., F.R.S.

symptoms	209
pathological anatomy	213
prognosis and treatment	214

DISEASES OF ORGANS OF DIGESTION AND RESPIRATION.

DISEASES OF THE TONGUE.

BY HOLMES COOTE, ESQ.

SURGEON TO ST. BARTHOLOMEW'S HOSPITAL.

ape and general anatomy of the tongue. Tongue-tie	215
hypertrophy and enlargement of the tongue	216

Acute inflammation (glossitis)
Abscess
Chronic ulceration
Nævus
Tumours of the tongue.
Ranula
Wounds, foreign bodies, &c.
Syphilitic affections
Cancer
Operations for cancer
Complete extirpation of the tongue.
Division of the gustatory nerve

DISEASES OF THE NOSE.

BY ARTHUR E. DURHAM, ESQ.

ASSISTANT-SURGEON TO GUY'S HOSPITAL.

I. Affections of the external parts
Acne rosacea (gutta rosacea vel rosea)
Integumentary hypertrophy (lipoma nasi)
Lupus
Lupus exedens (Noli me Tangere)
" non-exedens
" erythematosus
Cancerous affections of the skin, &c.
Rodent cancer
Epithelioma
Deformities and deficiencies
Absence of the nose
Occlusion and contraction of the nostrils.
Clefts and fissures
Deviations
II. Affections of the internal parts
Rhinoscopy
" anterior
" posterior
Nasal calculi; rhinolithes
Epistaxis
" plugging of the nares
Coryza
Catarrhal inflammation of the pituitary mucous membrane
" chronic
Hypertrophy of the pituitary membrane
Syphilitic coryza
" in the infant
Ozæna
The nasal douche
Syphilitic ulceration, caries and necrosis

	PAGE
Lupous and strumous ulceration with ozæna	282
Ulceration, caries, and necrosis from other causes	284
Idiopathic or constitutional ozæna	284
Glandular hypertrophy of mucous membrane	287
Affections of the septum	289
Blood-tumours	289
Acute abscess	290
Chronic abscess	291
New growths. Polypi	292
Papillary growths	292
Mucous, gelatinous or vesicular polypi	293
Method of removal	297
Fibrous polypi	305
Nasopharyngeal polypi	309
Operations for their removal	314
Cartilaginous growths	318
Osseous	319
' Osseous tumours of nasal fossæ and sinuses of face'	320
Malignant growths	322

SURGICAL DISEASES CONNECTED WITH THE TEETH.

BY S. JAMES A. SALTER, ESQ., F.R.S.

DENTAL SURGEON TO GUY'S HOSPITAL.

Subjects comprised in this essay	323
I. Alveolar abscess	324
Gum-boil	325
Pointing of alveolar abscess inside the mouth	327
" " remote from origin of abscess	327
" " on the face	328
Diagnosis of alveolar abscess	329
Treatment	331
II. Painful and difficult eruption of wisdom teeth	332
In the upper jaw	333
In the lower "	333
Spasmodic contraction of masseter	334
Suppuration and fistula from difficult eruption	334
Treatment	335
" of the spasmodic closure of the mouth	337
III. Tumours of the gum	
Epulis	338
Treatment	341
Congenital hypertrophy of the gum and alveoli	342
Treatment	344
Polypus of the gum	344
Treatment	345

	Vascular tumours	
	Treatment	
	Warty tumours of the gum	
IV.	Tumours of the tooth pulp	
	Polypus of the pulp	
	Treatment	
	Sensitive growth of pulp after fracture	
V.	Odontomes, or tumours of the hard tissues of the teeth	
	Enamel nodules	
	Exostosis	
	Hypertrophied, dilated fangs	
	Dentine excrescence	
	Warty teeth	
VI.	Abscess of the antrum	
	Varying extent of the antrum	
	Causes and symptoms of antral abscess	
	Treatment	
VII.	Dentigerous cysts	
	Pathology and mode of development	
	Causes of impaction of teeth	
	Summary of recorded cases of such cysts	
	Symptoms	
	Treatment	
VIII.	Alveolar or maxillary necrosis	
	Phosphorus disease	
	Causes	
	Symptoms	
	Treatment	
	Exanthematous jaw-necrosis	
	Treatment	
IX.	Hæmorrhage after extraction of teeth	
	Treatment	
X.	The application of obturators and false-palates in cleft-palate, &c.	
	Obturators	
	False-palates	

DISEASES OF THE MOUTH, PHARYNX, AND ŒSOPHAGUS.

BY GEORGE POLLOCK, ESQ.

SURGEON TO ST. GEORGE'S HOSPITAL.

OF THE MOUTH :

Quinsy ; acute inflammation of the tonsils	
Symptoms	
Causes	
Treatment	
Diagnosis from other inflammatory affections in this region	

	PAGE
Chronic inflammation and enlargement of the tonsils	404
Causes of enlarged tonsils	405
Forms " "	405
Effects " "	406
Pathological anatomy	407
Treatment	407
Excision of the tonsils	408
Malignant sore-throat (tonsillitis maligna)	410
Ulcerated "	411
Relaxed uvula	411
Removal of the uvula	412
Tumours of the palate	413
Cleft-palate (congenital)	414
Its mode of origin	414
Development of the mouth	414
Classification and description of different degrees	416
Varying width of cleft	420
Effects of cleft-palate	421
Treatment	422
Age for operation	422
Operation in cases of bifid uvula	424
" " cleft of soft palate	425
" " cleft of hard and soft palate	430
Treatment of acquired openings in the palate	436
Treatment after operation	438
Improvement in speech after operation	440
OF THE LIPS :	
Cracked lip	441
Aphthous ulceration	443
Nævus of the lips and cheeks	443
Cancer of the lip	446
Diagnosis from chancre	447
Treatment	448
Caustics in cancer of the lip	449
Cysts of the lip	450
Congenital cysts of the mouth	451
Other forms of cyst in this region	452
Treatment of these cysts	454
Solid tumours of the lips and mouth	455
TUMOURS OF THE JAWS :	
Cartilaginous tumours	457
Cystic "	457
Fibrous "	460
Recurrence of fibrous tumour	464
Myeloid tumours	465
Osseous "	466
Hypertrophy of the bones of the face	467
Vascular tumours	469
Mucous "	469
Cancerous tumours	469

OPERATIONS ON THE JAWS :

Removal of portions of the lower jaw	
" parts of its whole thickness	
" the whole of one side	
" the whole lower jaw	
" portions of the upper jaw	
" the whole upper jaw	
Osteo-plastic resection of the upper jaw	

OF THE PHARYNX :

Congenital defects	
Pharyngitis	
Abscess of the pharynx	
Ulceration	
Adhesion of the soft palate	
Dilatation of the pharynx	
Tumours	
" "	
Removal of such tumours	

OF THE ŒSOPHAGUS :

Dilatations and pouches of the œsophagus	
Contraction and stricture	
From fold of mucous membrane	
" cicatrization	
" simple chronic thickening	
" pressure of tumours	
" cancer	
Symptoms of stricture	
Treatment	
Passage of bougies	

DIPHTHERIA AND CROUP.

BY A. W. BARCLAY, M.D.

PHYSICIAN TO ST. GEORGE'S HOSPITAL.

DIPHTHERIA :

Relations of diphtheria to croup and scarlatina	
History	
Symptoms	
Albuminuria	
Paralysis	
Diphtheritic fever	
The fibrinous exudation	
Treatment	
Tracheotomy in diphtheria	
Ordinary sore-throat occurring in epidemics of diphtheria	

	PAGE
OUT :	
General characters	509
Diagnosis	510
Treatment	512
Tracheotomy in croup	513

DISEASES OF THE LARYNX.

BY ARTHUR E. DURHAM, Esq.

ASSISTANT-SURGEON TO GUY'S HOSPITAL.

THE LARYNGOSCOPE :

Invention of the instrument	514
Description of the laryngoscopes now in use	515
General arrangements for examination	517
The reversal of the laryngeal image	521
Management of difficulties in laryngoscopy	523
The general appearance of various parts of the larynx	527

GENERAL REMARKS ON THE TREATMENT OF DISEASES OF THE LARYNX :

Application of solid substances	529
" solutions	529
" atomised fluids	530
Spray producers	531
Various solutions in use in diseases of the larynx	532
Scarification of the mucous membrane	533
Galvanisation of the vocal cords	534

CATARRHAL LARYNGITIS :

Acute catarrhal laryngitis ; symptoms	534
" " laryngoscopic appearances	536
" " post-mortem "	537
" " diagnosis	537
" " treatment	539
Chronic catarrhal laryngitis	542
" laryngoscopic appearances	543
" treatment	544

CHRONIC GLANDULAR LARYNGITIS. FOLLICULAR DISEASE. DYSPHONIA CLERICORUM :

Causes of this affection	547
Symptoms and laryngoscopic appearances	548
Treatment	550

PHTHISICAL LARYNGITIS :

Frequency of this disease in phthisis	551
Symptoms and laryngoscopic appearances	552
Course and duration	554
Treatment	555

SYPHILITIC LARYNGITIS :

Various affections of the larynx in secondary syphilis
Diagnosis of "syphilitic" affection	.	.	.	"	.
Treatment

ERYSIPELATOUS LARYNGITIS

DIFFUSE CELLULAR LARYNGITIS. ACUTE ŒDEMATOUS LARYNGITIS

OTHER FORMS OF LARYNGITIS :

Exanthematous laryngitis.	From measles	.	.	.
"	"	"	typhoid fever	.
"	"	"	small-pox	.
"	"	"	scarlet fever	.
Gouty laryngitis

NEW GROWTHS. POLYPI, &c. :

Various forms of these growths
Papillomatous or warty growths
Fibrous and fibro-cellular	"	.	.	.
Adenomatous or glandular	"	.	.	.
Cystic tumours
Cartilaginous and osseous growths
Cancerous tumours
Symptoms of tumour in the larynx
Treatment	"	"	.	.
A. Operations through the natural passages :				
1. Caustics
2. Removal by forceps
3. By the wire snare
4. Galvano-caustic method
5. Removal by the knife
6. Puncture
B. Operations by external incision and section of cartilages

NERVO-MUSCULAR AFFECTIONS :

Nervous or functional aphonia
Atrophy, idiopathic, or from disease, of the laryngeal mus

SPASM OF THE GLOTTIS

DISEASES OF THE INTESTINES.

BY GEORGE POLLOCK, ESQ.

SURGEON TO ST. GEORGE'S HOSPITAL.

Abdominal abscess
Fæcal abscess
Treatment
Obstruction of the intestines
Causes of obstruction

	PAGE
Commonly acute. { Congenital malformation	603
{ Lodgment of foreign bodies	604
{ Twisting or 'dislocation' of bowel	604
{ Strangulation by loops, false membranes, diverticula, &c. .	609
{ " " peritoneal pouches, the foramen of Win-	
{ slow, &c.	610
{ Intussusception	612
{ Spontaneous cure of intussusception	617
{ Treatment	618
Commonly chronic. { Habitual constipation—accumulation of feces	619
{ Injury	621
{ Chronic peritonitis (tubercular) and abscess	621
{ Stricture from chronic disease of the intestine	621
{ " " cancer of the bowel or tumours in the	
{ neighbourhood	624
{ Relative frequency of these different forms	627
{ Diagnosis of the seat of obstruction and of its nature	627
{ Treatment of obstruction	630
{ Colotomy	632
{ Cases in which colotomy is justifiable	632
{ Not justifiable in intussusception	633
{ Insufflation and injection in intussusception	633
{ Indications and contra-indications to colotomy in	
{ its various forms	634
{ Opening the peritoneal sac (Littre's operation)	636
{ Lumbar colotomy (Amussat's and Calissen's operations) .	638
{ Paracentesis of the abdomen	641

HERNIA.

By J. BIRKETT, Esq.

SURGEON TO GUY'S HOSPITAL.

PART I.

PATHOLOGY AND TREATMENT OF HERNIA IN GENERAL.

General considerations.	Proportions of the two sexes	644
	Frequency of hernia at different ages	645
	Proportion of population affected with hernia	646
	Influence of occupation	647
	Influence of hereditary conformation and	
	other circumstances	648
	Immediate cause of hernia	650
Symptoms of hernia in general		650
Prognosis		652

Nomenclature of hernia
The essential parts of the tumour
The inciting causes of hernia
Congenital patency of the vaginal process of the peritoneum
Description of the changes which take place in this process
Condition of the vaginal process at birth
Of the gradual development of the hernial sac
The period of formation
„ organisation
„ contraction
The two kinds of hernial sac, congenital and acquired
The treatment of reducible hernia—palliative measures
Trusses
Radical cure of hernia
Morbid conditions of the hernia
Permanent irreducibility
Obstruction or distension
Inflammation
Strangulation
Immediate cause
Structural changes
Gangrene
Ulceration in the line of the stricture
Liability of recent hernia to acute strangulation
Morbid conditions produced by manual force
Artificial anus
Morbid conditions of the coverings of the hernia
Characters of the serum in the sac
Constitutional symptoms induced by the morbid state
of the alimentary canal above
Effects of obstruction
Constitutional symptoms of strangulation
Necessity of searching for hernial tumours
certain cases
Nature of the vomit
Symptoms of prostration
„ peritonitis
„ collapse
Pain as a symptom in strangulated hernia
Prognosis of strangulated hernia before operation
Necessity for liberating the bowel as soon as possible
Treatment of irreducible hernia
Injurious effects often produced by the taxis
Treatment before vomiting commences
Method of applying the taxis
Treatment during the stage of vomiting
The use of chloroform
Symptoms of rupture of the bowel by taxis
Cold applications
Purgatives and enemata
Other means of treatment
Injuries inflicted on the hernia—by the taxis

	PAGE
Injuries inflicted on the hernia—by accident	701
Injuries of the sac—by displacement ‘réduction en bloc’	703
” ” by laceration	704
Injuries of the coverings of the sac	705
Intra-parietal sacs	706
Operation for strangulated hernia—general considerations	706
Description of the operation	708
On opening the hernial sac	711
The instruments required in the operation	714
Treatment of the strangulated intestine	715
Perforation and wound of the gut	716
Difficulties in reduction—from bulk of intestine	717
” ” ” adhesions	717
” ” ” depth of mouth of sac	718
Treatment of the omentum	718
” cases of double hernia	720
” the case after operation	720
” the wound	722
Pregnosis of operations for strangulated hernia	723

PART II.

SPECIAL KINDS OF HERNIA.

Classification	725
Main in the epigastric region. Diaphragmatic	726
Epigastric	726
” mesogastric ” Ventral	727
Umbilical	727
Radical cure	730
Operation	730
Lumbar	731
” hypogastric ” Inguinal, its proportion to femoral hernia	731
Oblique, its varieties	733
Hernia into the vaginal process	734
Hernia into the funicular portion of the vaginal process	739
Frequency of these forms of hernia	741
Hour-glass contraction of the sac of oblique inguinal hernia	743
Infantile hernia	749
Inguinal-scrotal hernia of slow formation	750
Table of characteristics of the three varieties of oblique hernia	752
Diagnosis of tumours of the scrotum	753
Table of diagnostic characters	755

	Radical cure of reducible inguinal hernia
	The taxis
	The operation
	Oblique inguinal hernia in the femur
	Direct inguinal hernia
	Abnormal inguinal hernia
Femoral or crural hernia	
Varieties	
Development of its sac	
Diagnostication	
Trusses	
Frequency with reference to sex, age, and pregnancy	
Morbidity conditions	
The taxis	
The operation	
Wound of the obturator artery	
Obturator hernia	
Analysis of recorded cases	
Complications	
Diagnosis	
Treatment and operation	
Perineal hernia	
Pudendal hernia	
Vaginal, and ischiatic hernia	

DISEASES OF THE RECTUM.

By H. SMITH, Esq.

ASSISTANT-SURGEON TO KING'S COLLEGE HOSPITAL.

Hæmorrhoids	
External hæmorrhoids	
Their anatomy	
Treatment	
Internal hæmorrhoids	
Various forms, and structure of each	
Treatment of internal piles	
Ligature	
Clamp and cautery	
Nitric acid	
Ecraseur	
Prolapsus of the rectum	
Various conditions—complicated with piles	
Treatment	
Fistula in ano	
Formation of fistula	
Various forms of fistula—complete and incomplete	
Horseshoe fistula	

	PAGE
Causes and diagnosis of fistula	827
Treatment of fistula	828
Operation for fistula	830
Stricture of the rectum and anus	834
Treatment	835
Fissure of the anus	835
Stricture of the rectum (simple)	841
Anatomy	841
Symptoms	844
Diagnosis	846
Causes	847
Treatment	848
Dilatation	849
Incision	851
Ulcer of the rectum	852
Polypus of the rectum	858
Warty polypus	860
Villous tumour	862
Hæmorrhoids	863
Proctalgia of the rectum	865

DISEASES OF THE URINARY ORGANS.

BY SIR H. THOMPSON,

SURGEON TO UNIVERSITY COLLEGE HOSPITAL.

Division of the subject	868
Formations of the kidney	869
Acute and chronic nephritis	869
Symptoms	870
Diagnosis from	
1. Calculous nephralgia	871
2. Acute calculous pyelitis	871
3. Perinephritis: caries of the spine	871
4. Rheumatism or lumbago	872
5. Renal neuralgia	872
6. Acute cystitis	872
7. Calculus of the bladder	872
Treatment of acute nephritis	873
„ calculous nephralgia	874
„ chronic pyelitis	874
„ abscesses from nephritis	875
„ chronic nephritis	875
Haematuria	876
Diagnosis of its source	877
Treatment	878
Retention of urine	879
Obstructions of the ureters	880

Malformations of the bladder :	Deficiency	
	Multiplicity, &c.	
	Extroversion	
Synopsis of the diseases of the bladder		
I. CONDITIONS INVOLVING STRUCTURAL CHANGES :		
Acute cystitis		
Causes and symptoms		
Treatment		
Chronic cystitis		
„ with increased secretion (catarrh of bladder)		
Treatment of chronic cystitis		
Suppuration and ulceration of the bladder		
Vesico-intestinal fistula		
Hypertrophy and sacculation of the bladder, dilatation, &c.		
Tumours of the bladder.	1. Fibrous	
	2. Villous	
	3. Epithelial	
	4. Malignant	
Tubercle of the bladder		
Bar at the neck of the bladder		
Hernia and inversion of the bladder		
II CONDITIONS NOT NECESSARILY INVOLVING STRUCTURAL CHANG		
Paralysis of the bladder		
Atony from over-distension		
Irritable bladder		
Spasm of the bladder		
Perverted sensibility of the neck of the bladder, or neuralgi		
Incontinence of urine		
1. In childhood and youth		
2. In the adult		
Habitual engorgement of the bladder and overflow of urine		
Affections of the prostate gland :		
Malformations		
Acute prostatitis		
Chronic inflammation of the prostate		
Abscess of the prostate		
Enlarged prostate		
Symptoms		
Diagnosis		
Treatment		
Atrophy of the prostate		
Cancer „		
Tubercle „		
Cysts „		
Affections of the urethra :		
Congenital aberrations and malformations		
Hypospadias and epispadias		
Contractions of the urethra (stricture) :		
Organic or permanent		
Anatomical classification : viz.		
Linear stricture		

	PAGE
Annular	941
Indurated annular	941
Irregular or tortuous	941
Number in one urethra	942
Origin, nature, and seat of stricture	942
Obstructions from false membranes	943
Degree of contraction	943
Locality of stricture	943
Causes of organic stricture	945
Symptoms—Local	946
" Retention and engorgement	947
" Constitutional	948
Strictures classified according to their symptoms, as simple, sensitive or irritable, and contractile or recurring	949
Diagnosis of stricture	949
Method of passing the catheter	951
The endoscope	954
Treatment of stricture—gradual dilatation	956
False passages	959
Chloroform in dilatation of stricture	959
Continuous or permanent dilatation	960
Dilatation by mechanical means—forcible	961
Holt's and Thompson's instruments for this purpose	962
Constitutional treatment	965
Treatment by caustics	965
Incision. 1. Internal urethrotomy	967
From before backwards	968
From behind forwards	970
2. External urethrotomy	973
Perineal section	974
External division (Syme's operation)	975
Results of external urethrotomy	977
Spasm of the urethra (spasmodic stricture)	978
Inflammatory stricture	980
Tumours in the urethra	980
Urinary abscess	981
Urinary fistula. 1. Simple fistulæ	983
2. Fistulæ passing through indurated tissues	984
3. Fistulæ with destruction of the urethral walls	987
Treatment. Urethroplasty	988
Retention of urine	989
Treatment	989
Forced catheterism	991
Perineal incision	991
Puncture of the bladder by the rectum	992
" " above the pubes	993
" " by Dieulafoy's aspirator	994
" " through the pubic symphysis	994
Treatment of retention from enlarged prostate	996
Extravasation of urine	998
Treatment	999
Rupture of the bladder from obstruction	1000

URINARY CALCULI AND LITHOTOMY.

BY A. POLAND, Esq.

SURGEON TO GUY'S HOSPITAL.

The sources of morbid deposits in the urine	
Urinary deposits.	Uric acid
	The urates
	Uric oxide
	The oxalates
	Cystic oxide
	Phosphatic deposits
	Carbonate of lime
Formation of calculus	
Immediate causes of its formation	
Classification of calculi	
Physical characters of calculi	
Characters on section	
Chemical tests for calculi	
Kinds of calculus.	1. Uric acid
	2. Urate of ammonia
	3. Uric or xanthic oxide
	4. Oxalate of lime
	5. Cystic oxide
	6. Phosphate of lime
	7. Triple phosphate
	8. Fusible calculus
	9. Carbonate of lime
	10. Fibrinous calculus
	11. Urostealith
	12. Blood-calculi
Predisposing causes of calculus	
Sex and age. Statistics	
Other predisposing causes	
Calculus in the bladder. Symptoms	
Sounding for stone	
Prognosis of stone in the bladder	
The natural progress of a case to its fatal termination	
Treatment. General	
Lithontriptics	
Electrolysis	
Lithotomy. Perineal	
Instruments and preliminary precautions	
Steps of the ordinary lateral operation	
Aston Key's operation with the straight staff	
Accidents and difficulties of lateral lithotomy	
1. In cutting into the bladder	
2. In extracting the stone	
Causes of death after lateral lithotomy	
Statistics of	
"	
Mortality at different ages	

FOURTH VOLUME.

	PAGE
Table of cases of stone operated upon at the Norwich Hospital during 97 years	1068
Table of cases of lateral lithotomy at the same hospital during the same period	1064
Table of cases of recurrence of stone after lithotomy at the Norwich Hospital, with remarks by Mr. Williams	1065
Different modes of performing perineal lithotomy	1069
Boyer's method with the <i>lithotome caché</i>	1069
Dupuytren's bilateral operation	1069
Median lithotomy. The Marian operation; Civiale's medio-bilateral operation; Buchanan's operation; Allarton's operation	1070
Vidal de Cassis' quadrilateral section	1071
The recto-vesical operation	1072
The hypogastric, or high, operation	1078
Estimate of the different methods	1076
Statistics of the median operation at Norwich, with remarks by Mr. Williams	1079
Calculus in the prostate gland	1082
Symptoms and diagnosis	1084
Treatment	1084
Calculus in the urethra	1084
Treatment	1085
Calculus in the female bladder	1086
Various methods of treatment	1087
Lithotomy in the female	1089
Calculus in the female, obstructing labour	1090
Table representing the chief varieties of calculus	1092

LITHOTRITY.

BY CHARLES HAWKINS, ESQ.

INSPECTOR OF ANATOMY.

History of the invention of lithotrity	1093
Different forms of lithotrity	1095
Primary treatment	1096
Operation. Methods of seizing the stone	1097
Sir B. Brodie's method	1098
M. Civiale's method	1100
Completion of the operation	1104
Repetition of operation	1104
Removal of fragments after operation	1105
Complications after operation	1106
Use of chloroform in lithotrity	1109
Comparison of lithotomy or lithotrity	1110
Comparative mortality of lithotomy and lithotrity	1113
Lithotrity in hospital practice	1116
Sir H. Thompson's statistics of lithotrity	1118

	PAGE
Table of cases of stone operated upon at the Norwich Hospital during 97 years	1063
Table of cases of lateral lithotomy at the same hospital during the same period	1064
Table of cases of recurrence of stone after lithotomy at the Norwich Hospital, with remarks by Mr. Williams	1065
Different modes of performing perineal lithotomy	1069
Boyer's method with the <i>lithotome caché</i>	1069
Dupuytren's bilateral operation	1069
Median lithotomy. The Marian operation; Civiale's medio-bilateral operation; Buchanan's operation; Allarton's operation	1070
Vidal de Cassis' quadrilateral section	1071
The recto-vesical operation	1072
The hypogastric, or high, operation	1073
Estimate of the different methods	1076
Statistics of the median operation at Norwich, with remarks by Mr. Williams	1079
Calculus in the prostate gland	1082
Symptoms and diagnosis	1084
Treatment	1084
Calculus in the urethra	1084
Treatment	1085
Calculus in the female bladder	1086
Various methods of treatment	1087
Lithotomy in the female	1089
Calculus in the female, obstructing labour	1090
Plate representing the chief varieties of calculus	1092

LITHOTRITY.

BY CHARLES HAWKINS, ESQ.

INSPECTOR OF ANATOMY.

History of the invention of lithotritry	1093
Various forms of lithotrite	1095
Preliminary treatment	1096
The operation. Methods of seizing the stone	1097
Sir B. Brodie's method	1098
M. Civiale's method	1100
Completion of the operation	1104
Repetition of operation	1104
Removal of fragments after operation	1105
Complications after operation	1106
Use of chloroform in lithotritry	1109
Question of lithotomy or lithotritry	1110
Comparative mortality of lithotomy and lithotritry	1113
Lithotritry in hospital practice	1116
Sir H. Thompson's statistics of lithotritry	1118

LIST OF ILLUSTRATIONS.

FIG.	PAGE
306. Pendulous growths from synovial membrane	4
307. Soft anchylosis	17
308. Chronic osteo-arthritis of the knee— <i>Anterior view</i>	26
309. " " " " <i>Posterior view</i>	26
310. Fosse cartilage of elbow	37
311. The acetabulum almost filled up with bony deposit after hip-disease	45
312. Incipient ulceration of the cartilages	58
313. Ulceration of cartilage	59
314. Bony anchylosis of the hip	67
315. Necrosis of the acetabulum	83
316. Dislocation of the hip from disease	83
317. Chronic osteo-arthritis of the head and upper end of the femur	87
318. Disease of the tarsus, commencing in the calcaneo-astragaloid joint.	94
319. Angular deformity from spinal disease	110
320. Diagram of the formation of angular deformity	111
321. Caries of the spine, with anchylosis of its posterior portion	114
322. Atlo-axial disease	140
323. Re-pair of disease of the axis	142
324. De-formity of the neck from disease high up	143
325. Rivalve nasal speculum	256
326. Metz's nasal speculum	257
327. General arrangements for posterior rhinoscopy	258
328. Method of using the nasal douche	276
329. Rauehfuss's instrument for blowing powders into the nose	278
330. Hilton's snare for nasal polypi	301
331. Enamel nodule on a tooth-fang	350
332. Exostosis of a tooth	351
333. Magnified section of a dental exostosis	351
334. Dentine excrescence attached to a tooth	352
335. Dentine excrescence in the cavity of a tooth	352
336. A warty tooth	354
337. Several teeth enclosed in a warty mass	354
338. A large diffused tumour formed by a warty mass developed from the teeth	355
339. Fissure of the hard palate from syphilis	394
340. Obturator applied in the above case	394
341. Congenital cleft of hard and soft palate	395

FIG.	PAGE
242. Obturator applied in this case, lower surface	396
243. The same seen in profile	396
244. Inferior surface of total cleft of palate with separation of inter-maxillary bone	416
245. Double hare-lip with separation of inter-maxillary bone	417
246. Total mesial cleft of palate and lip	419
247. Model of complete cleft of palate	433
248. The same in section, showing the great inclination of its sides	433
249. Model of partial cleft of palate	434
250. The same in section, showing the much more horizontal direction of its sides	434
251. Cystic tumour of lower jaw	450
252. Drawing of patient with malignant tumour of upper jaw and deficiency of lower jaw	473
253. Pouch of the pharynx	487
254. General arrangements for laryngoscopy	520
255. Diagram illustrating the reversed image in laryngoscopy	522
256. Simple form of Siegle's spray-producer	531
257. The same instrument with Beigel's face-screen	531
258. Papilloma of larynx	575
259. Pedunculated fibrous tumour of larynx	577
260. Extroversion of mucous membrane, simulating laryngeal polypus	580
261. Mackenzie's laryngeal forceps	586
262. Laryngeal forceps, opening antero-posteriorly	586
263. Intussusception of gut, caused by the traction of a fibrous tumour	613
264. Hernia, at the femoral ring, of a part only of the walls of the bowel	673
265. Artificial anus after strangulated ileum, at the convexity of the tube	681
266. Artificial anus, the entire calibre of the bowel destroyed	682
267. Bowel injured from violent taxis	691
268. Double inguinal hernia	732
269. Diagram of two of the congenital imperfections of the vaginal process of peritoneum	735
270. Diagram of two other of these malformations	735
271. Direct inguinal hernia	76
272. Double femoral hernia	76
273. Condition of the ileum and sac in a case of strangulated hernia	76
274. Bubonocele and femoral hernia	77
275. Internal hæmorrhoids surrounded by external hæmorrhoids	79
276. Clamp for removing piles	80
277. Prolapsus recti, in the child	81
278. „ „ in the adult	81
279. Cancerous growth, protruding from the anus, surrounded by external piles	85
280. Warty polypi at the anus	86
281. Villous tumour of the rectum	86
282. Extroversion of the bladder	882
283. Sacculated bladder with enlarged prostate	887
284. Villous tumour of the bladder	900
285. Great hypertrophy of the prostate, with fibrous tumours	918

LIST OF ILLUSTRATIONS.

xxix

FIG.	PAGE
286. Vertical section of bladder and enlarged prostate	919
287. Typical example of enlarged prostate	920
288. Exploring sound with angular beak	926
289. Prostatic catheters	930
290. Long indurated stricture of the bulbous part of the urethra	942
291. Stricture of anterior part of urethra	942
292. Bulb-ended bougie and catheter	950
293. Diagram of a section of the urethra	952
294. Fac-simile of French and English scales for catheters	955
295. Holt's or Perrève's instrument for rupture of stricture	962
296. Sir H. Thompson's stricture dilator	963
297. " catheter-urethrotome	969
298. Civiale's urethrotome	969
299. Trélat's urethrotome	970
300. Uric acid deposit	1004
301. Urates of ammonia deposit	1006
302. Oxalate of lime "	1007
303. Cystic oxide "	1008
304. Phosphate of lime "	1009
305. Triple phosphate "	1010
306. Diagram of various sections of the prostate in different methods of lithotomy	1069
307. Sir B. Brodie's method of seizing the stone in lithotripsy	1099
308. M. Civiale's " " " "	1108

PLATE.

Plate of the chief varieties of calculus	1092
--	------

PART I.

DISEASES COMMON TO ALL THE JOINTS.

estimating the nature and importance of diseases of the joints, we have to bear in mind that we have to deal with structures, distinct in their formation, their functions, their pathology, but all so far united for a special purpose. Disturbance of any one of them is sufficient to derange the harmonious working of the whole. The study of diseases of joints, therefore, involves to a greater or less extent that of affections of the *osseous system*, the foundation, as it were, of the articulations; of *cartilage*, *fibro-cartilage*, or *membrane*, disposed between the bones to play the part of the buffers in a railway-train; of *fibrous tissue* or *ligaments*, the coupling-rods; of *synovial membranes* with their *synovia*, the lubricating-oils and grease by which the friction is diminished; as well as of *connective* and *adipose tissues*, which serve as packing material to fill up the hollows. External to the joints, moreover, but still playing an important part in their mechanism, are various muscles with their tendons, as well as synovial sheaths, the pathological conditions of which in their relation to joints cannot be entirely omitted or passed over.

It is only since the investigation of the diseases of the joints

which those tissues are found to have undergone. It must, however, be admitted that when disease has advanced to a certain stage, so many of these structures are likely to be involved, that an accurate allotment of the symptoms to their respective sources would be useless even if it were possible; so that, practically, we have often to disregard minute anatomical distinctions, and consider the case mainly as regards its severity and probable termination. It may be remarked, too, that the separation of joint diseases into those which affect the synovial membranes or ligaments, the bones, and the cartilages, is more obvious and more practically useful in adults, in whom these structures have arrived at their full growth, than in children in whom they are still in process of development. As Mr. Hilton justly observes, 'in childhood the separate and distinct disease of the articular structures can seldom be recognised with any useful precision during life.'*

The limits assigned to this essay are evidently insufficient to allow of any complete or comprehensive study of these varied diseases; no more than a sketch can be attempted; but that sketch, it is hoped, may be found sufficient for all practical purposes.

Anatomically all joints are not alike, being divided into the comparatively simple *synarthrodial* articulations, or those devoid of a synovial membrane, and the more complicated *diarthrodial*, in which a synovial membrane exists. Pathologically a similar distinction may to a certain extent be drawn, and the affections of the diarthrodial articulations may conveniently be separated from those of the synarthrodial, which will receive such notice as may be considered necessary in the latter part of the essay.

DISEASES OF THE SYNOVIAL MEMBRANES—SYNOVITIS.

Synovial membranes are liable to inflammation, forming what is usually, though not perhaps very correctly, termed *synovitis*.

The inflammation which attacks a synovial membrane may be of the ordinary character, or it may be modified by certain constitutional conditions. Common synovitis occurs, in many cases, as a local affection from the direct irritation of mechanical

* See also Holmes, *Surgical Diseases of Children*, p. 411.

injury, or results from undue exercise and exposure of the joint to atmospheric influences; it may also be induced by disease which has involved the adjoining tissues. This form of inflammation is generally confined to a single articulation.

The constitutional conditions which modify the character of the inflammation may be more or less permanent, and even hereditary. Sometimes we find the patient presenting evident marks of *scrofula*, and the local disease sharing in the peculiarities of that affection; in other cases, and principally in the poorer classes, who are ill-clad and unduly exposed to variations of temperature, the *rheumatic* diathesis is apt to prevail; whilst in those who live luxuriously, the local inflammation may be preceded or accompanied by various symptoms of disordered digestion, indicating the presence of an unusual quantity of uric acid in the system, or the existence of the *gouty* diathesis. The constitutional modifications, on the other hand, may result from causes which are obviously temporary. A peculiar form of synovial inflammation, for instance, may occur unexpectedly after parturition, or following some operation, in those in fact who are suffering from what is termed *pyæmia*; or it may take place suddenly during the progress of urethral discharge, when it is distinguished by the name of *gonorrhæal rheumatism*; or, finally, it may manifest itself during the existence of *venereal poisoning*. In all these modifications of inflammation, it may be remarked, there is a considerable chance of the topical disease manifesting itself in more than one articulation.

It will be convenient to commence with a description of the *simple* form of inflammation of a synovial membrane, and its consequences, and afterwards to notice briefly the peculiarities presented by synovitis when *scrofulous*, *rheumatic*, or *gouty*; *pyæmic*, *gonorrhæal*, or *syphilitic*.

Acute synovitis.—Synovitis may be acute or chronic. Acute synovitis, arising in the membrane itself from the causes already alluded to, occurs more frequently in adults than at an earlier period of life, in males than in females.

Pathological anatomy.—The simplest changes which result from inflammatory action consist in increased vascularity of the membrane, especially of its plicæ vasculosæ, followed by increase in the amount of the synovia, and some alteration in its healthy composition. A little later, we find the products of inflammation either effused into the synovial cavity, mingling with and

usurping the place of the normal secretion, or situated in the texture of the membrane and in the immediately contiguous tissues. The effusion into the joint-cavity may consist of serum, mixed with shreds of synovial epithelium, and a variable amount of organisable lymph, giving rise to false membranes connected with the walls of the cavity or floating in the secretion; some of the colouring constituents of the blood may also be present. In a large number of cases the inflammation is arrested before much mischief occurs; the vascularity of the membrane diminishes, the effused fluids are absorbed more or less completely, and the joint is restored to its original condition; or it may be left more distended than natural, and liable to fresh effusion; or a certain amount of thickening and loss of mobility may remain permanently.

In other cases, however, further changes occur within the

Fig. 206.



Pendulous growths from the synovial membrane. (From a preparation in the Museum of St. George's Hospital.)

joint, giving rise to villous or fringed processes, extending, perhaps, over the cartilages, which may still be found beneath them. At the same time a similar action takes place on the outer side of the membrane, increasing the amount of thickening and consolidation of the capsule. If the inflammatory action still proceeds unchecked, the fluid effused into the joint becomes turbid, and pus is formed, at first scantily, but afterwards in larger quantities; the other structures entering into the composition of the joint become involved in the diseased action; the articular cartilages

are affected, ulcerous excavations extend through their substance, and the articular extremities of the bones participate to a greater or less extent in the disorganising process. The parts outside the joint-cavity undergo similar changes; degeneration takes place in their newly-formed materials, and abscesses arise there, communicating with the joint by ulceration extending through the capsule, or making their way to the surface by means of sinuses. The ligamentous structures

become thickened or softened, or otherwise altered, so as, in some instances, to be no longer capable of holding together the bones, which may thus become completely or incompletely dislocated.

Symptoms.—The leading symptoms are those of inflammation generally, varying in degree with the amount of synovitis present, with the cause inducing it, with the particular constitutional condition, and with the individual joint affected. The amount of inflammatory fever depends upon the character and extent of the synovitis, and the importance of the articulation, being generally most marked when the hip-joint is the seat of the disease. Locally, we have *pain*, usually of considerable amount, increased by any movement of the part, and often characterised by a feeling of distension. It may be noticed also that the pain in articular inflammations, as in those of some other organs, is not always referred to the region in which the disease actually exists. In diseased hip, for instance, the suffering may be experienced mainly in the knee, giving rise occasionally to mistakes on the part of the careless observer. There is *swelling*, varying with the amount of effusion into and round the joint, and taking the shape not of the articular extremities of the bones, but of the distended synovial capsule, modified by the degree of resistance afforded by the surrounding structures. *Fluctuation*, at least in the more superficial joints, is usually distinct, becoming less so if the disease is prolonged and the inflammatory products acquire a more solid character. *Heat* and *redness* are more or less evident, according to the situation of the joint affected, its proximity to the surface, and the degree in which the tissues external to the synovial membrane are involved. The *position* which is unconsciously assumed in acute articular inflammation is often characteristic of the affection and of the particular joint implicated. To alleviate suffering, by securing rest, the limb becomes more or less rigidly fixed by involuntary contraction of the muscles in its neighbourhood, but, in addition to this rigidity, there is a strong tendency for the joint to become flexed, and for this flexion steadily, though gradually, to increase unless arrested by the surgeon. It has been supposed that the rigidity and flexion are due to over-distension of the joint-cavity by fluid; but the ease with which the joint can be moved, and the increased flexion relieved when chloroform is administered, points rather to undue muscular contraction.

The most rational explanation appears to be that given by Mr. Hilton, in his valuable work *On Rest*, viz. 'that the irritated or inflamed condition of the interior of the joint (say the knee-joint), involving the whole of the *articular nerves*, excites a corresponding condition of irritation in the same nervous trunks which supply both sets of muscles, extensors and flexors; but that the flexors, by virtue of their superior strength, compel the limb to obey *them*, and so force the joint into its *flexed condition*.'* At any rate the effect produced, that is, the position assumed by the limb, is practically of great importance, for it is capable of maintaining or aggravating the disease at the time, as well as of giving rise ultimately to great embarrassment in the event of ankylosis occurring.

In the majority of cases of uncomplicated synovitis, the inflammation speedily subsides, the fluid is absorbed, and the part returns to its normal condition. In others, however, the joint remains somewhat swollen, the synovial membrane is thickened, and the mobility of the part diminished, whilst the patient is left very liable to relapse from exposure to any of the exciting causes of synovitis. Or the acute symptoms may simply diminish, and without any interval of restoration to health the disease may assume the chronic form to be subsequently described. In a certain number of cases, unfortunately, the inflammation, instead of subsiding, increases in severity, and the disease is no longer confined to the synovial membrane, the cartilages and other structures becoming involved, in the manner to be more minutely described hereafter. Such extension is marked by fresh and excruciating suffering, by painful 'jumping' of the limb, as well as by crepitus on moving the joint, as the cartilages become destroyed, and is followed or attended in a large proportion of cases by suppuration.

Acute articular abscess.—*Suppuration in a joint*, however, occasionally occurs also *directly* as a consequence of acute synovitis, the effect frequently of some injury, or in children especially, as the result of acute inflammation supervening on chronic disease of the joint. In this acute form of joint-disease, whether we assign the synovial membrane proper, or some other of the varied textures with which it is more or less

* Hilton, *On the Therapeutic Influence of Rest*, p.156.

blended, as the starting-point of the disease, we may *suspect* the formation of matter, if the inflammation run very high; if one or more rigors occur, followed by exacerbation of the pain and general suffering; and if the tumefaction increase and take on a more decidedly phlegmonous character which extends to the surface. The constitutional disturbance will now be great, the pulse rapid, and the character of the fever will be altered, assuming somewhat of a typhoid form. When suppuration has occurred within the joint, and the case progresses unfavourably, the parts around become speedily affected. Abscesses are formed in the surrounding areolar tissue; at first, perhaps, separate from, but soon communicating with, the articular cavity. The fibrous tissues become softened and yield, the skin gets implicated, and openings form communicating with the joint, and discharging a more or less unhealthy pus. When much disorganisation has taken place within the joint, the most favourable issue to which we can look forward is discharge of the matter, and union of the opposed articular surfaces. The health may now improve. The swelling subsides, the discharge diminishes, and at last ceases, the sinuses heal up, the joint-cavity is obliterated, being filled with newly-formed connective tissue, and the limb is stiffened or completely ankylosed. If, however, the constitutional powers give way, the patient may be worn down by the severity of the pain and profuse discharge; the removal of the source of irritation, by excision or amputation, becomes necessary, or where this is impracticable, life is at last extinguished.

Chronic synovitis may be simply inflammatory in its character, though it occurs more commonly perhaps as a modification of the inflammatory process, dependent on certain constitutional cachexiæ, such as scrofula, syphilis, gout, or rheumatism. In some cases the distinctive characters of each of these varieties are clear and well defined; and no difficulty is experienced in assigning the affection to its appropriate place. In others, however, the boundary-line is not always drawn with precision; and the surgeon finds in these diseases, as in most others, that the definite descriptions necessitated in books are not always borne out in practice. Just as acute disease melts by insensible gradations into the chronic, so the constitutional modifications of inflammation are often imperfectly or scarcely marked. At the bedside of the patient, the surgeon must work out for himself many details; and every case will still offer some special

employment for his intellectual investigation. The author can only point out, as it were, the leading types; he who attempts more bewilders the reader instead of assisting him.

Acute inflammation of the synovial membrane, imperfectly treated, may by degrees assume the chronic form; or commencing without any great activity in its symptoms, the inflammation may pass into a chronic condition; in the latter case, however, the probabilities are very much in favour of the existence of some constitutional taint. The joint is sometimes much distended with fluid; and the synovial membrane itself, in protracted cases, loses more or less of its natural structure, is over-vascular, and permanently thickened, or converted into a firm and gristly substance. The disease may gradually yield, or it may remain stationary in this condition for a considerable time, and then, under the influence of fresh inflammation, the joint may ultimately become totally disorganised.

The symptoms are much the same as in the acute form of disease, only far less severe in their character. The constitutional disturbance is comparatively little felt. There is swelling, sometimes to a considerable extent, more or less pain is experienced, and the movements of the joint are often very greatly impaired. In some instances a peculiar crepitus or crackling is felt, or a sensation is communicated to the hand as if a number of loose bodies existed in the joint, depending, perhaps, on effusion of plastic material into the cavity, similar to that noticed in inflammation of the bursæ and tendinous sheaths. The simple form of chronic synovitis—the form, that is, not connected with any constitutional taint—usually terminates favourably unless neglected, as is occasionally seen in the labouring classes, when attack after attack may occur till at last the joint is destroyed, or life itself sacrificed. In cases, too, of long standing, where great induration of the soft tissues has occurred, complete restoration to a healthy condition becomes at best difficult, and relapses into a state of sub-acute inflammation are of frequent occurrence.

Treatment of acute synovitis.—The treatment to be adopted is partly general, partly local. If the inflammation runs very high and the patient is strong and plethoric, it may be necessary to take blood from the arm; at any rate, a brisk mercurial purgative, followed by salines, with antimony and, perhaps, with colchicum, will be advisable, or in some cases it may be expedient to place the system rapidly, but for a short time, under the

influence of mercury. Where the patient is less robust, any blood-letting had recourse to should be local; and in all cases, though it will be desirable to keep the liver acting by combining taraxacum with the saline medicines, or by administering occasionally some preparation of mercury, violent purgatives should be avoided, on account of the disturbance to the limb which they necessitate.

The primary and essential condition of the local treatment is complete *repose*. When the inflammation is of moderate extent, mere rest in bed, with the joint supported by a partially distended air or water cushion, will be sufficient. Where the inflammation is severe, and danger of permanent mischief threatens, more decided mechanical support is required. The position to be adopted varies with the individual joint, but in all cases must be that which keeps no ligament, and no part of the synovial membrane, on the stretch, and which, in the event of ankylosis occurring, will allow the limb to be used to the greatest advantage. Supposing the articulation to have already assumed an improper position, it is better to put it right *at once*, even if the existing inflammation is acute; for in unnatural positions the different parts of the joint are kept in a state of undue pressure or of undue tension, either of which interferes with healthy nutrition, and so opposes the curative process. Under the influence of chloroform, the replacement can, at this time, be readily effected.* The splint used should

* As Mr. Holmes Coote, at page 85 of his work *On Joint Diseases*, has quoted these remarks for the purpose of adding, 'To this practice I most strongly demur,' I may be allowed to cite the very judicious and forcible observations of Mr. Hilton respecting the mischievous and destructive effect he has over and over again seen produced by allowing the joint to assume or retain the fixed and flexed position induced by involuntary muscular contraction, 'resulting not seldom, I may say, from procrastination, or inexcusable want of courage on the part of the surgeon to rectify it, or from his yielding too easily to the importunities of the patient or his friends, in order to avoid giving pain to the patient. From the time when the limb begins to be flexed starts the mischief, which by-and-by cannot be controlled.' And again, after relating two cases in which amputation had to be performed: 'It seems to me that it is a great mistake not to correct it *at once* when they see an inflamed knee or other joint beginning to be contracted or flexed. I believe their duty is to prevent the increase of the flexion, and so avert the condition which, at all events, led to the necessity for amputation in these two cases.'—Hilton, *On the Therapeutic Influence of Rest*, pp. 161-3. I need not say how heartily I concur in these remarks, in which, it may be noticed, the very words employed in the text about correcting the improper position 'at once' are adopted by Mr. Hilton.

be well adapted to the limb, and be of *sufficient length*; in many cases it is absurdly short, giving no real rest to the joint, and allowing the weight of the parts below to drag upon the ligaments, and tend to produce deformity. In disease of the knee-joint, for instance, to give effectual support, the splint should extend very far up the thigh, and be furnished below with a foot-piece, to prevent the tendency of the foot to fall outwards or inwards, and so produce a corresponding twist of the affected articulation. The points here insisted upon are, of course, applicable not merely to cases of synovitis, but of all diseases of the joints, in which absolute rest is demanded.

Local depletion may be required to an extent determined by the acuteness of the inflammation and the vigour of the patient. If leeches are applied, it must be remembered that they act not merely by the abstraction of blood they give rise to, but also as derivatives, in consequence of the irritation produced by the bites. The application of leeches immediately over a *superficial* joint may tend, consequently, to increase the inflammation instead of diminishing it. In acute synovitis of such a joint as the knee, therefore, it is better to apply leeches at some short distance *above* the joint, so as to empty the distended vessels below. The same observations would, of course, apply, should cupping be employed instead of leeches.

After, or in many cases in place of, the local depletion, various emollient applications may be used. As a general rule, warm fomentations and poultices are the most successful in relieving pain. Occasionally, however, cold is more grateful to the patient. Whichever is employed, uniformity of temperature is much to be desired; the joint should not be exposed to frequent changes, but the applications should be maintained constantly at or near the same standard. It is on this principle that wrapping the joint in cotton wool, or covering it with ointments which exclude the air, is often of service. If cold be applied, it may be used as directed by Dr. Esmarch,* in the form of ice contained in bags of vulcanised india-rubber (far preferable for many reasons to bladders), separated from the limb by a piece of lint. If this prove too heavy, a simple apparatus may be contrived, by suspending, at a suitable angle, from the cradle placed over the affected limb, a bottle containing cold water, in which a few threads are partially immersed, with their ends

* *On the Use of Cold in Surgery*, New Syd. Soc. 1861.

hanging out of the mouth of the bottle; these, by capillary attraction, will keep up a constant drip upon a piece of lint covering the inflamed joint, and the supply of fluid may be so regulated, by varying the number of threads, as just to compensate for the loss by evaporation, and prevent the rag from ever getting dry. Such an arrangement I frequently used with advantage at the Hospital for Sick Children, both in cases where acute inflammation already existed, and in others where its occurrence might be anticipated in consequence of operations. The cold should be maintained without intermission, as long as any abnormal increase of temperature exists, provided it appears grateful to the feeling of the patient.

Treatment of acute abscess of joints.—If there is reason to believe that acute inflammation of the synovial membrane has proceeded rapidly to suppuration, the fact may be verified, at least in the more superficial joints, by the introduction of a needle-trocar. Supposing matter to be found, if situated outside the joint an early incision may prevent the pus making its way into the joint-cavity; and if inside the articulation, the urgency of the symptoms will seldom allow of much delay. It is better, therefore, when the situation of the joint admits of it, to evacuate the matter as early as possible by a free incision, which may be made under the protection of lint steeped in carbolic oil, and the wound afterwards dressed on the antiseptic principle, as practised by Mr. Lister. The opening in such cases is usually attended with marked relief; but fresh incisions may afterwards be made, or a drainage-tube introduced if any obstacle to the exit of pus shows itself. During this time the most perfect rest must be observed, and the limb maintained accurately in position by splints of sufficient length, moulded, if necessary, to the requisite form. Great attention must also be paid to the general health, and the strength supported as the activity of the inflammatory fever subsides.

If the matter within the joint decomposes without escaping freely, absorption of some of its putrid constituents may occur, showing itself by the irritative or typhoid character of the fever. In many cases life is now so evidently at stake, that removal of the diseased part either by amputation of the limb, or, in exceptional instances, by excision of the joint, becomes imperatively called for. In other cases, after providing as far as possible for the free evacuation of the pus, weak iodine injections may be used, not merely to modify the action of the secreting

membrane, but also to improve the character of the matter. With care and attention on the part both of the surgeon and the patient, acute suppuration of joints from synovitis may often be expected to terminate favourably, especially in children; but as more or less stiffness of the joint must be expected, special attention should be paid from the first to the position of the limb, instead of waiting, as is sometimes done, till ankylosis has taken place, to remedy the deformity which has been allowed gradually to arise.

Treatment of sub-acute and chronic synovitis.—Instead of giving rise to the formation of abscess, the acute inflammatory symptoms usually subside. As this change occurs, the antiphlogistic treatment is modified; some blood may still be abstracted by leeches applied directly over the part, and cooling applications may be used, or counter-irritation may be had recourse to, in the form of blisters, to produce a discharge of serum from the vessels of the skin. Blisters may be used earlier in inflammation of the deeper-seated joints than where the superficial articulations are implicated; for in the latter the small amount of intervening tissues allows the irritating action to be propagated to such an extent as sensibly to affect the synovial surface, acute inflammation of which may be again excited under the influence of too severe or too early a stimulus. In synovitis, therefore, of the superficial joints, blisters should not be applied too hastily, and when used may be placed some little distance above the inflamed spot; it is better also to employ a succession of ‘flying’ blisters rather than keep *one* open by irritating applications. Following the blisters, or in place of them, various stimulating liniments may be used, or the parts painted with concentrated tincture of iodine. As the inflammatory symptoms subside, in those cases especially where some thickening of the membrane has occurred, we combine a moderate amount of compression with our support of the part; this may be done by applying alternate layers of bandage and adhesive plaster with or without camphorated mercurial cerate next the skin, in the way described as ‘Scott’s bandage.’ Sometimes the bandage is starched, but care should then be taken, if any fresh outburst of inflammation occurs, that the bandage can be promptly removed, if necessary, by the patient. In some cases it is useful to employ splints made of cow-hide, softened in warm vinegar, moulded to the form of the joint, and fastened on the limb after having been lined with chamois leather.

Many similar contrivances will probably suggest themselves to the surgeon's mind.

If rest be a primary necessity in the acuter stages of inflammation, a time comes at last when this has to be discontinued, and passive motion of the joint substituted in its place; but the exact time when this change is to be made, often requires great judgment and discrimination. The mere presence of pain on motion is not always decisive of the necessity of rest, for in some cases, after long confinement, pain and aching are actually relieved by free and even strong movements.

A distinguished artist, for instance, came under my care for disease of the right wrist-joint following exposure to cold and wet. The inflammation had been severe, and the joint had been kept at perfect rest for several months, but still great aching and pain were complained of, the part was puffy, stiff, and sensitive, though not warmer than natural, and the patient in despair of ever again being able to follow his profession. With some degree of caution I removed the splints and employed passive motion, gradually increased in force, strong friction to the part, and localised galvanism to the individual muscles of the fore-arm, which had become rigid from confinement like the joint itself. The pain subsided under this change of treatment, and by degrees perfect motion was regained, so that the artist has been enabled to resume his profession with no impairment of his technical skill.

Perhaps the best indications for substituting motion for rest in those cases in which pain continues severe after active disease has subsided, are the comparative low temperature of the joint, and the tendency of the integuments to become dusky and puffy.

The treatment sketched out has been that proposed for disease of the synovial membrane itself; the treatment of those cases in which the bones and cartilages become involved will be alluded to subsequently, in the sections treating of diseases of those tissues.

Dropsy of joints.—We occasionally meet with cases in which the synovial membranes are very greatly distended with fluid, whilst few or no other symptoms of inflammation are present. The affection, therefore, has been compared to dropsy of serous membranes, especially that of the tunica vaginalis testis forming hydrocele, and has received the special name of *hydrops articuli*. I have placed it, however, immediately after ordinary synovitis, because in almost all the cases where an opportunity has been afforded of examining the joint, the lining membrane has been found somewhat thickened and unusually vascular,

especially in the situation of its vascular processes. The fluid differs from ordinary synovia, being transparent and of a yellowish colour, but presenting no appearance of 'threads,' though it coagulates on the application of heat; in fact, it closely resembles the fluid met with in ascites or in hydrocele. All synovial membranes are not equally subject to dropsy, which usually affects those of considerable extent, and in which the secretion is naturally abundant. It is most frequently found in the knee, sometimes in the elbow, very seldom in any other articulation. Hydrops articuli is scarcely ever a primary affection. It may follow an acute attack of synovitis, and is apt to recur in persons of weak and irritable constitutions, when once the membrane has been over-distended. In such persons the immediate attack may be induced by exposure to cold or very slight violence, or even by some less obvious constitutional disturbance.

The effusion takes place very rapidly, and forms a tumour, whose limits are those of the synovial capsule. Its character may generally be easily recognised by the shape of the swelling and its ready fluctuation. There may be some slight symptoms of inflammation present; but usually, though the limb is somewhat stiff, it may be used with scarcely any pain, even from the first, and the parts external to the capsule show no traces of redness or œdema. From thickening of the joint due to effusion of plastic lymph, hydrops articuli is distinguished by the fluctuation which, as M. Bonnet remarks, is best recognised by placing one hand above and the other below the tumour; if they are placed transversely, the displacement of the muscles or tendons may be mistaken for the movement of fluids. From abscess of the joint, it may usually be distinguished by the condition of the soft parts around as well as by the constitutional symptoms, though it occasionally happens that the true nature of the effusion can only be determined by puncture with a needle. The prognosis in these cases is most favourable; at least, the other structures of the joint seldom become involved, though the disease itself may be difficult of cure and very prone to relapse, especially in cases of some standing.

Treatment.—If any inflammatory symptoms are present, these must first be subdued by the ordinary constitutional remedies; by leeches and fomentations, or by cooling lotions. When all traces of inflammation are subdued, and the affection has assumed the chronic form, the joint may be blistered, or painted

frequently with a strong solution of iodine; this may be followed by compression by means of adhesive plasters and bandages, accompanied by stimulating and mercurial applications directly to the skin, for the purpose of promoting absorption. If the case is recent, under such treatment the effusion may be expected to subside; but it will still be necessary to support the relaxed membrane by wearing a laced or elastic knee-cap for an almost indefinite period.

In old and neglected cases, on the other hand, the affection resists all ordinary treatment, and must either be abandoned as incurable, or further and operative measures had recourse to, the most promising of which are iodine injections, employed as in cases of ordinary hydrocele. Should it be considered advisable to have recourse to this plan, and the knee be the joint affected (as is most commonly the case), the spot selected for the puncture is that part of the synovial membrane which is situated above and external to the patella; the limb should be in the extended position, and the opening rendered valvular by pulling aside the skin before introducing the trocar, which should be only of moderate size. The fluid injected may consist of tincture of iodine more or less diluted. In a case of hydrarthrosis of both knees in which a permanent cure was effected by this injection, after all other treatment had failed, Mr. Goodall* used equal parts of the tincture and of water, whilst Schuh,† who has employed this treatment successfully in three cases, made use of one part of tincture of iodine to four parts of water. The quantity injected may vary from half an ounce to two or three ounces, the latter quantity being that recommended by Schuh. As much as possible of the fluid is allowed to escape, after being retained in the joint for about five minutes, and care taken to prevent any admission of air to the articulation. When the trocar is withdrawn, the cutaneous wound is coated with collodion. The immediate result of the injection is to produce a certain amount of inflammation; to prevent this from going too far, the limb must be retained in a state of immobility after the operation by means of a properly adapted splint, and the case watched carefully, so that antiphlogistic measures may be had recourse to if necessary. This treatment is only suitable to cases of simple dropsy, uncomplicated with diseased conditions of the bones and cartilages, and

* *British Medical Journal*, No. 367. † *Wien. Zeitsch.* N. F. vol. iii. p. 5.

would only be advisable when all other measures had failed, and much inconvenience was occasioned by the distended capsule.

Scrofulous synovitis: pulpy degeneration.—There is a peculiar form of slow inflammation of the synovial membranes, which is sometimes described under the name of *gelatinous* or *pulpy degeneration of the synovial membrane*, but which appears to be inflammation of a chronic character, modified, possibly, by the scrofulous condition of the system. If we have an opportunity of examining the joint when the disease has advanced to a certain extent, the principal points which attract attention are the thickening of the synovial membrane, and the gelatinous pulpy appearance which it presents. The vascularity of the membrane is increased, and the mass of its substance appears to be converted into a soft yellowish or light-brown gelatinous material, frequently of very considerable thickness. The articular cavity becomes diminished as the thickening extends upon the *internal* surface of the membrane; and as this extension goes on most freely in the loose synovial folds which surround the cartilages, these are often partially or completely buried and concealed beneath the new formation. A somewhat similar process occurs at the same time in the areolar tissue at the *outer* surface of the membrane, which becomes condensed and infiltrated with a corresponding pulpy substance.

The joint may remain in this condition for a considerable period, and then, under favourable circumstances, a process of restoration occurs, the new elements shrink or are absorbed, and the articulation is restored more or less perfectly to its former condition. More frequently, however, the disease makes further progress; the gelatinous thickening of the synovial membrane extends further over the cartilages, which become themselves secondarily affected in spots; though these spots are not, necessarily, at first, in *direct* relation with the altered membrane. The precise nature of the change which occurs in the cartilage will be considered in the description of the diseases which affect that structure; it need only be said here that the cartilage becomes more and more altered at the points first affected, and that intimate adhesion soon takes place between the granulation-tissue which these points present and the corresponding growth from the synovial membrane. The change in the cartilage is attended or followed by inflammation of the bone on which it rests; the articular lamella gives way in places, a communication between the osseous tissue and the interior of


the joint is formed, and granulations spring up from the inflamed membrane lining the cancelli, which join with those already existing within the articulation. In the mean time matter has most probably been formed, which may be retained within the articular cavity, or may make its way to the surface by means of sinuses.

If the disease still makes progress, the gelatinous growth continues to extend in an outward direction, or it degenerates in patches, giving rise to fresh formations of matter in its own substance or in the articular cavity; the constitutional powers now probably fail, the tissues of the joint, including the osseous, are still further destroyed, and amputation or excision is had recourse to, or the patient dies hectic. When, on the other hand, an improvement takes place, the new growth is directed inwards towards the articular cavity, which it tends to fill up. As the constitutional powers improve, the gelatinous material becomes firmer and more fibrous, it continues to contract, the cavity within it diminishes and is obliterated, the bones are united by a firm and solid substance, and the joint is ankylosed, as in the accompanying figure.

Symptoms. — Pulpy degeneration makes its appearance most frequently in early life, at or before the time of puberty, though no period of life can be said to be absolutely exempt from risk of its occurrence. In many instances the immediate attack can be traced to some slight accident or other local source of irritation; in others no such cause can be discovered, the attack appearing to commence, as we may say, spontaneously. It is characterised by a want of acuteness, even at its commencement, the earliest symptom noticed (supposing the disease to be in the lower extremity) being slight limping, followed by some swelling about the joint.

The nature of the affection is most clearly recognised in those

FIG. 207.



Section of a knee-joint, the bones of which are united by ligamentous tissue after scrofulous disease. (From a preparation in the Museum of St. George's Hospital.)

joints which are the most superficial. In its early stages it is characterised by stiffness of the joint, and the presence of a soft, elastic, colourless swelling, occupying the whole extent of the articulation. The absence of genuine fluctuation should prevent the disease from being mistaken for dropsy or abscess of the joint; the size and doughy feel of the swelling distinguish it from simple synovitis; whilst its whiteness is opposed to the darker tint which is usually met with in malignant disease. The early stages of the swelling are usually accompanied with comparatively little pain, the disease in many cases being very insidious in its progress; in this respect, as well as in the shape of the swelling, which extends equally over the whole joint, instead of being specially manifest in the situation of one of the bones, differing from disease commencing in the cancellous structures. If the progress of the case is towards recovery, the swelling ceases to increase, solidifies slowly, and gradually diminishes in size, the pain at the same time yielding, and the condition of the general health becoming much improved. Even if arrested in this early stage, a considerable amount of loss of free movement of the joint must be expected, and great care taken that no recurrence of the symptoms is produced by too early exercise.

When the disease advances, the enlargement and stiffness of the joint increase; the pain, which was previously perhaps little more than a sense of aching or feeling of fulness, becomes more marked and altered in its character, being now described as a 'jumping,' 'gnawing,' or 'starting' pain, and felt with special acuteness at night-time. This change may be considered as a sign that the disease is no longer confined to the synovial membrane; that the cartilages are becoming ulcerated, or rather, more strictly speaking, that the bone on which those cartilages rest is itself participating in the inflammatory action. Another symptom, which would indicate a considerable amount of destruction of cartilage, is the existence of crepitation on any movement of the joint by the patient himself or by the surgeon. This crepitation, however, it should be recollected, may disappear or be masked at any time in consequence of soft and extensive granulations rising from the bones, and preventing their articular surfaces from rubbing on each other. The limb is now in danger of becoming rigidly fixed by spasmodic contraction of the muscles, causing it to assume positions varying in the different articulations, but all tending to produce future embarrassment, unless carefully watched and guarded against.

Before the disease has advanced to this point, in the majority of cases suppuration has occurred, either infiltrating more or less extensively the gelatinous mass, or limited so as to form circumscribed abscesses in its substance, or accumulated in the remains of the synovial cavity. If collected within the joint, there will be a general increase of the swelling, with throbbing pain, and perhaps fluctuation, attended by febrile disturbance to a greater or less extent. After a time the matter makes its way through the softened walls, and 'pointing' takes place at spots determined by anatomical peculiarities referable to the individual joint; the sinuses so formed presenting subsequently large and flabby granulations. When the abscess is situated in the gelatinous mass itself, the increased swelling instead of being general, takes place at any part of the mass which may happen to be the seat of the degeneration, and the opening forms at or near this point. In most cases, however, even when the joint-cavity was originally free, it becomes involved subsequently in the parietal abscess.

In proportion to the deterioration of the constitution will be the extension of the destructive action. The ligaments and neighbouring parts become softened and destroyed, the altered extremities of the bones are no longer firmly united to each other, and in this condition a partial or complete dislocation is occasionally produced under the influence of spasmodic muscular action; this dislocation being often attended with decided relief to the severity of the symptoms. If the constitutional powers improve, it is still possible for a process of repair and recovery to occur. The discharge in such cases diminishes, the openings contract and ultimately close, the skin resumes its natural appearance, while the swelling decreases and becomes more solid. The diminution in size of the joint continues till it becomes even smaller than it was before the attack, and the skin presents a peculiar puckered appearance around the articulation and at the seat of the cicatrices. The restorative process is of course attended with ankylosis, fixing the limb in the position it may have been allowed to assume.

This form of disease has generally been supposed to be strumous in its character, and connected with the scrofulous diathesis. Mr. Holmes,* however, has questioned its constitutional origin, and advanced many excellent reasons for doubting

* *Surgical Treatment of Children's Diseases*, pp. 424-7.

whether, in a large proportion of cases at any rate, chronic joint-disease may not simply be the result of some local injury or irritation, and essentially independent of any general cachexia, though strumous persons would be at least as liable to it as others. At any rate, as Mr. Holmes justly remarks, 'whatever our views may be as to the pathological question, I think we must all agree that practically we have no such proof of the constitutional nature of any of these chronic joint-affections as should lead us to refuse to entertain the general question of operative treatment. Individual cases must be judged of by their own symptoms.'

Treatment.—In our treatment we have to bear in mind that we are dealing with an affection originating in inflammation, but that inflammation of a chronic character and not improbably modified and altered by the strumous diathesis. Our efforts consequently have steadily to be directed to the improvement of the constitutional condition. Good air and plenty of it, warmth and light, attention to cleanliness and proper diet, are in all cases clearly inculcated. Where the tendency is to *scrofulosis*, the liver and bowels are usually sluggish, and require to be stimulated occasionally by tolerably brisk purges, combined with quinine; whilst in *tuberculosis*, purgatives must be given with caution, and be of the mildest character. In both cases, tonics and especially cod-liver oil, are of service; though these tonics should be administered with judgment, and in relation to the presence or absence of inflammatory fever.

The essential part of the local treatment consists in the joint being kept in a state of perfect and continuous *rest*, by means of splints, which should be well and properly made, and constructed so as completely to prevent any motion of the limb, and at the same time admit of any topical applications which may be deemed advisable. As the chance of ankylosis occurring is more considerable than in cases of ordinary synovitis, the *position* of the limb becomes of even greater importance, though it is to be regulated on the same principles. The splints may be made of leather or of pasteboard, or constructed of wire in the mode recommended by Bonnet and by Syme. If made of iron wire, a strong piece is bent to the shape of the limb, extending on both sides and united transversely to such an extent as to give the requisite degree of strength. The splint so formed is carefully lined with soft leather, or fitted with a kind of cushion, and will often prove highly useful. Too much attention can hardly be paid to the selection and

adaptation of the splint in the first instance, and to the necessity of its being worn uninterruptedly for a very long period of time. If inflammatory symptoms are present, it may be necessary to employ a few leeches, or to use cooling or warm applications, as may seem expedient. As the inflammatory symptoms subside, counter-irritants may be had recourse to; these, however, should be employed with some degree of hesitation, and with the cautions previously pointed out as specially applicable to their action on the superficial joints. At the Hospital for Sick Children I became latterly more sparing in my use of them. Mr. Barwell,* however, strongly recommends the employment of the actual cautery in these cases, when the first symptoms of inflammation have somewhat subsided, but when some such action still continues, and enlargement is still going on. The iron is to be white hot, and applied in lines, about an eighth of an inch in breadth and three inches in length, parallel to the axis of the limb. Four such lines are recommended for the knee, two on each side of the patella, at least an inch apart; dry lint may be applied immediately after the cauterisation. The advantage of this plan, according to Mr. Barwell, consists not merely in its acting as a counter-irritant, but also in the pressure produced subsequently on the subjacent parts by the contracting cicatrices. Extension by means of weights, as described in a subsequent part of this essay (see p. 49), is also of considerable service in many cases where the joint affected is in the lower extremity.

Where the disease is confined to the synovial membrane, and has assumed an entirely passive condition, measures may be properly adopted to produce absorption of the indolent gelatinous mass. Such measures would obviously consist in *friction*, *passive motion*, and accurately applied *pressure to the part*. These means, as pointed out by M. Bonnet, may be freely adopted, but still with care and caution, and at once abandoned if symptoms of renewed activity in the disease show themselves. If any camphorated mercurial application is used to the part, along with the pressure, in the way described as 'Scott's dressing,' it should be employed in these cases very sparingly, and in a very diluted form; the joint, too, should be examined frequently, to make sure that no fresh attack of inflammation has occurred. As the case progresses towards recovery, friction and passive motion are more freely used, and the treatment assimilates to

* *On Diseases of the Joints*, chap. v.

that of ordinary chronic synovitis. Should suppuration take place within the joint, the matter must be evacuated by free incisions made in depending positions, followed subsequently by slightly stimulating lotions to the sinuses, and moderate pressure to the joint, which is still to be maintained in a state of perfect rest, as well as in a suitable position. If the health give way, or the strength appear unequal to bear a long-continued discharge, and the circumstances of the case be such as not to admit of a prolonged use of the treatment which may be desirable, amputation of the limb or excision of the joint will often become necessary. Should improvement, on the other hand, take place, and cure by ankylosis be effected, we must be careful to employ passive motion in due time to prevent ossification, and maintain the uniting material in its fibrous condition.

Rheumatic synovitis.—In the description of synovitis, it was stated that this frequently arose from exposure to cold, or in connection with other causes loosely spoken of as rheumatic. The synovial membranes and parts about the joints are also liable to suffer in a special condition of the system by which the inflammatory process is materially modified, and which constitutes 'rheumatism' properly so called. In acute rheumatism or rheumatic fever, the articular affection, although it may be severe and troublesome enough, is evidently dependent on constitutional derangement, culminating in a kind of inflammatory paroxysm assumed to be connected with excess of lactic acid. Unlike ordinary synovitis, the constitutional disturbance in this case is primary instead of secondary; and though a joint, when attacked, may display signs of severe and acute inflammation, yet these signs will often disappear with extraordinary rapidity, whilst other articulations become as suddenly affected. Acute rheumatism, therefore, being essentially a constitutional malady, falls naturally to the care of the physician, and its description must be sought for in works on medicine.

Chronic osteo-arthritis; chronic rheumatic arthritis; rheumatic gout.—The synovial membranes, as well as the other structures entering into the composition of a joint, are liable to a chronic affection, which has long been practically recognised, though varying names have been assigned to it at different times, or according to the class of joints which it happens to implicate. The term *nodosity of the joints* was the one proposed by Dr.

Haygarth, whose attention was attracted to the hard swellings presented by the smaller articulations of the hands and feet, which at the present day more commonly receive the appellation of *rheumatic gout*. When the larger joints, such as the shoulder, the elbow, or the knee, are implicated, the disease is often termed *chronic rheumatism*; whilst a similar affection of the hip has been described by some authors as the *morbus coxæ senilis*. Dr. Adams of Dublin, however, in his valuable work on the subject, has arranged and classified the varieties of the affection, and proposed the term *chronic rheumatic arthritis*, for which, however, the name *chronic osteo-arthritis* has lately been substituted in the official *Nomenclature of Diseases*. It will be seen that the disease implicates other tissues besides the synovial membrane; and, indeed, the changes which occur in the bones in these cases are so specially marked, that Mr. Barwell considers the morbid action to be in reality a rheumatic osteitis. The first obvious symptoms during life, however, are those in relation with the synovial membrane, and the earliest traces of disease after death have been found to be connected with the same tissue. Cruveilhier, Dr. Robert Adams, and Sir Benjamin Brodie, coincide in the opinion that a chronic inflammation of the synovial membrane is the earliest local result as far as the articulation is concerned; and I have therefore thought it best to notice the affection in the present section.

Chronic rheumatism is met with in two distinct conditions: as a palpably constitutional affection, implicating a multiplicity of joints, and either succeeding to rheumatic fever or following exposure to the usual exciting causes of rheumatism; and, secondly, as a local affection attributed to over exertion, or injury, or arising without any assignable cause. The two sexes are, perhaps, nearly equally liable to be attacked; but Dr. Adams is of opinion that the larger joints, especially the hip, are more frequently affected in males, whilst females are more subject to the disease in the smaller articulations, such as those of the fingers; a fact which was originally pointed out by Dr. Haygarth. Though usually met with after middle age, chronic rheumatism is by no means confined to elderly people, for it has been seen in patients under the age of twenty, affecting simultaneously a considerable number of joints. Nor is it confined to any special rank; for though it prevails extensively among the labouring poor, it is also frequently met with in the wealthier classes, in those who have led indolent and luxurious

lives, and in whom it is apt to assume a more decidedly gouty character. Though chronic rheumatism materially diminishes the *comfort*, it does not appear to shorten the actual *duration* of life, at least in those whose means allow them to be supplied with all requisite appliances; the very poor, however, are exposed to accidental or extraneous maladies, from the helpless condition to which they are reduced, and so with them the chances of life are diminished. When once the disease has become established, its tendency is to progress, slowly but surely, until the joints affected become ultimately disorganised to a greater or less extent; sometimes the affection remains quiescent, or stationary, for an indefinite period; but very seldom indeed can anything like a cure be anticipated, and then only when the case has been recognised and the treatment commenced before the disease had made much progress.

As far as the local affection is concerned, if we have an opportunity of examining the joint at a very early stage of the disease, we find redness of the synovial membrane, which becomes thickened and fibrous, whilst its fringe-like processes are much developed, and more vascular than usual, the internal surface of the capsule presenting somewhat of the appearance of a villous mucous membrane. There is also effusion of a moderate amount of fluid within the joint. At a later period, the fluid is absorbed; but the ligaments, which had been distended by the effusion, recover themselves but slowly, laying the foundation, perhaps, of the partial or even complete dislocations which are occasionally produced subsequently by the spasmodic muscular contractions which are apt to occur in certain stages of the disease. The capsules become very considerably thickened, and sometimes a bony deposit may be met with in their substance; in the interior of the joint, too, there may be found one or more of those foreign or extraneous bodies which are noticed separately in this essay under the head of Loose Cartilages. The articular cartilages by this time are probably affected; at certain points they assume a yellowish hue, and become fibrous. As the disease progresses, they are destroyed to a greater or less extent; sometimes presenting patches of dense bone, which are smooth and glistening; whilst at other times the cartilaginous tissue disappears entirely, and in its place we find a compact ivory-like bony material, which soon becomes highly polished under the influence of the attrition to which it is subjected, or is marked with linear furrows, result-

ing from the increased wear to which it is exposed in certain situations in the movements of extension and flexion. In the hip-joint, the round ligament, and in the shoulder the tendon of the long head of the biceps, or the glenoid ligament, will often be removed by the process of absorption. The proper osseous structure becomes condensed beneath the articular surfaces and bony vegetations are thrown out around them, forming buttresses which interfere with the movements of the joints by their mutual apposition. The shape of the articular surfaces is altered; the globular heads of the femur and humerus are flattened, and, as it were, crushed, and the cavities for their reception exhibit a corresponding change, resulting in varying amounts of deformity. These alterations have sometimes been mistaken for fracture of the neck of the humerus, or of the cervix femoris, in which bony union had taken place. It may be mentioned, too, that the abnormal condition is not always confined to the articular extremities, but may involve the shaft, or even the entire bone.

The general symptoms in these cases present no very special character. In some instances, as Sir Benjamin Brodie remarks, the disease appears to be connected with over-indulgence in animal food, and the patient experiences some of the usual effects of dyspepsia, such as a tendency to acidity and flatulence. In other cases, and especially, perhaps, where the larger joints are implicated singly, no particular constitutional derangement can be detected, and, indeed, it is remarkable to how slight an extent the general health is affected by the great amount of topical mischief which may have occurred.

The local symptoms are the most characteristic, and consist essentially of *pain, enlargement and ultimate deformity of the joint, rigidity or stiffness*, and a peculiar *crackling noise*. The pain experienced is principally felt at night-time, or when the joint is set in motion after a period of rest. The amount of pain, which is usually of an aching character, is far less than might be expected from the amount of structural alteration which occurs, and very different from that experienced in other affections of the articular ends of the bone implicating the joints. When only a single articulation is involved, the general suffering is not great, and the patient's condition is very tolerable; even when several joints are attacked, the amount of pain experienced may be only trifling, though it may become aggravated, especially under the influence of cold and humidity,

and of atmospheric changes, to such an extent as to rest almost impossible. In the earliest stages, when enlargement depends probably on increased effusion, the synovial membrane or the bursæ in its neighbourhood swelling is soft, and fluctuation may sometimes be detected. At a later period the swelling becomes hard, and bony growths may often be felt connected with the periosteum and extremities of the bones, which lead to alteration of the form of the articular surfaces, and give rise to

FIG. 208.

*Anterior view.*

FIG. 209.

*Posterior view.*

Chronic osteo-arthritis of the knee-joint. (From a preparation in the Museum of the Middlesex Hospital.)

and peculiar deformity, varying in the individual joints, and characterised in the smaller ones, such as those of the fingers, by the appearance of irregular knots; whence the term *osteophytes of the joints*, proposed by Dr. Haygarth. The accompanying illustrations from a preparation in the Museum of the Middlesex Hospital, show the changes in the shape of the bones, the eburnation of the articular surfaces, the copious deposit

bones in the neighbourhood of the joint-ends and the numerous additamentary bones which are met with in old cases of osteo-arthritis. The case is related by Mr. De Morgan in vol. ix of the Pathological Transactions, where also a drawing will be found showing the extraordinary change of shape which had occurred in the upper end of the femur.

It may be noticed that there is little or no tendency to suppuration in these cases; abscess of the joint rarely, if ever, taking place. The amount of articular rigidity which occurs is peculiarly striking, rendering the affected joints practically useless, and simulating ankylosis, though true ankylosis rarely, if ever, occurs. In the advanced stages of the disease, stiffness or even immobility of the articulations may be present, depending on the deformity which has occurred in the articular surfaces, and on the bony outgrowths which have formed; and where many joints have been affected, the unhappy patient may be confined to his bed in a state of utter helplessness. Though the muscles are not immediately implicated in the rheumatic affection, at some period of the disease, probably when the bone itself begins to suffer, painful spasms are apt to occur, contributing materially to the discomfort of the patient, and giving rise in some cases to the partial or complete luxations which are occasionally met with. When the joints have become rigid and practically useless, the muscles in the neighbourhood degenerate and waste. The sensation of decided crackling experienced in the movements of the joint is evident both to the patient and to the surgeon, and is usually most marked when the articulation is first used after repose; it is especially noticed, therefore, in the morning. It depends upon the mutual attrition of the altered articular surfaces, and may be detected in most of the joints, but especially in the hip, and only disappears when complete rigidity has supervened.

Treatment.—Dr. Haygarth, to whom we are indebted for our earliest notices of this disease, lamented the scanty means of treatment which he could suggest, and hoped, when time had been afforded for investigation by the profession, that fresh and more successful curative measures might be discovered. These hopes, however, have as yet been only partially realised; and in the advanced stages of the malady, when the joint has been essentially disorganised, we can scarcely anticipate much improvement. Our treatment, therefore, to be of real service, must be adopted at the commencement of the affection; at a

later period we can but hope to mitigate the symptoms, to relieve the sufferings, and, at best, to retard the progress of the disease. In the early stage, when inflammatory symptoms are present, and any movement of the joint is attended with acute pain, it may be necessary to employ cupping or leeches, to keep the part at rest, and support it by means of bandages or splints. Care, however, must be taken not to prolong too far the period of immobility, for in these cases especially, want of exercise of the joint has ere long a deleterious influence on its structure. When the acuter symptoms subside, therefore, and rigidity increases, the patient should be encouraged to use the limb, to promote polishing and induration of the articular surfaces. Warmth and friction to the part are usually grateful. Flannel should be worn; and shampooing may be employed, or local douches or hot-air baths had recourse to. At certain stages of the affection counter-irritation may be used in the form of blisters or of painting with iodine; and in the nodosities of the fingers, the local application of a solution of iodide of potassium under oiled silk is sometimes of service; or cod-liver oil may be applied externally, as recommended by Sir Benjamin Brodie.

The general or constitutional treatment must be adapted to the individual case, and to the stage at which the disorder has arrived. Where any gouty element exists, where dyspeptic symptoms are present, and the case appears connected with too full a habit, great attention to the diet becomes of importance; stimulants should be abstained from; sugar, fruits, and raw vegetables avoided; and a moderate quantity of potash or magnesia be administered three or four hours after each principal meal. In other cases, however, the system appears impoverished rather than too full, and here it may be necessary to employ a better and more nutritious diet. In the earlier stages of sub-acute inflammation, it may be advisable to give a few alterative doses of blue-pill, combined, perhaps, with acetous extract of colchicum, and to administer Gregory's powder at night-time, with an active aperient at regular intervals. Perspiration should be encouraged, too, by the use of the Turkish or hot-air bath, or by taking Dover's powder, or other medicines to act upon the skin. When the urine is clear, Mr. William Adams recommends the use of sulphur externally and internally, and also the free use of dilute phosphoric acid. At a later period, the iodide of potassium may be administered and continued for several weeks at a time, if it appears to suit the

atient; for its action in these cases is somewhat uncertain. Various stimulating internal medicines may now be tried, such as turpentine or guaiacum; or cod-liver oil may be administered internally as well as externally; opiates also are usually of service for the purpose of procuring rest. Warm clothing should be used; and, where circumstances admit, residence in a warm climate during the winter months materially increases the comfort of the patient. Warm bathing is usually attended with relief; and if practicable, recourse should be had to the mineral springs, such as those of Bath or Buxton in this country, the hot sulphur baths of Luchon in the Pyrenees, Aix-la-Chapelle, Vichy, Ems, Wiesbaden, or Carlsbad on the Continent.

Gouty synovitis.—The gouty diathesis presents some points of resemblance to the rheumatic; in fact the chronic form of alteration of the joints just described is often termed rheumatic gout. The speciality of the gouty inflammation, however, consists in the sudden or 'explosive' appearance of uric acid in the part affected, giving rise to more or less local disturbance. Gouty inflammation of the joints occurs in an acute or in a more chronic form. The acute form of gout requires no description here, as it is essentially a medical complaint, and is amply treated of in medical works. Where gout has assumed a chronic form, however, the local affection may predominate over the constitutional, and such cases may at times fall under the care of the surgeon; they require, therefore, to be briefly noticed.

When gout first attacks a patient, the joint usually recovers its original mobility as the 'fit' passes off; but when repeated attacks have occurred, the restoration becomes more and more imperfect, and at last the articulation loses entirely its capacity for motion. Opportunities of examining the joint at an early stage are not common; but Dr. Garrod is of opinion that gouty inflammation is *invariably* accompanied by deposition of its peculiar salt. Unquestionably, as the disease advances, a deposit of the white, chalky-looking urate of soda takes place, in the substance of the ligaments and in the surrounding areolar tissue, as well as in the neighbouring muscles; or it may occur in the interior of the joint, thickening the synovial fluid, and studding the synovial membrane with small white masses; or in the substance of the articular cartilages, in the form of white spots, resulting from the presence of the crystalline salt in the hyaline

structure; or, finally, the urate may be deposited in the bone and in the periosteum. Resulting from this deposit we may expect to find, sooner or later, the tissues themselves undergoing changes; the synovial membrane will be thickened and vascular; the ligaments and areolar tissue condensed; and the cartilages subsequently become softened and are removed, so that the articular surfaces of the bones are exposed and altered.

Whenever a joint is subjected to protracted attacks of chronic gout, it is certain to be permanently injured, either by becoming so rigid that its functions are practically destroyed, or from the formation of *chalk-stones* around it. Chalk-stones—or *tophi*, as they are sometimes termed, from a Hebrew word signifying *concretion*—vary much in their consistence; they may be soft and semi-fluid, or hard as the chalk from which they derive their name. Under the microscope they present bunches of needle-like crystals of urate of soda; and chemically they are composed of that salt, united with a certain amount of animal or earthy matter. A concretion taken from the metacarpus was found by Lehmann to contain—

Urate of soda	52.12
Urate of lime	1.25
Chloride of sodium	9.84
Phosphate of lime.	4.32
Cellular tissue	28.49
Water, loss, &c.	3.98

100.00

Chalk-stones are generally found in the smaller articulations of the hands and feet. They form protuberances, distorting and crippling the articulation, which approach the surface, till the skin over them is thinned, and allows their white colour to become visible. Sir T. Watson relates the case of a gentleman who was in the habit, when at cards, of scoring the game on the table with his chalky knuckles. It may be mentioned, as sometimes facilitating our diagnosis, that there is another situation in which deposits of urate of soda are even more commonly visible than in connection with the joints, namely, in the cartilage of the external ear, in the form of little pearl-like bodies, varying in size from a split-pea to a mere white point, and situated usually about the fold of the helix. Out of seventeen cases in which Dr. Garrod detected gouty concretions, in seven they existed in the ears *alone*, in nine in the ears *as well as* around the joints, and only in one case were they found in other parts,

but not in the ear. The gouty deposit, which is at first liquid, becomes harder in time from absorption of its fluid parts, and accumulates in many cases to such an extent as to render the joint completely fixed, whilst the neighbouring structures are also inconvenienced by the distension to which they are subjected. The mere presence of the salt in connection with the joint does not appear usually to produce much irritation; after a time, however, inflammatory action, perhaps from a fresh gouty attack, may be set up, the integuments ulcerate, and a discharge, containing the urate of soda mixed with blood-globules, takes place through openings which are often most troublesome to heal. These abscesses, however, it may be mentioned, appear sometimes to act as safety-valves, and their cure is followed by renewed attacks of gout, which had been suspended during the time they remained open. The diagnosis between chronic gouty affections of the joints and the chronic osteo-arthritis already spoken of, is rendered easy when ulceration of the soft parts has occurred, and chalky matter exudes, which can be determined by chemistry to consist of urate of soda. At an earlier period some uncertainty may exist; though the case may be considered gouty, when it comes on after repeated and manifest attacks of gout, and when the gouty diathesis is plainly marked; it may be proved to be so, if a puncture of the swelling with a needle allows of the escape of fluid in which the presence of crystals of the urate of soda is revealed by the microscope: for the existence of this salt forms the specific characteristic of the *gouty* affection, and is never met with in rheumatism or in chronic osteo-arthritis.

Treatment.—As the disease is essentially a constitutional one, the treatment must also be mainly constitutional, and consists in impeding the undue formation of uric acid, as well as in getting rid of the surplus amount of the acid already existing in the blood. For the first purpose the diet has to be carefully attended to; the dyspeptic symptoms relieved as far as possible; the secretions, especially that of the liver, regulated; and the strength of the patient supported in those cases where there is a tendency to loss of tone. To free the blood, we may administer occasionally small doses of colchicum, when the strength of the patient admits of it; but our main efforts are directed to increase the action of the kidneys and skin by means of salines and alkalies. The salines, to be of service in chronic cases of gout, must be given in small doses and very

diluted; they should be taken on an empty stomach and some little time before food. The mineral waters, properly selected, offer many advantages for the administration of saline medicines, as well from the state of dilution in which they are found, as from the greater chance of the patient continuing to take them for a sufficient time, while the mind is occupied and the attention diverted by the change of scene and hopeful statements afforded at the various spas. The external use of the waters can also be conveniently and usefully adopted. The spa to be selected must depend to a great extent upon the special characters of the individual case. In some, Carlsbad, Wiesbaden, or Vichy may be recommended; whilst in other cases, Aix-la-Chapelle, Teplitz, Wildbad, and Buxton are more likely to be of service. Where the debility is great, or after a course of other waters has been taken, the ferruginous springs of Schwalbach or of Pyrmont are often beneficial. For those who are prevented from going to a great distance, or who are desirous of feeling their way and seeing which spring is likely to suit them best, a trial of the carefully prepared factitious mineral waters of the spa at Brighton may at times be deemed advisable, and the Turkish bath is often of much service.

It is, however, for the local mischief that the surgeon may principally expect to be consulted. The local treatment will vary according as it is directed simply to relieve the stiffness and rigidity of the joints, or to alleviate any active inflammatory action which may have been set up, with or without the formation of abscess and production of ulceration. To relieve the stiffness, one topical application has, indeed, been already alluded to, in the form of baths of the mineral waters. Where these cannot be had recourse to, weak alkaline lotions, with iodide of potassium, may be applied to the joint under oiled silk, or, supposing the smaller joints of the fingers to be affected, the whole hand may be wrapped in bandages saturated with water and then enclosed in a waterproof bag during the night. When a certain amount of low or chronic inflammatory action is going on in the ligaments and synovial membranes, small flying blisters are often useful, applied only for a short time, and with tissue-paper between them and the skin; in this particular condition, too, the internal administration of small doses of iodide of potassium is often of use. When acute inflammation occurs, as it may do when a new and severe attack of gout takes place in the seat of a chalk-stone, the

mount of swelling which ensues may cause the cutis to give way, whilst the cuticle remains entire. It is better in these cases to avoid, if possible, making any incision; for the salt is deposited not simply in the cellular tissue or collected as a mass, but also in the actual substance of the articular cartilages, and cannot, therefore, be expected to be entirely or freely evacuated. Soothing applications are of most service; but it may be remarked, that where an opening occurs naturally, or is made by the surgeon, the serious consequences frequently attendant on wounds of articulations are not to be expected, partly from the disorganisation which has occurred, partly also, no doubt, from the unimportant character of the joint which is usually involved. Severe operative measures are not called for in these cases. While the inflammation runs high, poultices and fomentations are of most service; afterwards, gentle pressure may be employed, and the sinuses which are left should not be irritated by probing or by stimulant applications, but allowed gradually to contract and ultimately heal. As already stated, when ulceration has once occurred, the case is almost invariably tedious, and may be rendered more so by too active surgical treatment.

Pyæmic synovitis.—Acute synovitis, frequently leading to suppuration in the joints, is not a very uncommon consequence of purulent or systemic infection, and is met with in phlebitis, after operations, and in connection with certain puerperal conditions, as well as in the progress of fevers, &c. For the description of the general condition giving rise to this complication, its symptoms, and its treatment, the reader is referred to the essay on PYÆMIA. It need only be stated here, that when this pyæmic condition exists, there is always danger of the joints, as well as of other parts of the body, becoming the seat of the local disturbance. Sometimes the attack is marked by the occurrence of violent pain, accompanied or followed by heat, redness, and swelling, affecting one or more of the articulations, which may present distinct fluctuation, provided they are superficially situated. In these cases more or less thin and liquid pus may be found in the interior of the joint, or, in some instances, diffused throughout the external areolar tissue. In other cases, however, large deposits of pus may take place in the joint, suddenly or insidiously, unaccompanied by pain, and perhaps unsuspected during life. The number of joints affected

may vary considerably. Only one may be attacked, or it may be difficult to find a single articulation which does not present pus when cut into. The inflammatory condition may remain confined to the synovial membrane, but frequently extends to the other structures, which become more or less disorganised. In a case of suppuration of the knee, following thecal abscess of the finger, Dr. Handfield Jones found increased vascularity of the synovial membrane, with villous projections from its surface; whilst the articular cartilage was ulcerated in places, the cartilage-corpuscles containing a greatly increased number of celloid masses in their interior, and the intercellular substance being fibrous, and deprived of its natural consistence.

The *treatment* in these cases is mainly directed to the relief of the constitutional infection, and its principles have been laid down in the essay on PYÆMIA. Locally, when we have reason to suppose that a joint is becoming implicated, it should be maintained in a state of perfect rest upon a splint, and the pain relieved by soothing applications. When suppuration has indubitably occurred within the joint, a free incision for the purpose of evacuating the matter is desirable, and the local treatment already recommended for abscess in the synovial membrane becomes applicable.

Gonorrhœal synovitis; gonorrhœal rheumatism.—A form of synovial inflammation is occasionally met with in connection with purulent inflammation of the urethra, or even with purulent ophthalmia. This complication appears to have been first described by Sir Benjamin Brodie, and usually passes now under the name of *gonorrhœal rheumatism*, though it differs in several respects from rheumatism, and is not necessarily connected with gonorrhœa. In these cases, during the existence of a urethral discharge, which may depend on gonorrhœa, or arise after the passage of a bougie, inflammation occurs in the synovial membranes of one or more of the joints, more particularly in the knee. Another peculiarity, which was pointed out by Sir Benjamin Brodie, is, that some form of ophthalmia very commonly accompanies or precedes the articular affection, or may alternate with it. All stages of gonorrhœa may present this complication, and usually no material change in the discharge is noticed, though sometimes it declines a little, when the synovial affection shows itself. The articular inflammation may last for a few weeks, or may continue for

months or even years. It is very apt to involve one joint after another, and presents a decided disposition to relapse. There is little tendency to suppuration, but sometimes the synovial membrane is thickened, and the movements of the joint permanently interfered with. In some few instances the case is much more severe; the cartilages become involved, and permanent rigidity may be the result.

The immediate cause of this affection is still a matter of dispute. By some it has been considered to be connected with the gouty habit;* whilst Mr. Barwell considers it to be 'in reality a slower form than ordinary of purulent infection, produced by inflammation of the prostatic veins.'† He allows, however, that he has 'no cases of dissection to prove this position,' but the following case related by Mr. Prichard certainly seems to favour Mr. Barwell's views.

A patient in the Bristol Infirmary for gonorrhœa and orchitis, was attacked with feverish symptoms, followed by great pain and swelling in the right knee. In a few days the lower part of the thigh was filled with matter which had escaped from the distended synovial membrane. The limb was subsequently amputated, and the joint found to be completely disorganised.

Treatment.—The local treatment in these cases is much the same as that recommended in ordinary synovitis. So long as acute inflammatory symptoms are present, the usual antiphlogistic measures must be had recourse to, and if the pain is very severe, as is sometimes the case, subcutaneous injections of morphia may be employed; when effusion remains, free blistering should be employed, and 'Scott's bandage' applied in those cases where there is thickening about the joint. At a subsequent period, friction, shampooing, local vapour baths, or the use of the mineral waters of Buxton or of Wiesbaden, will probably be of service. The general treatment must also be conducted on the ordinary physiological principles. As the inflammatory excitement passes off, the iodide of potassium, combined with colchicum and opium, may be administered, along with occasional active purgatives; to be followed at a still later period by mild tonics. In the more chronic forms of the disease the greatest benefit will be derived from change of climate, including a residence at the mineral springs recommended for chronic rheumatism, with the use of the waters both internally and

* Johnstone, *On the Genito-Urinary Organs*, p 317.

† *On Diseases of the Joints*, p. 101.

externally. The treatment of this affection by keeping boug in the urethra, or by the administration of cubebs and copai as once recommended, is hardly likely to be often had recourse to in the present day.

Syphilitic synovitis.—The character of the synovial inflammation in some cases is modified by the existence of constitutional syphilis. Occasionally, as Sir Benjamin Brodie remarks, the synovial membranes of one or more joints assume a sluggish form of inflammation in the early stages of secondary syphilis in connection with papular eruptions. More frequently, perhaps, the joint-affection comes on at a more advanced stage of syphilitic infection, in connection with affections of the bones, and especially of the periosteum. The inflammation in most cases is anything but rapid in its progress or acute in its symptoms, though the nocturnal pains may be severe; and it is readily distinguished by its occurrence during the existence of other and characteristic symptoms of venereal blood-poisoning. Its constitutional origin is also marked by its tendency to affect more than one joint. The general treatment is naturally directed to the cure of the syphilitic infection. Mercury may be used in the form of fumigations, in conjunction with the vapour-bath, in many cases where it cannot safely be taken internally. The iodide of potassium, in moderate or, if necessary, in very large doses (forty to sixty grains a day), combined with sarsaparilla or other medicines that tend to strengthen the patient, is often of the greatest benefit. Locally leeches are seldom required. Soothing applications may be had recourse to if the pain is severe, but rest of the limb by means of splints, and mild counter-irritation, are the remedies most frequently of service.

‘Loose Cartilages’ in Joints.

We occasionally find in some of the articulations small movable bodies, which may be quite free or still attached to the walls of the joint by narrow pedicles, and to which the name of ‘loose cartilages’ has been given. These bodies, which are usually met with in adults rather than at an earlier period of life, vary in their structure; some are comparatively soft and of a yellowish colour, and resemble masses of fibrine; in other cases, they are hard and glistening, rather white than yellow, and evidently composed of cartilage or fibro-cartilage, both

the surface and in their interior; frequently, too, they are partially, or sometimes almost completely, converted into bone. In many cases we find only one such body in a single joint, but it is by no means uncommon for two or three to be present, and occasionally they are met with in much higher numbers; Morgagni, for instance, discovered twenty-five smooth and polished globular bodies in the left knee of a woman who died of apoplexy. They vary considerably in size, being sometimes no larger than a barleycorn, whilst in other cases they may nearly equal the patella itself in their dimensions. When small, they are usually round or oval; when of considerable size, they may be elongated, flattened, or tuberos. The most common situation for these 'loose cartilages' is the knee, but they are not confined to that joint; they have also been discovered in the other articulations, including even the hip, which was long supposed to be exempt from them.*

'Loose cartilages,' in many cases, at any rate, appear to take their origin from the small irregular projections which naturally exist in connection with the 'vascular processes' of the synovial membrane. In certain abnormal conditions these processes increase in size and solidity, and may ultimately become detached from their connections, so as to form separate bodies, which are perfectly free in the joint, or they may still remain connected with the parietes by an elongated pedicle. They are composed of 'connective tissue with elongated nuclei, and coated with epithelium, and, though not always, contain a variable number of scattered fat- and true cartilage-cells; and they are not developed externally to the synovial membrane, but from an outgrowth of that membrane itself.'† Occasionally we find imperfectly formed bone in the substance of the masses, derived probably from a further stage of conversion of the cartilage, or they may

FIG. 210.



Cartilaginous growth connected by a narrow pedicle with the coronoid fossa of the humerus. (From a preparation in the Museum of St. George's Hospital. Series iii. No. 145.)

* See *Archives générales de Médecine*, tome xii. (1846), p. 363.

† See note to Kölliker's *Manual of Human Histology*, vol. i. p. 328.

be bony throughout. According to Dr. R. Adams, these foreign bodies, which may be supposed usually to take their origin from the synovial membrane, are especially connected with the condition constituting *chronic osteo-arthritis*, though it is not denied that they are seen in articulations which present no trace of that disease. It is probable, however, that the detached bodies found in joints may have more than one origin; occasionally, perhaps, they may result from a blood-clot which has lost its colouring matter, and now consists only of fibrine; and in many instances they can be traced to an accident by which, probably, a portion of cartilage or fibro-cartilage has been detached, and which may, after a time, become smooth from long-continued friction.

In support of this view, Mr. Brodhurst* relates an interesting case in which, six weeks after the accident, he removed from the knee-joint a 'loose cartilage,' which proved to be the anterior portion of the internal semilunar cartilage, retaining in every respect its normal appearance.

A 'loose cartilage' may exist in the joint for a length of time without producing sufficient disturbance to attract the attention of the patient, or to make him think that he is subject to anything more serious than occasional rheumatic twinges. At last, however, in some movement of the joint, the foreign body slips between the articular ends of the bones, and then a violent and sudden pain is experienced, with inability to use the limb, which remains fixed in the position which it may have happened to assume at the time. The pain continues until the 'cartilage' is dislodged by some means, when the immediate symptoms at once subside, though they are often followed by inflammation of the synovial membrane, accompanied with effusion. The character of the suffering in these cases is peculiar, being often sufficient to make the patient fall, or even to occasion syncope, and is probably due to the sudden stretching of the ligaments caused, in the case of the knee-joint, by the tibia and femur being forcibly separated when the loose cartilage gets between their articular ends during flexion and is accidentally caught during extension, before it has time to slip aside. The attention of the surgeon is now called to the part, and, on careful examination, he may succeed in discovering a firm extraneous body in the joint, which slips beneath the finger, to be lost beneath the patella or the tendinous

* *St. George's Hospital Reports*, vol. ii. p. 142.

structures above or below that bone. When the symptoms described have once occurred, they may be repeated at varying intervals; in some cases recurring on the slightest movement, and even during sleep, whilst in others they may be kept in check so as to give rise to but little practical inconvenience. If the attacks are frequent, however, the joint suffers sooner or later; the ligaments become relaxed, disease extends to the harder tissues, and the articulation is permanently injured. Many of the symptoms produced by a 'loose cartilage,' it may be mentioned, are occasionally met with as the result of partial dislocation of a fibro-cartilage in the knee, as well as, more rarely, in the articulation of the lower jaw, constituting what Mr. Paget terms *locked joint*.^{*} A description of this condition in the knee will be found in the essay on INJURIES OF THE LOWER EXTREMITY (vol. ii. p. 916).

The treatment to be adopted may be either palliative or radical. In the first place, however, any inflammation which may be present must be subdued, and any dropsical effusion got rid of by blisters or other appropriate remedies. When this has been done, a well-fitting elastic bandage or, still better, a back splint, which will prevent the joint from being flexed, should be constantly worn, whilst the loose cartilage may be fixed (by means of pads or strapping) to some part of the synovial membrane, in the hope that absorption may ultimately take place, as will sometimes happen. In many cases this treatment is sufficient; but in some inconvenience is still experienced to such an extent as to suggest further operative measures. When such is the case, the extraneous substance may be removed from the cavity of the joint, either by direct or by subcutaneous incision. Both these operations, however, are attended with a very decided amount of risk, and should only be performed when the palliative treatment has been fully tried and *failed*, and the foreign body continues to give rise to active mischief. The foreign body, too, should be clearly movable, and capable of being brought to a part of the joint in which it is easily accessible. The patient should be prepared for the operation, both by appropriate diet and regimen, and also by the limb being kept perfectly at rest for a few days. The loose cartilage is then guided to that part of the joint where it is most superficial, and fixed securely, so that it may not slip out of the way during the

^{*} *British Medical Journal*, January 5, 1867.

operation. If the *direct* operation is the one adopted, the parts over the foreign body are now divided, and a sufficient opening is made in the synovial membrane to allow the cartilage to be pressed out of the wound, without any unnecessary manipulation or disturbance of the joint. The edges are then accurately brought together by sutures or strips of plaster, and the joint is kept at perfect rest on a splint; constant cold may be applied to the joint, to prevent, if possible, the occurrence of inflammatory symptoms. In the *subcutaneous* operation, the cartilage having been securely fixed as before, a tolerably sharp tenotomy knife is introduced through the skin, at some distance, and passed through the cellular tissue to make a subcutaneous incision into the joint. Great care is taken in dividing the synovial membrane, that the foreign body does not slip aside, for which purpose it is better for the operator to secure it himself with one hand, whilst he makes the incision with the other. The loose cartilage is then pressed either through the aperture in the synovial membrane (as recommended by Mr. Square), or *through* it into the subcutaneous areolar tissue, and retained in either position by appropriate means in the expectation of its becoming ultimately absorbed.

Sir Benjamin Brodie states, in his work on the joints, that his experience does not enable him to determine which of the two operations (the direct, or subcutaneous), is to be preferred. According to M. Larrey, who refers to 167 published cases of the removal of loose cartilages by operation, out of 131 cases in which the old or direct method was employed, 98 were successful, 5 doubtful, and 28 died; whilst of 39 indirect operations, 19 were successful, 15 failed, and 5 died. From this it may be inferred, as M. Larrey states, that extraction by either method is attended with decided danger; but that in the *direct* operation, the risk to life is greater than in the *subcutaneous* operation. On the other hand, in the subcutaneous operation there is increased difficulty, and consequently less chance of success. On account of the difficulty which is experienced in getting the cartilage out of the joint into the cellular tissue without an undesirable amount of manipulation, Mr. Syme recommends another method, by which, he says, he generally succeeds without risk. This consists in 'making a free subcutaneous incision through the synovial membrane and cartilage, and applying a blister over the part where it is retained.'

DISEASES OF THE ARTICULAR EXTREMITIES OF THE BONES.

The diseases of the joints which originate in the articular extremities of the bones are next in importance and in frequency to those which commence in the synovial membrane. It will not be necessary, however, to enter much into detail in treating of them here, as the morbid processes have been already described in the essay on DISEASES OF THE BONES (vol. iii.). A brief notice of these affections in their relation to the joints will therefore be sufficient.

Simple inflammation of the articular ends of the bones, and its consequences.—The articular extremities of bones are liable to attacks of simple inflammation, which may result from injury, or depend on other causes. The changes which take place resemble those which result from inflammation of other tissues, when allowance is made for physical differences of structure. In favourable cases restoration to a healthy condition occurs, or the inflammatory products may give rise to deposition of increased ossific material, leading to hardening and increased weight or size of the part. In less favourable cases suppuration takes place, which may be circumscribed, forming a *local abscess* in the articular extremity, or diffused in the cancellous tissue, as in *osteo-myelitis*. *Local abscess of bone* is usually found in the head of the tibia, and, as a general rule, is recognised and remedied before the articulation is affected. If not, the synovial membrane becomes inflamed from time to time; and if the original mischief is still allowed to remain, the matter may at last make its way into the joint, giving rise to suppuration in that cavity. For a detailed description of the symptoms and treatment of *local abscess in the cancellous tissue*, as well as of *osteo-myelitis* (which may in like manner lead to a discharge of pus into the nearest joint-cavity), the reader is referred to the essay above mentioned.

The inflammation may lead to the death of the whole or a part of the articular extremity, constituting *necrosis*. If the dead portion is limited in its extent and situated near the surface, the articulation may escape with little or no damage, and the sequestrum separate, or be removed by operation. Sometimes the necrosis is deep-seated, presenting then many points of resemblance with local abscess of bone, like which it may lead to destructive inflammation of the joint. In some of these

cases the true state of things is only recognised when death or amputation allows the parts to be dissected; in others, however, a similar operation to that required for local abscess can be performed, the dead bone is removed, the pent-up matter evacuated, and the joint is saved. Sometimes the whole or the greater part of the head of the bone dies, giving rise, of course, to rapid destruction of the joint. As no hope can exist of saving the joint in such cases, its removal, when practicable, becomes the only course to be pursued. In early life, it should be noticed that the *epiphysis* may be separated from the shaft of the bone, and lie loose in the joint, as the result either of acute or chronic disease. In such cases removal of the detached epiphysis will probably often be sufficient, should the true state of the case be detected, which may sometimes happen, though there are no very positive symptoms to distinguish this from other forms of death of the articular extremity. In necrosis affecting the very small joints, such as those of the fingers and toes, the amount of constitutional disturbance involved is so slight that we may wait, if we think it expedient, for the natural separation of the parts. In this way a sufficiently useful thumb or great-toe may be preserved, the superficial position of the articulation facilitating the removal of the dead bone.

Finally, instead of death occurring in obvious masses, the inflammation may lead to a molecular destruction and removal of the osseous tissue, by a process analogous to ulceration, which is termed *caries*. Caries of bone in the vicinity of joints may arise from simple inflammation; it occurs, however, so much more commonly as the result of a low form of inflammation usually described as *scrofulous*, that it is convenient, practically, to defer its consideration, or to merge it in the description of the latter form of disease, to which we now proceed.

Scrofulous inflammation of the articular ends of the bones.—The pathological changes involved in this action, so far as the bone is concerned, are described in the essay on DISEASES OF THE BONES; they need not be recapitulated here. Suffice it to say, that we find the affected bone more vascular than natural (at least at first), then light and oily, as well as unusually soft, so as to admit of being cut with a knife, the cancelli being dilated and filled with a jelly-like substance. This form of inflammation is much disposed to lead to caries. Caries, when affecting the articular extremities of the long bones or the

uboid bones, such as those of the tarsus and carpus, commences frequently in separate points of the cancellous tissue, which gradually extend, and so approach the surface. At last the periosteum itself is reached, and implicated; it becomes detached from the osseous substance beneath, which is left rough and worm-eaten on the surface, bleeding freely, and giving exit to a foul discharge, in which are contained particles of disintegrated bone. The soft parts around share in the diseased action. Serum and lymph are effused into the neighbouring areolar tissue, giving rise to a characteristic firm and elastic swelling; at a later period suppuration occurs, and the tissues are traversed by sinuses.

When the disease is situated sufficiently near a joint, the articular cartilages in the immediate vicinity of the inflamed bone become affected; their nutrient supply, as far as the osteal vessels are concerned, is interfered with; their adhesion to the bone is impaired, and they become fibrous and ulcerate, the change commencing usually on their deeper or attached surface; or portions of them may become detached, and ultimately be found loose within the joint. The synovial membrane participates in the inflammation, which may either spread to it from the diseased cartilages, or may affect it more directly by extension from the periosteum to that part of the synovial capsule which invests the bone to a certain extent before it becomes united with the cartilage and perichondrium. It may be remarked that the affection of the articular ends of the bones sometimes assumes a peculiar form, which has received the name of '*spotted ulceration*,' but which is probably a variety of the strumous disease. When this is the case, several pits or depressions, containing a curdy matter, are found in the articular surfaces, the cartilages presenting a series of defined orifices, corresponding to the pits in the bone, their intervening structure remaining frequently unaltered. The diseased action, which has been supposed hitherto to be confined to one side of the joint, now becomes diffused; the inflamed synovial membrane goes through the series of changes which have already been described under the head of Synovitis; the articular cartilages on the opposite side of the joint become affected, they speedily ulcerate, and the cancellous structure of the bone on which they rest is exposed. The disease gives rise to the formation of matter within the joint; but suppuration, it has already been stated, takes place likewise in the contiguous soft

parts. As the case progresses, the matter makes its way to the surface, it may be directly, in the immediate neighbourhood of the joint, or forming tortuous sinuses, which open at a considerable distance, and give exit to a profuse discharge. When the long bones are affected, partial or incomplete luxations are apt to take place during the later stages of the malady. The disease may continue to progress till life ceases or the joint is removed; but sometimes, after dislocation has occurred, a reparative process takes place, healthy granulations arise in the bone, the synovial membrane goes through the changes tending to recovery, which have already been described, the discharge diminishes and ceases, ankylosis takes place, and the patient is left with a limb the utility of which varies much in different cases.

Scrofulous disease of the joints, commencing in the bones, as well as in the soft structures, is frequently met with in children of strumous habit, especially in those who have never thoroughly recovered from the depressing effects of some of the ailments to which they are subject. At the Hospital for Sick Children I was able, very often, to trace the attack to the weakened state of health which had remained after scarlatina or measles. In early life the disease may affect indifferently the extremities of the long bones or the cuboidal bones; but when it commences after puberty, which is sometimes the case, it shows a decided preference for the carpal or tarsal bones. The disease is very insidious in its earlier stages, occasioning little pain or inconvenience; in children, the first thing which attracts the attention of the parent in most cases is 'limping' when the joints affected are situated in the lower extremity, or unwillingness to use the arm when the upper limb is implicated. It is now noticed too, perhaps, that the child selects that position which will throw least stress upon some particular joint, or that he flinches when that joint is moved or pressed upon; and that the nights are somewhat restless. At this time the little swelling which exists is mainly situated over one of the bones which enters into the composition of the articulation. As the disease advances, the joint itself becomes more decidedly implicated; it is rigid and more or less flexed according to the care with which it has been treated; the swelling increases, and now takes more decidedly the shape of the synovial membrane, and there is usually effusion of serum and lymph into the areolar tissue over the affected bone, distending the skin, and producing

a 'white swelling.' If the case is neglected, painful startings of the limb begin to show themselves, aggravated at night-time, and causing the child to cry out in his sleep, or to wake up with a shriek of pain. When this occurs, it shows that the articular surface of the bone is affected, and beginning to be laid bare to the joint; the 'starting' becomes specially marked, too, when inflammation has extended to the opposite bone, and when, consequently, the two inflamed and sensitive osseous surfaces are subjected to mutual pressure by spasmodic contraction of the muscles. This muscular spasm is originally induced by the osteal disease; eventually, however, it reacts upon that disease, which it aggravates by the forced and violent apposi-

FIG. 211.



The acetabulum almost filled up with bony deposit after recovery from hip-disease, in which the head of the femur was dislocated. (From a preparation in the Museum of St. George's Hospital. Series iii. No. 90.)

tion in which it maintains the inflamed surfaces. Hence the great relief to suffering so often experienced when displacement of the bones takes place, under the influence of the morbid muscular contraction, after the ligaments and other retaining structures have become weakened. When this occurs, the mutual pressure of the inflamed bone-surfaces ceases, and from this period a curative process is often dated. Before this happens, however, matter has in all probability formed in or around the joint, and sinuses are produced, taking various directions in accordance with the anatomical arrangement of the part. If a probe is introduced, carious bone can sometimes be felt, but not always, in consequence of the irregular course which

the sinuses often take. The integuments around the wounds are thin and discoloured, the discharge copious, watery, and unhealthy. The general health by this time usually shows strong signs of breaking up, the child becomes emaciated, hectic manifests itself, profuse night-sweats occur, and, unless relief is afforded, some vital organ becomes implicated, leading to the death of the patient.

Prognosis.—The prospect of a successful issue to the case depends greatly on the treatment being commenced at a very early period, before the joint has become seriously implicated. It is most desirable, therefore, that any expression of pain or tenderness about a joint, or any hesitation in the manner in which it is used, should be at once attended to, and a careful examination made by a competent surgeon. In the wealthier classes this is usually done, and the progress of the case at this period can almost invariably be arrested, and a tolerably perfect cure insured. Even when the disease has arrived at a more advanced stage, when the joint has become implicated, and suppuration has taken place, care and perseverance on the part of surgeon and patient will usually be rewarded ultimately by success; for these cases occur for the most part in early life, and ‘in children the plastic powers of nature are so great that recovery may take place, when in the adult any such hope can scarcely exist.’* These observations, however, apply principally to the wealthier classes; for amongst the poor, who are struggling for their daily bread, early symptoms are generally unnoticed, and the disease has advanced to a great extent before surgical aid is invoked. Even if admitted into hospital, the patient can seldom be kept there long enough for complete recovery; moreover, the general health is apt to suffer, after a time, from the necessarily impure air of a sick-ward, and so the patient returns home, to be again exposed to the imperfect nutrition, the confined air, and the want of rest as well as of necessary comforts and appliances, which originally predisposed to the disease or accelerated its progress. Sooner or later, in all probability, he again applies to the same or some other hospital, in a still more advanced stage of the malady, from which he ultimately sinks, or has to submit to operation. If the prognosis, therefore, in the affluent classes is favourable, it is far less so among the poor and needy. Where circumstances are favourable, and the treatment com-

* *Lectures on the Surgery of Childhood*, by Athol Johnstone.

nenced at an early period, a few weeks may suffice for recovery with a useful and movable limb; even when portions of the cartilage have been destroyed, the functions of the joint may still be preserved, though the time required becomes considerably prolonged. When the disease has advanced to suppuration, and sinuses have formed communicating with the joint, recovery can only be expected at the expense of ankylosis, and after the lapse of many months or even some years.

Treatment.—If inflammation is present, we have always to bear in mind that it is of a low character; it is to the relief, therefore, of the constitutional condition that our general treatment has to be mainly directed. Everything which improves the general health—whether in the form of diet, air, cleanliness, or medicine—must evidently increase the power of resisting disease locally, or of remedying its effects. The best means of effecting this improvement, however, have already been so fully described in previous essays, especially that on SCROFULA, that it is unnecessary to repeat them here; I need only remark that whatever form of tonic may be selected, it must be given for a great length of time to be of service. As Sir Benjamin Brodie directs, the steel may be administered for three or four weeks at a time, and then suspended for a week or ten days; or some other tonic substituted, if the first should have ceased to agree. Even after recovery has taken place, as the constitution is in fault, it is necessary to persevere with the general hygienic rules prescribed in these cases, if we wish to prevent a recurrence of the disease in the same or some other part.

The *local treatment*, when the disease is seen in its *first stage*, is to a great extent negative. The disease is one of defective power; local depletion, therefore, should be avoided, as a rule, though occasionally a few leeches may be employed to relieve rapidly any outburst of acute or sub-acute inflammation. Cold or warm applications can be used at these times, whichever may be most agreeable and soothing to the patient. Counter-irritants, whether in the form of blisters, issues, or the actual cautery, unless in exceptional cases, are seldom required *at this period*, and, indeed, are more likely to do harm than good, by exhausting the patient. They are also objectionable for another reason: these cases usually occur in children, and repeated painful applications, with the frequent dressings subsequently required, induce fits of passion or of apprehension, which are

lowering to the patient and injurious to the malady. In fact, in the treatment of children, it is of great moment for the surgeon to acquire their confidence, to inspire them with a feeling of regard and affection; and this can never be done if his visit becomes frequently associated in their mind with pain and suffering. It may of course be necessary at times to have recourse to operative measures, or thorough examinations of inflamed and sensitive parts; but if practicable, these should be done *once for all*, and usually under the influence of chloroform. Is nothing, then, to be done locally? *Efficient and prolonged rest*, with the limb in a proper position, here as in scrofulous synovitis, is of the greatest moment, and absolutely necessary to prevent further mischief. The limb should be wrapped in a thick layer of cotton-wool, properly secured, pasteboard or other splints adapted to the form of the joint applied, and the whole secured by gummed or starched bandages, which, without being at all tight, should extend so far as to embrace the joints *above* and *below*, as well as the actual one which is diseased. It is necessary that the limb should be fixed to this extent in order that the contraction of those muscles which arise at some distance from the articulation may be prevented, and the joint kept *absolutely in a state of rest*. Support, to a sufficient extent, should be continued to the part for some time after apparent recovery. It is also desirable that the splints should be so adapted as to allow exercise to be taken, whilst immobility of the joint is maintained.

When the disease has been neglected or improperly treated in its first stage, it commonly happens that the limb has been allowed to become distorted, and more or less fixed in its abnormal direction. In such a case it is better at once to restore the limb to its proper position, of course under chloroform, and to adopt efficient means to prevent it from again becoming distorted. The expediency of this treatment depends not merely on the fact that ankylosis is to be expected (and for recovery by ankylosis to be desirable the limb must be fixed in a proper position), but also on account of its immediate influence on the disease. The painful startings are occasioned by pressure on the inflamed bone-surfaces; and so long as undue and irregular pressure is maintained by the muscles in a state of spasmodic contraction, pain must be experienced and irritation kept up. Dr. Bauer has proposed to divide the tendons of the contracting muscles independent of any malposition, as a means of relieving;

the starting pains; and such a plan might not unreasonably be adopted if all milder means of relief should prove unavailing. Supposing the disease to be situated in the lower extremity, a long splint is often employed with great advantage as far as mere rest is concerned, but with comparatively little power of relieving articular pressure. For this latter purpose Sir Benjamin Brodie suggested the application of a moderate but constant extending force, by means of a weight attached to a string, passing over a pulley fixed at the bottom of the bedstead, the other end of the string being secured to the limb, whilst counter-extension was kept up by straps, which prevented the body from slipping down in bed. Of late years extension by weight has come pretty freely into use both in America and this country, in consequence of the advantages which it presents, and the facility with which it can be applied. In Mr. Holmes' *Surgical Treatment of Children's Diseases*, and in some papers by Mr. Marsh in *St. Bartholomew's Hospital Reports*, the subject will be found fully discussed; but I may, perhaps, be allowed briefly to illustrate the plan of treatment by means of a case now under my care at the Brighton Hospital for Sick Children.

A girl about six years of age was admitted with neglected hip-joint disease in the second stage. Thigh much flexed and rigid. Knee turned inwards. Pelvis and spine much twisted. No sinuses or other indications of matter, but painful nocturnal startings and shrieks when the limb is touched. A broad strip of adhesive plaster was applied on either side of the leg from the knee downwards, forming a loop or stirrup some three or four inches below the foot; a flat piece of wood, broader than the foot, was inserted between the two lateral portions of plaster, so as to spread them out and prevent pressure on the malleoli. The plaster was further secured to the limb by two or three circular pieces of strapping, and a light bandage. A stout iron skewer was then passed through a common reel, and the skewer suspended by a couple of straps to the bar at the foot of the child's crib, so that the pulley formed by the reel was opposite the sole of the foot. A bag containing four pounds weight of lead was then fastened to one end of a small chain, the other end being free for the purpose of being fastened to the stirrup of plaster when required.* When all was ready, the child was placed under chloroform, and the pillows removed so that she lay flat on her back. On making gentle extension on the thigh, the muscles began to yield quietly, and then one end of the chain, passed over the reel, was hooked on to the stirrup, the four-pound weight hanging to the other end some distance from the floor. Straightening of the limb progressed steadily and visibly, and as this was going on, a long splint was fastened to the opposite or *sound* side of the patient, to correct the lateral twisting of the body and also to prevent any sudden jumping up in bed under the influence of fright. The hip was enveloped in cotton-wool; sand bags were

* Fig. 119, vol. ii. p. 869, will illustrate this apparatus.

placed on each side of the limb for support: the lower end of the bedstead was raised about three inches by pieces of wood in lieu of counter-extension, to obviate any chance of the body slipping down; and finally a cradle was placed over the foot to keep off the weight of the bed-clothes and prevent eversion. When the effects of the chloroform had passed off, the limb was nearly straight and the child made no particular complaint of pain, but was cheerful and took her meals. There was some 'starting' on the following night, but less than before, and next day the child was comparatively comfortable and happy.

In the second stage of joint disease, counter-irritation is more often required than in the first. The most generally beneficial form, I think, is the steady and continued use of warm applications over a large surface of skin. Iodine is often of service, and blisters occasionally, though these last had better be employed with care and caution in children. Severe pain, continuing after the limb has been properly secured and sufficient extension made, is often relieved by a few lines of the actual cautery.

In the third stage, when matter is forming, the limb must still be retained in a state of perfect rest and extension. Hot applications, or, perhaps, the actual cautery may be employed, but the pain often continues unrelieved till the matter is evacuated by incisions, or has made its own way through the ligamentous structures towards the surface.

If the case progresses favourably, and the sinuses contract and close, the treatment consists in the same local measures as those which have already been pointed out in the section on *strumous synovitis*; but the case may not progress favourably, and it may then become a question of resorting to operation. It has already been stated that in early life the limb may generally be preserved. With the children of the very poor, however, especially those in large towns, and, indeed at all periods of life amongst those who have to gain their livelihood by labour, this question often arises: Is the chance, or even the strong probability, of ultimate recovery with a stiffened limb, after many months, perhaps years, of confinement and forced abstinence from labour, preferable to a more immediate recovery with a loss of the limb or joint, but with all the risks attendant on a grave operation? Every case must be decided on its own particular merits; but having this consideration in view, it is evident that a surgeon is often justified in using the knife even where an operation is not absolutely necessary. When health is failing under the protracted confinement, suffering, and exhausting discharge, removal of the part often

becomes advisable to *preserve life*. Now in weakly subjects, anything which lowers the patient, or effects a drain upon the system, is prone to induce disease; the removal of the source of mischief, therefore, often saves life by preventing more important structures from becoming affected. On the other hand, it must be borne in mind that operations in strumous subjects are, in some cases, rapidly followed by the appearance of tubercle in internal organs, which did not previously exist, or, at any rate, had failed to manifest itself. In deciding the point, the surgeon will undoubtedly be influenced by the position and importance of the diseased part, and the consequent severity of the operation required. To expedite the patient's recovery, he would promptly remove one or several of the tarsal bones, when he would hesitate about amputating the thigh, or excising the head of the femur.

The comparative merits of amputation and excision will be discussed hereafter. (See the essay on EXCISION.)

The articular extremities of the bones are liable to *rheumatic inflammation*. In the essay on DISEASES OF THE BONES, the subject of *rheumatic osteitis* is described; and its relation to diseases of the joints is further discussed in the sections on Chronic Rheumatic Synovitis and Chronic Osteo-Arthritis, to which the reader is referred.

The joints may also suffer from syphilitic inflammation commencing in or extending to the bones. For this also the reader is referred to the essay on DISEASES OF THE BONES; and some further remarks on the subject will be found under the head of Syphilitic Synovitis.

Rachitic affections of the articular ends of the bones.—The bones are materially affected in the constitutional derangement giving rise to rickets. This condition, which is sometimes supposed to be of modern origin, though the lines of Martial* would seem to show that it could not have been unknown in his time, is described in another part of this work.† It is only necessary to remind the reader here, that though the alterations which occur are specially manifested in the *shafts* of the bones, yet the articulations do not entirely escape. In the early stage

* Cum sint crura tibi, simulent quæ cornua lunæ,
In rhytio poteris, Phœbe, lavare pedes.

† See SURGERY OF CHILDHOOD.

of rickets the 'knots' or swellings which take place in the osseous or cartilaginous ends of the bones are highly characteristic of the affection. At the anterior extremities of the ribs these enlargements are particularly obvious, affecting both the bone and the cartilage, and giving rise to an angular projection of the costo-chondral articulations. At a later period the articulations become much weakened, and permanent alteration of shape occurs. Sometimes the rib projects, and the cartilage is forced inwards; more frequently it is the rib which is driven inwards, and the cartilage is dislocated on its external surface, a connection between the two being still maintained by ligamentous tissue. If the chest is examined, the extremities of the ribs will be found presenting a row of nodosities, the sides of the thorax are hollowed, the thoracic capacity diminished, and the function of respiration interfered with to a more or less serious extent. In the limbs similar nodosities are met with at the articular extremities of the bones, first appearing in the lower ends of the radius and ulna, as well as in the malleoli, and afterwards affecting the knees and elbows. As the disease advances, the shafts of the bones undergo the alterations described elsewhere, the bending of the diaphyses occurring more frequently, as well as at an earlier period, in the lower than in the upper extremities. The articulations participate more or less in the deformity, being often twisted in an opposite direction to the curvature which the limb has assumed. The ligaments yield, and incomplete dislocations are apt to take place. At the knee, for instance, displacement may occur outwards or inwards, resulting from a partial yielding of the articular extremities of the femur and tibia, itself induced by the unequal distribution of the weight of the body on the surfaces at the joint.*

The nature of the affection, when the joints are implicated, can scarcely fail to be recognised, in consequence of the rickety condition of other parts of the osseous system, and the period of life at which it occurs. The treatment in these cases is mainly constitutional, to remedy, if possible, the specific diathesis; cod-liver oil, probably, being the medicine which is most generally beneficial. Locally, it is advisable in most instances, when the articulations are disposed to yield and become distorted, to apply mechanical support of such a nature as to counteract the tendency to displacement, and all

* See Bouvier, *Sur les Maladies chroniques de l'Appareil locomoteur*.

of exercise being freely taken. The articular deformity in this way is often relieved, and the results obtained are tolerably satisfactory.

Tumours of the articular ends of the bones.—In the essay on DISEASES OF THE BONES an account of the new formations which occur in connection with the osseous system will be met with. If the articular ends are involved, the joints may evidently become affected either by their movements being mechanically interfered with, or, in the case of cancer, by extension of the disease to other structures, and consequent disorganisation of the articulation. The symptoms, therefore, produced by these formations, especially the malignant, present certain points of resemblance with those of other diseases of the joints, which may be briefly noticed. In some cases, the nature of the tumour for various reasons is obvious enough, and no difficulty arises in the diagnosis, but this is far from being generally, or even usually, the case; the rules, however, by which we are to be guided are laid down in the essay to which the reader has already been referred. The form of malignant disease most frequently met with is the soft or medullary, and according to Rokitsansky it never commences, as a primary disease, within or upon the synovial membrane; the same remark applies with increased force to the cartilages, for the cartilaginous structures are sometimes found intact when every other part of the joint has been destroyed.* Whatever may be effected in the case of innocent formations, in malignant affections the only chance of relief consists in amputation, which should be performed, if possible, so as to remove the whole of the affected bone. Ever when this is done, the chance of permanent success is far from great; but amputation involving the retention of any portion of the cancerous bone is next to hopeless. Obviously, therefore, it is desirable to ascertain in which of the bones the disease began; and by attending to the point where the swelling first manifested itself, this can usually be determined.

ABNORMAL CONDITIONS OF THE ARTICULAR CARTILAGES.

The articular extremities of the bones are protected by a thin layer of cartilage, one surface of which is firmly connected with

* See Virchow, *Cell. Pathol.* Lect. xix.

the bone, whilst the other is free in the joint, except near the circumference, where it is covered by an extension of the periosteum, forming a special coat—*perichondrium*—which gradually terminates on the cartilage, without any well-defined border. It is to this perichondrium that the portion of synovial membrane which advances over the edge of the cartilage is inseparably united.

The articular cartilages, when fully developed, contain neither nerves nor blood-vessels, and yet they are living tissues, though simple in their structure. Nutrition in them, as in the cornea is not effected directly through the means of blood-vessels, but results from the conveyance of the nutrient materials from cell to cell. The cartilage-cells retain their powers of reproduction old cells disappear and are replaced by new ones, and so a round of nutritive changes takes place adapted to serve the wants and maintain the integrity of the tissue. The nutrient materials are obtained from the blood-vessels of the synovial membrane and of the bone; in both cases there exists a free vascular supply in close proximity to the cartilage, round the circumference of which are arranged the *plicæ vasculosæ* of the synovial membranes; whilst we find in the bones, beneath the cartilaginous surface (the *articular lamella*, however, intervening) vascular convolutions, dilatations, or plexuses, specially adapted for the reception of the blood and separation of its plasma.

The functions of articular cartilage are comparatively mechanical, intended as it is by its smoothness and elasticity to diminish friction and obviate the 'jar' or shock to the bones which would otherwise give rise to frequent fractures. Even for these purposes, however, it is necessary that a certain amount of vitality or nutritive power should exist, else the cartilage would gradually wear away in proportion to the amount of friction it has to sustain, and that we know is not the case; for if articular cartilages are somewhat thinner in the old than in the young, the change is not greater than that which is known to occur in other tissues of unquestionable vitality.

So long as the natural processes of nutrition and secretion, as well as the disturbed processes of inflammation, were attributed entirely to some mysterious action on the part of the blood-vessels, the healthy and diseased conditions of the non-vascular tissues scarcely allowed of explanation. But when we admit the nutritive power of the cell- or germ-constituents of the

tissue itself, and consider the vessels as intended to bring supplies of prepared nutriment into sufficiently near relation to them, much of the difficulty vanishes. In perfectly developed articular cartilage we have permanent nucleated cells contained in their capsules and imbedded in an inter-cellular substance; and if we have no vessels in the cartilage itself, they exist in sufficiently close proximity to the tissue for the elements they contain to be within reach of the more direct agents of nutrition. Such an arrangement would be insufficient for organs in which intense action is going on; but for the passive cartilages, the blood-vessels of the bones on which they rest, and of the synovial membranes which surround their circumference, afford a sufficient supply of nutritive material for the cartilage-corpuscles to maintain the integrity of the tissue. If this supply is interfered with, if the circulation in the bone or in the synovial membrane is disturbed beyond a certain point, we can understand that the nutrition of the cartilage should be prone to suffer, and so we find this tissue very liable to be affected *secondarily* in consequence of inflammation, common or specific, occurring in the synovial membrane or in the articular extremity of the bone. Its nutrition, however, may also become abnormal, directly or primarily, from some disturbed action in its own elements, and we have disease of the cartilage, attended with great alteration of structure, where the neighbouring tissues exhibit little or no sign of disease.

Articular cartilage then is so far under the influence of the laws which regulate vital tissue as to be at once capable of maintaining its own status during health, and of undergoing changes, many of which are analogous to those termed inflammatory, in a state of disease. Thus cartilage will maintain itself in its usual condition during a long life; it may be hypertrophied or atrophied; it may degenerate, may ossify, or perish; it may also ulcerate, and probably suppurate.*

Hypertrophy.—Articular cartilages, it is said, are occasionally found to present an increase in their substance without any obvious change in their actual texture. Such a condition, however, appears to be very rare; for, as a general rule, when the cartilage is thicker than natural, the tissue is soft and spongy,

* See Weber, *Ueber die Veränderungen der Knorpel in Gelenkkrankheiten*,—Virchow, *Archiv*, Januar 1858.

with a tendency to break up into fibres, exhibiting, in fact, obvious marks of disease.

Atrophy.—Simple wasting of the tissue in cartilage occurs more certainly than simple hypertrophy. In advanced life, and especially in those joints which are subjected to the greatest amount of pressure, such as the hip, we find the cartilages gradually thinned, their structure in other respects remaining healthy. The process is, of course, a slow one, advancing gradually with age, and constituting, in fact, little more than a sign of natural decay. True atrophy, in most cases, affects the whole surface of the cartilage; but a partial thinning is met with in comparatively early life, accompanied usually with sufficiently marked traces of previous disease to distinguish it from simple wasting.

Gouty deposits in cartilage.—In old and gouty subjects we sometimes meet with a peculiar change in the articular cartilages, resulting from the deposit in their substance of urate of soda. This salt is found not merely on the surface of the cartilage, but also in the inter-cellular substance, through which it is diffused, sometimes to such an extent as almost to supersede the normal structure, which then presents the appearance of a chalky mass. The predilection exhibited by the gouty material for this tissue is further shown by the frequency with which it is deposited in the cartilage of the external ear as well as in the articular cartilages.

Fatty degeneration of cartilage.—In advanced life, in joints which have long been disused, and under other circumstances unfavourable to healthy nutrition, articular cartilages sometimes undergo a process of degeneration corresponding to the analogous affection of the cornea, which passes by the name of *arcus senilis*. In such cases we find the whole or part of the articular surface presenting a series of fibres, free at one extremity where they project into the articulation, and attached at the other to the remaining cartilaginous tissue, exhibiting, indeed, somewhat the appearance of velvet. The cartilage is often thickened and pulpy, and much softer than natural, or it may appear rough and fibrous.

Ossification of cartilage.—Many cartilaginous structures have a natural tendency to ossify as age advances, and we find the

costal cartilages usually, and the thyroid and cricoid cartilages occasionally, undergoing this change, which represents in them a form of degeneration or decay. The articular cartilages, on the other hand, are by no means prone to such a transformation; for, when the developmental ossification is completed, any further tendency in this direction appears, as a rule, to be permanently arrested in the cartilage which remains beyond the articular lamella. Under certain circumstances, however, articular cartilage becomes converted into bone. This change, occasionally met with in the aged, is obvious enough at times during the progress of *chronic osteo-arthritis*; for in joints suffering from that affection, the place of the entire cartilage may be occupied by a particularly dense kind of bone, in which the Haversian canals are filled with calcareous matter; or portions of the cartilage may be replaced by this ivory or porcellaneous material, which remains on a level with the remaining cartilaginous structure.*

Ulceration of cartilage.—It has been already stated, in accordance with the opinion of Weber,† that articular cartilage, like the cornea, is very liable to an altered condition of nutrition, which may fairly be called inflammatory; a condition which shows itself principally in the form of ulceration. The researches of Goodsir and Redfern, followed by those of still more recent authors, have shown that the changes in ulceration arise immediately from a vital alteration occurring in the cartilaginous structure itself, and that the influence of the neighbouring blood-vessels is only indirect or secondary. When the disease is peculiarly active, almost the whole of the cartilage may disappear, a few isolated patches, which are usually reddened, alone remaining. In other cases, the cartilage is softened and abraded; or it appears as if portions of it had been scooped out, leaving smooth pits or depressions. Ulceration usually begins on the free surface of the cartilage; it may commence, however, on its attached surface, or even in its substance.

If we examine the changes as they occur on the free surface after synovitis, we notice that the colour of the cartilage is altered in spots, which either rapidly become holes, looking,

* See the section on Chronic Rheumatic Synovitis and Arthritis.

† See Weber's paper in Virchow's *Archiv* for January 1858.

to use Sir Benjamin Brodie's expression, 'as if they had been cut out with a chisel,' or the spots assume a fibrous aspect, becoming gradually excavated, so as to form pits or depressions with firm margins. Such a condition may be found by destroying the cartilage extensively, even laying bare the bone, without the presence of any membrane. In some cases, however, and especially in strumous synovitis, we find a thickened newly-formed vascular membrane extending from the diseased synovial cavity over the cartilage. Such an extension, indeed, may occur without any ulceration resulting; or even when the cartilage is affected, the excavations are not at all necessarily in connection with the membrane, for they may be found in situations to which the membrane has not extended. As the disease goes on, however, the cartilage, according to Mr. Barwell (in his description of the changes occurring after strumous synovitis), slowly undergoes a transformation into a form of granulation, between which and the synovial material growing from the synovial membrane adhesions form, resulting at last in absolute continuity. If the ulcer extends deeply, the bone beneath participates in the process, its vascularity is increased, and at last the cancellous tissue becomes inflamed. The articular lamella crumbles in masses, carrying with it portions of cartilage which may still be comparatively sound; and the granulations which spring from the exposed cancellous tissue unite with those of the synovial membrane. When the disease originates in the bones, and extends to the cartilage, the first process, according to Mr. Barwell, is one of simple degeneration, followed usually by true inflammatory ulceration. A portion of the cartilage, losing its nutrient supply, degenerates or perishes and is detached with its articular lamella from the inf

FIG. 212.



Incipient ulceration of cartilage. (Museum of St. George's Hospital.)

extends deeply, the bone beneath participates in the process, its vascularity is increased, and at last the cancellous tissue becomes inflamed. The articular lamella crumbles in masses, carrying with it portions of cartilage which may still be comparatively sound; and the granulations which spring from the exposed cancellous tissue unite with those of the synovial membrane. When the disease originates in the bones, and extends to the cartilage, the first process, according to Mr. Barwell, is one of simple degeneration, followed usually by true inflammatory ulceration. A portion of the cartilage, losing its nutrient supply, degenerates or perishes and is detached with its articular lamella from the inf

* Barwell, *On Diseases of the Joints*, p. 113.

surface of the bone, whilst the surrounding portions of cartilage undergo the changes described as occurring in ulceration after synovitis. Inflammatory ulceration of the cartilages, unaccompanied with disease of any other joint-tissue, may possibly take place, though cases of it are not easily met with; degeneration, on the other hand, being of comparatively frequent occurrence.

When the cartilages have been extensively destroyed and the cancellous bone-tissue on each side of the joint exposed, repair, if it occurs, takes place by granulation-tissue uniting the opposing surfaces, and producing ankylosis in the manner already described. Or the ankylosis may be incomplete, portions of cartilage still remaining unaltered. In slighter cases, a natural cure may occur after ulceration has gone on for some time, and affected the cartilage to a considerable depth. This takes place, as Dr. Redfern points out, 'by the formation of a fibro-nucleated membrane from the substance of the cartilage, without the occurrence of any new exudation.' Occasionally we find a patch of ivory deposit occupying the place of some portion of the cartilage, or there may be simply a scar left, marking the place where the ulcer had existed. To complete the sketch of the changes which this tissue undergoes, it may be stated that, in cartilage malignant affections are so rare that it is usually assumed to be altogether insusceptible of them.* For this insusceptibility Virchow gives reasons, founded on his views of the transference of the infection, by means of morbid juices, to the anastomosing elements in the neighbourhood.

Symptoms.—Lesions of cartilage may occur primarily or secondarily. As primary affections, we frequently find after death well-marked 'ulceration' where no symptoms had been noticed during life, excepting perhaps, occasionally, a certain amount of 'crackling' or 'grating' in the joints. These cases are of course chronic in their character, and the affection is of the nature which has been described under the head of degeneration. True 'inflammatory' ulceration is almost invariably secondary, the consequence of inflammation, common or specific,

FIG. 213.



Ulceration of cartilage. (From a preparation in the Museum of St. George's Hospital.)

* See Chance's edition of Virchow's *Cellular Pathology*.

occurring originally in the synovial membrane or in the cartilage. This affection may be acute or chronic in its progress. Lawrence * relates the case of a patient who died after amputation, in whom, as he believed, complete destruction of the articular cartilages of the femur and tibia took place in a few days. In most cases, however, it is much slower in its progress and in some, especially in old diseases of rheumatic origin, may be extremely chronic in its character.

Degenerative ulceration of cartilage is attended with a marked symptom, gives rise to no pain. Is pain produced by 'inflammatory' ulceration? The deservedly great authority Sir Benjamin Brodie for a long time connected with ulceration of the cartilages those severe pains which are increased at times, aggravated by any motion, and attended with involuntary startings of the limb. The fact, however, that no new disease has been discovered in articular cartilage, even with our improved powers of investigation, renders such a view inadmissible, unless we conclude that our means of investigation are still too defective to warrant our placing trust in them. In his last edition, however, Sir Benjamin Brodie avows that he is 'inclined to the opinion that the increased sensibility in these cases is in the bony plate beneath the cartilage rather than in the cartilage itself; and that the presence of these pains, with involuntary startings of the limb, is always regarded as a sign of the bone partaking of the disease. We may assume then, that disease of articular cartilages is attended with no distinctive or characteristic symptoms (if we except the 'crackling' on motion, mechanically attendant upon the roughness of their surface). In cases of disease commencing in the synovial membrane when starting pains come on, or may be taken indeed as evidence, not merely that the cartilage is affected, but that the disease has further extended to the bone. When the disease, on the other hand, has commenced in the bone and involved the cartilages, the joint may speedily become opened; and the suppuration and abscess which occur are attended with the symptoms which have already been described. The treatment, as well as the diagnosis of disease of articular cartilage, so far as they can be ascertained, are found in the description of the affections of the other structures of the joints.

* 'Lectures on Surgery,' *Lancet*, 1829-30.

on the other hand, may *originate* in the peri-articular
s, and either simulate disease of the joint or lead
o it. The variety of these lesions is so great that it is
le, within the limits of this essay, to attempt a detailed
tion of them, and this indeed would be unnecessary, as
cription will be found in other parts of the work. A
arks, however, upon the relation of each of these
es to the pathology or diagnosis of joint-disease may
haps, be considered out of place.

Diseases affecting the Ligamentous Structures.

Ligaments and fibrous structures around the joints are no
liable to inflammation, acute or chronic, in consequence
of injuries which they sustain from sprains or other acci-
(see the essays on injuries of the different regions). Slow
ly, the reparative process appears to be slow also; for it
frequently happens that the movements of a joint are hampered,
its use attended with great pain for a long time, after
a comparatively slight injury, in which the ligamentous or
fibrous tissues alone have apparently been injured. Oppor-
tunities of examining the parts in these chronic cases have not
often been afforded, or have not, at least, been often made use
of; but occasionally the ligaments have been found thickened
and pulpy, independent of any actual laceration of their sub-
stance. From the frequency with which chronic or saturnine

believe that the disease may in many instances originate in these structures.

In the sketch of the advanced stages of articular disease we have seen the ligaments ulcerating or yielding, and so permitting the exit of matter from the interior of the synovial membrane, or allowing complete or partial luxations to occur. Sometimes, on the contrary, we find layers of a fibrous tissue, having somewhat of a lardaceous appearance and occasionally attaining a very considerable thickness, developed in the ligaments or in the areolar tissue around them, and forming the immediate cause of what is termed extra-capsular false ankylosis. In rheumatic and syphilitic affections, it is probable that the ligamentous structures share in, or even originate, the diseased action, and at least contribute to the pain which is so severely felt. In gout they may form the seat in which the urate of soda is deposited. In chronic rheumatic arthritis they suffer to a great extent; the capsular ligaments may be greatly increased in thickness, or portions of bone may be developed in their substance; and in the hip the round ligament may entirely disappear, or in the shoulder no trace of the glenoid ligament be left.

The ligaments, it was said, are liable to become relaxed in the progress of articular disease; a similar looseness of them is sometimes found as a primary affection, or at least dependent only upon some constitutional condition. In persons of lax fibres, especially in delicate or hysterical young women, a preternatural mobility of the joints is not uncommon, so that extension of the fingers or thumb may be carried nearly as far as flexion, and the nails and the back of the hands be made to meet. In many cases this state is not attended with any practical inconvenience; and even where partial luxations occur, the bones return at once to their normal position. Occasionally, however, it happens, perhaps in those persons in whom the undue mobility of the joints is not attended with a corresponding relaxed condition of the muscles, that complete dislocation takes place, which may not be reduced spontaneously, but may require surgical assistance. Instances of this kind, involving an unusual liability to luxation, are frequently recorded. A similar condition of relaxation of the ligaments, involving loss of power of the articulation, and leading occasionally to luxation, may follow palsy of a limb, and is often met with as a result of infantile paralysis. In many children it is necessary

to have recourse to mechanical contrivances to obviate this weakness, whilst means are being adopted to increase the muscular power by which the joint may be moved or the limb exercised. The symptoms, progress, and treatment of articular disease, as it implicates the ligaments, will be found included in the preceding sections.

Diseases affecting the Muscles.

The altered conditions of the muscles and their tendons by which the joints are forced into certain positions, or retained in the positions they have assumed, may be primary or secondary in their relation to the articulation. In the latter case disease commences in the joint, and the muscles are affected subsequently, contracting spasmodically from reflex irritation, so as to aggravate the symptoms and give rise to increased mischief by the inter-articular pressure they produce, as well as by the tendency to partial or complete luxation caused by the predominance of one group of muscles over its antagonists; at a later stage, too, they become fixed and rigid, so as to impede functional exercise of the joint even when the original articular affection is subdued. Hence the importance, which has already been insisted on, of early attention to the position of the limb, and the necessity of counteracting the reflex muscular action by appropriate splints or by properly applied extension. The muscular lesion, on the other hand, may be primary, as far at least as the joint is concerned; from some disturbance in the nervous system, an undue or unbalanced contraction of certain muscles occurs, and the articulation is fixed in a particular position, which becomes permanent if the muscular structure degenerates and loses its proper contractile material. Such a condition of spastic rigidity is sometimes found in the adult, the result usually of serious disease affecting the cerebro-spinal axis. It is much more common in early life, as a consequence of the infantile paralysis which, if seldom fatal, is frequently quite incurable. In many of these cases the amount of deformity which occurs is deplorable, and the malposition of the joint, so long as it lasts, obviates any hope of stimulating muscles to contract, or prevents that exercise of the limb on which any prospect of improvement depends.

Rigidity of the muscles, terminating, perhaps, in atrophy or degeneration, is often seen when a limb has been maintained for too long a time in one position, and especially when pressure

has also been exercised on the muscles and their tendons, and their action arrested by the application of tight bandages. This condition of the muscles may also be accompanied by grave mischief within the joint. After fractures requiring prolonged immobility of the limb, or at least in which the limb has been kept fixed for a length of time, rigidity of the parts takes place to a greater or less extent, and is very obstinate in its duration. In some of these cases, and especially when the patient is advanced in life, or the constitution is otherwise predisposed to such alterations, we may find, as M. Tessier has pointed out, effusion of serum, mixed perhaps with blood, into the joint, the synovial membrane at the same time being injected and its vascular fringes swollen. In other cases the mischief may have advanced still farther, and the articular cartilages have materially suffered, becoming reddened from contact with the effused blood, and softened in their structure, or presenting actual loss of substance on their surface. These changes, it may be remarked, are not confined to the articulation in the immediate neighbourhood of the fracture, but extend to others which have been maintained in a similar state of fixedness. The importance of due action of the muscles need scarcely, therefore, be insisted on, or the propriety of having recourse to passive motion at the earliest period of which the case will admit.

Finally, it may be noticed that an articulation is sometimes found to be perfectly rigid, from muscular contraction, in that condition of the nervous system which we term hysterical (see Articular Neuralgia). By examining the limb, however, during natural sleep, or in the insensibility produced by chloroform, the relaxed condition into which the muscles fall, and the free movements which can be impressed upon the joint, leave no doubt as to the true nature of the case.

The subject of undue or impaired muscular contraction, and its results, will be found treated in the essay on ORTHOPÆDIC SURGERY (vol. iii.) ; but it plays so important a part in articular disease, that it was impossible to avoid all notice of it in this place.

Diseases affecting the Areolar and Adipose Tissues.

The areolar tissue around the articular extremities of the bones, or over the synovial membranes, commonly participates in the diseases of these structures, and is either simply cedematous, or, in severe cases, becomes the seat of abscess or permeated by sinuses. Acute inflammation, on the other hand, not

frequently attacks the areolar tissue in the neighbourhood of joints, as the result of blows or contusions, of wounds, or of erysipelas. It occurs, too, in many cases as a secondary consequence of inflammation of the synovial bursæ. The inflammatory action frequently runs high, and there is much constitutional as well as local disturbance. The swelling is considerable, the part hot and tender, the skin reddened, and any movement of the joint, including the articulation, is attended with pain. There is usually much tendency to suppurate, and when the synovial bursæ are implicated, matter forms rapidly and in considerable quantity, presenting some resemblance to acute abscess of the articular cavity. The diagnosis, however, between this affection and disease originating within the joint, is not difficult. The rapidity with which the symptoms have come on distinguish it from disease of the bones; whilst the position of the swelling, which extends over and conceals the osseous prominences, prevents it from being confounded with synovitis. By careful manipulation, too, it will be found that the joint itself may be handled without producing pain, provided that no pressure or tension is exercised upon the inflamed parts around. This kind of inflammation is not uncommon in delicate or strumous children, in whom it is very apt to involve the articulation, unless great care is taken. The knee and the shoulder are the two situations, perhaps, in which the affection is most frequently seen.

The treatment in these cases is sufficiently obvious. The limb must be kept at perfect rest; poultices and fomentations applied; and if matter forms, early and free incisions should be made for the purpose of evacuating it, lest the pus should find its way into the synovial cavity, and the case become one of serious consequence, resolving itself, in fact, into acute suppuration of a joint. The incisions, though free, should not, of course, be so deep as to open the articulation, under the mistaken idea that this is already implicated. In strumous children the general health has to be sedulously attended to, and care taken not to depress the constitutional powers.

With respect to the affections of the adipose tissue, there is little to be said; according to Rokitsansky, a branching growth of fatty tissue is occasionally met with, chiefly in the knee-joint, in the free part or in the duplicatures of the synovial membrane corresponding to the *lipoma arborescens* of J. Müller.

Diseases of the Synovial Bursæ.

A description of the pathological affections to which the synovial bursæ are liable will be found in the essay on AFFECTIONS OF THE MUSCULAR SYSTEM, to which the reader is referred. The situation of the subcutaneous bursæ, at least of those which are most liable to disease, is well known, and they are at once accessible to examination, and the symptoms attendant on their morbid conditions so characteristic, that mistakes concerning them are hardly likely to arise. In the case of the deep-seated bursæ, however, this is by no means so certain. Their diseases simulate to a much greater extent disease of the neighbouring joints, and, indeed, frequently lead directly to it; for in many instances these bursæ are continuous, or have some connection with the articular synovial membranes, so that direct extension of inflammation may occur from one to the other. In the description, therefore, of the special characters which attach to the diseases of the individual joints, I shall have occasion again to refer to the deep-seated bursæ, and to the means of distinguishing their affections from those of the articulations.

Altered Conditions of the Integuments around the Joints.

The diarthrodial joints are intended to allow of free movements of the bones which enter into their composition. If these movements are interfered with, the joint is truly affected, whether the seat of the obstruction is in one of the proper articular structures, or has only an indirect connection with them. Whenever the integuments have been extensively destroyed around a joint, the resulting cicatrix, and the accompanying induration of the subcutaneous areolar tissue, may cause the articulation to be partially or almost wholly rigid. The cases in which this usually occurs are those resulting from extensive burns; and here too we often find a limb rendered useless, and hideous deformity produced, by the original extent of the injury, or by the want of proper forethought during the healing process. In the essays on BURNS and on PLASTIC SURGERY, the importance of considering the position to be chosen for the limb, when the burn has invaded the circumference of an articulation, has been pointed out, as well as the best means to be adopted for preventing injurious contraction at the time, or for remedying it if it has unfortunately occurred. I have thought it right, however, not

entirely to omit in this essay any allusion to the fact that a natural condition of the surrounding integuments is of importance to the integrity of the joint.


ANCHYLOSIS.

A want of precision occasionally attaches to the sense in which the term ankylosis is used. Originally, as the word itself implies, it was applied solely to joints which had become fixed in an *angular* or bent position; rigidity when the limb was straight receiving another designation (*ὀρθόκωλον*). The position which the limb has assumed is no longer specially designated, but we speak at the present day, somewhat clumsily it is true, of ankylosis as being *true* or *false*; the latter, at any rate, having a somewhat vague signification in the minds of many authors. By *true ankylosis* (*synostosis*) is meant actual osseous union of the contiguous bones. This implies, in most cases, destruction of the entire joint, and constitutes a further or final stage of the process of junction by fibrous tissue, which has been described in previous sections, and which forms one variety of false ankylosis. A

form of osseous ankylosis, moreover, is sometimes met with in which the new bone is placed *outside* the articulation. This variety is most common in the spine, which occasionally presents superficial bridge-like bony processes extending from one vertebra to another, or even forming a kind of osseous sheath enclosing several vertebrae.

False ankylosis may be *intra-capsular*, when it signifies a junction of the articular surfaces of the contiguous bones by a comparatively soft or membranous tissue (the joint-cavity being partially or completely destroyed), or it may be *extra-capsular*, the union of the bones in this case depending on adhesions

FIG. 214.



Section of a hip, presenting complete bony ankylosis of the ilium and femur. (From a preparation in the Museum of St. George's Hospital.)

around the articulation, with thickening of capsular or ligamentous tissues in that situation. Extra-capsular false ankylosis, usually accompanied by a false joint, but it may exist alone, as was seen in a case of Bonnet, where the articulation was completely destroyed, section of all the muscles around had been performed, whilst the interior of the joint was found to be free. The ankylosis resulting from the presence of fibrous tissue in the ham uniting the bones. Impaired mobility of the joints may result from various causes. In chronic osteo-arthritis, for example, the articular surfaces are altered, or bony growths form around them, which mechanically impede the motion. A joint may be crippled by the contraction of muscles, or by injuries or diseases affecting the integument. Muscular retraction, too (so called contracture ankylosis), may alone interfere with motion. These cases, however, may be taken as examples rather than ankylosis.

Ankylosis is occasionally seen in adults, and is sometimes the cause of the minor joints without the existence of any other disease to account for its presence. As a rule, it is either true or false, results from a variety of causes, whether specific, which has affected a joint, or from a general amount of destruction of the parts, or from a defective reparative action. In the majority of cases, the difficulty in determining the fact that a joint is ankylosed is if not altogether lost. In the hip joint, it is necessary to be careful that increase of motion is not taken for action of the articulation. In the knee, the exact cause of the rigidity is not always clear. It may be osteo-arthritis, irregular osseous growths, or a false joint. When retraction of the muscles is the cause of the impaired mobility, it is not always apparent beneath the skin. If the areolar tissue is indurated, or at one part of it is indurated, we may suspect that there is ankylosis, which it may be not entirely absent. Subsequently, when the mobility of the joint is lost, the means at our disposal have been

* Bonnet, *Sur les Maladies des*

It may happen, the muscles only after chloroform. being restored as may be practised of the operation. Or, as M. has been practised restoration of the most practicable long but fine teguments punctured section of the on the admission; which are being

main which is extensive hours, but usually, of any very degeneration of the joint in the manner described effectually secured at the Hospital wrapped in cotton as recommended by been restored to the symptoms produced by it, if the case is one of such measures as the joints. For this, and steady and, which may be permanent (the child); or still other contrivances, which is a full description.

From these proceedings, of the lesion are mainly specially in childhood, at an early constitutional condition favourable. In middle life, or at least of much improvement, and as advanced age is

augmented by modern discoveries : by the employment of chloroform we annihilate pain, and get rid of the active muscular contraction which would greatly impede our efforts ; and by means of subcutaneous surgery we are furnished with a safe and ready method of overcoming the passive or permanently rigid state into which some of the muscles are likely to have fallen ; a state which adds materially to the resistance offered by the joint.

The replacement may be immediate or gradual. In a large number of cases, where the required amount of alteration is not very considerable, the immediate process is preferable, the chance of mischief arising in consequence of the sudden change from a position which may have lasted for a length of time, not being very great if the operation is properly performed, and the after-treatment careful and judicious. In many instances, however, the restoration of shape can be better effected gradually or painlessly by simple mechanical contrivances carefully adapted to the individual joint, and in such cases this plan of treatment may properly be adopted. (See the essay on ORTHOPÆDIC SURGERY.) If immediate reduction is determined on, it is desirable that the surgeon should have a clear notion of what he is about to do ; that he should have determined the position to be chosen, and calculated the obstacles to be overcome. The key to success in the process of immediate replacement consists in the previous manipulation. When the patient is placed under chloroform, therefore, the movements for the purpose of breaking down the adhesions which have formed should be properly adapted to fulfil their purpose. Instead of applying the force employed simply in one direction, by attempting at once to straighten the limb, the first process is to move the joint in every sense so as to render it as supple as possible. Accordingly the limb is to be alternately flexed and extended (or, it may be, abducted and adducted likewise) steadily and gradually with somewhat increasing force, till as much as possible of the natural extent of motion is obtained. When the adhesions are sufficiently broken down, the limb may then be placed, by appropriate traction and pressure, in a proper position, and secured on a splint.

Many cases of ankylosis, it was stated, are complicated by the existence of muscular retraction, which prevents the necessary movements for the relief of the ankylosis. If the existence of this retraction is evident, the muscles may be divided subcu-

taneously a few days before the main operation. It may happen, however, that a retracted condition of one or more muscles only becomes evident during the manipulation under chloroform. One of two courses may then be adopted. Having restored as much motion as possible to the joint, tenotomy may be practised on the opposing muscles, and the remainder of the operation suspended till the wounds have entirely healed. Or, as M. Bonnet recommends, after subcutaneous section has been practised, the movements may be persevered in till restoration of the articular functions is effected to the greatest practicable extent. If the latter plan is adopted, a very long but fine tenotomy-knife should be employed, and the integuments punctured at a considerable distance from the point of section of the muscle, to avoid if possible the risk attendant on the admission of air to the cut surface during the movements which are being performed.

As the effects of chloroform pass off, the pain which is experienced may be expected to be severe for some hours, but speedily subsides, without the development, usually, of any very severe inflammatory symptoms. Constant irrigation of the joint may be practised for the first day or two in the manner described in a preceding section, the limb being effectually secured on an open splint (the plan I usually adopted at the Hospital for Sick Children); or the whole limb may be wrapped in cotton wool, and the bandage at once starched, as recommended by Bonnet. When the proper direction has been restored to the ankylosed limb, and any inflammatory symptoms produced by the force employed have subsided, it is time, if the case is one which allows of any hope of success, to adopt such measures as may tend to develop renewed mobility of the joints. For this purpose, baths and friction may be employed, and steady and judicious passive motion had recourse to, which may be performed by the hand (usually sufficient in the child); or still better, in the adult, by various mechanical contrivances, which could hardly be understood from any verbal description.

In estimating the chances of success from these proceedings, the age of the patient and the duration of the lesion are mainly to be considered. In youth, and especially in childhood, at an age, that is, under sixteen, assuming the constitutional condition to be good, the results are usually favourable. In middle life, there is still a fair hope of success, or at least of much improvement. Beyond this period, however, and as advanced age is

attained, the chances are much diminished; and if the lesion is of old standing, if the adhesions have acquired great solidity, or if the anchylosis is complicated with other local mischief, not much hope of benefit can fairly be entertained.

When the anchylosis is *osseous*, the means already spoken of are obviously inapplicable. Supposing the position in which the anchylosis has occurred to be inconvenient, it has been proposed by Mr. Barton of Philadelphia to remove a wedge-shaped portion of bone from the projecting surface of the angle formed by the limb; the extent of the broad part of the wedge varying, of course, with the amount of the deformity. When the osseous wedge has been removed, either from the anchylosed part itself, or a little above or below this point, as may be considered expedient, the remaining thickness of the bone is fractured, and the two surfaces brought together so as to redress the angular deformity previously existing.*

Finally, cases of anchylosis are sometimes met with in which the limb is not merely useless, but an incumbrance, whilst the continuance of disease in the bones, or the existence of other complications affecting the surrounding structures, prevents replacement from being effected. It may then become expedient to remove the diseased joint, or even the entire limb. The propriety of such operations must obviously depend upon the merits or peculiarities of each individual case.

ARTICULAR NEURALGIA.

We frequently meet with cases where pains are experienced in the joints without any appreciable material lesion to account for them. In many instances these are termed, and no doubt correctly, rheumatic or gouty pains, for these two conditions may have manifested themselves unquestionably in other articulations, or may ultimately appear in those which have been the seat of the erratic suffering. Occasionally pain gives a brief warning of mischief about to be developed, which it precedes only by a short time. In the pyæmic condition, for instance, sharp pain may be felt in some joint, and though no other sign of diseased action is present, we are led to expect the formation of matter in this situation, and the progress of events common to it justifies our apprehensions.

* See also the essay on ORTHOPÆDIC SURGERY, vol. iii. p. 722, for an account of a less severe form of operation recently introduced by Professor Gross.

The pain may have a local origin, independent of any disturbance in the joint in which it is experienced. The pressure of a tumour on some nervous trunk may give rise to suffering referred to its articular branches; and all surgeons are familiar with the fact, that pain in the knee is one symptom of disease within the hip, and yet this sympathetic affection may so mask the original disease, that I have frequently had children with diseased hip brought to me, in whom blisters or other local remedies had been applied to the knee, under the idea that *it* was the seat of mischief.

Unquestionably, however, the majority of these cases are dependent on that abnormal condition of the nervous functions which is termed hysterical. The condition, indeed, is not absolutely confined to the female sex, as the original signification of the term would imply (*ὑστέρα*, womb), for it may occasionally be met with in males, though far less frequently than in the other sex. When this morbid condition of the nervous system prevails, a local direction is often given to the perverted sensibility, by some accidental concentration of the '*attention*' on the part. The physiological effects of undue '*attention*' on organs are of great interest, and serve to explain many curious phenomena exhibited in certain disturbed conditions of the nerve-centres, which may be induced artificially, as in hypnotism, or arise from less obvious internal causes, as in hypochondriasis. I am unable in this essay to devote more space to this subject, but must confine myself to a simple notice of the effects which may be produced by involuntary or automatic '*attention*.' A woman is brought into close relation with some one suffering from cancer of the breast; the attention is directed involuntarily to the corresponding organ in herself; the part becomes painful, swelling even occurs, and what is termed neuralgia of the breast is set up. A case is related of a gentleman who lost an intimate friend from cancer of the œsophagus: the sufferings which he witnessed made a strong impression upon his mind; he began himself to experience difficulty of swallowing, and ultimately died from the effects of spasmodic dysphagia.

In hysteria, when the attention has been fixed on an articulation from mental or from emotional causes, or in consequence of some slight injury, symptoms which might be supposed to indicate structural disease of the joint are apt to show themselves. There is great *pain*; pain out of proportion to the other symptoms, and frequently extending over a large surface,

but often suspended if the attention is diverted, and completely absent during sleep, which may be prolonged and sound. The *tenderness* is also exaggerated, and usually greater in the integuments than in the deeper structures; it is little felt, too, on firm pressure on the articular surfaces effected from a distance. There is sometimes a slight amount of *swelling*, which is diffused in the external areolar tissue, and arises from a turgid condition of the small vessels, or from increased effusion into the areolar interspaces. On handling the part, *fine crepitation* is often felt, evidently situated near the surface, and differing essentially from the grating or crackling which may be attendant on removal of the cartilages. Frequently *convulsive* or *spasmodic action of the muscles* is induced by any attempt to bend or straighten the joint; or sometimes there is a more permanent *rigidity*, by which the limb is fixed in some improper position. When volition is suspended, however, the rigidity and the spasm are suspended likewise. General symptoms indicating *hysteria* are often present; in many cases there is habitual coldness of the extremities, with other evidence of weak circulation; but articular neuralgia is sometimes seen in stout and florid girls. Some irregularity of the menstrual functions is usually present.

The diagnosis in these cases is generally not difficult, for it is commonly easy to detect symptoms which unequivocally vary from those which would be induced by true structural disease. It may be mentioned, however, that in old cases the patient may have acquired so much information regarding the symptoms which ought to be present, that less chance is afforded to the surgeon of detecting any palpable inconsistency. Where any doubt exists as to the true character of the affection, time should be taken before a positive opinion is pronounced; a careful examination, too, should be made, to detect and remove any local cause of nervous irritation; for these sympathetic pains may be maintained by disturbance of parts with which there is indirect nervous communication, and relief of the one will be followed by cessation of the other.

The general principles of the treatment have been laid down in the essay on HYSTERIA. In most cases tonics and antispasmodics are beneficial. Great attention should be paid to the due action of the skin and liver; and where a tendency to periodicity is present, quinine is of marked service. Change of air and occupation are beneficial, especially residence at the sea-side, where salt-water bathing, general or local, can be

easily enjoyed. The patient should be urged and encouraged to use the articulation. If the limb is contracted, and any attempt at restoring the joint to its natural position is strongly resisted, chloroform may be had recourse to, and the part maintained afterwards for a *short* time on a splint, with advantage; but exercise of the joint should speedily be enforced. The moral treatment to be adopted is of material importance. The attention of the patient must especially be directed elsewhere, every effort should be made to cultivate the power of 'self-control,' and healthy occupation given, if possible, to the mental as well as to the bodily functions. A judicious mixture of firmness with kindness must also be exhibited both by friends and surgeon, and the confidence of the patient acquired, which can never be done if any form of deception is had recourse to, as has sometimes been recommended.

INJURIES OF THE JOINTS.

Wounds of Joints.

The joints are liable to various and severe injuries. The ligaments and other soft parts in the neighbourhood may be strained or otherwise damaged; the bones may be suddenly forced apart, and their natural relation to each other altered; or a fracture may extend into and implicate an articulation. These injuries, however, have already been described, generally, under the heads of DISLOCATION and FRACTURE, or, as they affect the individual joints, in the essays which treat of injuries in the different regions. The subject of *wounds of the articulations* is the only one that remains, therefore, to be noticed in this place.

In treating of wounds of the joints, I shall confine myself to those in which the synovial membrane is actually penetrated, indicated by the presence in the blood of thin lines or streams of semi-transparent glairy synovia. The wounds may be punctured, incised, or contused. The gravity of a *punctured* wound is influenced to a certain extent by the course which it takes, as well as by the nature of the instrument with which it is inflicted. When oblique in its direction, of narrow dimensions, and made with a sharp clean weapon, it corresponds to the wounds produced in subcutaneous surgery, and has comparatively little tendency to give rise to suppuration; a direct wound, on the other hand, even of the same size, is more prone to produce serious mischief, especially if it is made with a blunt or jagged

instrument. To a certain extent, the same observations apply to *incised* wounds, but their greater dimensions much increase the chance of the admission of air, and of the supervention of destructive inflammation, whatever the direction they have taken. *Contused* wounds of a joint may be inflicted from within or from without. The former have been described under the head of Compound Dislocation; the latter are produced by heavy or blunt instruments, and are formidable from the injury of the soft parts they involve, and the consequent chance of mortification, as well as from the inflammation they may be expected to induce.

Under favourable circumstances, the edges of an incised or punctured wound, if maintained in perfect apposition, may unite speedily by the processes described in the essay on WOUNDS, and the articulation be restored to a sound and healthy condition. When the wound is much contused, the edges gaping, or when other circumstances are not favourable, severe inflammation takes place, suppuration rapidly ensues, and destruction of the joint may be expected to occur. The mischief may not even be confined to the articulation; pyæmia may be established; or the patient may sink, exhausted by the profuse discharge, and the fever excited by the local irritation; there is also a risk of tetanus making its appearance.

The causes of the grave character of these lesions are not, perhaps, difficult of comprehension. When early union does not take place, and repair has to be effected by what is called the 'second intention,' suppuration occurs; but the matter is formed not merely at the divided surfaces, as in ordinary wounds, but extends to, and accumulates in, a large and often intricate synovial cavity, which does not communicate freely with the surface, but allows of burrowing and confinement of the pus, and so favours its decomposition, with the consequent constitutional disturbances. The structure of some of the tissues which form the walls of the cavity is also most unfavourable when suppuration has occurred; instead of being vascular and full of life, soft and able to approximate and contract, the cartilaginous surfaces in the joints are unyielding, and so low in their vitality as to be little capable of efforts at reparative action; even when the cartilages disappear, the osseous structures which are exposed are themselves far less adapted for the process of granulation than the softer tissues, and obliteration of the pus-forming cavity becomes a slow and tedious process,

then interrupted by local or constitutional mishaps. Finally, even if repair does take place, it has been effected at the expense of the functions of the part; for union of the opposing surfaces of an articulation involves ankylosis, and consequent loss of mobility—the purpose for which the joint was constructed.

When a wound exists in the neighbourhood of an articulation, it may be of such a size, and have so freely exposed the interior of the joint, that no doubt can exist as to its extent. In certain cases, however, it is not always easy to determine whether the synovial membrane has been injured or has remained intact. We are guided in our opinion by the position of the wound, its direction, the degree of penetration, and the form of the weapon. Our diagnosis may be assisted by the presence in the flowing blood of semi-transparent glairy synovia in thin lines or streams; but the absence of synovia does not prove that the articulation is uninjured (for its escape may be prevented by various circumstances), and its presence is not conclusive in favour of injury, for the fluid may have been discharged from a synovial bursa or from a tendinous sheath. When any doubt exists, researches with a probe had better be avoided, and the case treated at first as if penetration had undoubtedly occurred.

The dangers attendant on a wound extending into an articulation are influenced by several circumstances. In early life such wounds are usually attended with less severity of symptoms than at a more advanced age, and a good constitution is more likely to do well than one which has been broken down by prolonged debauchery. A small joint will escape with impunity where a large one would be destroyed; and even if the articulation is destroyed, the general disturbance may be trifling. The upper extremity is more favourably situated than the lower, as far as the chance of recovery is concerned; and finally a wound inflicted with a sharp instrument is more likely to unite than one which has been attended with much violence to the softer tissues.

In managing a wound of a joint, it becomes evidently of the highest importance to procure, if possible, immediate union, and prevent the occurrence or check the spread of inflammation. Every wound involves a tendency to inflammatory action; but the smaller the amount which is excited, and the more complete the absence of inflammatory exudation, the better the form of healing which occurs. The size of the wound, and the degree

of violence with which it was inflicted, are elements in the production of inflammation over which we have no control; a third element is the duration or amount of exposure to the atmosphere; this is within our reach, and the importance which has long been attached to the prevention of access of air to a wounded joint is fully justified on theoretical as well as on practical grounds.

When there has been much laceration and contusion of the soft parts, little or no chance of immediate union exists. For the description of the management of this class of wounds, however, the reader may conveniently be referred to the essays which treat of Gun-shot Wounds and of Compound Dislocations; for contused wounds of joints usually fall within one or other of these two categories. In punctured or incised wounds, if seen at a sufficiently early period, the treatment is directed to procure rapid union, and prevent the occurrence of inflammation, or reduce it to a minimum. For this purpose, we retain the wounded surfaces in close apposition, and adopt such measures as will insure perfect immobility of the joint, whilst we exclude all exposure to the air. If any foreign or detached body is present—a piece of glass, for instance—it is carefully removed, and the wound, having been gently cleaned and thoroughly washed with a solution of carbolic acid, must then be closed effectually, and its surfaces, for their entire depth, kept in close and steady apposition. The mode of effecting this must vary; where only a small puncture exists, it may be sufficient to apply a piece of lint, which may be soaked in carbolic oil or blood, or covered with collodion; if the wound is of some extent, something more is required; on the whole, sutures are preferable to adhesive plasters, being less liable to be disturbed, and admitting of the application of moisture to the part; the twisted suture appears to be the most efficient, as it maintains the divided surfaces in contact to a considerable depth. The wound may then be protected from the air by a covering of stout tin-foil, previously washed with the solution of carbolic acid, and nicely adapted to the surface, this being again covered by a rather larger piece of lac plaster charged with carbolic acid, and a fold of lint dipped in carbolised oil (as described by Mr. Lister) applied over the plaster and renewed, by being moistened with the oil, once at least in twenty-four hours; the under dressings of course not being interfered with unless it becomes absolutely necessary to do so. The limb

must be carefully secured on a splint, or otherwise fixed in that position which will be most advantageous to the patient should rigidity ultimately occur. As perfect immobility is of the greatest importance, the splint must be carefully selected, so as effectually to prevent any articular movement. The immediate and continued application of cold to the part, in the form of irrigation or of iced water in vulcanised india-rubber bags, is usually attended with great advantage in preventing or subduing inflammation. The general treatment, at an early period, should be moderately antiphlogistic, without permanently lowering the strength, great demands on which may be expected should events take an unfavourable direction. Care should be taken not to disturb the antiseptic dressings so as to allow the wound to be unprotected by them, and the sutures may be left with advantage for many days, perhaps a week, or so long as there is any hope of union being obtained, unless their removal is required to give exit to matter.

In spite of the measures adopted, or in consequence of the case coming too late under treatment, inflammation may run high and suppuration occur. When matter has unquestionably formed, any attempt at union is abandoned, and free exit afforded to the pus, either by enlarging the original wound, or, if necessary, by making free incisions in a more depending position; but an attempt may still be made to arrest decomposition of the matter by the continued use of antiseptic dressings. The further progress of the case does not differ materially from that already described under the head of Suppuration in Joints (see p. 6), to which the reader is consequently referred.

PART II.

DISEASES OF INDIVIDUAL JOINTS.

THE limited space at my disposal requires that any observations I have to offer respecting the peculiarities presented by disease as it affects individual joints should be extremely brief.

Diseases of the Hip.

Lesions of the hip have attracted especial attention from the importance of the articulation involved, the depth at which it is situated, the mass of soft parts which surround it, and the little leverage afforded by its upper attachment in the pelvis.

At the same time, unfortunately, disease of this joint is met with only too frequently, and constitutes one of the most troublesome lesions to which the human frame is liable.

As in other joints, the structures in which disease for the most part commences are the synovial membrane and ligaments and the articular extremities of the bones. Mr. Aston Key was of opinion, from the amount of mischief which is sometimes found in the ligamentum teres at an early period of the disease, that this was the part in which inflammation usually commences. At any rate, there is reason to believe that the ligaments (including the capsule) are often among the earliest structures to be affected.

Acute arthritis of the hip, whether occurring primarily, or in the course of chronic disease, is very serious and often fatal. The symptoms are usually unmistakeable, and if matter forms, as is often the case, it should be evacuated early, and antiseptics freely used. The expediency of amputation or excision must be judged of by the circumstances attendant on each individual case.

Common synovitis, as a primary affection, is less frequent in the hip than in the superficial joints; though when it occurs, its symptoms are usually severe, and the suffering experienced considerable.

Morbus coxarius; strumous synovitis and ostitis.—The disease of the hip which is most common, and which is so often met with in early life, is chronic or strumous inflammation, sometimes commencing in the synovial membrane or ligaments, but frequently, also, in the articular extremities of the bones. In retaining the term *strumous*, however, we must bear in mind that, as already stated, it by no means follows necessarily that there is a special tendency to tubercular deposit in the joint or in any other part of the body, or that such general cachexia exists as will prevent reparation of the local mischief. If attention to the general health is *desirable*, attention to the local treatment (especially in the form of securing the joint in a most perfect state of rest), is *indispensable*, and any neglect of it on the part of the surgeon hardly admits of excuse.

It is of great importance to recognise chronic hip-disease as early as possible, before permanent mischief has occurred. The first symptoms to be noticed are, a slight limp in walking, with disinclination to bear the whole weight of the body on one particular limb; if attention is now carefully drawn to the

patient, the thigh will probably be found to be slightly flexed on the abdomen, and a certain amount of feverishness may be noticed towards evening, after the day's exercise, with a little restlessness during the night, accompanied, perhaps, with slight occasional twitches or jumps of the thigh, which will usually be somewhat flexed and adducted during sleep. If a careful examination of the hip is now made, it will be found to be held a little stiff, and some pain will be evinced on moving the joint or on pressing the articular surfaces together, either directly at the great trochanter or front of the joint, or indirectly through the foot.

If these symptoms are not attended to, and the disease proceeds a stage further, the lameness becomes very marked, the limb will be found apparently altered in length, flattening of the nates will be detected on examination, and very decided pain will be complained of. The causes of the apparent alteration in length are mechanical. The thigh is flexed on the pelvis, but combined with this flexion, which is very seldom *simple*, are two principal deviations; in the one case we find inclination and rotation *outwards*, attended with apparent elongation of the limb; in the other, there are inclination and rotation *inwards*, producing apparent shortening. Now, supposing the femur to be bent at an angle on the pelvis, and the foot turned out, when the patient tries to stand or walk the weight of the limb has to be supported, but in order that the foot may reach the ground, that side of the pelvis is depressed and advanced in front of the other; in this way the limb *appears* to be elongated. In the opposite condition of flexion combined with adduction and rotation inwards, the corresponding side of the pelvis, for a similar reason, is raised and carried backwards; a position which involves apparent shortening when an attempt is made to place the limbs parallel with one another. The position which is recommended as best suited for examination is that in which the trunk and the *sound* limb are perfectly straight; the nature of the alteration, as it affects the diseased side, is then more evident than if the two thighs are placed parallel.

The fixed and altered direction of the affected extremity soon entails secondary deformity in other parts, especially in the spine. When the thigh is fixed at an angle with the pelvis, in order that the weight of the extremity may be supported, the pelvis itself becomes inclined forwards; to compensate for this inclination and maintain the balance of the body, the vertebral

column is strongly curved, producing a deep hollow in the lumbar region, with corresponding projection of the abdomen anteriorly. The exact nature of this secondary deformity is readily seen, either in the recumbent or in the erect posture, on supporting the affected limb at the angle which it has assumed, when the pelvis and spine may easily be restored to their natural direction. Lateral deviation of the spine may also be induced, to compensate for the lateral inclination or twisting of the pelvis. The pelvis, it will be noticed, shifts or alters its position; and, indeed, increased mobility of the pelvic articulations often occurs to an extraordinary extent, especially in the later stages of hip-disease, when recovery is taking place, or ankylosis or rigidity of the hip itself. In such cases, in a life, the amount of motion which becomes developed at the sacro-iliac synchondrosis and other neighbouring joints closely imitate the natural mobility of the hip, and easily deceive a careless observer.

The condition of apparent shortening frequently follows that of apparent lengthening, though it may occur where no lengthening has ever manifested itself. Is there ever any *real* alteration in length? Though many causes have been advanced which might give rise to actual elongation of the limb, it is hardly probable that any of these should actually occur, and it may be said that the lengthening is, in truth, only apparent, depending on the altered position of the pelvis. In the early stages of hip-disease the same observation holds good with respect to the opposite condition. In other cases, however, a certain amount of genuine shortening really does take place. When the osseous structures are involved in the disease, the spasmodic contraction of the muscles maintains constant pressure of the head of the femur on the upper part of the acetabulum; if this is already softened, and the cartilages ulcerated or absorbed, the border or edge of the acetabulum yields to the pressure, it becomes deeply ulcerated and excavated, and the head of the femur itself, which has been more or less disintegrated, is drawn up into this excavation, which is enlarged to receive it, in the direction of the external iliac fossa. This condition constitutes what has been termed *spontaneous dislocation* from disease, which differs evidently from traumatic dislocation, in which the head of the femur clears the rim of the acetabulum which remains intact. When this false or incomplete luxation has occurred, the trochanter major will be found to be displaced

wards, and a certain amount of real diminution in the length of the limb will be present, increased, of course, if the head of the femur is destroyed to any extent. In some cases, however, luxation from disease actually occurs, as may be seen in the accompanying illustration, taken from a preparation in the Museum of St. George's Hospital, in which the head of the femur is situated very near the anterior superior spine of the

FIG. 215.



of the acetabulum. (From preparation in the Museum of St. George's Hospital.) *a.* Head of femur, extensively destroyed by abscess in the joint. *b.* Acetabulum containing the remains of the ligamentum teres. *c.* Necrosed portion of the acetabulum. *d.* Perforation of the acetabulum corresponding to the necrosed portion.

FIG. 216.



Dislocation of hip from disease. (From a preparation in the Museum of St. George's Hospital.)

1. In very exceptional cases, too, the head of the femur is dislocated through the acetabulum, so as to enter the pelvic cavity. Another, and a very striking change, which will be found at an comparatively early period, consists in a certain amount of atrophy in the fulness of the nates. These are often flattened and appear wider than natural. This alteration depends partly on wasting of the gluteal muscles from want of use, partly also, on the position of abduction, on the twist of the pelvis and consequent diminution in the projection of the tuber ischii.

The pain which is experienced is not confined to the hip, but is often referred to the inner side of the knee. It is due probably to pressure on, or irritation of the branch of the sciatic nerve distributed to the capsular ligament and ligamentum teres, referred sympathetically to the terminal cutaneous

branches of the same nerve. In the same way, if the articular branches of the anterior crural or sciatic nerves are more particularly involved, we may find the pain referred to the front or outer side of the knee, or even to the heel or ankle. From whatever cause it arises, this pain is sometimes the first symptom which attracts attention, and may lead to mistakes regarding the real seat of the disease. The chance of error is increased if puffiness over the knee accompanies the pain, as it does sometimes just as swelling of the testicle may be produced by the passage of a calculus along the ureter, from pressure on the spermatic plexus, and consequent nervous disturbance of the nutritive conditions of the parts supplied by its peripheral branches.

In the third stage of hip-disease, all the symptoms are much aggravated. The starting pains are very violent, rendering the nights much disturbed. The limb is still more distorted, as real shortening is often present from the incomplete luxation already described. There is much fever with increased local heat, suppuration has now taken place, and crepitus may be detected when the articular ends of the bones are moved against each other. When abscesses form, they may point in almost any part of the limb, either in the vicinity of the joint, or after having burrowed among the muscles to a great extent, especially when the erect posture has been long maintained. If the sinus is situated some distance down the thigh, near the inferior attachment of the tensor vaginæ femoris, the probabilities are in favour of the femur being the seat of mischief. When the acetabulum has originated or participated in the disease, and matter has formed in the pelvis, the abscess often shows itself at the outer part of the groin in the neighbourhood of the antero-inferior spine of the ilium; occasionally it passes down by the rectum, into which it may burst, or it may reach the surface close to the anus.

There are one or two points in the diagnosis to which I may briefly direct attention. I have seen congenital dislocation mistaken for hip-joint disease, and the child subjected to much unnecessary confinement and suffering. A little attention readily distinguishes the two affections. The history of the case, as the existence of evident limping from the first moment when any weight was allowed to rest on the limb, the absence of pain and the extent of real shortening, aggravated when the limb is pressed upwards, diminished when extension is made, are symptoms of congenital dislocation which differ widely from the

which would have been presented by severe disease of the joint. We occasionally meet with cases in which a difference in the length of the two lower limbs exists, resulting in some instances from congenital malformation, in others from infantile paralysis, which has led to wasting and defective growth of one extremity. Such a condition, if existing to any extent, causes limping in the act of walking, and leads to secondary deformity of the spine, in the form of lateral curvature, to counteract the tendency of the body to fall towards the affected side. This state might easily be mistaken for diseased hip, as well as for diseased spine, until a careful examination of the whole limb is instituted, when the cause of the symptoms is readily detected.

Psoas abscess, with or without disease of the spine, may be mistaken for hip-disease. A case of this kind, where the psoas abscess was connected with extensive tubercular disease in and around the kidney, came under my care at the Hospital for Sick Children, which had been considered and treated previously as disease of the hip. The limb, in such a case, may be flexed on the pelvis, and any attempt at extension give rise to great suffering; it will be found, however, that, provided the psoas muscle is kept relaxed and no traction upon it allowed, free movement of the joint may be effected, and its articular surfaces rotated on one another or pressed together with much force, but without suffering. The situation of the abscess, also, serves to assist in the diagnosis, for pelvic abscess, depending on hip-joint disease, is almost invariably complicated with sinuses in other situations. Inflammation of the bursa under the psoas-iliac may simulate hip-disease, and be with difficulty distinguished from it. Sometimes, indeed, the bursa communicates with the joint, and disease beginning in the one involves the other in its course. If such is not the case, and the inflammation is confined to the bursa, there will be pain and tenderness on pressure at the front and inner part of the joint, with, it may be, some indistinct swelling in that situation; the mass of muscles, however, obscuring the bursal tumour. The limb is usually instinctively flexed on the pelvis, as in hip-disease, and any attempt to extend the limb, or any movement which involves traction on the psoas muscle, is attended with suffering. The diagnostic marks correspond to those just mentioned. If care is taken to avoid any pressure on the bursa, or action of the psoas, the joint may be freely handled without suffering; moreover, no tenderness or swelling can be detected behind the trochanter.

The treatment of strumous disease of the hip does not differ materially from that already recommended for strumous inflammation of the joints generally. If detected at the earliest stage, complete rest of the joint, at first in bed, afterwards by means of a splint, which must be long enough to secure perfect immobility of the hip, will be sufficient. But the rest must be continuous and maintained for an adequate period, the general health being at the same time properly attended to. In the second stage, when the limb is apparently altered in length and some starting pains have manifested themselves, confinement to bed for a long period is necessary, and extension by means of weights, to relieve the muscular spasm, may be used in the way already described. If the thigh is flexed and adducted, it must be straightened, which can be done readily enough under chloroform when the disease is recent, and with care and caution even at a later period when adhesions have taken place.

When suppuration has occurred, if the symptoms are not very urgent it is usually better to abstain from opening the abscess, as absorption may be expected occasionally to take place. If there is great pain, and the matter is too deep-seated to be reached conveniently, relief is sometimes obtained, temporarily at all events, by a few lines of the actual cautery. If the pain and distress are still very great, an opening may be made either with a knife or a trocar. In some cases where it is necessary to let out deep-seated matter, in order to avoid a large vessel, the mode recommended by Mr. Hilton may be employed, viz, after making a small incision through the superficial structures, to force a probe-director through the soft parts into the cavity of the abscess, and when pus flows along the groove of the director, to introduce a pair of fine dressing forceps into the abscess, and enlarge the opening by forcibly separating the blades.

If the case has gone on unfavourably, the bone become carious, and extensive suppuration taken place, resulting in sinuses, the health suffers, and the only chance of recovery, or even of preserving life, may consist in excision of the head of the femur, or removal of the diseased bone. Such an operation, in children at any rate, is more formidable in imagination than in reality and I have had occasion to perform it without any injurious consequences. The subject of excision, however, will be treated of in a separate essay.

Morbus coxæ senilis.—In advanced life another form of disease

in the hip, which is known by the name of the *coxitis*, or *chronic osteo-arthritis*. Unlike strumous disease, this disease seldom occurs under the age of forty, and appears as a local affection, without any visible impairment of the general health. In chronic arthritis of the hip, stiffness of the articulation, and a certain amount of tenderness extending down the *front* of the thigh; the pain, though augmented after much exercise, diminishes at night, and is not increased even by rough pressure of the articular surfaces against one another. The mobility of the joint is much interfered with, and a loud crackling can be heard when the limb is exercised. When the disease is established, the foot is much everted, the nates are flattened, and there is *apparent* shortening of the extremity to a considerable extent, in consequence of elevation of the right side of the pelvis; a certain amount of *real* shortening may also take place after a time, from the changes which take place in the femur, the head of which may become altered and flattened, whilst the neck may assume a horizontal position, so that the upper extremity of the bone may be below the level of the lesser trochanter.

At this stage of the disease, too, bony protuberances may perhaps be felt around the head of the bone. The characters of the disease are so distinct from those of the strumous disease of the hip, that the stakes between the two are well marked. From sciatica the disease is distinguished by the character of the pain, as well as by the presence of shortening of the limb, real or apparent, and by the position of the foot. The appearance of the bone somewhat resembles that of the neck of the femur, which it might possibly resemble if any injury or acci-

dent previously occurred. The symptoms, however, and the slow and gradual progress of the disease, as well as the osseous growths which are formed, are sufficient for *accurate diagnosis*. Where

FIG. 217.



Head and upper part of the shaft of the femur affected with chronic osteo-arthritis. (From a preparation in the Museum of St. George's Hospital.)

constitutional symptoms are present, the disease extends to other and smaller articulations, and the character of the affection cannot fail to be recognised. The treatment of this affection is noticed in the section on Chronic Rheumatic Arthritis.

Neuralgia of the hip.—This joint is frequently the seat of hysterical pain, closely simulating structural disease. The diagnostic characters of this affection have already been pointed out (see Articular Neuralgia). It may be remarked, however, that apparent shortening of the limb may be present in these cases, owing to alteration in the direction of the pelvis and its relation to the vertebral column, induced by long continuance in one unnatural posture. Lateral curvature of the spine also is frequently noticeable; but the nates are not flattened, and abscesses do not form.

Diseases of the Pubic, Sacro-iliac, and Sacro-coccygeal Joints.

These articulations, in consequence of their anatomical arrangement, are less exposed to disease than the proper diarthrodial joints of corresponding size.

As a consequence of hip-disease, especially in early life, the pelvic articulations become unusually movable; to such an extent indeed, that rigidity, or even complete ankylosis, of the diseased hip-joint may be attended with but little loss of the mobility of the limb.

Disease of the symphysis pubis is very uncommon.

A case, however, of extensive ulceration of its fibro-cartilage, with superficial caries of the articular surfaces of the contiguous bones, accompanied with a large abscess in the sheath of the rectus, occurring in a female, will be found in the fifth volume of the *Pathological Transactions*. Dr. Humphry also quotes a case related by Otto, in which the symphysis, as well as the right sacro-iliac joint were completely ossified in a middle-aged woman. Two other cases of ossification are also referred to, but ankylosis is undoubtedly very rare.

Acute inflammation of the sacro-iliac joint may result directly from injury. A case is related by Louis in which a sack of corn fell on the loins of a man who was stooping at the time. The accident gave rise to some swelling of the part, followed after a time by violent pain and much febrile disturbance, and death ensued at the end of the twentieth day. Very extensive inflammation, attended with suppuration, was found in the right

sacro-iliac articulation, and the bones were separated to a certain extent from one another.

Disease of the sacro-iliac joint may arise in connection with pregnancy or parturition. During pregnancy the union between the pelvic bones is said to become less firm, and in certain cases the relaxed condition of the sacro-iliac joint is so considerable as to give rise to pain, as well as to difficulty in standing or walking; during exercise, too, the patient may be sensible of a certain amount of motion between the sacral and iliac bones, and sometimes a distinct 'crackling' can even be heard. In most of these cases the ligaments resume by degrees their normal condition, and the pain and limping gradually subside. Occasionally, however, the relaxed condition may last for a considerable period, and the limping may continue for months or years, or even, according to Ludovic, for the remainder of life. It may happen, too, that actual inflammation of the joint is set up; the pains become violent; there is swelling about the part; and if suppuration takes place, the case is most likely to terminate fatally.*

Sir B. Brodie relates the case of a married lady who was unable to walk without crutches, and who experienced pain after taking exercise, referred to the right groin as well as to the right sacro-iliac articulation, where a projection existed, as if the ilium had been displaced and drawn upwards; the affected limb was two inches *shorter* than the sound one. Pains resembling sciatica had been complained of for many years, but the first distinct symptoms of the affection appeared to be referred to her pregnancy, four years previously. Recovery took place, though the limb remained permanently shortened. Mr. Hilton also relates a case of sacro-iliac disease, occurring after parturition, attended with suppuration within the pelvis, in which the abscess subsequently became absorbed and a cure was ultimately effected after prolonged mechanical rest.

The chronic form of *sacro-iliac disease* is often traceable to some local injury, though Mr. Erichsen (to whom we are indebted for a very excellent description of this affection), considers it to be usually strumous in its origin. The symptoms consist mainly in pain in sitting and standing, or even lying on the affected side, as in any of these positions the weight of the body tells upon the inflamed part. The pain may be localised by direct pressure on the affected joint, or by pressing the bones together or the sacrum forwards, and in many cases a little stiffness with some elevation of temperature may be detected

* See Desormeaux, *Sur les Relâchements, Écartements, etc. des Symphyses du Sacrum*.

in the sacro-iliac region. Frequently, however, as Mr. Hilton points out, the actual seat of the disease may be masked by the pain being referred (through the nerves in relation with the joint), to the hip, knee, or back part of the calf, whilst the psoas muscle may be kept spasmodically contracted from nervous irritation, giving rise to a flexed condition of the hip. There is an absence, however, of any increased warmth in these parts, and by taking care to keep the pelvis quite at rest, the other joints may be moved freely without producing pain. When suppuration takes place, and in most cases it only does so at a late period of the disease, the matter may point posteriorly in the immediate neighbourhood of the articulation, or it may extend outwards to the great trochanter, or upwards to the loin; if, on the other hand, the matter accumulates in front of the joint, it may pass into the perinæum or even open into the rectum, or it may leave the pelvis by the great sciatic notch and point in the gluteal region.

The affections with which sacro-iliac disease is most likely to be confounded are diseases of the hip and spine, but by careful attention to the points indicated, the real seat of the mischief can generally be made out satisfactorily, except, perhaps, when the disease is situated at the junction of the last lumbar vertebra and sacrum, and affects mainly one side.

When sacro-iliac disease is in an advanced stage, and especially when matter has formed to any extent, the prognosis is by no means favourable. In an earlier stage, however (and hence the importance of accurate diagnosis), a cure may not unreasonably be expected, as Mr. Hilton relates several cases, occurring at varying ages, to show that by giving perfect and prolonged rest in the recumbent position, and by steadying the pelvic bones by means of a strong and broad belt so as to favour ankylosis, recovery will often take place.

Disease of the sacro-coccygeal joint is occasionally met with as the result of injury, and, for anatomical reasons, more frequently in females than in males, in early life than at an advanced age. The symptoms, as described by Mr. Hilton,* are tenderness with increased heat about the joint; pain on defæcation as well as on rising from a seat or sitting down, and on walking or running rapidly, the pain in these movements being caused by the action of the corresponding muscles attached to the coccyx

* Hilton, *On Rest*, p. 424.

There is also pain on direct pressure on the joint, and on moving the coccyx when grasped between the thumb and the fore-finger introduced into the rectum. Rest in the recumbent position, and the prevention of constipation by appropriate diet and the use of laxatives, constitute the treatment recommended.

It may be remarked that symptoms very nearly resembling those mentioned are often met with in hysterical females. In these cases, however, the temperature about the part is not increased, and there is an absence usually of pain when the coccygeal bones are moved in the way previously described.

Diseases of the Knee.

The diseases of the knee have been studied with extreme care, owing to the frequency of their occurrence, as well as to the size and accessible position of the articulation. Hence it is that the affections of this joint are often taken as typical of articular disease generally. The knee is liable to the various morbid changes which have been described in the first part of this essay; to one of them, indeed—chronic inflammation of the articular extremities of the bones—it is specially exposed, for *local abscess* is more common in the head of the tibia than in any other situation; and necrosis of the lower end of the femur is often met with, at first simulating, but subsequently inducing, suppuration of the joint.

The position which the limb habitually assumes in severe disease of this articulation, whether simply inflammatory or strumous in its origin, is one of considerable flexion, frequently combined with torsion, in consequence of eversion or inversion of the foot. If the ligaments become extended or destroyed in the progress of disease, the spasmodic contraction of the flexor muscles acts upon the head of the tibia, which it tends to dislocate backwards into the popliteal space. In our treatment, therefore, we have to guard against this tendency; the foot must be supported to prevent torsion, the limb must be maintained uninterruptedly in a nearly *straight* position, and extension by weights may be used as in the case of hip-disease.

In *chronic osteo-arthritis*, the knee-joint soon acquires a strong inclination *inwards*, whilst the tibia is usually rotated outwards and the foot everted. At a more advanced period, if the limb becomes rigid in a semi-flexed position, the patella is occasionally found resting on the outer condyle of the femur, or even completely dislocated outwards. The amount of synovial

it, rather than in the articulation itself. In this way the whole thigh may be inflamed and swollen, and the appearances somewhat closely resemble those of acute periostitis of the femur. The depth at which the matter is situated prevents it from coming rapidly to the surface, and does not readily allow of fluctuation being detected. Much constitutional disturbance is, consequently, usually present, and the amount of pus which forms before an outlet is obtained is often very large. In such cases early and deep incisions are obviously called for; incisions which shall extend through the whole muscular layers, if the seat of the suppuration is to be reached, and the symptoms produced by the pent-up matter abated.

Diseases of the Ankle.


The ankle-joint is liable to the various diseases which have been already described, and is often involved in the progress of caries affecting the irregular bones of the tarsus; occasionally, indeed, it is not easy to decide whether the disease actually affects the ankle, or is confined to the astragalus or calcaneum; the point may, however, be determined by the free motion in the latter case of the ankle-joint when the patient is placed under chloroform. When the joint is distended with fluid, the tumefaction and fluctuation are most evident between the two malleoli, on either side of the extensor tendons; in extreme cases, the effusion is perceptible also posteriorly, at the inner and outer borders of the tendo Achillis. When suppuration occurs, the matter may make its way to the surface in the immediate vicinity of the joint, or it may ascend the leg, or descend into the foot in connection with the tendinous sheaths around the ankle, so as to open at a considerable distance. The direction which the foot is prone to assume in disease interfering with the use of the joint is that of flexion, combined generally with varying amounts of inversion or eversion. The position to be selected, on the other hand, to avoid present stretching of the ligaments, and as most useful in the event of ankylosis, is that in which the foot is in the same plane with the leg, and forms with it an angle only slightly exceeding a right one. It may be remarked, that disease of this joint is more favourably situated for recovery than disease of the thigh or knee, because, by the employment of a wooden leg, whilst the ankle is carefully supported by splints, exercise may be taken, and the general health consequently maintained.

Chronic osteo-arthritis seldom affects the ankle, but when it does, the tarsal joints are usually implicated at the same time. Increase in the breadth between the malleoli, which are preternaturally prominent, combined with projection inwards of the scaphoid and depression of the instep, are the appearances commonly noticed.

Diseases of the other Articulations of the Foot.

The tarsal bones are so small, and the articulations between them so numerous and in such close proximity, that distinctions between the morbid conditions of the two can with difficulty be drawn. Disease in the tarsus, in the majority of cases, arises in strumous inflammation of the bones, which

FIG. 218.



Disease of the tarsus, commencing in the joint, between the os calcis and astragalus.
(From a preparation in the Museum of St. George's Hospital.)

soon extends to the cartilages and to the synovial membranes. Occasionally, however, an articulation is the starting-point, as may be seen represented in the accompanying illustration (taken from a preparation in the St. George's Hospital Museum), in which the disease is found commencing in the joint between the astragalus and calcaneum. Some importance must be attached to the original seat of the mischief, in consequence of the varying degrees of simplicity of the synovial membranes; of these, the median, or *scapho-cuneiform*, is by far the most complicated, as it extends between so many tarsal bones. When disease involves only the simpler synovial membranes, or is confined to single bones in con-

with them, the probabilities of a successful issue are actively great; and if operative measures are necessary, of individual bones may be had recourse to with happy results. I have several times removed the calcaneus, the cuboid, or even the internal cuneiform, and re- has occurred with a very useful foot. When disease of the astragalus or the scaphoid, on the other hand, and the ligamentous synovial membranes are implicated, the chances of curing the mischief become much diminished; though one of these bones can often be removed successfully, and a very effective foot.* The middle and external cuneiforms, also, are unfavourably situated, though less so, perhaps than the astragalus or scaphoid. It may be remarked, however, that the repair which takes place after excision in these cases is usually most complete, for a large portion of the periosteum is left behind in the operation; reproduction of bone consequently occurs to a greater or less extent, and restores to the foot all the solidity which is required. After removal of the whole of the calcaneum, I have found the heel still rigid, and the amount of deformity very trifling.†

Inflammation of the complicated *scapho-cuneiform* synovial membrane may be looked upon, according to Mr. Erichsen, as a special affection of the foot, commencing usually in disease of the scaphoid. At first there is pain and swelling in the region of the joint between the scaphoid and cuneiform bones. The swelling is increased by bending down the foot, and extends over the tarsus to its outer side, as the articulation between the internal cuneiform and the cuboid becomes involved. At a later period 'the foot assumes a remarkable bulbous or clubbed appearance; the symmetry of the heel and the outline of the foot are unimpaired, but the fore part and dorsum of the foot are greatly swollen, glazed, and possibly perforated by sinuses discharging their unhealthy pus.'‡

Diseases of the metatarsal and phalangeal articulations call for special observations. They are frequently *gouty* in their character, or, in some cases, are examples of *chronic osteoarthritis*. It is worthy of notice, perhaps, that in the metatarso-

* Holmes, *Surgical Treatment of Children*, p. 502 et seq.

† See also *Pathological Transactions*, vol. xi. p. 217, for an account of the removal of the parts in a foot from which I had removed two of the tarsal

‡ Erichsen's *Science and Art of Surgery*, 3rd ed. p. 715.

phalangeal articulation of the great toe, the deformity induced by the latter affection has been mistaken for dislocation; a condition which it may closely resemble.

The subject of *bunion* is considered in the essay on SURGICAL AFFECTIONS OF THE SKIN.

Diseases of the Sterno-clavicular, Acromio-clavicular and Sternal Joints.

A certain amount of enlargement of the *sterno-clavicular* as well as of the *acromio-clavicular* articulations may sometimes be seen in persons who have long been in the habit of working hard with their upper extremities. Acute inflammation of the *sterno-clavicular* joint is also met with occasionally. Such a case in an adult came under my care, in which suppuration rapidly took place, with destruction of the fibro-cartilage, inducing spontaneous dislocation of the inner extremity of the clavicle. The shoulder and arm having been fixed, and the dislocation reduced and prevented from recurring by appropriate splints, recovery took place with a useful extremity.

Disease of this joint is easily recognised, but, owing to the great inobility and peculiar anatomical characters of the articulation, it is difficult to secure perfect rest and, for like reasons, ankylosis is seldom met with. Cases of *sterno-clavicular* disease will be found related in Mr. Hilton's work on *Rest*, and in the *Lancet* of April 1864.

Chronic osteo-arthritis of these articulations may also occur, producing enlargement of the articular extremities of the clavicle, stiffness and pain on movement of the joints, as well as weakness of the upper extremity.

Disease of the articulation between the first and second pieces of the sternum, where no synovial membrane exists, is rare, but an interesting case is related by Mr. Hilton, in which the principal symptoms were pain about the sternum, especially on coughing or talking or making any respiratory exertion, with difficulty of breathing, and great tenderness on direct pressure on the joint. Recovery ultimately took place.

Diseases of the Shoulder.

The shoulder-joint resembles the hip in its anatomical characters, but is, fortunately, far less subject to disease, and when disease occurs the results are usually much less severe, the per-

tence of pain and the production of rigidity constituting the conditions most to be apprehended. If suppuration takes place, which is seldom the case, the matter often follows the course of the long tendon of the biceps, and opens at the front of the arm below the deltoid. Not unfrequently, too, it passes out of the articulation by the opening which exists in the synovial membrane beneath the subscapularis, and follows that muscle, arriving at the surface near the back and lower part of the shoulder. The most favourable position in which a diseased shoulder can be placed for treatment, is that which is almost naturally adopted; the arm should be parallel to the trunk, and the elbow slightly separated from the chest, the fore-arm being at the same time supported by appropriate mechanical means. If ankylosis takes place, owing to the great mobility of the scapula the arm may still be moved very freely, and the limb consequently remains very useful.

The condition of parts induced by *chronic osteo-arthritis* of the shoulder has, no doubt, been frequently mistaken for the result of injury. After the stage of synovial distension has passed, wasting of the soft parts occurs; rigidity, as well as crepitation on movement, are met with, and the head of the humerus appears to be elevated and carried forwards. When the disease is of long standing, the deltoid is atrophied. The capsular ligament is usually thickened, and has sometimes osseous particles in its substance, or it may be perforated at its upper part. The synovial membrane shows marked signs of inflammation. The glenoid fossa loses its ligament as well as its cartilage, and presents the characteristic ivory-like material; occasionally, too, irregular osseous growths are found around its edges, increasing its capacity. The under surface of the acromion, and even of the acromial end of the clavicle, may be hollowed out by pressure of the humerus upon them, and are often covered with a polished porcelain-like material. Separation of the acromion into two portions is also met with sometimes. The head of the humerus is often much enlarged, and its surface eburnated. The tendinous structures around the joint may appear partially disintegrated, and the intra-articular part of the long head of the biceps is usually destroyed, the remains of the tendon adhering to the edges of the bicipital groove.

Disease of the *deep-seated bursæ* around the shoulder-joint is occasionally met with, and may prove somewhat perplexing. The bursæ beneath the *subscapularis* and the *infra-spinatus*

habitually communicate with the synovial membrane, so that their diseases are in fact those of the articulation itself. The bursa beneath the *deltoid* is distinct; and when that bursa is inflamed, the muscle is rendered unusually convex, and its contraction of it is attended with pain. The articular surface, however, can be freely pressed together, and the joint exercised without suffering, provided the *deltoid* is not used, and no pressure is made on the tumour. A sensation of fine crepitation can often be felt in these cases on moving the shoulder.

Diseases of the Elbow-joint.

The diseases of the elbow resemble those of the knee, although they are less severe in their character, and more favourable in their results. This remark, indeed, applies, for obvious reasons, to all the articulations of the *upper* as compared with those of the *lower* extremities. When effusion takes place into the synovial membrane of the elbow, the swelling is most evident at the back of the arm, on each side of and above the olecranon; it is also perceptible at the outer side of the fore-arm, beneath the extensors. If suppuration occurs in the joint, the direction which the matter can make its way with the greatest ease is upwards and backwards between the humerus and triceps; abscesses point, therefore, usually at the borders of this muscle, the outer being the border more commonly selected. The synovial capsule may also give way in front, in which case sinuses will be met with at the side of the brachialis anticus. If the bones are diseased, however, the fistulous orifices may lead directly to the part which has become necrosed or carious. In cases of *strumous osteitis*, which is somewhat common in this region, complete death and separation of a portion of bone in the olecranon, or in one of the condyles, frequently *without* the existence of caries, is by no means unusual. A comparatively slight operation, therefore, is occasionally successful in cases where excision of the articulation might at first sight have appeared necessary. The radius, it may be remarked, is less prone to suffer than either the humerus or the ulna.

The position in which the joint should be placed when splints are required, is that in which the fore-arm forms a right angle with the arm, and is in a state intermediate between pronation and supination. As the joint naturally falls into this position

occurrence in this disease, taking the form occasionally of 'additamentary bones.' Detached or pendulous are often met with, sometimes in great numbers.

Disease of the bursa over the olecranon, constituting 'miner's elbow' can scarcely be confounded with disease of the joint. Inflammation attacks the bursa beneath the triceps, however care is necessary. According to Mr. Barwell, the condition here closely resembles that which is met with in the case of a synovial bag existing between the triceps and the olecranon, which may be a distinct bursa, or, as is more commonly the case, be a prolongation of the articular synovial membrane. Mr. Barwell remarks that, in inflammation of the bursa when it does not communicate with the joint, 'there will be no swelling or tenderness between the inner condyle and olecranon process when the arm is bent at right angles; but the infallible test is, that the line of junction between the head of the radius and the olecranon is as clear and well-defined as ever. If, on the other hand, the inflamed structure communicates with the joint, these parts will participate in the general puffiness and swelling.'*

Diseases of the Wrist, and other Articulations of the Hand.

The complicated relations of the various synovial membranes of the wrist and around the carpal bones facilitate extension of disease from one to the others: the anatomical arrangement of



hand usually lies on a cushion in a state of complete pronation; a condition which places the posterior ligaments on the stretch, and favours luxation backwards, the only form of spontaneous dislocation to be apprehended. In acute inflammation of the wrist, therefore, relief may often be afforded by so adapting the mechanical support as to obviate this tendency to pronation. If luxation of the ulna has already occurred, and the triangular fibro-cartilage been destroyed, the hand must be secured in the position recommended, and pressure exercised on the back of the ulna to reduce its dislocation, and maintain the bone in its normal position. As stiffness of the fingers is very apt to occur from want of exercise in cases of diseased wrist, passive motion of their articulations must be had recourse to as early as possible.

Chronic osteo-arthritis of the wrist is most common in females beyond the middle period of life, though it is met with occasionally at an early age in both sexes. The back of the wrist in this form of disease exhibits usually a peculiar projection in consequence of enlargement and partial luxation backwards of the ulna, and, though to a much less degree, of the radius. The fingers are frequently distorted, becoming permanently flexed as well as adducted; the *second* phalanges, however, of the fingers, as well as of the thumb, are usually *extended*, giving to the hand a peculiar appearance highly characteristic of the disease in its advanced stage. Enlargement of the articular extremities of the fingers is very common, representing the 'nodosities' of Dr. Haygarth. Partial or complete luxations of them may also be present.

Synovial tumours, or 'ganglia,' about the wrist are treated of in the essay on AFFECTIONS OF THE MUSCULAR SYSTEM.

Diseases of the Temporo-maxillary Articulation.

The only affection of this joint to which it is necessary specially to direct attention is *chronic osteo-arthritis*. This disease usually attacks the jaw in old persons, though it has been seen in early life; generally speaking, too, it is constitutional rather than local, and shows a marked disposition to symmetrical disturbance. The condyle and ramus of the lower jaw are usually enlarged and elongated; in a case related by Cruveilhier, however, the condyle was found to be entirely absent. The glenoid cavity is more or less altered, being frequently en-

ations are diseased, the whole jaw is carried forwards. Symptoms are commonly less urgent than those presented by the same disease in other joints, but Dr. Humphry refers to a case in which this disease caused so much discomfort and distortion of the face, that he was led to excise the condyle of the affected joint, and with very good result.* A case, too, is recorded by Mr. Hilton in which osseous ankylosis of this articulation had occurred to a considerable extent on both sides, in a patient in whom ankylosis was also found between the skull and the jaw as well as between some of the succeeding cervical vertebrae.†

ATHOL A. JOHNSTONE.

* Humphry, *On the Human Skeleton*, p. 303.

† Hilton's *Lectures on Rest*, p. 105.

DISEASE OF THE SPINE.

THE spine is subject to a disease, commencing in its bones and joints, so formidable in its results, that it is emphatically called Disease of the Spine. From Caries being the kind of morbid action which invades the bony structures, that term is sometimes used to designate the disease. Owing to an angular projection or hump being formed in the spine at the chief seat of disease, and to distinguish it from Lateral Curvature, the name Angular Deformity is sometimes applied. The term Malady of Pott, derived from the first standard description of the disease having been given by our countryman, is still occasionally met with, but only in foreign works.

Disease of the spine is correctly included among those of the joints generally. The morbid action which leads to the destruction of parts of the bodies of the vertebræ and fibro-cartilages, is identical with what attacks other articulations. But as the spine combines several offices essentially distinct from each other, such as being a flexible column and at the same time the guardian of the spinal cord, its diseases present special characters, which justify their being treated of separately.

Caries: ulceration.—The disease called caries is generally considered to be a morbid process confined to bone. The part of the bone attacked, commonly circumscribed within a small area, at first, is gradually consumed by disintegration and absorption. The fibro-cartilages contiguous to the diseased bone undergo what appears a similar and simultaneous destruction of substance. And during the process, there is a constant discharge of pus, both from the bone and the cartilages. Hence it may be concluded that the diseased action in bone called caries, is the same that, witnessed in the fibro-cartilages, is called ulceration.

The question has been mooted as to whether the morbid

tion exists as to the segment of the spine to which the
laction confines its attacks. It is a remarkable patho-
fact, that spinal disease selects the bodies of the vertebræ,
no-cartilages, and the accessory ligaments, forming the
segment of the spine, as the particular objects of its
a; to the exclusion of the pedicles, arches, transverse,
and spinous processes, together with the ligaments and
ations, which form the posterior segment. That is,
and ulceration commence in the anterior portion of the
destroy it extensively from front to rear, and proceed
pwards and downwards; but they do not go backward, or
be line of partition between it and the posterior portion.
be deemed singular that, numerous as are the small
formed by the opposing surfaces of the oblique processes,
posterior segment, disease is scarcely ever witnessed in
It will presently be seen that the structures of that
it have a remarkable disposition to take on reparative
in antagonism, as it were, to the morbid action in

ase of the spine occurs with equal frequency in the two

It may commence at any period of life; but it is
ely rare past the middle term; it is most rife from early
ood to adolescence.

disease is generally admitted to be one of the class called
one: yet it is often met with in persons who do not other-

surgeon, is a prominence of one, two, or more spinous processes at some part of the column.

Unprofessional persons will speak of that projection as a 'growing out' of the back. To the surgeon it is significant of a certain portion of the front of the spine having been destroyed by disease. He infers that an excavation there has caused the superior part to fall down at an angle to the inferior; and concludes that as the bend is directly forward, the loss of substance must be in the bodies of the vertebræ and intervertebral cartilages. The apex of the angle, it will likewise be perceived, will be formed by the spinous process of the vertebra in which the destruction has been greatest.

In the advanced stage of the disease, no symptom can be more easily detected than the prominence of the spinous processes; for the muscles in each trough will have become atrophied from want of use; and the ridge will stand out conspicuously, like a keel, with rugged, serrated edge, and knob distinct; the transverse and oblique processes may even be well-marked. But in the recent stage, when the angle has just begun to appear, there are sources of obscurity, which sometimes make the diagnosis difficult. The protuberance can be more easily discerned in some of the regions of the spine than in others. It can be recognised in the dorsal before it can in the lumbar; and the reason is obvious. The natural curve in the dorsal region, has its convexity presenting backward; the ridge of spinous processes is, therefore, comparatively superficial; hence a trifling increase of the natural bend caused by the destruction of even a small part of one of the bodies, will be distinctly apparent. But it is different in the lumbar region; the natural curve there has its convexity presenting forward, and owing to the hollow in the back, the tips of the spinous processes are situated comparatively deeply. When there is loss of substance, therefore, in the front or convex aspect of the curve, consequent on carious ulceration of the bodies and intervertebral cartilages, it must be long before the spine will fall forward, and bend in the reverse direction, so that the ridge of spinous processes shall form a convexity and visible projection posteriorly. Moreover from the great depth and magnitude of the lumbar vertebræ, it is obvious that the destruction of their substance must be proportionally extensive, for the column to be bent at an angle. In the cervical region, the normal curve is similar

in direction to that in the lumbar. When the disease, however, attacks the vertebræ there, the effects differ from what are observed elsewhere; the portion of spine above the carious alteration does not fall forward; the weight of the head overcomes that tendency; and the result is that the head subsides vertically, so as to approximate to the upper part of the chest, and obliterate the appearance of the neck. Accordingly, the occiput comes in the way of an examination, preventing the tips of the spinous processes from being felt by the fingers.

But in reference to these normal curves of the spine, it is to be remembered that they are not properly established till about puberty; and that up to that age, the column is very flexible. A difficulty of diagnosis is, therefore, sometimes found in young patients, as a consequence of this suppleness; especially if they have been previously debilitated by illness. When a weakly child sits, he usually stoops; and if the spine be examined, the ridge of spinous processes will probably project conspicuously and alarm the parents; it is not unlikely, moreover, that the chief protuberance will be about the middle of the lumbar region, where a hollow might be expected in the adult. If it should be apprehended that disease of the spine was commencing, the point of greatest interest would be to observe whether there were any distinctly abrupt projection of one particular spinous process, to mark it as the apex of an angle, however obtuse. Should doubt still remain, the child may be laid on its belly, and the hips elevated to a higher level than the spine; if the structures be sound all appearance of the abnormal protuberance will then be removed; if, on the contrary, the projection be consequent on disease, it will continue to be perceived.

Rigidity of the affected portion.—For the detection of disease in the spine at an early period, no symptom is more valuable than diminution of the natural flexibility of the column at the suspected part. And the reason why that defect should indicate disease at its incipient stage, may be stated. Pathology teaches that as soon as caries and ulceration attack the bodies of the vertebræ and intervertebral substances, adhesive inflammation is set up in the adjoining structures, more particularly in the posterior segment of the spine. Coagulable lymph is thrown out in the interstices of the laminae, joints, and processes situated at the back; and as the lymph solidifies the vertebræ become agglutinated and deprived of their natural mobility. Parallel instances are met with in diseases of the

joints elsewhere: for example, an early symptom in hip disease is interruption of the free motion of the articulation. The rigidity of the spine may be made evident by the simple expedient of directing the patient to stoop forward, and rise alternately; or to incline his body first to one and then to the other side; if the column be carefully watched during these movements, and caries has commenced, a distinct contrast will be perceived between the natural flexibility of the spine throughout its principal extent, and an abnormal inflexibility in the portion occupied by about four or five adjoining vertebræ. The latter bones will appear to be moved as a united, compact mass. Or if the palm of the hand be planted broadly and flatly over the vertebræ suspected to be diseased, including some of the sound adjoining parts, and if the patient be asked to stoop and rise as before, a distinct sense of movement among the various processes will be communicated to the fingers at the sound part while there will be an absence of such a feeling at the diseased part.

Heat and swelling.—When the disease is acute and the ulceration in active progress, both these symptoms will be present. The increase of temperature may be ascertained by the touch but it will be certified better by the thermometer. The swelling is seldom such as to be prominent and distinct; it consists rather in a general fulness which partially obscures the forms of the bones. When considerable, it indicates an actively progressing condition of the caries. When absent, and the points and surfaces of the bones are visible and defined, it furnishes an argument for supposing that the disease is dormant and that ankylosis may have taken place.

Pain.—This, which is a faithful guardian in general of the textures of the frame, is a fallacious monitor in regard to disease of the spine. It fails to warn when danger is imminent, and it alarms needlessly. As a proof that carious ulceration of the vertebræ is not always attended with pain, it may be stated that numerous patients have gone through the whole course of the disease, so as to have angular deformity of the worst description, without having been confined to bed a week or day, from inability to go about. Again, other patients almost always females, have suffered sharp pain in a part of the spine, from supposed disease, while every vertebra was perfectly sound. Nevertheless, in acute cases of real disease of the spine, pain in the affected part of the back is a leading

symptom. It will be varied sometimes by the feeling of painful constriction about the epigastrium, as if a cord were tied round the waist. The act of raising the patient, or turning him round, in bed, will cause great agony. If his limbs be affected with convulsive startings the shock communicated to the spine will be attended with much pain. His nights will be sleepless. Such attacks do not commonly last long, but they are apt to recur; they indicate accessions of the acute morbid action.

In a large proportion of cases, pain is a subordinate symptom; and some devices are necessary to make it be felt. While the patient is upright, a shock may be given to the spine, by placing the hands on both his shoulders, and suddenly depressing them; the jerk will probably cause pain in the affected part. Or he may be asked to jump from a low stool upon the floor. Percussing each spinous process in succession is another mode. The plan of carrying a hot sponge down the spine, in the expectation of the warmth bringing out the pain, has not much to recommend it. When dull aching pain, always returning to the same part of the spine, follows exercise, jolting in a carriage, or sitting for a long time, it betokens latent disease.

Neuralgia of the spine.—It is necessary, for assisting diagnosis, to notice here a kind of pain which fixes itself in some particular spot of the spine, and is liable to be mistaken for a symptom of organic lesion. It differs from those pains in the back which can be traced directly to rheumatism, dyspepsia, diseases of the kidneys, or of the uterus, or ovaries. The affection is nervous and functional. It may be considered to have the same relation to the spine which *clavus hystericus* has to the head. Pains resembling it visit different parts of the body; but they have one common character in affecting females prone to hysteria almost exclusively. Sir B. C. Brodie gave to them the name ‘*hysterical neuralgia*.’ They are not uncommon in the joints generally, particularly the knee. When that articulation is affected, the pain may be of long duration, exhausting to the constitution. It has fallen to the lot of the writer to have witnessed two cases of that hysterical neuralgia of the knee-joint, in both of which the patients, young females, consented to lose their limbs on account of the severity of the pain. They were under the care of the same surgeon, who conceived that their sufferings were owing to acute ulceration of the cartilages of the joints, requiring amputation: and chloroform had not then been introduced. In both cases, the cartilages and

all the other structures of the joints were found, after the operations, to be perfectly healthy. The lesson taught by these examples, and enforced by other considerations, is that, when the spine is affected with similar pains, they may be acute and long persistent, without boding actual disease.*

Every part of the spinal column would seem equally prone to be affected with this neuralgia. Perhaps it may be met with more frequently at the site of the 'vertebra prominens,' than elsewhere; and when seated there, the protuberance of the spinous process increases the fear that disease may be its origin. A peculiar feature of the pain is that it is circumscribed within a small area, so that the part might be covered with the point of the finger; and the description often given is that it resembles what might be produced by a nail being driven into the side of the spine. The actual suffering to which the pain gives rise, is undoubtedly great; but the effect on the patient's mind is not the least of the evil consequences. She becomes the victim of unfounded apprehensions as to the nature of her complaint—being commonly possessed with a conviction that a frightful corroding ulcer, of some kind, is destroying her spine. Forthwith she is furnished with an invalid bed; and months or years may elapse before she rises from it.

The main points which deserve attention, in forming a diagnosis, are: 1, that no appreciable angular deformity exists; 2, that there is no perceptible rigidity in the affected part; 3, that

* The subsequent history of one of these patients was not a little remarkable. Although the stump was apparently sound, it soon became the seat of pain, equal to what had been felt in the knee. Thereupon the surgeon repeated the amputation higher up. Healthy cicatrisation took place; but ere long pain, as bad as at first, returned in the new stump. The next proceeding was to excise a large portion of the sacro-sciatic nerve. But this operation had no better result than the preceding. Neither surgeon nor patient, however, lost heart. As there still remained a portion of limb capable of removal, amputation at the hip-joint was performed. The large wound healed favourably. Yet the pain did not depart; it continued in the cicatrix as great as ever, and of the same kind as at first. The young woman was eventually sent to the sea-side. Then she became pregnant. Upon her return it was ascertained that as soon as she felt herself in the family-way, the neuralgia ceased. Having subsequently become the mother of a large family, she has been frequently seen on account of her children; and according to her statement, she has still occasional attack of the old pain in the stump.

The writer has been informed of a similar case which occurred recently. The surgeon was induced, chiefly by the solicitations of the patient, a woman of hysterical temperament, to amputate her limb for supposed ulceration of the cartilages of the knee-joint. The structures were found healthy.

while undergoing the examination, the patient will wince and jerk the body, each time the spine is touched; which is contrary to what she would do if the pain depended on disease: lastly, in reference to those cases in which the pain in the back has lasted over many months, and the lady has been confined all the while to her couch, and yet no deformity is apparent—which is the history of almost all these neuralgic cases—it may be argued that, if caries had been engaged, all the while the pain was felt, in destroying the vertebræ, deformity of an unmistakable kind would have been produced.

It is not uncommon for hysterical patients, especially those who have been long confined to the recumbent position, to lose control over their lower extremities—to have ‘hysterical paraplegia.’ When neuralgia of the spine is added, the complication no doubt increases the difficulty of diagnosis. It will, however, be perceived further on, that when paraplegia occurs as the result of carious ulceration, it is preceded by strikingly marked angular deformity of the column; a condition that does not belong to neuralgia. Yet, in certain rare cases, deformity may not be wanting to increase the perplexity.

The writer had under his charge a governess, twenty-two years of age, affected with paraplegia; she suffered at the same time from neuralgia in the spine. On examining the seat of the pain, which was in the lumbar region, a projection of some of the vertebræ was distinctly visible. The patient had been obliged to relinquish her situation in a family, as it had been stated by the medical attendant that she had spinal disease, accompanied with affection of the spinal cord. From inquiry, however, into the history of the deformity and examination of the whole spine, it was made out satisfactorily that the bulging of the spine at the loins, was a consequence merely of lateral distortion; that the pain situated there was nervous; and that the dragging of the limbs and inability to walk were symptoms of hysterical paraplegia. Under treatment by physical exercises, shampooing, shower baths, tonics, &c., she perfectly recovered in a few weeks.*

Angular Deformity.

When the substance of one or more bodies of the vertebræ has been removed by caries, it may easily be understood that, from the superimposed portion having been undermined, it will fall forward and form an angle with that below. The pressure of the superincumbent weight, however, is not the only

* See article HYSTERIA, vol. i. p. 379. *Lectures on Local Nervous Affections*, by Sir B. C. Brodie. Also a masterly essay *On the Moral Constitution in Females of Hysterical Diathesis*, by Dr. Robert Ferguson, prefixed to Dr. Gooch's Works, published by the New Sydenham Society.

agent in producing the bend; when a patient is confined to the recumbent position, the angle continues to become acute, which shows that the action of the abdominal muscles shares in causing the deformity.

FIG. 219.*




From the bodies and fibro-cartilages in which the substance takes place, being in the median line, the descent of the upper portion is generally in a direction straight forward. That, at least, is invariably the case in the dorsal region. In the dorsi-lumbar and lumbar regions, at which great freedom of motion exists, a lateral deviation will sometimes be combined with the movement forward. The surgeon may be deceived by that inclination to one side; and commit the dangerous error of supposing a case of angular deformity to be one of lateral distortion.

* The figure, copied from a photograph, shows the appearance presented in a case of angular deformity from spinal disease. The delicate girl, fourteen years of age, had become deformed to the extent shown in the drawing, shortly before her admission into the hospital. She was generally able to go about and attend school, during the progress of the deformity. Latterly, she failed in her power of walking. But it was ascertained by frequent observation that this defect did not proceed from paraplegia; from inability to balance herself in the newly-deformed condition of the region of her body. She became strong on her limbs, after a short stay in the hospital.

The writer was consulted about a girl, aged fifteen, who had a projection at the dorsi-lumbar region, with a distinctly-marked deviation of the spine to one side, simulating closely lateral curvature. The medical attendant had considered the case to be of that kind, and put the patient on a course of calisthenic exercises. It was distinctly ascertained, however, that the prominence was the effect of caries; and the principal diagnostic sign was the directness with which the spinous processes stood out backwardly, against the skin; for it is a never-failing observation in regard to lateral distortion, that, owing to the rotation of the column on its long axis, which always accompanies incurvation, the spinous processes are pointed laterally, toward the concavity; and that to such a degree that they are nearly hidden from view by the overlapping of the edge of the longissimus dorsi.

But angular deformity does not consist merely in a falling forward of the undermined portion of the spine, while the inferior remains erect. It is not the same as when a rigid pole, like a flagstaff, is broken by the wind, and the upper part falls over. Were that the case, a patient in whom the spine had become bent would not be able to preserve his equilibrium; his head would be thrown inordinately in front of the line of gravity, and if he attempted to walk, he would be precipitated forward. When a person, therefore, has spinal disease, yet is able to be upon his legs, he is instinctively led to make certain efforts for restoring the balance; the nature of which the accompanying diagram is intended to illustrate. In his first endeavours, he grasps both thighs above the knees with his hands, thus making supports of his arms; he tries at the same time both to elevate the head, and to carry the upper part of the body back, obviously

FIG. 220.*



* A. B. Spine in profile. c. Seat of disease. c. d. Upper portion fallen forward, d^I d^{II} d^{III} d^{IV} upper portion both elevated and carried back to the line of the centre of gravity. E. Apex of angle moved backward.

with the design of bringing them more directly into the centre of gravity. Eventually he succeeds; and it is obviously by causing the portions of the column above and below the angle to arch backward. But these incurvations will have the effect of carrying the angle itself also backward. Hence the prominent 'hump,' which is such a conspicuous feature in the deformity from spinal disease, is produced. It may also be perceived that the apparently proud air with which the patient carries himself—his face turned upward and his body thrown back—is merely an effect of his continued efforts to restore the head to its proper vertical position above the base of the spine. When the disease is in the lumbar vertebræ, and a deep chasm is formed just at the base, the column falls down in its whole length; there are therefore no means of transporting the angle backward in the manner described above; and the patient is in the pitiable condition of being obliged to carry his body horizontally, *ventre à terre*, progressing on his knees, or feet, and hands.

Changes in chest and abdomen.—When the disease is seated in the dorsal region the whole thorax is included in the deformity, and the abdomen partially; the trunk being at the same time diminished in height, as well as rendered broad and square, without distinction of waist. Owing to the falling forward of the portion of spine above the diseased part, the antero-posterior diameter of the chest is increased in length; the sternum also protrudes, being sometimes bent at an obtuse angle; and the projection there forms a kind of counterpoise to the posterior protuberance. The viscera both of the thorax and abdomen accommodate themselves, by alterations in their forms, to the changed condition of the walls of their respective cavities, neither their internal structure nor their functions appearing to suffer. The œsophagus and trachea are simply shortened; the aorta, having to give off its intercostal branches, follows the course of the spine, and is, therefore, about its natural length.

The disease sometimes attacks two different parts of the column, either at once, or at distinct periods; thereby causing two angular projections of the spine in the same patient. In the last case witnessed by the writer, the boy had just recovered from disease in the dorsi-lumbar region, when it appeared in the cervico-dorsal vertebræ; there it extended to the spinal cord, and the patient died in a few months with paraplegia.

Spinal Abscess.

The importance of the subject of abscesses connected with disease of the spine, may be judged of when it is stated that they are the most frequent causes of death, in this complaint. Yet happily in a large number of cases the disease runs its whole course, producing extreme deformity, and ending in ankylosis, without evidence of any abscess having formed.

It has been shown above that the destructive effects of spinal disease are confined to the anterior segment of the column; and that while the morbid processes are committing extensive ravages there, an opposite reparative or defensive action is going on in the posterior segment. Now, the different courses allowed by spinal abscesses cannot be properly understood unless the influence of these distinct agencies be kept in view.

And here a remark concerning the constitution of abscesses generally may not be out of place. It belongs to all abscesses to have an element of self-cure as well as of increase. In the walls of a simple abscess two different processes are observed to be going on. The deeply-seated part of the parietes carries on a formative action, opposed in its nature to an absorptive one in the more superficial part. And it is by the concurrent growth at the bottom or basis, and removal and thinning of the structures toward the point of acumination, that the contents are eventually brought to the surface and discharged. On the pus being evacuated the cavity is filled up by granulations springing from the under surface, on the same principle as a wound is healed by the second intention. When the abscess is cured by resolution, it is from the adhesive, formative, or reparative process overcoming the suppurative and absorptive. The pus is then taken up and carried into the system; the walls proportionately thicken by new growth till they close in and obliterate the cavity.

The mode in which an abscess is first formed, in disease of the spine, appears to be as follows. The morbid process being a species of ulceration, the act of removal of substance is accompanied with suppuration. In an ordinary ulcer, say of the skin or mucous membrane, the pus thus discharged is either collected under the dressings or washed away. But in the spine the matter is caught by the connecting tissue; that becomes thickened and condensed; and at length an abscess, with proper pyogenic mem-

brane, is constructed. When a collection of pus is thus formed, various causes may lead to its progressive increase. From the denudation of the bone and fibro-cartilages, and the ex-

FIG. 221.*



of their surfaces to the pus, the ulceration may be extended and spread. Owing to the natural flexibility of the spine, the

* The patient from whom the specimen represented in the figure was removed, was a blacksmith, twenty-eight years of age, and was admitted to the hospital for a psoas abscess, about the size of a large orange, situated in the left groin. He had an abrupt angular deformity in the back with protrusion of the sternum; and his muscles, particularly those of the upper extremities, were largely developed. The deformity had taken place when he was a boy, and it had not increased since. His father being a blacksmith, he had been early put to the same work, and had continued uninterruptedly till three weeks before his admission, when he first noticed the swelling in the groin. On the abscess being tapped, a straw-coloured fluid was discharged; the opening did not heal; ordinary pus continued to flow; and he died exhausted with hectic fever. It will be remarked in the figure that the posterior arches and articular processes of the vertebræ, at the seat of the disease, have been firmly united together by bone.

tion between the opposing diseased surfaces will be great; but it will be made worse by the additional mobility consequent on the destruction of the fibro-cartilages and other ligaments which connect the vertebræ. Again, in proportion as the excavation proceeds, and the spine becomes more bent, all the adjacent structures in front of the column will become further relaxed, and so facilitate the accumulation of pus in the abscess.

Yet experience shows, as already remarked, that in a large number of cases the disease proceeds to the effect of producing extreme deformity, even combined with paraplegia, without any signs of abscess presenting themselves. But it is not to be concluded on that account that abscess has not existed in these cases. Owing to the inaccessible situation of the diseased parts, a collection of matter may have formed in connection with the carious vertebræ, and its presence would not be known if it remained of small size. An abscess, therefore, arrested in its progress before becoming sufficiently large to show itself outwardly, might have been lodged for an indefinite length of time in the angle of the diseased bones, and been eventually removed by resolution, without its existence having been suspected.

Let it therefore be asked, under what conditions it may be expected that an abscess in course of formation, in contact with diseased vertebræ and fibro-cartilages, may be arrested in its progress and dispersed? That inquiry introduces the subject of ankylosis.

Ankylosis.—When inflammation of the adhesive type affects a bone, the first change observed is a thickening of the periosteum, extending more or less to the connecting tissue around; by that means the adjoining structures are agglutinated and a bed prepared for the deposit of ossific matter. As stated before (supra, p. 105), one of the consequences of caries attacking the bodies of the vertebræ is, that adhesive inflammation is set up in the posterior segments of the bones. It may therefore be assumed that shortly after the commencement of the disease, the connecting tissue in the interstices of the vertebræ behind, will be consolidated, and by taking on the office of additional ligaments, will bind the arches and different processes together.

The immediate effect of this cohesion of several vertebræ will be to diminish, if not altogether stop, the jarring and friction between the diseased surfaces, which tend so greatly to aggravate the morbid action. An incipient abscess will, therefore,

be in a comparatively more favourable condition for undergoing cure by resolution. Accordingly, it may be supposed that the diseased process will sooner or later cease; that the abscess will be obliterated; and that the confronting surfaces of the partially destroyed vertebræ will come into apposition. The consolidated posterior segment will preserve the vertebra at rest, like splints applied to a fractured bone; the surfaces will therefore be expected to unite; with, perhaps, remnants of the bodies of several adjoining vertebræ all fused into one. Such is a description of *true* anchylosis.

Spurious anchylosis.—This term is applied in reference to the joints in general, to that kind of union between the surfaces after disease, in which the bond consists either wholly of fibrous structure, or of that combined with osseous matter. In the spine it denotes that the posterior surfaces of the vertebrae alone, including the laminae and various processes, are soldered by bone, while the surfaces of the bodies, partially consumed by the caries, are apart; and are connected above and below by the remains only of the walls of the obliterated abscess.

The writer had an opportunity of exhibiting, at a meeting of the Pathological Society,* a series of five different specimens of vertebræ united by bone at caries, and in which the anchylosis was partial. The appearances were essentially the same in all. Down the whole length of the posterior segments the affected vertebræ, perfectly solid ossific union could be traced between the margins of the laminae and the oblique processes, including their articular surfaces: and it was so firm and general, that all mobility was destroyed. No similar union had taken place on the fore part. The bodies had been destroyed to different degrees, and gaps of various width separated them; that the continuity of the spine as a column, depended wholly at the diseased part on the posterior segment. The amount of separation between the surfaces of the bodies, was owing to two distinct causes; in some of the specimens, distance at which the surfaces were apart was to be accounted for principally by the quantity of substance lost through the carious ulceration; in others was obviously consequent mainly on the too great obtuseness of the angle the seat of flexion, and the inability of the surfaces, from the spine having become fixed, to approach closer to each other.

The importance, in a practical view, of keeping this distinction between the two forms of anchylosis in mind is obvious.

1. It bears on the question, as to which position of the spine during the treatment, is most favourable for obtaining solid union between the surfaces at the close of the disease. The fri-

* *Path. Trans.* vol. i. p. 328.

A young patient are naturally much distressed when they see the rapidity with which the spine becomes bent, and they would induce the surgeon to employ measures to keep the column straight. They are not aware that a certain amount of deformity is a necessary condition of cure in this dangerous case. Loss of substance in the fore part of the column is inevitable: the destroyed portion cannot be replaced: the only way by which a cure can be effected, is by the upper and lower portions coming into apposition, and uniting as a broken bone would do; but that approximation and union can only take place by the spine becoming bent at an angle more or less acute. Hence, even if the treatment by extension were safe—which it is not—a sufficient argument against it would be, that in the event of its being carried out, the only union obtained would be in the posterior segment, and a permanent gap would be left in front.

2. It is sometimes advantageous for the surgeon to be acquainted with the proceedings of quacks. Bone-setters and others frequently undertake to straighten the spine in cases of angular deformity; and as they occasionally have a certain temporary success which gains them reputation, it is more necessary to understand what they do, and at what a risk. They will take, for example, a young man who has a prominent hump in the dorsal region, consequent on disease which he may have had in childhood. By confining him in the recumbent position, employing mechanical means to stretch the back, and obliging him on physical exercises, not forgetting to manipulate the projecting vertebræ sufficiently often, the spine may be made straighter, and the hump less distinct.* The quack will probably represent that the cure has been effected by his re-joining the dislocated bones at the angular projection. But it is obvious that the apparent improvement has been brought about wholly by the extension of the sound portions of the spine above and below the apex of the angle. Each of these portions, besides being inclined from before backward, is curved; and it can be easily understood that, by restoring them to a vertical line, the whole column will be rendered straighter, and the protuberance less distinct. But it is not

See a series of casts taken from patients who had undergone treatment by mechanical extension for angular deformity, contained in the Museum of the Essex Hospital.

to be expected that the improvement will last. As no can be made in the relations of the vertebræ at the ankylosis, and their surfaces, in conformity with the they form, are placed obliquely, the longitudinal axis of the and lower portions respectively must likewise be bent in - ence to each other. Hence the spine will soon return former condition. But it is the danger of the treatment constitutes the chief objection. And that arises from its impossible to know beforehand what is the exact kin degree of union that may have taken place at the seat of dis or what may be the state of the abscess. If the cavity or abscess has been obliterated, and the former communica between it and the carious bodies closed, and if the anchy has been perfect, so that no remnant of the disease has been the treatment, however rough, might do no harm, if it di good. But the results would be disastrous, if the anchy were spurious, and the walls of the abscess were the only nection between the upper and lower portions in front. O to the union between the vertebræ at the back, the mol would be destroyed, and it might be inferred that the anchy was perfect; yet the osseous bond might not be sufficiently st to resist the violence sometimes employed in endeavouri straighten the spine, by ignorant practitioners.

A friend related to the writer the following case, and also showed hi preparation connected with it. A young woman had caries of the cervical and upper dorsal vertebræ, from which she recovered, so as to b to resume her employment. The angular deformity being considerabl applied to a quack, who promised to make the spine straight. The trea consisted chiefly in employing mechanical means to stretch the neck. increasing the power one day, there was an audible snap, with sudden The patient was found immediately afterwards paralysed from the neck (wards, and in a few days she died. It was found at the post-m examination, that there was an abscess, nearly obliterated, in front of e vertebræ; and that the walls of the abscess had been extensively torn their connections with the bones. The spinal cord was diffuent near th of injury. It cannot be doubted that, when the force of extension wa creased and the snap heard, the bony connections of the posterior se of the vertebræ had been broken, the walls of the abscess detached fro bodies in front, and the spinal cord ruptured.

Psoas and Lumbar Abscess.

The abscesses hitherto treated of have been suppose have stopped growing before they attained a large size; have remained dormant, or been removed by absorption.

next to be considered are those which have progressively enlarged, and travelled long distances to evacuate their contents externally. Owing to the great dimensions which the latter commonly acquire, and to their cavities communicating with the vertebræ while still in a state of disease, they are especially dangerous in their consequences; they are, indeed, the most frequent causes of death in spinal disease.

A spinal abscess, at its commencement, lies upon the carious vertebræ, filling up the angle in front. It is favoured in taking that position by the bend in the column which has the effect of relaxing the structures on the fore part, and facilitating the accumulation of fluid. For the same reason abscesses will sometimes increase to a large size in the dorsal region without producing undue pressure on the roots of the lungs. As the pus accumulates, it is prevented by the portion of spine which forms the upper side of the angle overhanging it, from ascending; it cannot proceed backwards, owing to the thickness of the structures;* it would be difficult for it to make its way down the front of the bodies: accordingly, the ordinary course which an abscess takes when it enlarges is downward, by the sides of the bodies of the vertebræ. Sometimes the main sac bifurcates, one branch descending along the right and the other along the left side. In the thorax, the walls of the abscess are covered in front by the costal pleura; in the abdominal region, by the parietal peritonæum. When the abscess is connected with diseased dorsal vertebræ, it encounters in its descent the diaphragm. But that barrier is overcome by a particular process. As the abscess comes into contact with the diaphragm and compresses it, adhesive inflammation is set up in their respective surfaces; the consequence is, that they become united over a considerable area; an opening is next formed by absorption within the boundaries of the adhering structures; the abscess then protrudes; and extravasation of pus at the margins is prevented from taking place by the firm union of the parts encircling the opening.

Psoas abscess.—When the abscess has thus perforated the

* The writer does not remember more than one case in which a spinal abscess in the dorsal region opened directly backwards. There were two orifices near the projecting angle. The pus flowing from them had bubbles of air mixed with it; and it was noticed that these were drawn within the openings in inspiration, and expelled in expiration.

diaphragm and gained its abdominal side, it comes into union with the heads of the psoas muscle. That muscle is formed by one set of fibres from the sides of the bodies of the vertebrae, and by another from the roots of the transverse processes. The ligaments stretched across both origins in front, are the ligamenta arcuata. As the abscess, therefore, travels downwards, it has to pass through a narrow strait; it is prevented from enlarging before part, by the resistance of the ligamenta arcuata, and the back, by that of the spine and lowest rib; hence, in order to proceed it has to force its way in the line of the psoas muscle. That, however, can only be done by penetrating its interior. It accomplishes this, in the first place, by inserting its most advanced part, like a wedge, between the origins; it then splits and distends the fibres, so as to form a cavity for the reception of the pus; the muscular fibres become incorporated with the walls of the abscess; and the psoas at length is converted, more or less thoroughly, into abscess. But the muscle charged with pus does not expand equally in every direction. The fascia iliaca forms a kind of sheath for it; and this, being particularly strong on the inner side and united firmly to the brim of the true pelvis, prevents the growth of the abscess inwardly. On the outer side, however, the connections are loose; and there enlargement takes place freely. The abscess now chiefly occupies the hollow between the united fibres of the iliacus internus and psoas muscles on the inside, and the crest of the ilium on the outside. When the advanced part reaches the level of Poupart's ligament certain retardation occurs; and then a bulging will be observed along the line of the flexure of the groin. The abscess perforates the abdominal walls in the same way as it formerly did the diaphragm. And the opening is invariably at the same place: namely, behind Poupart's ligament, between the united tendons of the iliacus internus and psoas muscles, and the anterior inferior spinous process of the ilium. The situation corresponds to the point of junction of the outer with the middle third of Poupart's ligament.

Before the protrusion takes place, there will be a certain amount of thickening and condensation of the connecting tissue in the groin—a consequence of the inflammation excited by the near approach of the abscess—and a provision at the same time for strengthening its walls when it does emerge. On the abscess actually escaping, it will be suddenly relieved from

terable compression to which it had been subjected in the interior; and it is from that cause probably that it increases in size as soon as it appears outwardly. The enlargement is sometimes greatest in the groin, and for a short way.

But it is more common for the abscess to descend for a distance on the thigh; and the direction then varies. It follows the walls abutting at the commencement of its course against the origins of the sartorius and tensor vaginae femoris, and then, by their opposing its progress outwardly, the line which it most frequently takes is inward and downward, along the sartorius; and it often accompanies that muscle to near the knee. Sometimes, however, soon after its appearance in the groin, it turns sharply inwards to occupy a space over the adductor muscles. In a comparatively rare set of cases it bends outwardly. The least frequent direction is straight downwards. Occasionally, the abscess divides at the groin; one portion going inwardly and the other outwardly.

Neck of the abscess.—It seems appropriate, on the analogy of a hernia, to call the narrow part of the abscess where it passes through the abdominal parietes, its 'neck.' It may be that, when an abscess has been formed in the thigh, dependent on matter descending from a diseased portion of the spine, it is free, like others arising differently, to secrete additional pus from its own pyogenic membrane, or to absorb it; but it would appear that abscesses originally spinal, are mainly dependent on the parent one. Now, the channel of communication between the well from which the fluid is drawn, and the reservoir below, is constricted at the neck more than elsewhere. It will depend, therefore, greatly on the width of the aperture as to whether the abscess in the thigh be abundantly or sparingly replenished from above. In general there is a disposition in the opening to contract: although it has never, according to the writer's knowledge, become entirely closed. The degree to which the narrowing has gone, will commonly be judged of pretty accurately by the feel of the parts when the contents of the abscess have been squeezed upwards, and then when compressed, and comparing them with the condition of the corresponding parts on the sound side. Occasionally both the well and the channel leading from it to the abscess are constricted to such a degree that no traces of them can be found: the collection appears circumscribed; and a question arises as to whether it be really connected with the spine. The surgeon

may, therefore, have recourse to another mode of examining if he bandages the abscess firmly for a few days, it will be to have had the effect of reducing the swelling considerably; let him next leave the abscess without bandages permit the patient to be in the upright position; he will find that the swelling has returned to its original dimensions—thus showing that the channel was pervious, and the pus could be passed along it, to and fro.

Lumbar abscess.—It was stated formerly, when describing the process by which a spinal abscess, originating in the thoracic region, made its way through the diaphragm, that the pus was effected through a narrow opening, bounded in front outwardly by the ligamenta arcuata, behind by the lower ribs, and on the inside by the bodies of the vertebræ; and added, that the abscess continued its descent by penetrating the psoas muscle. At the point where this obstruction occurs, with, and where the ribs cease to form an obstacle to the progress of the abscess backward, it occasionally perforates the peritoneal parietes, and presents itself in the loins as a ‘lumbar abscess.’ In some cases the whole abscess takes that form, but in others only a part; so that the lumbar is merely a shoot of a psoas abscess. Again, a lumbar abscess may be in connection with diseased upper lumbar vertebræ; just as may be the case with a psoas abscess. The lumbar abscess penetrates the quadratus lumborum, and it appears under the skin on the outside of the sacro-lumbalis muscle. In its progress posteriorly the abscess meets with opposition from various fasciæ and tendinous expansions; which deflect its way, and cause it to assume outwardly a broad and flat appearance with a slight elevation.

Spinal abscess in the neck.—Disease affecting the articulation between the two uppermost vertebræ of the neck, the atlas and axis, is characterised by certain peculiarities which entirely deserve to be treated of separately: its consideration will, therefore, be postponed to the end of the article. When any of the cervical vertebræ are the subjects of caries, a collection of pus will form on their fore part, as in the spine generally, instead of the matter accumulating there, and forming a prominent swelling which might compress the trachea or phagus, or dropping downwards, it is deflected laterally. In the neck the abscess appears outwardly on one or both sides of the spine, inclining somewhat posteriorly.

varieties in spinal abscesses.—It is proper to mention some options or anomalies met with in abscesses in different parts of the spine.

In describing the first stage of the formation of psoas abscess, it was stated that the sac occasionally bifurcated; one branch descending along one side of the bodies of the vertebræ, the other along the other. Cases in which both have moved and pointed at the groin are extremely rare. Dissection sometimes reveals the fact, not ascertained during life, that the abscess, on one side, ran its whole course and discharged its contents on the thigh, another, on the opposite side, stopped short about midway. In every case an examination ought to be made to ascertain whether there be two abscesses or only one.

It is not uncommon, in a case of psoas abscess, for the collection of pus to be arrested in its progress downwards, just above Poupart's ligament, or when it is lying on the venter ilii. It may then increase in size and become a prominent, globular swelling, with distinctly defined boundaries.

A young girl, fourteen years of age, had disease in two or more of the dorsal vertebræ; it had caused great angular deformity, and for a certain time, paraplegia; at length, a globular swelling, visible to the eye, formed in the right iliac region. After remaining stationary for a short time, it began to diminish; then it got gradually smaller, till, after the lapse of several months, it had entirely disappeared, so that no trace of it could be found. The treatment consisted in the continued application of the tincture of iodine. Finally the patient's health was quite restored; the deformity alone remaining.

When a spinal abscess, in descending, reaches the hollow of the ilium, instead of pursuing its course downwards, it occasionally turns sideways, mounts over the crest of the ilium, discharges its contents, by one or more openings, over the iliac muscles. The same thing is sometimes met with in connection with lumbar abscess. The occurrence is more frequent in children than in adults: a difference which may be accounted for by the relatively small size of the pelvis in the former.

Sir B. C. Brodie has described certain cases in which part of the contents of a psoas abscess, when lying in the iliac space above Poupart's ligament, has made its way into the inguinal canal, and so appeared at the external abdominal ring like an inguinal hernia.*

* *On Diseases of the Joints*, p. 267.

The writer has witnessed a similar case, in which the tumour had some resemblance to a reducible hernia. Along the flexure of the groin, and for some distance above, there was general fulness and induration of the structures. At the external abdominal ring, and for a short distance below, in the line of the spermatic cord, there was a circumscribed swelling, which became more prominent when the patient coughed. The contents of the supposed abscess, however, were quite soft and compressible. The man was not aware that he had disease of the spine, having come to the hospital for some comparatively trivial complaint. On turning to the back, there was an absence of any direct projection of the spinous processes; all that could be perceived was a slight arching backward of the vertebræ in the lumbar region, which was in contrast with the hollow naturally existing there. The chief diagnostic symptoms were well-marked rigidity of the column at the loins, and natural flexibility elsewhere. Shortly after his admission he rapidly got worse; the abscess in the groin broke; and on his death, the bodies of three lumbar vertebræ were found carious and surrounded by pus.*

5. *Contents of spinal abscess discharged by the lungs.*—It has been already stated that when abscess forms behind the posterior mediastinum in connection with diseased dorsal vertebræ it is generally prevented, by the bending and receding of the spine, as well as by the cavity emptying itself downwards, from enlarging to such a degree as to press on the roots of the lung and impede their functions. But a danger of another kind is to be apprehended from the proximity of the collection of pus to the lungs: inflammation is apt to spread from the pleura costalis which covers the walls of the abscess, to the pleura pulmonalis and substance of the lungs: pleurisy or pneumonia, or both combined, may then be excited, and death may ensue. A different result, however, may be looked for, in a few exceptional cases: namely, the evacuation of the abscess through the lung. The process by which that is effected is the same that has been described in explaining how an abscess perforates the diaphragm in passing from the thoracic into the abdominal region. It is the same also as the process by which an abscess of the liver discharges its contents through the lungs. Firm adhesion is first established between the walls of the abscess and the surface of the lungs; next, in some central part within the area of these adhering structures, an opening by absorption is made, having the margins secured by the surrounding adhesion; this opening brings the interior of the abscess into communication with lung substance, which becomes condensed in the neighbourhood; a tunnelling is commenced through the latter un-

* See preparation in the Museum of the Middlesex Hospital.

large bronchial tube has been reached and opened; then the sputum is coughed up and expectorated.

The writer was consulted in the case of a boy affected with angular deformity, the apex being formed by the spinous processes of the fourth and fifth dorsal vertebrae. He had been confined to bed at the commencement of the disease, five years before; he had since then been strong and active; and the object of the consultation was merely to ascertain whether anything could be done to improve his figure. During the interview, the patient took from his pocket a small box containing fragments of bone; and these he said he had coughed up, together with a large quantity of matter, when his illness was at the worst. The structure of the pieces resembled that of the bodies of vertebrae; they were generally cancellous; but had here and there portions of cortical layers upon them; they were four in number, irregularly cube-shaped, the sides measuring, on an average, from three to four lines. It was supposed that they had lain loose in an abscess at the seat of the deformity, and been expectorated.

6. *Abscess opening into intestine.*—It is not uncommon for abscesses in the abdominal region, with whatever viscus they may be connected, to empty themselves into some part of the intestinal canal if they come into contact with it. For example, an abscess of the liver frequently terminates in that way.

The following is a brief notice of a case in which a spinal abscess communicated with a portion of intestine, and in which not only did the matter of the abscess pass into the bowel, but the contents of the bowel discharged themselves by the abscess. The patient was a girl, thirteen years of age, who had an acute angular deformity at the centre of the dorsal vertebrae, and a psoas abscess of several months' duration on the left side. The abscess reached down to the knee; the external opening being over the inner condyle of the femur. As it was proved that the abscess admitted some of the contents of the intestines into its interior, the patient, during the writer's visit, picked from the orifice the contents of one or two peas, and a bit of undigested stalk of greens, which she had been lately eating.

7. *Spinal abscess discharging its contents by the urinary bladder.*—

An under gardener, twenty-four years of age, had angular deformity; the most prominent spinous process being that of the seventh dorsal vertebra. He could not tell when the disease began; and as he had never missed his day's work on account of it, he had reluctantly left his situation to become a patient. By carrying the fingers deeply in the direction of the psoas muscle, a large abscess could be distinctly felt. He remained in the hospital two months, during which no perceptible change occurred. At the end of six months he returned; having worked at the spade the greater part of the intervening time. He stated that a week before, he had slight trouble in passing his water; and shortly afterwards, he voided a large quantity of matter of a yellow colour, along with the urine. That he had continued to do

since; but latterly, the proportion of the pus to the urine had diminished. During his stay in the hospital, about a third of what was consisted of pus. No trace of the abscess in the iliac region, which had been repeatedly felt before, was discernible. The patient left the hospital on his own accord, a fortnight after his readmission; and he could not be traced afterwards.

Diagnosis of spinal abscess.—One of the most frequent sources of uncertainty in distinguishing abscesses connected with the vertebræ from other swellings in the same locality, is the insidiousness with which spinal disease, as already stated, commences and proceeds.

The following is an example, being that of a psoas abscess mistaken for a hernia. The patient was a young man, son of a large farmer, fond of hunting, and he had regularly followed the hounds till a few days before coming for consultation. He had observed, for a week, in his left groin, a swelling, which his medical attendant told him was a rupture requiring a truss. On flexure of the groin there was an oval tumour, of well-defined shape, resembling a hernia, except in the point at which it emerged. It was behind Poupart's ligament, an inch and a half externally to the part where a femoral hernia descends; in other words, it protruded at the place where a psoas abscess appears when it passes out upon the thigh. Above the tumour, and in the iliac region generally, there was fulness and tenderness. On examining the back, no distinct protuberance of any of the spinous processes was visible. But in the loins, there was a perceptibly greater bulging forward of all the vertebræ than is natural, without undue prominence of any individual bone; the lumbar vertebræ were likewise abnormally rigid; free mobility being in the dorsi-lumbar region. The patient owned that some months before, he had received an injury in the lower part of his back, which was so slight that he had not given up any of his active pursuits for some time. The dangerous nature of his illness was explained to him. He was loth to forego his hunting. Ere long, the abscess enlarged, broke, and discharged a large quantity of pus. When last heard of, his recovery was hopeless.

An abscess originating in caries of the lower lumbar vertebra and extending it may be to the sacrum, is liable to extend laterally on the wing of the ilium, and to rise above the crest of the ilium. A swelling of that kind, if it be on the right side, is often confounded with a cæcal abscess. Or if it surmounts the crest of the ilium and discharge its contents backward and downward, the symptoms of the hip will be disturbed; and it will be supposed to have originated in morbus coxæ. The diagnostic sign most to be relied on in these doubtful cases, is the diminished flexibility observable in the lumbar region.

A tumour of medullary cancer is not unfrequently met with according to its situation, for either a lumbar or psoas

The uncertainty will be increased if the cancer have invaded the spine, and by converting part of it into its own morbid structure, has caused the column to become bent. Again, it is the peculiar nature of this kind of carcinoma to throw out globular projections on the surface, resembling greatly the pointing of an abscess; moreover, the feeling of elastic resistance at the apices of these lobes, is so much like that of undulation of pus within an abscess, that a mistake in diagnosis is very apt to be made.*

The writer had lately under his care, in the hospital, a female, of middle age, who had lateral curvature of the spine, from girlhood; and in whom the lump was so prominent and abrupt, that it might be mistaken for angular deformity from caries of the vertebræ. She was admitted for a tumour that occupied the inner side of the left wing of the pelvis, and was on a level, at an exterior part, with the crest of the ilium. An oval-shaped, projecting lobe, of a course of time, formed on its most depending face, near Poupart's ligament; and the feeling communicated to the finger, when examining that part, bore the closest resemblance to what is conveyed by pus, when near the surface of an abscess. The subsequent progress and termination of the case, showed that the tumour was one of medullary cancer.

Treatment of external spinal abscesses.—In discussing the question of the two kinds of ankylosis of the spine, the true and false, reasons were advanced for believing that, when an abscess at the seat of disease subsided, to give place to ossification, it was to the arrest of motion caused by the consolidation of the posterior segment of the spine that the beneficial effects were principally to be ascribed. In such cases, the abscess is confined within the range of the rigid portion of the column. But when unfavourable influences interfere with the termination in ankylosis, and the abscess enlarges by extension of its walls downward, it gets beyond the bounds of the inflexible part, takes its place in movable structures, and consequently loses the advantage which rest is calculated to confer. Thus progressive increase goes on, and the abscess eventually presents superficially, where, of course, exposure to the bad effects of motion is worse. These considerations enforce the propriety of employing in the management of such abscesses, every available means for procuring perfect rest.

Another general observation applicable to the subject is, that in common with other collections of pus having their origin in diseased joint structures or diseased bone, it is the continued

* See Article CANCER, vol. i. p. 565.

irritation kept up in the walls of the abscess and adjacent parts by the presence of the carious bones and ulcerated surfaces which chiefly leads to the fresh formation of pus; whence measures designed to forward amendment in the abscess, ought to be accompanied with others directed to improve the condition of the spine at the original seat of disease.

An additional remark may be made, which still applies to large abscesses whether originating in the spine, hip, or other joints, when diseased. It relates to the contrast observed in the condition of the patient, before and after the evacuation of the contents: either when the abscess has burst spontaneously, or been opened by the lancet. It is constantly seen in practice that an abscess of extraordinary size will often remain for many weeks together perfectly quiescent; perhaps enlarging a little at one time, and diminishing a little at another; the patient being all the while free from constitutional disturbance. But if the abscess point and break, or the surgeon open it, and the puncture does not heal, but the matter continues to run, an important change will soon occur. The unfavourable symptoms come on at various periods in different cases. It may be expected that within a few days the quantity of pus from the abscess will greatly increase: that it will become thin, and perhaps variously discoloured. At the same time also, the patient will sicken, his pulse will be quickened, his skin hot with occasional profuse perspirations; the fever will be hectic, attended with gradual wasting and increasing debility; the liver will probably enlarge by fatty degeneration; and he will die exhausted by the combined effects of the purulent discharge and the fever. (See article HECTIC, vol. i. p. 287.)

Now it is the anticipation of that change being brought on by evacuating a lumbar or psoas abscess, that chiefly leads to hesitation and delay in the treatment. The surgeon may rest, for a while, on the hope that he may possibly succeed in dispersing the collection by resolution: he therefore keeps the patient rigorously at rest; supporting the strength by tonics and diet; and he applies assiduously tincture of iodine to the back, at the seat of disease; to the side, in the course of the abscess; and to the swelling itself; he thinks that if the abscess be not altogether removed by such means, its size will be diminished, and the walls converted into a fitter condition than before for the operation of puncturing. His expectations, however, will probably not be answered favourably. He may

perceive that at some particular part there is an appearance of accumulation, and he is desirous of averting spontaneous opening. Necessity, therefore, seems to compel him to adopt the plan of tapping the abscess and drawing off the pus. It may be supposed that the proceeding has been carried out in the most approved way, with all the precautions commonly urged (see article ABSCESS, vol. i. p. 115); yet the almost inevitable termination will be, a bursting forth of the matter, from the temporarily closed openings artificially made, and an established discharge from the abscess, with all the uncontrollable ill-effects apprehended as the sequel. These considerations prompt us to greet with welcome the valuable labours of Professor Lister, and of others, in their present active endeavours to improve his part of surgery.

Affection of the Spinal Cord from extension of the Disease.

When we consider the delicacy of structure and constitution of the medulla spinalis, and its contiguity to the centre of so destructive a morbid action as caries, we may be surprised that its functions are not lost in a greater number of cases than experience shows to be the fact. Yet, as the surgeon has no means of foretelling, in any particular patient, whether the cord will be affected or not, he is kept in constant apprehension.

Several circumstances combine to protect the cord from injury. First, in proportion as the spine falls forward, and forms an angle at the seat of disease, the abscess is propelled to the front, and the chasm between the bodies is closed behind, so as to prevent the pus from having access to the vertebral canal. Secondly, it is within the posterior segment of the spine that the cord is contained; and it has been formerly shown that, while the anterior segment, composed of the bodies, is the special seat of caries, the posterior is exempt from disease, and has a reparative or defensive action carried on within it: the cord, it may therefore be inferred, will not be injuriously affected by such kind of action. Thirdly, it might justly be feared that when a part of the spine had been destroyed by caries, and a deep gap formed between two or more of the bodies, the column would be so much weakened that it would be liable to be broken by any slight injury; and that the cord would therefore be crushed and deprived of its functions. But it has been already pointed out (supra, p. 115), that when caries attacks the vertebræ

in front, a process of agglutination and consolidation, terminating in firm osseous union, is immediately set up in the posterior surfaces of the same bones. Hence, in proportion as the spine yields to the pressure of the superincumbent weight, and the upper part tends to fall forwards, that movement is restrained and moderated by the opposition of the structures behind, which have thus been endowed with increased strength. Accordingly the angle at the seat of disease is formed in a slow, gradual manner, without jarring or abruptness; and the cord becomes accommodated to its novel position without loss or impairment of its functions.

Before describing the morbid condition of the cord from which paraplegia ensues, it may be interesting to notice certain considerable changes in its structure which are connected with the deformity, but which do not involve any deprivation of its functions. When a large portion of the front of the spine has been removed by caries and the surfaces have united, not only will the characteristic angle be formed, but the column will be considerably diminished in length at the seat of disease. This shortening is to be attributed to the loss of substance which the affected vertebræ will have sustained, and to the having collapsed into a comparatively small size before union. The vertebral canal and the cord contained in it, will necessarily participate in this bending and shortening, and be accommodated to both. Now it may be inferred that, to bring about the adaptation adverted to, a succession of minute interstitial changes in the substance of the cord must have been wrought during the progress of the deformity; that certain portions of the material will have been abstracted and their places not filled up; and that other portions will have been arranged in a different order from at first. It is not found, however, that either the atrophy, or novel arrangement of the nerve substance, has a prejudicial effect on the functions; for these will remain entire, however extreme the deformity. The explanation is founded on a well-known law in the pathology of the brain and cord; namely, that changes of extraordinary magnitude may be effected in their structure by encroachments of various kinds, without the sacrifice of their functions, on condition that the intrusions are made slowly and gradually. Remarkable examples of the kind, connected with atlo-axial disease, will presently come before us (*infra*, p. 143). Moreover, every case of lateral curvature furnishes examples.

Disorganisation of the cord ; paraplegia.—From what has just been stated, it will be perceived that the change of structure in the cord which causes paralysis of the lower extremities, is not the mere alteration of its figure or size consequent on the angular deformity. It consists in a conversion of a portion of its substance into a new morbid tissue. A certain part of the organ, adjacent to the seat of caries, undergoes a process of breaking-down of its texture; the most conspicuous effect of which is a softening, approaching to liquefaction, of the white columns and grey matter. If the diseased part be tapped with the finger, it will readily yield to the slightest pressure; if a stream of water be allowed to fall gently upon it, the soft substance will be washed away, leaving the connecting tissue. When the cream-like matter is examined in the microscope, it is observed to be made up of innumerable fragments of nerve-fibres, granules, oil-globules, and amylaceous cells, the débris of medullary tubes and ganglion-cells. A tinge of yellow pervades the softened and neighbouring parts. The membranes are not commonly present any decided signs of having been irritated, nor is the vascularity much altered.

It will depend on the depth to which the morbid change extends, in any particular case, as to the functions of the cord which will be abolished. We owe to modern discoveries the knowledge of the fact, that to the anterior column, from which the anterior roots of the spinal nerves arise, belongs the power of controlling the muscles; and that to the lateral column, from which the posterior roots arise, belongs sensation. Hence if the disease penetrate only so far as to destroy one column, leaving the other sound, we may anticipate that the function pertaining to the disorganised column will alone be lost.

Experience proves that, in paraplegia caused by disease of the spine, motor power is much more frequently destroyed than sensation: and so general is this fact, that it may be regarded as a comparatively rare thing for both motion and sensation to be lost simultaneously. As to the reverse kind—abolition of sensation with motor power remaining—it is doubtful whether it has ever been met with.

When we consider the relative position of the different columns of the cord to the centre of disease in the spine, we shall at once perceive why motion should be first destroyed. The anterior column, on which motor power depends, is separated from the bodies of the vertebræ only by the theca

vertebralis and the other membranes of the cord : hence it may be expected that, when the morbid action proceeds backward, this column will be deprived of its function before any other. And its liability to be so involved will be increased by the compression to which it is peculiarly subject from the bending of the spine, as it forms an acute angle at the part.

Moreover, if we trace a filament of an anterior root to its origin, we shall find that, as it is about to join the cord, it subdivides into numerous minute radicles, and that these lose themselves almost imperceptibly on the *surface* of the anterior column. If, on the other hand, we take a filament of a posterior root, we shall observe that it continues as a single fibril up to the cord ; that when arrived there, it dips bodily without subdivision, into the fissure between the posterior and lateral columns ; that while in the fissure, it pursues an unbroken course to the bottom ; and that its actual termination is in the *deepest* part of the cord, where the central cineritious matter is situated. It appears, therefore, that the part of the medulla appropriated for sensation must be in the internal well-protected recesses of the organ ; where it is less likely to be reached by disease invading the exterior, than the column of motion, situated more superficially.

It will depend on the height in the spine at which the cord is diseased as to the extent of the body that will be paralysed. If the disease be in the lower lumbar vertebræ, that is below the termination of the cord, there may be an absence of paralysis : unless, as occasionally happens, the products of inflammation have extended upward, to involve the medulla above. In not a few cases, one extremity is more completely deprived of motor power than the other. Comparing the paraplegia resulting from disease with that from injury, it is remarkable that the former is seldom accompanied with loss of command over the bladder or rectum ; and that when these organs are affected, it is commonly in a slight degree. Retaining sensation, the patient may have an urgent desire to pass water, but he will be unable to do so, and will require the assistance of the catheter. Again, the stools may pass at one time involuntarily, at others consciously. In general, the torpidity of the bowels renders it necessary to give aperients frequently.

The cases are comparatively rare in which the loss of power in the limbs leads to the formation of bed-sores. If the patient be an adult, who suffers much pain at the seat of disease,

making it difficult to shift his position, there will be danger of such sores; but not so great as in cases of paraplegia from fracture of the spine, when the fæces and urine pass involuntarily, and the hips are constantly sodden in the filthy moisture. In a young person, on the other hand, whose body is light, in whom sensation has not been lost, and who can feel the irksomeness of lying for a long time in the same posture, we do not look for sores; he can turn himself by the use of his arms, change the position of his hips, and thus avoid the bad effects of pressure.

Spasmodic actions of the paralysed limbs.—When the lower extremities are entirely bereft of voluntary power, they are subject, nevertheless, to convulsive actions; through some obscure morbid influence acting on the distal portion of the spinal cord cut off from connection with the brain. The spasms are the same that are witnessed when the cord has been destroyed by violence, or by the encroachment of tumours, ramollissement, &c. A frequent form of the attack consists in a sudden jactitation of one or both legs. While the patient is lying quiet, the limbs will suddenly be drawn up; they will be bent both at the knees and hips—so that the heels will come in contact with the back of the thighs, and the knees with the abdomen: the flexor muscles will then relax, and the limbs will fall straight, as before. In such cases the spasms are of short duration; but they may be repeated with great frequency, so as to cause much distress. A different form consists in the legs being drawn up, and then remaining bent. In these cases, the limbs and body will be so rigidly fixed, that the patient may be moved about in bed, in one mass, like a log of wood. Cases will be met with in which both the above forms are presented, at different times, in the same patient. The spasmodic actions, whether clonic or tonic, are apt to be excited by certain irritants; thus tickling the soles of the feet, pinching or pricking the skin, the act of moving, will bring on the sudden startings, or will aggravate the contractions of the rigidly bent limbs. Micturition and defæcation sometimes, but rarely, excite attacks. Yet they frequently come on without any perceptible cause of irritation. When the patient is drowsy and about to drop asleep, he is peculiarly subject to be woke up by the limbs jumping.*

* 'Contributions to the Pathology of the Spinal Cord,' by William Budd, M.D. *Med.-Chir. Trans.* vol. xxii. p. 153.

The pain occasioned by the spasmodic affections is sometimes excessive. In certain cases it equals, if it does not surpass, the agony of tetanus in its worst form: happily, in the majority the suffering is moderate and endurable. Sometimes when the convulsions are strong, they shake the whole body, and communicate a painful jar to the diseased vertebrae. Again, the patient have bed-sores, the concussion produces friction, the hips against the bedding, and that aggravates the ulceration or sloughing. When it is attempted to subdue the spasms, by applying splints to the limbs, or by bandaging the leg that is affected to the one that is quiet, and fastening them, when secured, by india-rubber tubing to the bedposts, the plan commonly fails; the skin is liable to be frayed, and the pain is not diminished. There is no particular medicine that can be relied on for overcoming the convulsions. Good effect sometimes follows injecting morphia under the skin.

Diagnosis.—As intimated before, paraplegia consequent on caries of the vertebrae does not differ materially from that which may be caused by morbid affections of the cord, of various distinct kinds. The principal diagnostic symptom in spinal disease is, doubtlessly, the angular deformity. Observation shows that paraplegia of this kind does not occur till late in the disease, after the bodies have been partially destroyed. Consequently, if there be no distinct projection of the spinous processes, and rigidity of the vertebrae, to indicate excavation of the column, it is probable that the paraplegia depends on some other morbid affection. Again, palsy arising from caries is commonly distinguished by motor power being abolished while sensation is unimpaired. If the reverse be found, namely, loss of sensation with retention of motor power, it may be inferred that the cord is encroached upon by some morbid growth advancing from behind forward.

Prognosis in paraplegia.—Experience proves that in paraplegia from spinal disease, restoration of the lost powers of the limbs is not uncommon. The mode in which recovery takes place appears connected with the bending of the spine at the seat of the caries. It has been previously stated that in the majority of cases, the paralysis consists in the loss of function by the anterior column alone, motor power being the only property destroyed: whence it was inferred that but a comparatively small part of the cord was disorganised. It may therefore, be conceived that, in proportion as the spine sin-

seen place before the diseased surfaces of the bodies come in contact, and when a permanent gap has been left between them, there will be little chance of recovery. Prognosis is more favourable in children than in adults; it may be ascribed in considerable degree, at least, to the presence of the sound portions of the cord adverted to being readily effected in the former than in the latter, owing to greater flexibility of the spine in early life. The patient had lately under his care a boy, six years old, with a deformity in the dorsal region. Within a period of years, his lower limbs were twice completely paralysed: at first time, for six months; the second, after an interval of months, for four. He recovered from the paraplegia, but afterwards died from the effects of extensive abscesses. Restoration of the power of the limbs, in some cases, is partial. Certain groups of muscles continue paralysed. Unequality in the condition of the muscles leads to paraspasms of contractions and deformities, chiefly in the hands and toes.

Treatment of Spinal Disease.

General treatment in spinal disease is founded on the principles that guide us in hip-disease, scrofulous synovitis knee-joint, disease of the ankle, &c. The remedies are

fixing the diseased vertebræ, we endeavour to prevent jarring and attrition between the rugged opposing surfaces of the carious bodies, and, by removing that source of irritation, to encourage the morbid action to cease. Secondly, by keeping the bones at perfect rest, we afford the parts the most favourable opportunity for becoming consolidated by ossific union.

The most effectual method of arresting motion between the diseased vertebræ, in aid of nature's efforts, is to place the patient in the recumbent position on his back; and to give him the greatest advantages whilst confined, it is desirable that he should be furnished with an invalid-bed, provided with conveniences for enabling him to lie upon it, day and night, without rising. If an ordinary couch be used, the patient will lie with greater security and comfort if the frame for the horsehair mattress, instead of being horizontal, be gently inclined downward from the top to where the hips come, and then be elevated (but to a slighter degree), with a double incline, for the knees to be bent over the angle. A hollow or socket for the reception of the angular projection of the spine ought to be made in the mattress. Or a substitute may be contrived by filling an india-rubber pillow with sand, or suitable grain or seeds, and placing it under the patient's back, protected with blanket and sheet. In young persons, further security will be required, by enclosing the body in a stiff corset; and, perhaps, by strapping them down to the couch. It will add to the convenience if the bed-frame be portable, and capable of being easily transferred to the body of an invalid-carriage, or to a bench in the garden, with a view to the patient's getting fresh air.

It is an important duty of the surgeon to be peremptory in enforcing strict confinement in the recumbent position. He will find this task, of course, more difficult in children than in adults; and the principal cause is the freedom of the patient, in the majority of cases, from pain. Nevertheless, however moderate the symptoms with which the disease is ushered in may be at first, the incipient stage is a period during which a rigid observance of perfect rest ought to be insisted upon. And the time over which that care should extend ought not, even in the most favourable cases, including children, to be less than six or ten months. In patients approaching adolescence, the confinement must be longer.

If the case should proceed favourably, without pain in the back or abscesses coming forward, for the number of months

specified above, the next practical question will be, concerning the propriety of liberating the patient from his long durance. That is equivalent to asking, how it may be ascertained that ankylosis has taken place; for it cannot be considered safe to allow the patient to get into the upright position till the bones have become strongly knit. Accordingly, in examining the angular projection, what has chiefly to be made out is, whether there be any motion between the affected bones; for if the part be perfectly stiff, it may be presumed that they are united. Atrophy of the muscles may assist in confirming the point. But it will not be prudent to rely on the solidity of the ankylosis altogether. The patient ought to be supplied with artificial supports. Space does not allow of a detailed description of the best kind of these instruments to be used in different cases; for, of course, they must vary, not only according to the situation of the disease, but the age of the patient. One general remark may be made, that the corsets are to be constructed much in the fashion of women's stays, so as to embrace the body above and below equally and generally; they ought also to be strengthened as much in front as practicable, to counteract the tendency to stoop forward. For giving them the proper stiffness, steel ribs are preferable to whalebone. In young, feeble children, whose bodies are extremely flexible, while their hips are too narrow for being used as a base of support, it may be found best to encase them in separate pieces of prepared leather, moulded to the body, and afterwards inserted in corsets; the lateral pieces or splints may be prolonged downward, to rest on the seat of the chair, and brought high enough to reach the arm-pits, so as to act as crutches while sitting.

Prone-couch.—A mode of reclining in the prone position, instead of on the back, has been much recommended. The method consists in making the patient lie upon his chest and abdomen, on an elevated frame, not unlike a desk, with an inclined board upon which his legs may hang: he is prevented from sliding by pegs at the top received into the arm-pits. The object of the position is, by fixing the upper part of the trunk, and allowing the hips and legs to glide down the incline, to keep the spine on the stretch, and so diminish the acuteness of the angle at the seat of disease. But such a design is opposed to the principles advocated above (see p. 117). It is evident that if the angular projection be diminished, the surfaces of the diseased bodies of the vertebræ, which ought to meet, will

be kept apart, and ankylosis will be prevented. Moreover, the patient cannot sleep, and cannot empty the bladder or rectum while confined on the apparatus, it will be necessary to move him frequently, to the great injury of the back. But will, perhaps, be said, that although the employment of the prone-couch be not adapted to caries in its active stage, it may be advantageously used to improve the figure, when ankylosis has been effected. That desire on the part of patients, who have narrowly escaped death at the price of being deformed, to have their cure made absolutely perfect by getting rid of the hump on the back, may be natural; but it is scarcely necessary to say how utterly vain such hopes are.

Counter-irritants.—It is not long since that, in cases of angular deformity, issues were almost invariably ordered, on each side of the protuberant vertebræ, whether the patients were young and weak, or the disease acute or chronic, or there were abscesses or paraplegia. They had a debilitating effect, and were attended with much constitutional irritation. According to the practice of the present day, issues are reserved for subduing casual accessions of the morbid action, when the pain is great, or when there is a threatening of paraplegia. They effect all their good in a short time, so that it is not necessary to keep them open. Blisters give rise to so much general inconvenience that they are seldom had recourse to. Compound tincture of iodine, as a local application, has supplied us with an efficient substitute for these coarser remedies.

DISEASE OF ATLAS AND AXIS.

Disease affecting the two highest vertebræ of the spine has certain peculiarities which entitle it to separate consideration. The same cause which makes accidents to those bones more dangerous than elsewhere,* renders morbid action more perilous. The portion of spinal cord contained within the ring of the atlas is the most vital in the nervous system. Hence, whatever tends to weaken the joints in that part, exposes the patient to the hazard of sudden death, or of paraplegia including the whole body below the head.

Both atlas and axis, especially the former, differ from the other vertebræ in their anatomical structure. The atlas is

* See INJURIES OF THE BACK, vol. ii. p. 394.

efficient in the part that corresponds to a body; and there is a fibro-cartilage either between the axis, or the occipital bone and it: hence, as these structures are the proper seat of caries, it may be expected that the morbid process which attacks the two bones will differ from the ordinary disease of the spine.

The joints particularly subject to be affected are those between the atlas and axis. And a reason may be assigned for their being peculiarly prone to disease. The term 'axis' implies that the vertebra so called is the centre of the rotatory movements of the head; and as the spokes in a carriage-wheel inserted into the axle act upon that central point with great lever power, so that when a break-down occurs it is the axle that gives way, a corresponding impulse is directed upon the pivot of the axis, by sudden violent rollings or twistings of the head. These shocks and sprains, often repeated, predispose the articulations, in a delicate patient, to disease.

The morbid action thus set up in the atlo-axial joints is similar in pathological character to what commonly occurs in the wrist, elbow, knee, &c. Proceeding generally in a chronic form, it gives rise to a gradual softening and spongy thickening of the synovial and fibrous tissues that connect the vertebrae; and subsequently to ulceration of the cartilages, and superficial corrosion of the bony structures. The disorganisation will go on in most cases to an extreme degree without signs of abscess.

That which contributes principally, as intimated before, to the importance of disease in this locality, is the danger threatened to the spinal cord at its most vital part. Owing to the head being sustained on the atlas, and the ligaments which bind the latter and axis ceasing to be able to retain them in position, the weight of the head causes the atlas to glide forward and downward upon the axis. This movement implies that not only are the articular surfaces separated from each other, but that the odontoid process of the axis is detached in an equal degree from the surface of the atlas which rolls upon it. Now, it is from this last-mentioned change that the risk to the spinal cord chiefly arises. The anterior arch of the atlas cannot be separated from the odontoid process, except by the stretching and elongation of the transverse ligament which embraces the neck of the process, and of the accessory restraining ligaments. Accordingly, if the weight directed on the atlas should be suddenly increased while this *extension is in progress*,

and the ligaments are correspondingly weakened, the danger is imminent of the latter being ruptured. And should that event occur, there would be nothing to prevent the atlas, with the head as its freight, from sliding abruptly forward on the surface of the axis to its utmost reach. The consequence would be, that the portion of spinal cord contained within the ring of the atlas would be compressed by the posterior arch of that

FIG. 222.*



bone against the odontoid process in front, and be thoroughly crushed, and the patient's death would be instantaneous.

But, happily, these cases are not always so formidable. In not a few, the transverse and restraining ligaments undergo extensive stretching of their fibres without being ruptured.

* Atlo-axial disease. Figure from specimen in Middlesex Hospital Museum. *a, a.* Portion of occipital bone, in proper relation to atlas. *b, b.* Section of atlas. From destruction of the ligaments of the atlo-axial articulations, the atlas has slid forward, borne by the weight of the head, and has carried the odontoid process of the axis with it. *c.* Odontoid process of axis: loosened at its base by disease, it has been carried forward in connection with the atlas. *d, d.* Section of axis: its odontoid process detached: its articulations with the atlas, and with the body of third vertebra, and its body, all much diseased: the great projection backward of the spinous process in reference to the occiput and atlas, is a result of the gliding forward of the latter. Within its canal the spinal cord is seen bent at an obtuse angle: the consequence of the portion contained in the foramen magnum and atlas being carried forward, while that in the axis has been stationary.

Case: The patient, a butler, past the middle age, and corpulent, had for some

They continue, therefore, to retain a hold of the atlas; and to keep a check upon it, as it glides forward upon the axis. Besides, the connecting structures in general surrounding the diseased vertebræ are agglutinated by deposition and induration of coagulable lymph, and the shifting movement proceeds slowly. Thus displacement may take place to an extraordinary degree. In many cases the atlas will be transported to the front to such a distance, that less than one-half will remain resting on the axis, the anterior part protruding unsupported beyond its level. The posterior arch of the atlas will then cease to be in relation or coaptation to that of the axis; it may be transported so far forward, that it will even bisect the vertebral canal into equal parts. Hence about a moiety of the area of the canal appropriated for holding the spinal cord and its membranes may be absolutely cut off. It might have been conjectured that a loss of substance, and alteration of figure so great as this description implies, would have been altogether incompatible with the cord retaining its functions. But experience shows that if the morbid action be chronic, and the dislocation take place slowly and gradually, the disease may run its course without paraplegia ensuing, and the patient will recover with deformity of the neck alone. Again, it is found that in those formidable cases in which paraplegia both of the upper and lower extremities has occurred, the patients may regain the power over their limbs, and be restored to perfect health; the spine, of course, remaining distorted. The phenomena are to be accounted for according to the pathological principle already adverted to (p. 134). The diminution in the substance

time stiffness in the neck, with pains in the head and arms, supposed to be rheumatic. When seen by the writer, a large swelling extended from the upper third of the neck behind, to the highest part of the occiput, concealing the forms of all the bones which it covered, except the spinous process of the axis, which appeared unduly prominent. He could perform the nodding, but not the rotating motion, of the neck. The vertical axis of the head was advanced perceptibly too far forward, in comparison with the vertical axis of the neck; the effect apparently of the head having been transported to the front slightly beyond its proper plane in relation to the spine. The disease was seen to be atlo-axial; and he was removed to the hospital. Progressive paraplegia soon commenced, and it proceeded downward; first, one arm became weak and then paralysed; shortly afterwards, the other also became weak and then paralysed; and the lower extremities followed the same successive course. But before the legs were totally paralysed, he died; his death having been accelerated by a severe attack of bronchitis.

and the disfigurement of the cord generally have been sequent on a slow process of interstitial change; and it has been seen that deviations from the normal size and shape

FIG. 223.



The specimen from which the figure was taken was removed from the body of a man found dead in the streets. There was fracture at the base of the skull, and at a short distance from the foramen magnum; the boundaries of the fracture being seen in the figure. *a, a.* Portion of occipital bone, insulated by the fracture. *b.* Anterior tubercle, on anterior arch of atlas. *c.* Posterior tubercle, on posterior arch of axis. *d.* Odontoid process of axis. Its situation corresponds to the centre of the vertebral canal, at the level of the foramen magnum. *e.* Body of spinous process of axis. About three-fifths of the articulating surface of the atlas have been displaced forward on the corresponding surfaces of the axis, thereby accounting for the projection anteriorly of the front segment of the axis. *f.* Spinous process of axis. The posterior segment of the ring of the atlas is displaced forward, the canal of the axis nearly in its middle. The distance between the surface of the odontoid process, and anterior surface of the posterior arch of the atlas (the antero-posterior diameter of the vertebral canal), measured third to a quarter of an inch; that between the odontoid process and the anterior arch of the atlas, at the original sites of their articular surfaces, five-eighths of an inch. The space between these points is traversed by an irregularly cylindrical beam of bone which unites the two surfaces: the growth of bone is not visible in the drawing, but it had obviously been during the process of displacement of the atlas upon the axis, and at a slow rate at which it had taken place. For a more detailed descriptive specimen, accompanied with a drawing representing it in a different position, see *Med.-Chir. Trans.* vol. xxxi. p. 289.

the most surprisingly great kind, can be made in the spine, as well as the brain, without sacrifice of their contents, if wrought in that gradual manner.*

* In a specimen of consecutive dislocation of the atlas upon the axis, described by Mr. Paget, the displacement exceeded what has been stated in the text. Fully two-thirds of the atlas must have been entirely thrown



is deformity from caries in the dorsal vertebræ. The most common age at which it commences is near puberty; but infants nor adults are exempt.

symptoms.—A general swelling around the nape of the neck,

articulating surface of the axis. On measuring the part of the vertebral column available for holding the spinal cord, it was found that the distance was two lines from the odontoid process. and three lines. at the most. from the

DISEASE OF THE SPINE.

attended with stiffness, and pain in motion, are the earliest indications of the disease. The patient will prefer the recumbent position to being erect; keeping his head sunk in a soft pillow. When upright he will steady his head by placing a hand on each side; and if asked to rotate it, he will stiffen his neck, and turn the whole body round. The nodding motions he will be able to perform. A sudden jolt, or a tap with the hand on the crown of the head, will give pain. As the swelling of the upper part of the neck may probably be mistaken at first for enlargement of the lymphatic glands, or deeply-seated abscess arising from some other cause, the most certain mode of deciding the question as to the real nature of the disease, is to observe accurately the relative position of the spinous process of the axis to the occiput. Notwithstanding its large size, that process is not readily felt in the normal condition; it lies sunk in a hollow, arched over by the ligamentum nuchæ. But in atlo-axial disease the spinous process becomes distinct, not only to the touch, but to the eye. That change, however, is not consequent on the axis itself protruding; for this bone is stationary; the spinous process appears to project merely because the atlas, carrying the skull upon it, glide away to the front, thereby leaving it more exposed and defined. In conformity with this explanation, it will be perceived that the head droops, as it were, upon the neck; that the chin and face generally are advanced preternaturally forward in comparison with the throat, while the back of the head is deficient in rotundity and fulness in reference to the line of the spine. It may happen that one side of the atlas will move forward more

figure. The disease had apparently commenced four months previously; but it had caused so little pain or inconvenience, that she had not been obliged to give up her work during its progress. She complained of pain darting from the swelling in the neck upward to the crown of the head, and downward over the shoulders, and along the arms. She had also slight incontinence of urine. The most prominent part of the swelling in the neck was formed by the spinous process of the axis; above which, and between it and the occiput a distinct hollow could be felt. Chiefly by rest and counter-irritants, she improved so much that in six weeks she was discharged. But three weeks afterwards, she was readmitted on account of an aggravation of the former symptoms, and there being a weakness of the lower extremities, which caused them to sink under her when she attempted to walk. In a month she was sufficiently recovered to be sent to a convalescent hospital at the seaside. She subsequently called various times at the hospital, to show that she had regained her strength completely. A cast from the neck is in the Museum of the Middlesex Hospital.

rs and clavicles, perhaps give the first alarm. The latter
ons are sometimes prolonged to the finger-ends, accom-
with slight muscular weakness. Such symptoms alone
be explained by supposing that the roots and trunks of
ves given off from the cord in the neighbourhood of the
, had become affected by the process of displacement and
action external to the spine, without the medulla itself
involved. But prudence will not allow that view to be
ch relied upon. Should there be added any symptom of
wer extremities becoming paralysed—as a tendency to
walking, or a sense of pricking or of numbness in the feet
conclusion will at once be drawn that the morbid action
etrated to the medulla. In certain cases the paralysis
upper and lower extremities will be but imperfectly pro-
ed; and it will pass off. In a single case seen by the
, the muscles became affected with tonic spasm; the whole
was so rigid that on turning the patient in bed she rolled
be stiffness of a corpse having rigor mortis; that condition
for a month; it gave place gradually to ordinary paralysis,
which she eventually recovered. When the paraplegia
plete, the patient is in a singularly helpless state, being
ted of voluntary power and sensation both in the upper
lower extremities at once. Yet, notwithstanding the
ently hopeless condition of patients in these cases, ex-
nce proves that they frequently recover. The prognosis
be, of course, more favourable in young persons than in
above the middle period of life. With his colleague

DISEASE OF THE SPINE.

Treatment.—In disease of the atlo-axial articulation utmost danger would be incurred if the patient fell forward his head; in its stretched and attenuated condition the transverse ligament would readily give way, the atlas thus be free to slide forward and compress the spinal cord against the posterior arch. Special care is therefore required to keep the head and neck of the patient in a state of perfect rest. Accordingly, constant uninterrupted confinement in the bent position is necessary. And the patient's head is to be laid on cushions, which will prop it equally on all sides. For that object, the india-rubber pillow, stuffed to a proper degree of firmness with grain, so that it may be pressed into a hollow which shall fit the back of the head and nape of the neck, is to be used. Or the sand-bags recommended by Mr. Hilton will be more easily procured. A band ought also to be passed over the forehead, from the sides of the pillow, to restrain motion. In the event of the patient being restless and liable to lift the head inadvertently, as during or sleep, it is a useful precaution to have a shield made of either in gutta-percha, or prepared leather, to the back of the neck; this ought to take in the shoulders below, and reach the vertex above, so that it may be secured by bands both to the shoulders and head, to arrest movement.

Necrosis of atlas.—Before quitting the subject of disease of the upper vertebræ, a rare affection of the atlas may be noticed. This consists in necrosis and exfoliation of the anterior arch of the bone which corresponds, in the lower vertebræ, to the body. The disease is connected with syphilis of the back of the pharynx; and it has been commonly observed in syphilitic patients. In some recorded cases the whole thickness of the bone has come away, as shown in the accompanying illustration. The articular surface of the odontoid process of the axis was exposed and visible. From the patient's surviving, it cannot be doubted that the insertions of the transverse ligament had been preserved entire; and that it had retained its hold of the p

Pain and the Therapeutic Influence of Rest,' by Mr. Hilton, Lecturer on the Diseases of the Spine, contains a very remarkable case of disease of the superior cervical vertebræ, with protrusion of the atlas and axis, through the occipital bone, into the foramen magnum. This case has been communicated by Dr. Lochee and Mr. C. H. Moore to the Medical Society of London. See number for November 23, 1867. The preparation is in the Museum of the Middlesex Hospital.

oughing ulcer in the posterior fauces, with occasional attacks of difficult deglutition and breathing, have been the most prominent symptoms. When the fragment has been picked away, the sore has healed, and the patient recovered. It need scarcely be said, that a patient who has such a threatening disease, in a condition of so much danger to life, is in a critical condition as the dead portion of bone remains as a source of irritation, and that the greatest caution is demanded to protect him from sudden bending forward of the head.*

ALEXANDER SHAW.

*See case narrated by Mr. Keate in *Medical Gazette*, vol. xvi., 1835. 'A Case of Exfoliation of the Anterior Arch of the Atlas,' by Mr. R. Wade, *L-Chir. Trans.* vol. xxxii. p. 65. Lectures by Mr. Hilton, op. cit. p. 99.

DISEASES AND INJURIES OF NERVES

PART I.

NERVE-LESIONS AND THEIR MORE IMMEDIATE EFFECTS.

DISORDERS OF NERVES.

NERVES are subject to many disorders and injuries, most of the other parts of the body. They are liable to inflammation, suppuration, ulceration, and gangrene; to atrophy and enlargement; to softening and induration; to the formation of cysts, fibrous or neuromatous tumours, and cancerous growths along their course; to compression or stretching; to contusions, lacerations, punctures, and partial or complete division.

Inflammation of Nerves.

Inflammation of nerves is less common than that of other soft tissues. It attacks in preference the sanguiferous, robust, and adults rather than children, in whom neuritis is rare. It may be either acute or chronic, and either idiopathic or traumatic.

Acute neuritis is generally preceded by the usual prodromata of inflammation. Its characteristic symptoms consist in tearing, darting, or lancinating pain along the course of a nerve trunk or its branches, attended with a sensation of tingling, formication, or numbness, which are sometimes followed by partial or complete paralysis of the muscles supplied by the portion of the nerve which is below the seat of inflammation. The pain never becomes suddenly severe or ceases suddenly, intermits, as in non-inflammatory neuralgia; for although frequently remits or abates, it is a continuous pain which gradually increases in intensity and gradually subsides. It is always aggravated by pressure on the inflamed portion of

erve, by contraction of the muscles to which it distributes branches, or even by the slightest touch along its peripheral course; while, on the other hand, it is relieved by firm pressure above the seat of inflammation. Sometimes the nerve is swollen at the part affected, resembling a cord stretched along the surface of the limb. Usually the movements of the limb are much impaired, and there is often a convulsive agitation of its muscles. The pulse is frequently strong and quick, the temperature of the skin over the inflamed part is much increased, and there is generally a variable degree of sympathetic fever, in proportion to the size of the affected nerve and the intensity of the inflammation.

The seat of inflammation in neuritis is the neurilemma and the connective tissue between the bundles of fibres; and whatever changes the fibres may undergo are the consequences of deranged nutrition. When the inflammation has been of moderate intensity and not of long duration, the nerve has been found to be more or less altered in colour, varying from a pale rose to a deep red, according to the greater or less amount of vascularity in the neurilemma and interfibrillar connective tissue. For the same reason, also, there is a variable increase in the diameter of the nerve, its component bundles being separated from each other more widely than usual; but the fibres retain their natural structure, or at least do not appear to have suffered any remarkable change.

The inflammation may either terminate in resolution or reach its greatest intensity. In the latter case, the nerve is extremely vascular and of the deepest or violet red throughout its entire thickness. Its volume is still further augmented, sometimes to twice or even three times its natural diameter. This is the result partly of the increased hyperæmia, and partly of the inflammatory products that are poured out. These products consist of sero-fibrinous or sero-sanguinolent fluids; of pure blood, or blood mixed with pus, infiltrated into the neurilemma and between the bundles of the nerve-fibres, which undergo disintegration or degeneration, in consequence of impaired nutrition. Sometimes the nerve is softened and easily torn, or even reduced to a pulp; at other times it is indurated and dense from coagulation and hardening of fibrinous exudation. In some cases it is interspersed with minute ecchymoses, or with hard grains along its inflamed course; in other cases it is covered with small abscesses, or enveloped in pus or puro-

sanguinolent fluids, or fibrinous exudation; or it contains adhesions to contiguous parts.

Chronic neuritis may either result from incomplete resolution of acute inflammation, or be the continuation of a milder. In either case it is a frequent cause of certain kinds of neuralgia, of neuroma, and painful subcutaneous tubercle. Its morbid anatomy consists of an increased vascularity of the affected nerve; sometimes of a varicose state of its blood-vessels, of a thickening and induration of the neurilemma, in consequence of coagulable exudations. In a variable degree the nerve assumes a somewhat slate colour, loses its characteristic opacity, and when examined under the microscope, the nerve fibres are found to have fallen, to a greater or lesser extent, into a state of disintegration.

Idiopathic neuritis, although much less common than the former, is, I believe, more frequent than it is generally considered to be. Its chief exciting causes are exposure to cold, to wet, and particularly to the combination of both; the suppression of profuse perspiration; the arrest of hæmorrhoidal discharges; excessive bodily fatigue; strumous or tubercular affection of the system; sometimes visceral inflammations and suppurations.

Martinet relates the interesting case of a deserter—an athletic young man who after an obstinate chase through a wood, was captured by the gendarmes, covered with sweat, and in a state of wild excitement. On the third day after his capture, he was unable to sustain himself on his legs, and experienced excruciating pain along the back of his thighs, in the course of the sciatic nerves. Soon after, the patient was attacked with pneumonia, and died on the eighth day of his illness. On post-mortem examination, both sciatic nerves were found to be as thick as the index finger, and hard and brittle. All their component bundles were separated from each other by a sanguinolent fluid, and were penetrated by a multitude of blood-vessels, imparted to the nerve a deep-red colour.* Gendrin mentions the case of a similar case of a man who walked from Lyons to Paris. During the last days of his journey, he experienced a dull pain in the course of the sciatic nerve. The right leg became entirely useless, and the patient died after a fortnight. On examination, the right sciatic nerve was found to be enlarged, of a rose colour at one part, and of a deep red at another. It was infiltrated with serous fluid, and three times its natural diameter.† Martinet has mentioned nine other cases of idiopathic inflammation of nerves, the most important of which I will very briefly describe.

Case 1: Violent and continuous pain in the course of the left

* L. Martinet, 'Mémoire sur l'Inflammation des Nerfs,' *Revue médicale*, tome 2^{me}, obs. v.

† Gendrin, *Histoire anatomique des Inflammations*, tome ii. pp. 143

formed on the sacrum, and death ensued a few days later. On post-mortem examination, the spinal cord was found healthy, but the median nerve on the left side, at its separation from the other branches of the brachial plexus, was of a deep red colour *throughout its entire thickness*, for about two inches in length. A nearly similar condition was found in the anterior branch of the seventh cervical nerve which contributes to form the median. The sciatic nerve, enveloped in a large quantity of gangrenous areolar tissue, was red, at its superior part, and for a length of about two inches and a half, of a brown colour through its whole thickness, but without any alteration in its volume or consistence.*

In this interesting case, we learn that nerves situated at a distance from the seat of inflammation, may consecutively and independently become affected by idiopathic neuritis.

Earle,† Sir Charles Bell,‡ Barensprung,§ and others, have recorded several cases of idiopathic inflammation of nerves. Barensprung's case was of a scrofulous child, one year old, in which both roots of the sixth, seventh, eighth and ninth intercostal nerves were found enlarged and red. Martinet's eighth case was also a child, twelve years old, who had for some time shown the characteristic symptoms of phthisis, and in whom there was violent inflammation of the left sciatic nerve. The child died in about seven weeks in a complete state of marasmus. On examination, numerous tubercles were found in both lungs, the greater number in a state of suppuration. The left sciatic nerve was considerably increased in volume, but the whole of its upper third was filled with sanious pus, which separated it from the surrounding muscles. A kind of pus was found between its bundles. This case and the foregoing one are particularly interesting from the fact of their occurring in children, in whom neuritis is rare.

traumatic neuritis is much more common than idiopathic. The term implies, it is the result of some external injury, such as wounds, contusions, fractures, dislocations, surgical operations, &c. According to my own observations, it is less fatal than the idiopathic form.

purgatives; tartarated antimony, opiates, belladonna, and the preparations of aconite. When there is reason to believe that the inflammation is of a gouty, rheumatic, or syphilitic character, colchicum, iodide of potassium, or perhaps mercury, should be employed. In cases where it is complicated with serious internal disease, or associated with a low cachectic condition, the treatment must be modified accordingly.

In *chronic* neuritis repeated blisters along the course of the nerve sometimes afford great relief. Anodyne fomentations, and especially liniments of aconite, belladonna, chloroform, and opium are frequently of signal service. At the same time attention must be directed to the state of the general health. Vegetable and mineral tonics, iodide of potassium, and mild aperients are sometimes indispensable.

Ulceration of nerves, as a primary affection, appears to be unknown; but nerves in the neighbourhood of ulcers are liable to become involved in the morbid process, and then they are generally the cause of intense and protracted suffering.* In such cases, the surrounding soft parts are often enlarged, the skin increases in thickness, the muscles and tendons share in the ulceration, and even the bones increase in dimensions. On either side of their ulcerated portion the nerves are generally thickened, sometimes to nearly double their natural diameter.

When these ulcers are exceedingly painful, Mr. Swan recommends the use of ointment made with powdered opium, or lotions made by mixing well-powdered opium with water, or lime-water. These should be applied on lint, and then a folded cloth moistened with water, or laudanum and water, should be placed over the surrounding skin; at the same time the digestive organs should be regulated. When other remedies have failed and the disease is confined to the soft parts, excision of a portion of the nerve will be advisable, but if the bone be affected by caries or necrosis, amputation may be necessary. Mr. Swan recommends, that in excising a portion of the nerve, the operation should be performed as far as possible from the ulcer because there is then a much greater probability that the

* Nerves sometimes ulcerate in consequence of the pressure of tumours, &c. A striking case of ulceration of the sciatic nerve, produced by pressure of an aneurism, which extended backward from the right groin, is recorded by Morgagni, *De causis et sed. Morb.* epist. 50, p. 11.

external wound will heal by the first intention, and that consequently the cut ends of the nerve will escape inflammation and ulceration. It is also advisable to divide the nerve as near the upper part of the wound as possible, as the end of the nerve will thus retract from the wound, and consequently be less liable to become inflamed, should the external parts have assumed this disposition.*

Tumours of Nerves.

Of all the morbid growths that are found in connection with nerves, tumours are by far the most frequent. They vary considerably both in size and structure, but are commonly a source of acute pain and of the most distressing nervous affections. The smallest kind are known as *painful subcutaneous tubercles*; the larger kind, as *neuromata*. The former are either more or less spherical, or oval, or fusiform, and sometimes rather flattened. Their colour is generally white, but occasionally there is a brownish tint on their surface or in their interior. They are always firm, sometimes hard, and have a fibrous or fibro-cartilaginous structure. Their size varies from that of a millet-seed to that of a pea or a grain of oat. They are situated in the subcutaneous areolar tissue, and are generally more or less embedded between the fibres of the nerves, which are separated and stretched over them. Sometimes, also, they have a close connection with a subcutaneous vein in the neighbourhood of the nerve. They are generally solitary, and mostly confined to the upper or lower extremities, especially the former. Women are more subject to them than men; of eighteen cases collected by Wood, fourteen were women.

It is not easy to determine the cause or mode of origin of these tubercles. They appear in many instances to be the result of chronic inflammation, and have been found to follow certain injuries of nerves, especially contusions. When once deposited, they seem to grow with some rapidity, but may then remain stationary for a number of years.

This small subcutaneous tubercle is a source of the most severe pains, which dart like electric shocks along the course of the nerve, sometimes in both a peripheral and central direction. The pains recur very irregularly, and last, at each attack, from ten

* Swan, *On Diseases of Nerves*, &c., p. 83, 2nd ed.

minutes to two hours or more. They begin gradually, increase in intensity, and gradually decrease, leaving the tubercle and its surrounding parts more or less tender to the touch. They are frequently excited or aggravated by changes in the weather from hot to cold, or the reverse, by storms, &c., and sometimes by mental emotions. The paroxysms may be repeated several times a day, or there may be remissions of days or even weeks. In many instances the tubercle is not tender to the touch except during the attacks, which are much aggravated by pressure.

In all cases of obstinate neuralgia, especially of the extremities, a careful search should be made for the possible existence of tubercles.

Valleix relates the case of a lady who suffered for fourteen years from acute pain in one of her knees, which no kind of treatment succeeded in relieving, until M. Cabaret, on carefully examining the part, discovered a small tubercle about the size of an oat attached to a cutaneous nerve.* After its removal the pain entirely ceased. Marjolin furnished Descot with notes of a remarkable case of painful subcutaneous tubercles of the scrotum. A robust man, thirty years of age, had complained, for more than a year, of sharp, lancinating and intermitting pains in the right side of the scrotum and inner side of the corresponding thigh. On examination, he found, in the areolar tissue immediately beneath the skin of the scrotum, several small, smooth, lenticular, movable, hard, and isolated bodies. When compressed between the fingers, they caused the same kind of pains as the patient was accustomed to feel. A small incision of the skin was made over each of them, and they were easily extracted by compressing the edges of the incision. The operation was followed by a complete cessation of the pains.†

Neuroma.—Neuromatous tumours, although they differ from each other considerably in size, are larger than the subcutaneous tubercles, and are more variable in structure. They have been found of every intermediate magnitude between a small grain of wheat or oat and a large melon.

Dr. Smith of Dublin, found on the left sciatic nerve of a man who died in the Richmond Hospital, a neuromatous tumour which measured ten inches in its transverse, and eleven inches in its long diameter. The fibres of the nerve were separated from each other, and studded with small oblong tumours.‡

Neuromata are either round, oblong, oval, or fusiform. They are situated either between the neurilemma and the nerves, or in the connective tissue between bundles of the nerves. In the former case, the bundles, although pressed together and flattened, are not separated from each other; while in the latter case they

* Valleix, *Traité des Névralgies*, p. 563.

† Descot, *Dissertation sur les Affections locales des Nerfs*, p. 245.

‡ *Treatise on the Pathology, Diagnosis, and Treatment of Neuroma*. 1849. Plate ii.

and wavy fibrous tissue, in connection with a variable number of nuclei and small cells. In an interesting case of paralysis of the third cerebral nerve with slow hemiplegia on the same side, from supposed syphilitic disease of the brain, under the care of Dr. Hughlings Jackson, a neuroma of the third nerve, among other lesions, was discovered after death. The description which I gave of the structure of this tumour is applicable generally to neuromata of this kind. 'On examination under the microscope, it was found to be everywhere composed of a multitude of nuclei, with some nucleated cells, and an abundance of fibrous tissue. Some of the nuclei were more or less oval, but the majority were round, and measured about the $\frac{1}{32000}$ th of an inch in diameter. Amongst these were also bodies of the same nature, but of all degrees of smaller size, and either elongated, angular, or in the form of granules. The nucleated cells were of very small size, and very granular in structure. A few of them were sparingly scattered through the tissue, but, for the most part, they were collected here and there into small, oval, or circular groups. Between these elementary bodies there was an abundance of ordinary fibrous tissue. The fibres were generally more or less tortuous, but straightened, and in their course each divided again and again into finer filaments, which were connected with the cells and with all the other elementary bodies; the granules and angular or elongated nuclei filling up the interspaces between those that were oval or round; and thus contributing to form with these and the fibres, a close, uninterrupted, and

Occasionally the tumour has been found to be lobular; the lobules having been connected with branches of the nerve on which the mass was seated, as in a case that will presently be described.

Unlike the subcutaneous tubercles, neuromata grow somewhat rapidly, and steadily enlarge, although they are liable to increase and diminish alternately in size. Their number is exceedingly variable. Sometimes they are solitary, one only being found on a nerve-trunk; sometimes several form on the same trunk, or on the same branches of a nerve; and in rarer instances they form on all or nearly all the spinal nerves in the same individual. Dr. Hughes Bennett states that in the Museum of the Richmond Hospital, Dublin, he examined a most remarkable series of preparations, taken from two individuals in whom almost every nerve of the body presented knotty swellings.* In the case already quoted from Dr. Smith of Dublin, in reference to the enormous neuroma on the sciatic nerve, there were also found on the anterior and lateral regions of the right *lower* extremity, no less than 150 tumours, and about 100 on the nerves of the right *upper* extremity.† In another case described and figured by Dr. Smith, the abdomen and lower part of the thorax were covered with a vast multitude of small superficial neuromatous tumours, which might easily have been confounded with malignant tubercle.‡

Neuroma may be either traumatic or idiopathic. In the former variety the tumours are caused by wounds, blows, pressure, or by foreign bodies in contact with nerves, or lodged within them. They are generally solitary and solid in structure.

The idiopathic variety appears to be the result of several different causes, viz. chronic inflammation, rheumatism, gout, syphilis, or some other peculiar constitutional vice. Some pathologists believe that the disease is of the nature of scirrhus, but this opinion is not supported by facts. As in traumatic neuroma, there is generally only one tumour, but in some cases, as already shown, the number may be immensely multiplied.

When the tumours are very numerous, there is little or no

* Dr. Hughes Bennett, *Principles and Practice of Medicine*, 4th edition, p. 195.

† *Treatise*, plate ii. This is apparently one of the cases mentioned by Dr. H. Bennett.

‡ *Treatise*, p. 17, plate vi.

pain; but a solitary neuroma, whether traumatic or idiopathic, is a source of the most violent agony, which shoots along the nerve like sudden electric shocks; the pain is sometimes continuous, but is always aggravated by pressing or touching the tumour, which seems, during the paroxysm, to be in a state of erythism. Mental emotions and atmospheric changes are also as unfrequently the exciting causes of an attack. After the disease has existed for a length of time, the pain which at first occurred only in paroxysms may become continuous. In idiopathic neuroma, it is more frequently limited to parts below the tumour, while in the traumatic form it may shoot both upwards and downwards. In a few cases on record, the patients were subject to epileptic convulsions, and in two or three they were cured by removing the tumour.

John Hunter removed from the musculo-cutaneous nerve, one about the size of a hen's egg, over which the fibres of the nerve were spread out. Sir E. Home removed from the axilla of a Frenchman, a hard oval tumour about three inches and a half long, and two inches broad. It was enclosed between fibres of one of the large nerves of the axillary plexus, and was the cause of terrible suffering. Louis removed from the fore-arm of a young lady, a tumour which made its appearance fourteen years back, as a small hard substance midway between the bend of the arm and the hand. At the time of the operation it had so much increased in size, that it extended over the whole front of the fore-arm, and was about six inches in diameter in the middle. The pain increased in severity in proportion to its growth. On examination after its removal, it was found to consist of several lobes, which were connected by pedicles to branches of the radial nerves, like grapes to their stalks.*

The exciting cause and mode of origin of these tumours are often exceedingly obscure; but not unfrequently they may be traced to external injuries, particularly to blows and contusions. Cases of this description are recorded by Sir C. Bell,† Dupuytren,‡ and others.

When the neuroma is superficial, the diagnosis is by no means difficult. In general the tumour is movable laterally, but cannot be moved in the direction of the nerve without causing great pain. In most cases the skin which covers it remains unaltered, but when the tumour is very large the cutaneous vessels, especially the veins, are sometimes dilated.

The only effectual remedy for these morbid growths is the knife. The operation may consist either of excision of the

* Delaroche et Petit-Radel, *Encl. méd.*, art. 'Tumeur.'

† *Operative Surgery*, vol. ii. p. 161.

‡ *Apud Valleix*, loc. cit. p. 328.

tumour, with or without a portion of the nerve, or of amputation of the affected limb. When the neuroma is very intimately connected with the nerve, and there is reason to believe that it cannot be extricated from the nerve without great disturbance or injury, it is better to remove a portion of the nerve with it. With regard to the sciatic nerve, which supplies the whole of the limb, excision of a portion of its length would be attended with serious consequences, and therefore unless the tumour be confined to that portion of it which forms the posterior tibial nerve, and the peroneal nerve can be saved, it would probably be advisable to amputate the limb.

Tubercle. — Dupuytren and Cruveilhier have spoken of tubercle in nerves, but there appears to be no evidence to prove that it is ever found in this situation as a primary disease; although nerves are sometimes destroyed by the softening of tubercles in which they are involved.

Cancer, unlike tubercle, is frequently met with in nerves, both as a primary and secondary affection. The form of the cancer is in general either the white medullary, or melanosis. It may occur at any part in the course of a nerve, but it is mostly found near its peripheral extremities. It generally grows to a considerable size, as in medullary cancer of the retina. As a secondary disease it is communicated to the nerve by a contiguous cancerous growth.

Dubois, on several occasions, found cancerous growths in the substance of nerves of both the arms and legs.* Moutard-Martin saw one on the median nerve, and its removal was followed by cancer of the brain. Dupuytren removed one from the posterior tibial nerve, which, moreover, presented a series of nodulations, like grape-stones. It was the cause of acute lancinating pains. In another case he found the trifacial nerve transformed into encephaloid substance.† Cancerous growths have been found, also, on separate branches of the trifacial, on the spheno-palatine ganglion, on the optic nerves, and on the phrenic nerves.‡

Hypertrophy of nerves, in the proper sense of the term, has no actual existence as a *morbid* condition. It is true, as we have already seen, that the inflamed portion of a nerve fre-

* Viel-Hautmesnil, *Considérations générales sur le Cancer*. Paris, 1807.

† Breschet, *Dictionnaire de Médecine*, art. 'Cancer.'

‡ Descot, *Sur les Affections locales des Nerfs*.

the result of several different causes, viz., chronic
nation, stretching, compression, and even severe contu-
Of these the most common is compression caused by
aneurisms, enlarged glands, &c. Nerves of special
often waste after loss or wasting of the central organs to
they belong, as the optic nerve after wasting of the
of the eye. In some instances the nerve-fibres only are
ted, and nothing may be left but the neurilemma; in
the neurilemma itself is involved in a greater or less
. During the process of atrophy, the nerve-fibres be-
even; their white substance is broken, at intervals, and
d from their axis-cylinders in masses, which again break
smaller fragments that are mingled with fatty particles.

INJURIES OF NERVES.

lusions of nerves are frequently followed very insidiously
most serious consequences. If the blow be not very
, the principal change produced in the nerve is the extra-
on of blood and other fluids into the connective tissue
in its bundles. There are pain, numbness, tingling, or
ation in the parts supplied by its branches, followed, in a
r or less degree, by partial and temporary paralysis, and
y of the muscles. If the contusion be very violent, the
of the nerve may be lacerated or crushed, and then the
y and paralysis are of a much more serious character, and
allow the injury either immediately or not till after an

of a great variety of causes. Of these, dislocation of the joints, and the attempts made at reduction are among the most common. The effects are more serious in proportion to the length of time before the dislocation is reduced. Sensibility suffers much less than the power of motion. If compression of a nerve be severe and long continued, it may give rise to incurable paralysis and wasting of muscles. When the injury is not so severe, it causes pain of a more or less acute, more or less constant, character, attended sometimes with feeling of numbness or tingling. Nerves may be compressed also, by aneurisms; by tumours of different kinds formed either upon them, or in their immediate neighbourhood; by enlarged glands or bursæ; by abscesses; by cancerous growths; by distension of the rectum and sigmoid flexure of the colon; hæmorrhoids and other causes.

Sir B. Brodie relates the interesting case of a man who had severe pain on the inside of his knee. No marks of disease could be detected in the knee; but in the thigh there was an aneurism of the femoral artery. Sir E. Jenner applied a ligature round the artery in the upper part of the thigh; the tumour immediately ceased to pulsate, and the pain in the knee ceased also.* A remarkable case related by Morgagni, of aneurism of the right groin pressing on the sciatic nerve and producing ulceration with intense pain, has been already quoted. Scarpa also describes an interesting case of aneurism of the abdominal aorta, causing disorganisation of the lumbar nerves and injury to the anterior crural and obturator nerves.†

Tumours in the immediate neighbourhood of nerves frequently, by the pressure which they exert, the cause of much suffering as those which are formed upon them.

Sir B. Brodie mentions the case of a gentleman who suffered severe increasing pain in the left leg, from the foot to the knee, in the course of the peroneal nerve. As the limb presented no morbid appearance, the disease was diagnosed by the name of neuralgia. After a considerable time the patient died of dropsy, and on opening the abdomen, a large solid tumour was found attached to the left side of the lumbar vertebræ. It was evident that this tumour had pressed on the origin of the sciatic nerve.‡ Mr. Travers published an interesting case of compression of nerves by a medullary tumour in the brain, and Sir William Lawrence mentions the case of a tumour in the forearm of a gentleman, situated over the course of the ulnar nerve, and causing excruciating pains, like electric shocks, upwards and downwards, in the direction of

* *Works*, vol. iii. p. 135.

† *Treatise on Aneurism*, translated by Wishart, p. 99.

‡ *Works*, vol. iii. p. 135.

§ *Med.-Chir. Trans.* vol. xvii. p. 389.

of the nerve.

enlarged glands are not unfrequently the cause of compression and distension of nerves; and there is no nerve so frequently in this way as the facial. Several cases of this kind were published by Sir Charles Bell.†

Woddie mentions a case in which 'two lymphatic glands, enlarged to the size of large walnuts, were found situated beneath the skin, on the anterior surface of the thigh.' A considerable branch of the lumbar nerves lay over each gland, being thus kept stretched like strings of a violin over its surface, giving rise to violent pain and convulsive movements of the leg.§ The author mentions a case of severe neuralgia in the foot, caused by the weight and pressure of internal hæmorrhoids, after each evacuation of the bowels.|| Romberg calls particular attention to neuralgia of the sciatic nerve caused by crural hernia, and gives a long but interesting description of it.¶

Nerves of a very acute kind are more often excited by small local growths in nerves than by large tumours over which the nerve-fibres are stretched; for nerves may be stretched to a considerable extent without exciting any painful sensation, but they become inflamed, or their nutrition is impaired, when the slightest touch causes acute pain. It therefore seems highly probable that the painful effects of compression and distension, however means they may be produced, are in many instances the result of inflammation or some peculiar kind of irritation which has been excited. When nerves are bruised or otherwise injured at their exit from a foramen or bony canal, they are more liable to cause suffering, because in the event of inflammation following, they are sure to be compressed.

them. It is somewhat surprising, that, except in par are vitally important, laceration of nerves is not follow serious consequences so often as might be expected.

Béclard has related many cases of this description.* Flaubert m case in which the last four nerves of the brachial plexus were torn : spinal marrow, by violent extension in attempting to reduce a dialo the shoulder-joint. The patient died at the end of eighteen days.† George's Hospital Museum,‡ there is a preparation showing the ant of one of the cervical nerves torn from the spinal cord, by dislocati vertebræ in a heavy fall.

Ligature of nerves.—It sometimes happens that in t artery a nerve is accidentally included in the ligatu certain cases this accident has been followed by severe lo constitutional disturbance, and even by fatal effects. instances of tetanus resulting from the same cause, after tation, have been recorded by several writers. When referred to the amputated limb along the distributi particular nerve, there is reason to believe that this ne been included in a ligature; and if on pulling at the the pain experienced be aggravated, there can be no lon doubt as to its cause. Under these circumstances, if or any other alarming constitutional disturbance be thr the ligature should be immediately removed.

Hennen gives a most interesting account of the effects of ligatures in the case of a general officer who suffered amputation of an arm, d stroyed by a bullet in action. After the occurrence of fever and sloughing, an attempt at clearing the ligatures was attended with excruciating pain. 'He has,' says Dr. Hennen, 'frequently, after th of dressing was over, with great accuracy pointed out on my arm th the internal cutaneous nerve, as the site of his ideal pain; often he ha that of the external; and on one occasion, I, with utter astonish the general neurology of my arm and fingers traced by him. Once ever know him to refer his pain to the sensorium itself. On the from using an artery forceps to the ligature, on which the slide m stiffly, I exerted a greater force than I intended. He convulsiv hand to his head, expressed a sense of exquisite pain in his brain, tears dropped from his eyes, a paralytic contraction momentarily mouth, an universal paleness spread over the uncovered part of his b uttered a piercing cry, exclaiming that the agony of his head and insufferable. The state of collapse was so great, that I was obliged

* Béclard, apud Descot, *Affections locales des Nerfs*, p. 41.

† Flaubert, 'Mémoire sur plusieurs cas de Luxation, etc.;' *d' Anat. et de Phys. pathologiques*, 1827, vol. iii. part i. p. 102.

‡ Series viii. No. 131.

affected her health, that she became extremely nervous. 'There was a constant state of convulsive action of the muscles of the stump. Everything that could be done to improve her health and relieve pain was fairly tried for several months, without the least good effect. She was hysterical, and the paroxysms were frequent.' After the stump had healed, a pulsation almost aneurismal could be felt in the situation of the ulnar and radial arteries. Having seen, in several cases, symptoms similar to those caused by the extremities of the nerves becoming ganglionic, and a second operation rendering requisite, Mr. Langstaff removed the arm above the elbow-joint, and proceeded to securing the arteries, drew out each nerve to the extent of half an inch from the surface of the stump, and cut through them to prevent their interrupting the progress of cicatrisation of the integumental parts. The patient was relieved of all her painful sensations, a good stump was made, and her health improved. On examining the amputated part, the median, radial, and ulnar nerves were found remarkably large. The extremities of the two last were greatly increased by deposition of organised lymph.

Sometimes, after the complete cicatrisation of the external wound by which a nerve has been divided, the cicatrix becomes inflamed, and the nerve participating in the inflammation may give rise to the most acute pain, particularly when the part in which it belongs is put in motion. In other cases one or both ends of the divided nerve may be involved in the cicatrix and be the cause of the most severe suffering. The position of the cicatrix is sometimes important, as when it happens to cross at a right angle the tract of a large nerve which is so related to it as in certain positions of the limb to be pressed against, thus causing pain and even loss of motion.

Incomplete division of nerves is, in general, productive of more severe consequences than when the division is complete. If nearly the whole thickness of the nerve be divided, the two portions will retract and put the undivided portion on a stretch; but except in peculiar constitutions, under unfavourable conditions, or when the nerve becomes inflamed, this tension does not in general appear to give rise to serious results. Astley Cooper, in removing a tumour, cut away two-thirds of the thickness of the median nerve. Tingling of the fingers, and some partial numbness, ensued, but nothing more.* But partial division of nerves occurring in persons of a peculiarly nervous and irritable, or hysterical constitution, is frequently a cause of the most distressing local and general symptoms.

Punctures made with different kinds of pointed instruments are the most common instances of this species of injury

* *Lectures by Tyrrell*, vol. iii. p. 171.

res. The first symptom which ensues is generally acute pain in the injured part, coming on most frequently at the time of the accident, and darting along the course of the nerves either towards their peripheral extremities or in the opposite direction towards the spinal cord and brain. It is sometimes periodical in its attacks, and attended with redness, tingling, or more or less numbness and swelling of the part. Contractions of the limb, violent spasms or tremors of the muscles, trismus, and even epileptiform convulsions, are not uncommon, and in females hysterical excitement is a frequent consequence. After a time, if the symptoms continue severe, the spirits become depressed, there is more or less prostration of strength, and, in some cases, delirium and coma supervene. The pain is often reflected on other nerves with which the injured nerve is connected only through the nervous centres, and this sympathetic influence is sometimes experienced to a surprising and terrible extent.

Dr. Wardrop has related the case of a woman who pricked the forefinger of her right hand, near the point, with a gooseberry thorn. The wound was immediately followed by great pain, swelling and redness, which in a few days extended along the forefinger and adjoining phalanx of the middle finger. At the end of three months the pain and swelling disappeared, except from the first phalanges of the wounded finger, which remained extremely painful. The patient's general health suffered considerably, and she had severe nervous exsiccations two or three times a day, during which the pain extended along the finger to the back of the hand, and between the two bones of the fore-arm, passed through the elbow-joint, and up the back of the arm to the neck and head, producing a sensation at the roots of the hairs as if they had become erect. To these symptoms succeeded dimness of sight, and subsequently the pain extended to the stomach, producing nausea and vomiting. She had constantly the feeling of a lump in her stomach, and vomited after taking food or drink. At the end of seven months, three incisions were made at the end of the finger, but they gave her no relief. She was afterwards completely cured by amputation of the finger.*

A case in which the most violent effects arose from the puncture of a nerve during the operation of bleeding, was related to Mr. Swan by Dr. Wilson of Lantham. The patient, who was a woman, had been bled by a gardener. On the second day she fell into strong convulsions, and some time after became comatose. The median vein in which she was bled had not healed, and was somewhat inflamed. An incision was made just above the orifice of the vein, when the patient immediately cried out, 'I am well; I can stir my arm.'

Even fatal effects occasionally follow the puncture of a nerve during the operation of bleeding. Such a case is recorded by Bonetus.†

* *Medico-Chirurg. Trans.* vol. viii. p. 246. Sir B. Brodie has described a case of the same kind, occurring in a young lady eleven or twelve years of age (*Works*, vol. iii. p. 170).

† *Sepulchretum*, tom. ii.

Incised wounds of nerves are frequently the cause of the kind of symptoms as those produced by puncture.

Hamilton relates the interesting case of a girl, aged seventeen, a nervous temperament, who, while cutting bread, wounded the septum between the thumb and forefinger of the left hand. The pain from the first was severe, became still more so, and extended to the thumb, forefinger, the hand, up the forearm, inside of the arm to the axilla, shoulder, and to the neck. These and other symptoms continued for nearly three months, when the external wound had healed, when she got a fright in the street, as into an hysterical fit. After two days of hysterical symptoms, all pain swelling left the arm and never returned.*

This interesting case is a good illustration of the remarkable influence of hysteria in aggravating the effects of local injury and affords an example of the singular manner in which effects are removed by sudden emotions of the mind. Brodie, who paid so much attention to this important subject, justly observes, that when the patients are subject to hysterical paroxysms, sometimes the paroxysms cease on the appearance of the local symptoms, and sometimes, on the contrary, the appearance of the former is followed by an abatement of the latter by complete recovery from them.

He mentions the case of a young lady, who having long suffered hysterical neuralgia of the hip-joint and thigh, was entirely cured of symptoms on being thrown from a donkey.

I knew the case of a girl, who, while suffering from complete paraplegia, jumped up in a fright, on seeing a mouse run across the road, was cured from that moment.

For an important case of incised injury of nerves, of which the fact is the substance, we are again indebted to Mr. Wardrop.† A young gentleman cut the distal phalanx of the left thumb obliquely across the radial side of the nail. The wound readily healed, but on the patient living free a few days the thumb became painful, and although no change in its appearance could be perceived, and the cicatrix seemed perfectly natural, the pain extended to the forefinger, the radial side of the middle finger, and up the arm to the neck and side. The pulse was frequent and tense, the face flushed, the tongue white and frothy. Copious general bleeding gave almost immediate relief. The symptoms, however, returned and yielded to another bleed with copious purging. The paroxysms of pain were several times produced by mental excitement, and on some occasions were brought on by taking even a small portion of animal food. The wounded thumb, while at all times painful and extremely tender to the touch, was sometimes

* *Dub. Med. Journ.* vol. xiii. case iii. p. 42.

† 'On Local Nervous Affections, and Local Hysterical Affections,' vol. iii.

‡ *Medico-Chirurg. Trans.* vol. xii. p. 205.

interesting case has several points worthy of particular attention. It affords a good example of acute inflammation in a recently injured nerve by a too early use of the part it supplies, by an incautious mode of living, or by mental exertion. We also see not only that the pain—or rather the pain exciting the pain—was reflected from the injured nerve to other and even distant nerves with which it is connected only through the medium of the nervous centres, but even after the connection of the injured portion of the nerve with its nervous centres had been interrupted, pain was still felt in the other nerves by accidental causes. Indeed, in severe or long-standing cases of traumatic neuralgia, it is reasonable to believe that the nervous centres in which the nerves are implanted, are sometimes secondarily and entirely deranged or diseased, and in this way become the seat not only of neuralgia, but even of paralysis in other and distant parts. Among other cases that might be mentioned, one recent case of complete division of a nerve, recorded by Mr. [Name], appears to be of this nature.

A lady, aged twenty-three, on December 20, 1822, wounded the ulnar nerve of the second finger of the left hand, near the middle of the second phalanx, while squeezing an orange. The pain extended to the centre of the left breast, and from the side of the neck to the face, along the branches of the facial portion of the seventh (?) nerve. On the sixth day after the accident the wound had healed, but was extremely tender. The pain was excruciating when she attempted to move the arm with the hand in a state of pronation. Whenever pain was produced in the superciliary nerves of the left side, after

At the place of the division of the nerve at the first operation, both its extremities were incorporated with the cicatrix, as were those of the dorsal branch, which had also been divided. Although her local and general symptoms were much relieved by the operation, she continued to feel pain in the hand, arm, neck and face, and about four months later (in July), she began to complain of her spine. About the end of November she complained very much of pain in her back, with tingling in her arms, and a difficulty of supporting herself erect. Percussion along the spine produced uneasiness in every part, and much pain about the lower dorsal vertebrae. Some time after, she was seized with violent pain in the left knee, which lasted for two or three days. Pressure on each side of the spinous processes of several of the vertebrae produced pain, and percussion with a key made it very severe. She had difficulty in voiding her urine. In October 1824, she complained of numbness and loss of sensation in the left hip and shoulder, and of pain at the back of the neck, with a feeling as if the neck could not support the head. At the latter end of the year 1825 and beginning of 1826, she was affected by dizziness, in fits of which she fell down, but never entirely lost her consciousness. Her left arm and leg were weaker than the right. When the membrane lining the left external auditory meatus was touched, cough was produced. Up to March 1829, she had varying degrees of pain and weakness, soreness in the throat with an appearance of venous congestion, and disorder of the digestive organs, with great tenderness in a spot about the size of half a crown at the pit of the stomach. She had also some pain and swelling about the uterus and vagina. When sleeping on her left side she very frequently awoke with pain in the amputated finger. *Tic douloureux* was brought on by either exciting or depressing emotions, by exposure to strong light, and by fits of sneezing, with which she was frequently troubled. When Mr. Swan last saw her on October 30, 1833, she was nearly in the same state as she had been in for a long time; but we are left in ignorance of the subsequent course of the disease. The patient's father had a paralytic attack.*

Here, then, we have an instance of very extensive neuralgia, of incomplete but extensive paralysis, and other severe symptoms extending through a period of ten years, as the effects of simple division of a nerve of one of the fingers. That these morbid symptoms arose out of some peculiarity of constitution, there can be no doubt; and it seems highly probable that this peculiarity was of an hysterical character.

Foreign bodies embedded in a nerve or pressing against it are often the exciting cause of violent pains and other symptoms similar to those which result from wounds.

Jeffries relates the case of a girl who suffered from violent and almost continual neuralgia of the face for fourteen years. A hard, pointed substance was felt under the skin of the right cheek; and at this point the slightest touch

* Swan, *Diseases, &c. of Nerves*, 2nd edit. p. 129. Of this remarkable case I have given the briefest possible abstract of the most salient and important points, but the details, which are very full, deserve attention.

brought on an attack of pain. An incision was made, and a triangular piece of a china cup removed, with complete cessation of the neuralgia.

(One of the most interesting and remarkable cases of this description is related by Denmark.* A young soldier was wounded by a musket-ball which pierced the triceps extensor cubiti, about one inch and a half above the outer condyle of the os humeri, grazing the inner side of that bone, and passed obliquely downward through the brachialis internus, and out anteriorly through the bend of the arm. The wound soon healed without exciting any particular morbid symptom. The man was subsequently admitted into Basilar Hospital, with excessive pain, frightful dreams and startings. The fore-arm was always bent and in the supine posture. A small tumour could be felt in the site of the wound on the anterior part of the arm, which he could not bear to be touched. The pain began at the extremities of the thumb, and all the fingers except the little one, and extended up the arm to the part wounded. It was a burning pain, and so violent as to cause a continual perspiration from the face. Neither opium nor any other means employed assuaged his intense sufferings. At his own request, the arm was amputated, with immediate relief, and he was discharged cured in three weeks. On examining the amputated arm, the radial nerve seemed to be divided with, and intimately attached to, the wounded parts, for the space of an inch. It had itself been wounded, and at the place of the injury, was thickened to twice its natural diameter, and seemed as if contracted in its length. On further examination, Dr. Denmark was surprised to find, in dividing the fibres on the posterior part of the wounded nerve, that there was a small portion of the ball firmly embedded in it, which had been driven off by crushing the bone.

Treatment of injuries of nerves.—When an incised or other open wound is accompanied and followed by very acute pain, by more or less numbness, and by partial or complete loss of sensation and motion in the part beyond the wound, we may conclude that a nerve has been either injured or divided. Under these circumstances it is of primary importance to remove all dirt or other foreign bodies that may be present, to promote healing by the first intention, and enjoin perfect rest. If these precautions are not adopted and the external wound inflames, the nerve is unable to participate in the inflammation, and may then become the source of the violent and distressing symptoms already described. When this is the case it may be necessary to apply leeches to the neighbourhood of the wound, to use either cold lotions or warm fomentations of poppies, and poultices, and to act briskly on the bowels by means of purgatives. When the inflammatory symptoms run high and the constitutional disturbance is considerable, I have seen the most decided benefit derived from a combination of tartarated antimony, opium, and

* *Med.-Chirurg. Trans.* vol. iv. p. 48.

hyosciamus, given at short intervals, the antimony being in sufficient quantity to keep up a state of great nausea.

In some cases, after all inflammatory action has subsided and the external wound has completely healed, painful and distressing symptoms may still persist at intervals for a very considerable period; and in other cases where the external wound has healed at once in the kindest manner by the first intention, similar symptoms may supervene. Most of these latter cases are found in persons of a more or less hysterical constitution. But, as we have already seen, there are other causes besides hysteria, such as pressure, dragging, or inflammation of a cicatrix; a premature use of the limb—that is, before the nerve is sufficiently repaired or united, and while the nerve is in this state, it is in some cases very susceptible of irritation by errors in diet, intemperate habits, mental emotions and influences of the most trivial nature. One part of the treatment will of course consist in avoiding or removing the exciting cause, if this can be ascertained. When the cicatrix presses on the nerve, active frictions with moderate exercise of the part will relieve the pain. If it be simply painful or tender, the application of belladonna will frequently allay the irritation; or if it appears to be inflamed, the removal of a little blood by means of two or three leeches may be necessary. If pain and tenderness be experienced in the course of the nerve, repeated leeching, or blistering, and sometimes actual cautery will afford relief. In cases of that intense *burning* pain which is sometimes so agonising in nerve-wounds, there is no treatment so successful as repeated blistering. The painful state of the parts to which the wounded nerve is distributed may be treated by the local application of anodynes, such as chloroform, aconite, opium, and belladonna; and the most effectual mode of using the two latter substances, is by subcutaneous injection. Mr. Pearson * has recorded a very remarkable case of neuralgia, which, after resisting every kind of treatment, yielded at length to the application of a stimulating embrocation that was rubbed into the part for ten minutes twice or three times a-day.†

In some instances when every other remedy has failed, the happiest effects have resulted from the use of mercury exhibited for a period sufficiently long to affect the gums. Mr. Hamilton

* *Medico-Chirurg. Trans.* vol viii. p. 252.

† R. ol. olivæ, ʒijss.; terebinth, ʒjss.; acid. sulph. ʒjss. M.

as recorded two cases of this description.* In many instances, however, in which this remedy has been employed, it has completely failed.

In extreme cases, when all other resources are exhausted, it becomes a question whether the nerve should be divided or a portion of it excised, or whether the member which it supplies should be amputated. If the constitutional disturbance is of an alarming character, there should be no hesitation in once dividing the nerve, or excising a small portion; for these operations have been frequently followed by an immediate arrest of all the symptoms, as in Dr. Wilson's case already mentioned.†

In a case of contused thumb followed by violent symptoms, Sir Astley Cooper removed five-eighths of an inch of the radial nerve with complete success. Mr. Sherwin relates the case of a servant-maid, who suffered from the most alarming symptoms after the operation of venesection. At the end of a fortnight she was cured by a deep incision above the cicatrix.‡ A similar case with the same result is recorded by Dr. Watson.§ Mr. Earle excised about half an inch of the radial nerve, for punctured wound of the thumb followed by pain, spasm, temporary trismus, &c., which had lasted for two months. The patient was soon relieved, and after some months completely cured.

In many instances in which division of the nerve has failed, amputation has been performed with complete success, as in the case of Mr. Wardrop.|| A still more decided example is recorded by Mr. George Bell.¶ A lady, twenty-six years of age, while cutting a loaf of bread, injured the nerve of the thumb, on the radial side. The wound healed kindly, but was followed by excruciating pain and other violent symptoms. Two years after, on two different occasions, an incision was made through the soft parts to the bone, in the neighbourhood of the wound, but with only temporary relief. Mr. Bell removed the thumb at the second joint, and in five or six weeks she was restored to perfect health.

However, in some cases, especially of females of marked hysterical constitution, neither the division of nerves nor amputation is of any permanent benefit; while in others which have resisted every kind of treatment except by operation, a spontaneous cure has occasionally been effected.

When nerves have been contused or compressed, without coming inflamed, the best kind of treatment consists of the repeated use of exceedingly hot fomentations, frictions with

* *Dublin Medical Journal*, vol. xiii. pp. 41, 48.

† See p. 165.

‡ *Duncan's Medical Commentaries*, vol. iv.

§ *Medical Communications*, vol. ii. p. 251.

|| See p. 165.

¶ *Edinburgh Journal of Medical Science*.

stimulating embrocations, and Faradisation. But if tenderness and pain be felt in the course of the injured portion of the nerve, it will be necessary to apply leeches, cold lotions, and blisters, and to insist on perfect rest.

NEURALGIA.

It has been already seen that a large proportion of neuralgic cases are due to traumatic irritation or inflammation; the some are the result of idiopathic inflammation, in either acute or chronic form; that others are the consequences of ulceration; of compression and irritation caused by aneurism, by enlarged glands, by tumours, &c., either in the neighborhood of nerves or in their substance. Moreover, neuralgia of the tri-facial nerve has in some cases been found to depend on disease of the bones of the head and face. In the case of the late Dr. Pemberton, the frontal bone was thickened, and in the falx cerebri, there was an osseous concretion. Sir H. Halford found hypertrophy and other morbid states of the cranial and facial bones, such as exfoliations of the antrum, considerable thickening of the frontal, ethmoid, and sphenoid bones. Dr. Bright relates an interesting case of facial neuralgia in which he found fungoid tumours of the brain and disease of the cranial bones.† In some instances the disease of the bone is the result of syphilis, which also attacks the nerves themselves, not only in the head but in all other parts of the body.

Many cases of neuralgia have a gouty or rheumatic origin; but there are others that do not depend on any of the causes above mentioned, and in which, when there has been an opportunity of post-mortem examination, no appreciable alteration has been discovered in the nerves. Yet that some peculiar, but perhaps only periodical, change takes place in the tissue either of the nerve itself or of the nervous centre from which it arises, there can be no question; for to speak of functional disorder without some material change, is to speak of an effect without a cause. In some instances, the exciting cause appears to act directly and immediately on the nerve-tissue, as in cases of neuralgia resulting from atmospheric changes, particularly from electric states of the atmosphere. In other instances, the effect

* Travers, *A Further Inquiry concerning Constitutional Irritation, and the Pathology of the Nervous System*, p. 351. 1835.

† *Reports of Medical Cases*, p. 506.

may be produced through the medium of the circulation, by malaria and blood-poisoning of different kinds. Anæmia, again, especially in females, is among the most common causes of neuralgia; as is also hysteria, even when unattended by anæmia. Disorders of digestion, constipation, the sudden suppression of secretions, or of habitual hæmorrhages, such as hæmorrhoids, the catamenia, &c., may all be productive of neuralgia in persons already predisposed.

The points of diagnosis between this form of neuralgia and that which arises from neuritis or inflammation of nerves, are very important, because the treatment required is very different in the two kinds of cases. In non-inflammatory neuralgia, although the pain may be excited or aggravated by the slightest touch, it is generally relieved by firm pressure, which is not the case in neuritis. Moreover, the pain is always interrupted by emissions of variable length in different cases, and is extremely varied in its character. It begins and ceases suddenly, and is apt to shift from one branch of a nerve to another, with a rapidity and intensity which are never experienced in neuritis. Antiphlogistic treatment, which relieves the latter, frequently aggravates non-inflammatory neuralgia.

NUTRITIVE AND OTHER CHANGES RESULTING FROM INJURIES AND DISEASES OF NERVES.

(1) *Changes due to Injuries of Nerves.*

In the earlier part of the present century a few observers occasionally and incidentally noticed the morbid effects of injuries of nerves on the nutrition of certain tissues; but it is only within the last ten or twelve years that the subject has been systematically studied. These lesions of nutrition consist chiefly of,

(a) Diminution of temperature in the parts to which the injured nerve is distributed.

(b) Cutaneous eruptions, ulcerations, and alterations in the colour and texture of the skin, and its appendages.

(c) Increase or diminution, and alterations in the quality, of its secretions.

(d) Periodical swelling of joints and other parts of a member.

(e) Atrophy and contraction of muscles supplied by the injured nerve, with distortion and stiffness of joints.

nger. I merely mentions a case of anæsthesia, in which similar effect produced by warmth.†

Mr. Jonathan Hutchinson, in a valuable and interesting article on 'Injuries to Nerve-trunks,'‡ observes that the parts paralysed by division of the nerve which supplied them, never, even in the warmest temperature, be raised to the temperature of adjacent, unparalysed parts. 'Nor does even the existence of active inflammation raise the part to the normal state, although it much increases it.' 'It would appear, then, that while a paralysed part can be cooled to almost any extent, it cannot be raised by artificial heat beyond a certain point, that point much below the maximum of its uninjured condition.'

(b) Romberg, more than thirty years ago, very fully described some of the most important nutritive changes that the skin and its appendages undergo, when deprived of the influence of the supply of nerve-influence. There is found, he observes, increased branny or scaly desquamation of the epidermis, accompanied by an absence of transpiration. The nails curve and become rough and ragged, and fall off.

He mentions a case of Dieffenbach's, in which there had been violence for ten years in the right foot. On the outer and posterior side of the foot, near its middle, there was a tumour about five inches in circumference. It was also painful, and, when pressed, augmented the pain in the foot. The tumour was removed in 1836, and Romberg saw the patient two weeks afterwards. There was complete anæsthesia in the parts supplied by the peroneal nerve. The muscles of the leg and foot were atrophied. It

during the operation of blood-letting, was followed by the most severe neuralgia and other symptoms, and the wounded arm became covered with hair.* Larrey mentions the case of a soldier who received a kick from a horse on the right eyebrow. The frontal sinus was fractured, and after some days, cutaneous hyperæsthesia and tetanic symptoms of the same side supervened. To Larrey's great surprise, the hair and moustaches of the injured side matted up and became exquisitely tender to the slightest touch. About a week later the nails became ragged and rough, and fell off.† Similar effects were observed by Bellengeri from a contused wound above the supra-orbital foramen. The hair on the injured side not only became stiffer, but grew with much greater rapidity.‡ Pouteau records a remarkably interesting case, with symptoms similar to those of Larrey and Bellengeri, and resulting from a contused wound of the supra-orbital nerve.§ In a young lady, who came to me with painful neuralgia of a circumscribed portion of the scalp, the hair over the painful part was not only coarser and stiffer, but perfectly white. Reading, or playing the piano for a few minutes only, brought on the most insupportable and insupportable feelings, radiating as it were from the affected part. In a case of temporal neuralgia mentioned by Trousseau, the hair on the affected side turned white, and acquired considerable stiffness in a very short time.

With regard to cutaneous eruptions resulting from injuries of nerves, numerous interesting observations besides the one already quoted from Romberg, have been recorded by Rayer,|| Earle,¶ Ennet,* Opplolzer,†† Charcot,‡‡ Raynaud,§§ Kuhl, ||| Thomas,¶¶ Paget,*** Brodie,††† and others. These cutaneous affections have been shown to consist chiefly of erythema, and of vesicular eruptions allied to eczema, erysipelas, and herpes. Mr. Paget has given an excellent description of the erythema observed in cases of injured nerves. 'In well-marked cases,' he says, 'the fingers which are affected are usually tapering, smooth, hairless, almost void of wrinkles, glossy, pink or ruddy, or blotched as if with permanent chilblains. They are commonly also very painful, especially on motion, and pain often extends from them up the arm.'

* *Dublin Med. Journal*, 1838, vol. xiii. p. 46.

† *Clin. chirurg.* tome i. p. 200.

‡ *Archiv. gén. de Méd.* 1835, tome vii.

§ *Œuvres posthumes*, tome ii. p. 92, obs. iii.

|| *Traité des Maladies de la Peau*, 1822, tome ii.

¶ *Medico-Chirurg. Trans.* vol. vii.

** *Journal de Physiologie*, 1859.

†† *Allgemeine Wiener medicinische Zeitung*, No. 48, Nov. 1866.

‡‡ *Journal de Physiologie*, 1859.

§§ *Thèse de Paris*, 1862.

|| Apud Samuel, *Die trophischen Nerven*, p. 189.

¶¶ *Archiv. der Heilkunde*, 1866, p. 153.

*** *Medical Times and Gazette*, 1864.

††† *Works*, vol. iii. 'Local Nervous Affections.'

It is to Drs. Mitchell, Morehouse, and Keen, of the United States, that we owe the most precise account of the different morbid changes in the skin, and other tissues, resulting from particular injuries of nerves.* They have shown that the changes in the skin which follow wounds of nerve-trunks are of two kinds. The first is the result of entire division of the nerves of the part, with palsy of the whole limb. In many instances there is early oedema of the part. The skin thickens and dries; the epithelium hangs in patches here and there, and is yellow and even pale-brown. These peculiarities, however, are partly the consequence of mere disuse. But the nails become curved as in tubercular disease.

The second kind of pathological change is the result of a partial division of the nerve. The skin is deep red or mottled, or red and pale in patches. The cuticle seems partially lost, so that the cutis is exposed in places. The sub-cuticular tissues are nearly always shrunk. The surface of all the affected parts is glossy, generally devoid of wrinkles, and perfectly free from hair. Sometimes the fingers are dotted with islets of thin and red and glossy skin. This glossy appearance of the skin is always accompanied by pain, generally of a burning character.† 'The particular time can be named as the period at which these changes in nutrition first show themselves. This alone can be said, that they do not belong to cases of complete destruction of the nerves. They may begin within a few days, or at any later date; but usually they arise while the wound is healing.'

If the nerves of the hand be injured, the hair, after some months, disappears from the affected fingers, and the nails undergo remarkable alterations. These alterations consist of a curve in their long axes, an extreme lateral arching, and sometimes a thickening of the cutis beneath their extremities. In other instances the skin at that end of the nail next the third finger-joint becomes retracted, leaving the sensitive matrix partly exposed. At the same time the upper line of union of skin and nail retracts into or under the latter part, and in place of a smooth edge is seen through the nail as a ragged and notched border. 'No deformity of the nails in tubercle at all approaches that which nerve-wounds occasion. Indeed, we think it would be possible for one familiar with these cases to diagnose the

* *Gunshot Wounds and other Injuries of Nerves.* Philadelphia, 1864.

† Page 80.

presence of nerve-lesion from the form of these protuberant oddly-curved nails.' When the nails of the toes have been hacked, the curving is as marked, but a distressing ulceration apt to occur at their angles, and to break out again and again. The best remedy then is excision of the outer edge of the nail, matrix and all.*

In a case mentioned by Mr. Jonathan Hutchinson of division of the ulnar nerve, the nail of the little finger, about two months after the accident came off together with the skin, 'like the end of a glove.' It grew again, but not to more than half its original size.†

(c) *Alteration of secretions after injury of nerves.*—We have already seen that after the nerves of a limb have been completely divided, the skin is generally dry; but injuries of nerves without complete section modify the secretions in regard both to quality and quantity. The secretion is generally abundant, and sometimes intensely acid, so that an odour like vinegar can be felt in the neighbourhood of the patient. In one instance the odour of the sweat was 'disgustingly heavy,' and resembled the smell from a bad drain.‡

d) *Periodical swelling of joints, &c.*—This peculiar affection, which is particularly interesting to the surgeon, is the result of idiopathic neuritis and traumatic affections of nerves. It closely resembles ordinary rheumatism or gout that it is not easy to point out the distinguishing characters of the two kinds of disease. Remak calls it *arthritis spuria*.§ Dr. Mitchell has recorded some very interesting cases of this affection, in the *American Journal of the Medical Sciences* (1831, vol. viii.). It may occur at any time after the first few days of injury to a nerve, and may attack any articulation, or all the articulations of a member. Once fully established, it keeps the joints stiff and sore for weeks or months. When the acute stage is past, the articulations become hard, and partial ankylosis results. Of all the agencies which impede movement, it is the most difficult to relieve.

e) *Atrophy and contraction of muscles from injury of nerves* have been for many years carefully investigated by Dr. Duchenne

Page 82.

London Hospital Reports, vol. iii. p. 308.

Mitchell, Morehouse, and Keen, *op. cit.* p. 86.

Allg. med. cent. Zeitung, Berlin, 1863. *Ueber den Einfluss des Nervensystems auf Krankheiten der Knochen und der Gelenken.*

DISEASES AND INJURIES OF NERVES.

anlogne), whose work (*Électrisation localisée*) contains
 of the most valuable cases. Complete division of
 nerve is always followed to a certain extent, in the course of
 time, by atrophy of all the muscles to which that nerve is dis-
 tributed; and the wasting is much more rapid when the main
 artery has been divided or interrupted. But in *partial* division
 or injury of the nerve, only those muscles are liable to suffer
 which are supplied by the injured fibres, and in this case the
 wasting is more rapid and decided than when the nerve has
 been *completely* divided. Thus, a whole group of muscles may
 be affected, or only part of a group; a single muscle only, or
 even only part of a muscle. Moreover, a muscle may be affected in
 either a lateral or a medial direction. So soon as the
 muscular fibres begin to suffer in nutrition, their tension or
 tonicity diminishes, they become softer, more flabby or relaxed.
 Sooner or later they begin to waste, and after a time, in many
 instances, they acquire a kind of morbid tension, and contract
 or shorten permanently to a greater or less extent, causing a
 variety of deformities, alterations in the form of the joints, and
 loss of natural movement. This morbid tension, however, is
 not always in proportion to the degree of atrophy, for in some
 instances the wasting is slight while the muscular contraction
 is extreme. Moreover, a muscle may become shortened from
 paralysis of its opponent, and this condition must be dis-
 tinguished from the shortening which is due to atrophy.

Early loss of tone in a muscle, from injury of its nerves, is a
 bad sign. Rapid diminution of size and contraction, when
 due to nutritive changes, is still worse. All these changes are
 ominous of ultimate deformities and permanent loss of power.
 In the diagnosis and prognosis, no less than in the treatment
 of injuries to nerves and their consequences, electricity plays
 the most important part; and for its first application to these pur-
 poses, we are chiefly indebted to the remarkable sagacity and
 skill of Dr. Duchenne (de Boulogne).† The way in which he has

* Mitchell, Morehouse, and Keen, p. 120.

† The following statement is made by the most competent judges on the
 subject. 'As respects its value in traumatic lesions of nerves, we feel com-
 strained to state that it has been understood and rightly appreciated by Dr.
 Duchenne alone. After a year of great experience in the use of electricity, we
 are still satisfied of the essential correctness of almost every proposition on the
 subject which this distinguished physician has laid down.' (Mitchell, More-
 house, and Keen, op. cit. p. 136.) I am glad to learn from Dr. Duchenne, that

made it an instrument of diagnosis and prognosis, in diseases and injuries of nerves, is no less important than beautiful in a practical point of view.

When a healthy muscle is made to contract by means of electricity, it becomes the seat of a peculiar pain, which is called electro-muscular *sensibility*. This pain, however, is not caused by the contraction, for it is still experienced in muscles which have lost the power of contracting under the influence of electricity. The property of contracting under the same influence, is called electro-muscular *contractility*. Now it has been laid down by Dr. Duchenne as a law which is very remarkable, but which subsequent experience has fully confirmed, that the persistence of this property of electro-muscular contractility is not necessary for the exercise of *voluntary* contraction.* On the other hand, muscles that are partially or completely paralysed, with regard to the *will*, may sometimes retain their normal degree of electro-muscular contractility; while in certain cases, muscles which have lost, to a very unequal degree, the power of contracting under the stimulus of electricity, may seem to be all equally paralysed as regards the *will*. There is another remarkable and important practical point to be remembered, viz. that although the retention of electro-muscular contractility is not necessary for the exercise of voluntary contraction, yet that whenever it is lost, or even impaired, the prognosis is unfavourable in reference to the return of *voluntary* control, and indicates that the affected muscles will suffer in their nutrition and become atrophied. It is curious, however, and very important, that notwithstanding the loss of electro-muscular contractility, the *therapeutical* power of Faradisation may still exercise its influence on the paralysed muscles. If electro-muscular sensibility be retained, the chances of recovery are more favourable; and whenever it is impaired, if pain be excited by the action of electricity, the effect is a favourable sign.

Different forms of paralysis which depend on traumatic affections of nerves, and which may appear to be exactly alike—which cannot, indeed, be distinguished from each other by

somewhat abridged English translation of his laborious work is in course of preparation. It ought to be in the hands of every surgeon and physician. To one, judging from its unhappy title alone, would expect to find in it so much a mine of pathological facts.

* *De l'Électrisation localisée*, 2^{me} éd. p. 350.

the ordinary modes of diagnosis—may nevertheless differ essentially from each other with regard to their progress, their gravity, or their termination. The electro-muscular test alone enables us to discriminate those cases in which muscles will remain paralysed and become atrophied in the course of a few months, from other cases in which it may be predicted with certainty that the paralysis, without atrophy of the muscles, will disappear, either spontaneously or by means of Faradisation.

A man under the care of the late eminent surgeon M. Roux, at the *Hôtel-Dieu*, had dislocation of the humerus forward. The reduction was deferred till the next morning, when it was found that the head of the bone had changed its place to the axilla. The muscles of the arm, fore-arm, and hand were completely paralysed to the will, but sensibility was retained. This state of things having continued for nearly a month, with gradual wasting of the muscles, Dr. Duchenne was requested by M. Roux to examine the arm by means of Faradisation. All the movements of the upper extremity were found to be completely abolished, and it would therefore have been natural to conclude that all its muscles were equally affected, or rather that all the nerve-fibres that supplied them were equally injured. But it was found that some of the muscles had completely lost their electro-muscular contractility, and subsequently wasted rapidly; while the other paralysed muscles, which had not suffered in this way, not only escaped atrophy, but recovered their voluntary power after a few applications of electricity. The deltoid had partially lost its electro-muscular power, and therefore required for its restoration a little longer time than those which wholly retained it.

In paralysis resulting from *cerebral* disease, the electro-muscular contractility remains unaffected, and therefore Faradisation is of great importance in the differential diagnosis of paralysis which follows injuries of nerves or affections of the spinal cord.

A man was admitted into the *Hôpital de la Clinique*, for paralysis of the right arm. According to his own account, some days before his admission he woke in the morning with his arm paralysed, but perfectly free from pain. At first he only felt formication in the fingers, with a sensation of numbness and coldness in the whole limb; but a few days after, he began to experience a lancinating and intermitting pain of no great intensity in the right cervical region. The suddenness of the attack, without any preceding pain, occurring in a young man who had not suffered from any traumatic lesion of nerves, or been apparently exposed to any other influence that might occasion paralysis of the arm, would reasonably suggest a cerebral cause. But on examining the limb by means of Faradisation, it was found that the majority of the muscles had lost, in a greater or less degree, their property of contracting under the stimulus of electricity—a property which, as already stated, is retained in paralysis resulting from cerebral causes. Judging from these facts, Dr. Duchenne at once came to the conclusion, that the real cause of the paralysis was to be referred to some alteration either of the brachial and cervical plexus or of that portion of the spinal cord from which they derive their

origin. On further examination a month after by M. Nélaton, who then for the first time took charge of the patient, he was found to be affected with tertiary symptoms of syphilis. That distinguished surgeon thought that some kind of exostosis might be the cause of the paralysis, and accordingly prescribed the proto-iodide of mercury. After some days the pain ceased, and then the paralysis gradually disappeared from the muscles which had not lost their electro-muscular contractility.

Treatment.—For restoring the paralysed and wasted muscles to their original condition, electricity is unquestionably the most important agent. It should be employed daily for at least ten or fifteen minutes, and each muscle should be separately Faradised. The period when the treatment by electricity should commence must vary according to the peculiarities of the case. If the electro-muscular contractility remain intact, Faradisation should be employed as soon as possible. But when this property is lost—that is, when the nerve-influence is no longer supplied to the muscles, and more particularly when the muscular sensibility is impaired—the treatment by electricity is of no service, and should not be employed until some months after the injury, or until there is good reason to believe that the nerve-lesion is repaired. In addition to electricity, friction and shampooing should be employed, with passive motion, and voluntary exercise of the muscles, as soon as voluntary control is in any degree restored. The douche is often of great service, particularly if hot and cold water be alternately used. Care should be taken to preserve warmth in the affected part. When flexion or extension of a limb occurs in consequence of contracted muscles, splints, bandages, and other apparatus will be necessary.

When the effects of the nerve-lesion are aggravated or prolonged by some peculiar diathesis, or when the health of the patient is suffering from severe pain and want of sleep, general as well as local treatment must be adopted. Narcotics, iron and quinine, cod-liver oil, and phosphorus are the most worthy of recommendation. The hypodermic method is the best mode of administering narcotics. If morphia be employed, the first injection should not exceed the sixth of a grain.

(2.) *Changes due to Idiopathic Affections of Nerves.*

Although traumatic lesions of nerves give rise to the most serious nutritive changes in the tissues to which those nerves are distributed, it is not uncommon to find that idiopathic affections of nerves are followed by changes of a similar nature.

The principal alterations in the nutrition of the skin resulting from idiopathic neuritis and neuralgia are manifested in the form of herpetic eruptions. Rayer, I believe, was the first to point to neuralgia as the cause of these eruptions, while other writers, including even Valleix and Cazenave, regard the neuralgia as a consequence of the cutaneous affection. Rayer's opinion, however, has been fully confirmed by subsequent observation. The pain is frequently found to precede the eruption, while the latter is sometimes seen to follow the course of the affected nerve.

Barensprung has recorded a very valuable case, to which I have alluded, and which proves in the most convincing manner, the close relation between idiopathic inflammation of nerves and herpes zoster. The patient was a child one year old. The eruption extended round one side of the trunk from the sixth to the ninth rib, and after death, which resulted from tubercular disease, the roots of the sixth, seventh, and eighth intercostal nerves, especially the seventh, were increased in size and of a red colour, in consequence of the presence of enlarged and tortuous vessels in the neurilemma. The diameter of the seventh intercostal exceeded by more than one-half that of the fifth or the ninth. The corresponding intervertebral ganglia were firmly adherent to the intervertebral canal; their connective tissue presented an inflammatory redness, and they were decidedly increased in size.*

MM. Charcot and Cotard have related the equally interesting case of a woman, aged seventy-eight, from whom a cancer of the right breast was removed in August 1865. In October of the same year, she complained of sharp pains in the right shoulder, and in the right half of the neck. These pains were continuous, but increased in the severity at intervals, during which the patient appeared to suffer the most acute pain. About December 15 of the same year an herpetic eruption made its appearance over the whole right half of the trunk, limited exactly, both before and behind, to the median line. The eruption occupied all the other parts of the skin which are supplied by branches of the cervical plexus on the right side, but was not followed by any diminution of pain. The patient died on December 26, after symptoms of effusion into the pleura on both sides. On post-mortem examination it was found that both the anterior and posterior roots of the spinal nerves in the cervical region were perfectly healthy, the intervertebral ganglia, as well as the communicating tracks formed by the union of the two roots, presented a slight tumefaction and a vascularity which was manifested by a bright red colour. The difference was striking when the corresponding parts on both sides were examined.†

Among the most remarkable cases of neuritis giving rise to herpetic eruptions of the skin, are those which M. Leudet, of the Hôtel-Dieu of Montpellier, has shown to be caused by the inhalation of carbonic acid.

* Barensprung, *Annal. Charitékrank. zu Berlin*, vol. xi. part ii. p. 96.

† *Mémoires de la Société de Biologie*, 1866, p. 41. From such observations, Barensprung seems to be quite satisfied that the intervertebral ganglia on the posterior spinal roots, give origin to the so-called trophic nerves; but it appears to me that facts do not warrant the conclusions at which he arrives.

When the symptoms of asphyxia have subsided, it is not common, after a variable number of days, to find certain orders both of sensibility and motion. The lower extremities the parts which are usually affected, although different parts of the nervous system are sometimes involved. In one case, which proved fatal, M. Leudet found the right sciatic nerve at least one-third thicker than the left; its neurilemma injected, thicker, and more indurated than on the opposite side.

In these singular cases, the skin was affected by eruptions similar to herpes. These eruptions appeared almost immediately after the subsidence of the symptoms of asphyxia, and generally, lasted only a short time.*

Herpes is not the only cutaneous eruption that has been found to follow neuralgia and neuritis, whether traumatic or idiopathic: lichen has been observed by Canuet in cases of neuralgia; † acne and erysipelas have been seen by Hasse ‡ and Bernberg § to accompany the same affection.

J. LOCKHART CLARKE, M.D.

Recherches sur les troubles des Nerfs périphériques, et surtout des Vaisseaux, consécutifs à l'asphyxie par la vapeur de charbon. Archiv. de Médecine, 1865.

Thèse de Paris, 1855, p. 29.

‡ *Nervenkrankheiten*, p. 62.

Op. cit.

DISEASES AND INJURIES OF NERVES

PART II.

REMOTER CONSEQUENCES OF NERVE-LESIONS.

INTRODUCTION.

TWO distinct groups or classes of symptoms may be caused by a lesion of a nerve : in one class the symptoms are the effect of the *loss of function* or *cessation of action* of the nerve ; in the other, on the contrary, they are produced by *an action* of the injured or irritated nerve. In the first class, therefore, the symptoms depend on *lack of action*, while in the second, they are due just to the reverse, i.e. *existence of action*, in a nerve.

Each of these two classes of symptoms may be subdivided. As there are at least four different kinds of nerves, four distinct kinds of symptoms of loss of function or cessation of action may be observed after a lesion of a nerve :

1st. There may be paralysis of motion.

2nd. There may be paralysis of centripetal nerves (sensitive, incito-motory, &c.).

3rd. There may be paralysis of the vaso-motor nerve-fibres in consequence of which the blood-vessels may be distended and full of blood.

4th. There may be paralysis of the special nerve-fibres, the normal influence of which consists in giving activity to nutrition and secretion.

Leaving aside these four kinds of symptoms, I will confine myself here to the study of some of the effects of irritation of nerves. These effects may be grouped under two heads—the *peripheric*, or *direct* ; and the *remote*, *indirect*, or *reflex*. Of the peripheric, or direct effects, I will simply say that they give origin to five kinds of symptoms :

mentioned kinds of changes in the circulation of blood, are various alterations of nutrition or secretion in the skin, the joints, &c.

These five kinds of effects of irritation of nerves, four, viz. first, third, fourth, and fifth, may also be caused either by the reflex action, or by some peculiar influence upon or in the nervous centres. I propose here to treat only of our last kinds of symptoms.

the functional affections or disorders, and most of the diseases are frequently produced by an influence upon the nervous centres by an irritation of any part of the length of a nerve. This irritation may occur in the ramification of a nerve in a mucous membrane or the skin (and to worms, a calculus, &c., or to cold); or it may depend upon an organic or a functional affection of the trunk of a nerve (as of wounds, burns, tumours, or neuralgia). In most cases an inflammation of the various viscera owes its origin to a reflex influence on the organ which becomes inflamed, arising from the irritation of some cutaneous nerve-fibres brought about by cold air. I will not say more here on this origin of visceral inflammation caused by cold acting on the skin, my object being only to give an outline of the various effects of injuries or diseases of all parts of nerves excepting the work of their terminal ramifications.

the various reflex and other remote effects of irritation of

nutrition and secretion. After having mentioned clear and positive facts, showing that all these affections may be caused by an injury to, or a disease of, a nerve, I will briefly give the rules concerning the diagnosis and treatment of these effects of injuries and diseases of nerves. This essay will therefore consist of two sections: the first, relating to facts demonstrating the existence of a great variety of reflex and other remote effects of irritation of centripetal nerves; the second, giving the principal features and rules of diagnosis and treatment of these effects of diseases and injuries of branches and trunks of nerves.

SECTION I.—AFFECTIONS OF THE NERVOUS CENTRES AND OTHER ORGANS, CAUSED BY AN INJURY TO, OR A DISEASE OF, A NERVE.

Epilepsy.—Of all the nervous and other complaints that may be due to an irritation starting from the trunk, branches, or ultimate ramifications of nerves, very few, if any, are more frequent than epilepsy. Diseases of all the mucous membranes, or their irritation by worms, diseases of the cerebral or spinal meninges, dentition, &c., are known to be frequent causes of this convulsive affection. But it is not so well known that an injury to, or a disease of, a nerve, not rarely produces epilepsy. Indeed, even a man of great authority as regards epilepsy, Dr. Herpin,* has lately maintained that in cases in which this affection seemed to have been caused by an external injury, it was, in reality, due to a disease of the nervous centres. It is most important that medical practitioners should not be misled by such an opinion. The rational treatment of that form of epilepsy which appears after a wound, or an organic affection, of a nerve, would certainly be neglected if such a view were admitted. It is, therefore, necessary to demonstrate that in a number of cases an epileptiform affection and also the most genuine epilepsy have been caused by a lesion of a nerve. This demonstration is given by the following kinds of proof:—

1st. In a great many cases, epilepsy has appeared in persons in whom there was no other cause for its production but a wound, a burn, a tumour, an inflammation, or a neuralgia.

* *Des accès incomplets d'épilepsie*, par Th. Herpin, p. 36. Paris, 1867.

2nd. In a number of the above cases a peculiar sensation, generally mis-called aura epileptica, arose from the seat of the irritated nerve before all or most fits.

3rd. In many of the same cases, the application of a ligature round a limb above the seat of the irritation often prevented the occurrence of fits.

4th. In some of the same cases a pressure on the seat of the external irritation invariably brought on a fit.

5th. In many cases of epilepsy apparently due to an irritation of a nerve, the section of that nerve above the seat of irritation, or the amputation of the limb; the extirpation of a tumour, of a cicatrix, of a decayed tooth, of a carious bone, &c., have cured the patients.

6th. I have discovered, in certain animals, that the irritation of the sciatic nerve by a broken bone, or by some other causes (crushing, tying, or cutting), invariably produces a temporary or persistent epilepsy. This convulsive affection never disappears without treatment, unless the irritation has ceased in the sciatic nerve.

These various facts clearly prove that epilepsy may be due to an irritation of a nerve, and exist without any serious organic change in the nervous centres. For the details of cases like those I have mentioned I will refer the reader to the new edition of my work on epilepsy, which will soon be published.

I will only say here, that some of these cases have been observed by perfectly reliable men, such as Sir Astley Cooper, Sir Benjamin Brodie, Dieffenbach, Baron Larrey, &c. I will give one of those cases as a good specimen of reflected epilepsy. It was put on record by Dr. W. Laing of Aberdeen.

M. D., aged twenty-one, had the left hand lacerated by machinery. She went on well till the night of March 6, when she was seized with convulsions, and after a day or two, with trismus and other tetanic symptoms. On April 7 she was dismissed, *cured*; but on June 24 she returned to the hospital. About a fortnight before her readmission she suddenly fell down in an epileptic fit; and since, the attacks have become more and more frequent, recurring five or six times a day, and lasting about five minutes, after which she remained a considerable time in a state of stupor. On the 25th the fits were so severe that she was put in the strait-waistcoat. When the fits were slight, they were confined chiefly to the injured arm. On touching the fingers smartly, the arm was convulsively withdrawn; and when this was done while she was lying in a state of stupor, violent convulsions of the arm were produced. The patient often felt a sensation arising from the injured hand, previous to her fits. As the remainder of the hand was of little use, the fore-arm was amputated: the patient never had the slightest appearance of epilepsy after the operation, and was dismissed

cured, a month afterwards. The digital branches of the median nerve, and branch of the ulnar, were found enlarged to four or five times their size, and their extremities bulbous, and firmly embedded in a hard cicatrix.

Tetanus.—Referring to the article on TETANUS in the volume of this work, I will only say a few words on the important questions relating to the nature and to the local treatment of this affection.

I am really surprised that some persons still doubt that, owing to a peculiar influence exerted, on the circulation of blood and the nutrition of the spinal cord and the medulla oblongata, by the irritation of a centripetal nerve, that tetanus arises from a traumatic lesion. I hope that the following facts and reasoning will show that this convulsive affection is dependent upon an irritation arising from the injured nerve, and not—as Dr. B. W. Richardson, Roser, Billroth, and others are inclined to admit—from toxæmia.

The relation between the wound and tetanus seems to be positively established—at least, in those cases in which the muscles attacked with spasms are on the side injured. Letourneau, Sir Gilbert Blane, Swan, Dupuytren, and Mr. Currier, who cites the preceding authors, have seen such cases. My friend Dr. G. H. B. Macleod † relates two cases of fatal tetanus in which the tetanic spasms were almost entirely limited to the side injured. Baron Larrey states that when the wound causing tetanus is in the anterior part of the trunk, emprosthotonos is the form generally observed.

Other strong arguments in favour of the view that traumatic tetanus is caused by a peripheric irritation, are: 1st, that very frequently the muscles in the neighbourhood of the wound are either the first attacked or the most affected with spasms; 2nd, that in many cases it has been observed by myself and others, that even a slight pressure on the wound or the cicatrix has increased the spasms or produced them, in periods of relaxation (after chloroformic anaesthesia, for instance).

It may seem strange that tetanus will sometimes follow even the slightest wound, and that it will come at any period of inflammation or cicatrization, and even when there is no pain at all in the wound or its neighbourhood. But this last fact is

* Aberdeen Infirmary Reports, in *Lond. Med. Gazette* for Dec. 25, 1840.

† *A Treatise on Tetanus*, pp. 87, 174. London, 1836.

‡ See his excellent work, *Notes on the Surgery of the War in the Crimea*, pp. 155-161. 1858.

ses of wounds of all varieties, there were 18 cases of ; giving a proportion, when the nerves were simply l by a sharp knife, of one case of tetanus out of 1,364 s; and of one case out of 55 patients, after wounds and es, when the nerves were bruised or much irritated. In stical table given by Dr. Friederich,* the influence of rritation of nerves in causing tetanus is also demon- : out of 176 cases of that affection, only 11 occurred mputation, and 33 after gunshot wounds, while 61 were contusion or comminutive fracture, and 71 to wounds eturing instruments, or dilaceration and bruising of by nails, pieces of wood, &c. Dr. Lawrie's statistics with those of Friederich.†

cases of cure of tetanus either by an amputation of a r by section of a nerve, clearly prove the dependence of fection on an irritation starting from some peripheral f a nerve. The same conclusion flows out from numerous ke the following, to which I might add many others. as is pointed out to have been caused by a small splinter e sticking in the radial nerve (Hennen), by a portion of p embedded in the cubital nerve (Dupuytren), by an ap- ion of caustic potash on the coraco-brachial nerve (Frere), t corns in the tibial nerve (J. Hutchinson), by a splinter od in the radial nerve (Morgan), by neuritis (Curling, letier, Ervion, and others) by ligatures on nerves of

fragments of broken bone (Alquié), and by partial division tearing of nerves (Swan, Liston, Billroth, &c.). In cases in which the spinal cord was either inflamed or rendered tremely excitable, the section of a nerve or an amputation been of no avail; and in some cases, also, in which an inflammation had been propagated high up in the trunk of a nerve towards its roots, these operations have been useless; such failures might have been avoided had these more treatment been applied earlier, and had all the nerves divided higher up than they have been. I need not say the simple division of a nerve should always be preferred to an amputation, unless there are some special reasons for the last operation.

Hysteria.—The extreme frequency of this affection in wounds renders it difficult to prove that it may be due to an irritation of a nerve. However, there are cases in which it seems clear that hysteria was really caused by a wound, or the formation of a tumour.*

Two very interesting cases are reported by Dr. Parsons; † one observed himself, the other by Dr. S. P. Hildreth.

Morgagni mentions the case of a young girl, who, after a wound to the neck by the biting of a sparrow, was attacked with fits of trembling and recurring sixteen or eighteen times a day. ‡

Raynaud relates the case of a woman, who, after having received a wound of the breast, had a first attack of hysteria. Two small tumours soon appeared at the injured place, and for seven years hysterical attacks occurred several times every day. These tumours were removed by Boyer, and immediately after the operation the attacks ceased, and did not recur again. §

Three years ago, in a patient of mine of a highly nervous temperament who never had had any marked symptoms of hysteria, convulsions, delirium, with some degree of lock-jaw, frequently appeared and during three or four days, after a slight wound by a needle in the knee-joint, just below the patella. The needle broke at the time of the accident, and a small part of it remained under the skin. As soon as the needle was removed from the little wound, after the extraction of the point of the needle the patient got well, and has had no return of hysterical symptoms since.

* An interesting case, published by Hamilton (*Dublin Journal*, 1838, p. 42), shows how careful we must be as regards the signification of hysteria in hysterical patients. All the most characteristic features of neuritis in a patient after a wound, but they disappeared at once after a violent attack of hysteria.

† *American Journal of the Med. Sciences*, April 1851, pp. 307, 312.

‡ *De Sedibus et Causis Morborum*, Lutetiæ, 1822, vol. vi. p. 61. liv. § 45.

§ Raynaud, in *Archives de Médecine*, 1829, vol. iii. p. 434.

these facts, and the cases I have already mentioned, that hysteria, like epilepsy and tetanus, may be caused by irritation of a nerve. I must say, however, that a persistent hysteria is very rarely obtained. Various operations, as extirpations of cicatrices, amputations, &c., to cure local spasms due to an irritation of nerves, have proved only for a time, in several cases of Sir Benjamin Brodie, of Mr. Hancock, Tyrrel, and Bransby Cooper. ||

Catalepsy.—I have seen a case of this affection in which it was brought on at once by even a slight pressure on the spots between the shoulders. Catalepsy, therefore may, like the neuroses, be produced by a peripheric irritation. ¶

St. Vitus's dance.—St. Vitus's dance may be caused by an injury to a

case of Dr. Borelli, of Turin, chorea was caused by a neuroma of the arm of a child thirteen years old. This convulsive affection was at once cured, and in four days cured, after the extirpation of the neuroma.** Dr. Borelli mentions a case of chorea caused by the irritation of a finger by a splinter of a red nail.†† Dr. J. Malden has cured a woman of choreic movements after the extraction of a decayed tooth.‡‡

Drophobia.—I will try to show elsewhere that the symptoms

traité de l'Hystérie, 1847, p. 253.

Lectures illustrative of certain Local Nervous Affections, 1837, pp. 40–46.

Leçon inaugurale, soutenue le 20 novembre, 1855. Paris.

oc. cit. p. 83.

of this terrible affection depend on a local effect of the virus on the wounded nerves, and that some chance of cure might be obtained by division of the irritated nerves. Already in the last century, G. Hicks proposed this means of treatment against hydrophobia.* I owe to Dr. Stokes of Dublin, the mention of a most important case, showing that there is good ground to hope that hydrophobia might sometimes be cured by amputation or division of a nerve.†

Tremulous movements.—The so-called *trembling palsy*, which so often consists simply in involuntary tremulous movements, without any palsy, may be caused by an irritation starting from a nerve.

Sabatier‡ relates the case of a young man, who, after a wound of the saphenous nerve, near the knee, was attacked with violent trembling of the leg and thigh, which lasted many months. In a patient sent to me by Mr. Erichsen, an injury to a nerve of the left arm has produced trembling in both arms. In another patient, for whom I was consulted by Mr. M. H. Collis of Dublin, shaking palsy began in a fractured limb, and thence extended to the other limbs. Several cases are on record showing that the shaking due to an external injury may become general.§

Rotatory convulsions.—I have found that an injury to the auditory nerve in animals is at once followed by rotatory movements. I do not know of any case of wound of that nerve in man having produced the same symptoms; but several cases are on record in which these movements have been observed in man when the auditory nerve was irritated by an inflammation or some other cause (an injection of caustic, &c.).

I would refer for these cases to my work on the *Central Nervous System*, p. 195, only adding here, that since that publication I have seen three cases of that peculiar kind of involuntary movements, caused by an affection of the internal ear. An irritation of other nerves may produce the same effect. Dr. Krieg relates the case of a patient wounded on the forehead, and on whom a touch of the injured skin produced attacks of exceedingly rapid rotatory movements.||

Local convulsions.—It is well known that wounds of the branches of the fifth pair of nerves may by a reflex influence pro-

* *Lond. Med. and Phys. Journal*, vol. xvii. p. 277.

† The case above mentioned, and the reasons I have for the hope I have expressed, will be found in the Appendix to my work, *Lectures on the Physiology and Pathol. of the Central Nervous System*, p. 261 et seq. 1860.

‡ *Médecine opératoire*, vol. i. p. 254.

§ *Inquiry concerning Constitutional Irritation*, by B. Travers, p. 115, 1836; and *Treatise on Diseases and Injuries of Nerves*, by J. Swan, p. 124, 1834.

|| *Histoire de la Musculation irrésistible*, par le Dr. Roth, p. 78. Paris, 1806.

ther nervous affections a reflex vascular contracture frequently occurs.

A prominent American clergyman has given me the details of a most remarkable case of prolonged reflex contracture of blood-vessels, he himself being the

After having violently struck his leg (two inches above the knee) a piece of paling-fence, he was soon apparently cured of the local injury; influence upon the blood-vessels of the limb showed itself; so that for a winter it remained extremely cold, and ever since, for twenty years, its temperature has been lower than that of the other leg. Swan † has also observed a case of permanent reflex contracture of blood-vessels. I will revert to this subject in speaking of reflex muscular atrophy.

Paralysis.—I will not enter here into the discussion of the question of the mode of production of paralysis when caused by irritation of a nerve. I will only say that there are two entirely different kinds of paralysis produced by such an irritation: in one kind a congestion or even an inflammation takes place in a part of the nervous centres by an influence derived by the peripheric irritation, and the paralysis then proceeds accompanied by the usual symptoms of congestion or inflammation of some part of the brain or spinal cord; while in the other kind the paralysis exists without any symptom of congestion or inflammation of the nervous centres. This second kind of paralysis caused by an irritation of a nerve is most likely due, at least in some cases, to a reflex contracture of blood-vessels ‡

of my researches on this subject, with my friend's Dr. Tholozan (*Journal*

in a part of the nervous centres. In some cases, if not in all, it may also depend, however, on a peculiar influence exerted on nerve-cells, in these centres, by an irritation starting from peripheral nerve-fibres, producing what has been called *inhibition*, or, at any rate, changing the condition of activity of these nerve-cells.

Paralysis of the various muscles of the eye, including the iris, is pretty often observed in cases of wound of the infra- or supra-orbital nerves, or in cases of neuralgia. I have seen several cases of that kind of paralysis (caused by a neuralgia), and all characterised by their evident relations with that cause.

In a case, which I have carefully watched, a sprain of *one arm* at the elbow-joint soon produced a paralysis of *both arms*, but more marked in the uninjured arm than in the other. Every change in the degree of pain in the injured elbow was accompanied by a corresponding change in the degree of the paralysis. The pain has now ceased for many years, and the paralysis, which ceased with it, has not reappeared, the two arms having at present as great power as they ever had before the injury.

Baron Larrey states that almost all the men who received slight wounds of the shoulder in the Syrian campaign, were attacked with paralysis of the injured limb. These patients were cured in Egypt, where the air is purer than in Syria. In some, at least, of these cases, the paralysis was clearly due to an influence exerted on the nervous centres by the irritation of superficial nerves.*

Facial paralysis from a neuralgia is not rare; in one case this paralysis and neuralgia were both caused by the irritation of the infra-orbitalis nerve, and cured by the extirpation of the irritating cause, a piece of porcelain. (Jeffrey, quoted by Tillaux, *Des Affections chirurg. des Nerfs*, p. 15; Paris, 1800.) Fabricius Hildanus, quoted by J. Barthez (loc. cit. p. 83, vol. ii.), relates a case of paralysis of one arm, caused by a piece of glass in the ear. The arm may be paralysed by an influence arising from very distant nerves. Drs. S. W. Mitchell, Morehouse, and Keen,† give two interesting cases, in one of which the irritated nerve was the sciatic, and in the other the crural.

Parsons mentions a case of paralysis of the face and arm, caused by a prick.‡

Roche, in an able dissertation, relates the case of a physician, in whom general convulsions and afterwards complete paralysis of sensibility and motion

nervous centre, he vainly looked for blood-vessels on the surface he had under his eyes. Had he waited some time, he would not only have seen blood-vessels appear, where he at first had not seen any, but he might have had the proof that there is a contraction of the blood-vessels of the surface of the spinal cord, powerful enough to render them almost invisible, and occurring when the incito-motor nerves of the skin and other parts of the back are strongly irritated in the operation of laying bare the cord.

* *Mémoires de Chirurgie militaire*, 1812, vol. ii. p. 153.

† Circular No. 6, *Reflex Paralysis*. Philadelphia, 1864. See also a paper on 'Paralysis from Peripheral Origin,' by Dr. S. Weir Mitchell. New York, 1866.

‡ *Amer. Journ. of the Med. Sciences*, April 1861, p. 310.

or more extensive paralysis are also reported by the three American I have named. Schenkus (Barthez, loc. cit. vol. ii. Notes, p. 41), has case of general paralysis produced by a wound of the eyebrow. I have consulted by an American officer, who became paralysed in a slight of the four limbs, chiefly the right arm and the left leg, from a gunshot of the cervical plexus, and partly also of the brachial plexus of the

seen a most interesting case of general paralysis, caused by an of the nerves of the penis, cured by the successful treatment of and the operation for phimosis.

iplegia from peripheric irritation is less frequent than paralysis or paraplegia.

are, however, some remarkable cases on record. In one of them, I by Dr. Shearman, there was hemiplegia of the right limbs, caused ouloureux of the right inferior axillary nerve. Tonics and galvanism e patient.† In another case, reported by Baron Larrey, a lady was with hemiplegia on the same side where she suffered from a facial t, the hemiplegia being more evident during the attacks of neuralgia; ctions were cured by moxas.§

collected thirty-seven cases in which there was hemiplegia, due to tion of either the auditory or the trigeminal nerves near their origin, e crus cerebelli. I have tried to show elsewhere that, in those cases, dyis which exists in the limbs on the side of the irritated part, is pro- to a reflex influence.||

they Cooper mentions the following fact: 'Mr. Toulmin attended a account of her suffering severely from a diseased tooth, and she d also to be afflicted with hemiplegia. Mr. Toulmin extracted the and in a short time the paralytic affection entirely subsided.' ¶

* *Accidents nerveux traumatiques*, p. 65. Thèse. Paris, 3 janvier 1861.
Annales médicales de Paris 1834 n. 358

Anæsthesia.—A reflex anæsthesia is not rare in cases neuralgia.

Many cases of reflected anæsthesia from a wound and other injuries of nerves are on record. A remarkable case of Baron Larrey (loc. cit. vol. v. p. 1821), and an important one of Roche (quoted in my *Lect. on the Centr. Syst.* p. 131), particularly deserve to be mentioned. I have observed anæsthesia of the arms in an able lawyer of London, which was caused by a blow on the back of one knee. Several interesting cases have been published by Drs. S. Mitchell, Morehouse, and Keen. In one case a shell-wound of the left arm produced anæsthesia of the right thigh.* I have several times seen anæsthesia of the whole of one side of the face, in cases of neuralgia of one of the trigeminal nerve on the same side. I have seen also a case of anæsthesia of a part of the forehead and face, in consequence of the irritation of a branch of the fifth pair, on the cheek-bone, by a bruise. In those cases the anæsthesia subsided when its cause was cured.

Amaurosis.—The cases of amaurosis due to an irritation of one of the branches of the trigeminal nerve are frequent enough for my dispensing with quotations of cases. In the first edition of this work I quoted cases of Wardrop,† Notta,‡ Dr. Noyes of New York, Mr. J. Hutchinson.|| This last observer has since published several able papers on this subject. I have myself seen five or six cases of amaurosis evidently due to an irritation of the supra-orbital nerve. Cases of amaurosis due to an irritation of a nerve of the trunk or limbs are not frequent. Dr. Colhoun has reported a remarkable case of sudden and complete amaurosis of both eyes, caused by a gun-shot wound of the scapula.¶

Deafness.—A neuralgia of the face sometimes produces deafness; other kinds of irritation of branches of the fifth nerve may also cause deafness. Dentition and decayed teeth have been pointed out as having had the same effects. Pearson has reported a case of wound of the thigh in which deafness was one of the symptoms produced (*Medical Facts*, vol. vi. p. 109).

* See an excellent paper by Dr. S. Weir Mitchell, on 'Paralysis from Peripheral Origin,' reprinted from the *New York Medical Journal*, 1860. I have lately found that the irritation of the sciatic nerve in the lower extremity produces a slight anæsthesia of the lower limb on the opposite side, at the face and neck on the side of the irritation.

† *Med.-Chir. Trans.* vol. xii.

‡ *Archives gén. de Med.-Chir.* etc., juillet 1854, pp. 12-21.

§ *American Med. Times*, March 15, 1862.

|| *Med. Times and Gazette*, May 7, 1859.

¶ *The Medical Examiner*, p. 806, vol. ii. Philadelphia, 1839.

d by a peripheric irritation on the nervous centres of the able to act on circulation, secretion, and nutrition, through that influence, producing a cessation of some of dinary interchanges between the blood and the tissues. pposite conditions of the blood-vessels may exist in this form of collapse; these contractile tubes may be dilated racted. But in one or the other of these conditions, the instead of being black, in the veins, is reddish and some- arterial-looking, and the production of heat ceases in the ries. This third form of collapse, consisting, as I have a a cessation of the ordinary chemical interchanges be- the tissues and the blood, is the most prominent in a number of cases, and it is also the most dangerous of the of shock or collapse.*

of smell, taste, and hearing.—The celebrated experiment of die, showing that a section of the trigeminal nerve may a loss of the five senses in the head, is, most likely, explained by a reflex influence. At any rate, an important f Dr. Blondlot shows that an irritation of the infra- lis nerve (wounded) may produce, by a reflex action, loss ring, taste and smell.†

ralgia.—The frequency of *tic douloureux*, caused by an ion of a small part of the dental nerves, is such, that it is nly useless to mention cases. But *tic douloureux* may be d by irritations of other nerves, while also an irritation of ss of the jaw may cause a neuralgia elsewhere than in

pressure on the tumour brought on a severe attack of tic. patient was cured by the removal of the tumour.*

In the first edition of this work, I gave cases of neuralgia reported by Mr. Gay,† Dr. R. Rowland,‡ Parse Tavignot,|| Dr. Greene of New York,¶ Wardrop,** Maréchal Dr. Castle of New York,†† Marchal de Calvi,§§ and Romberg. The view I then maintained, that a neuralgia may be caused by a reflex action, being now pretty generally admitted, I give here a reference to these cases, and to important publications made on this subject by Ch. Londe,¶¶ J. Mason Warrar and Ch. Mauriac.†††

Delirium.—I will simply mention here three cases which show quite decisively that delirium may be caused by an irritation to a nerve.

A boy, aged fourteen, trod on a piece of glass, which penetrated the foot but was removed. Four years after, he began suddenly to talk in a strange, wild way; true delirium set in, and nothing appeased the patient. Near the ball of the big toe a small reddish elevation was found. The pressure was made upon it, the seizure returned with violence. An incision was made, and a trifling piece of glass was removed. Much as the patient had during the operation, with equal suddenness did all the symptoms vanish. He was surprised on being told of all the senseless things he had uttered.

I have published a case very similar to this, which I owe to the kindness of Mr. Campbell de Morgan, and in which the attacks of delirium took place when pressure was made on a wound of a toe containing a foreign body. Soon as the irritated part was taken away by the cut of a bit of skin

* *On the Nature and Treatment of Tic Douloureux, &c.*, by Dr. Henry Williams, p. 114. 1854.

† *The Lancet*, 1846, vol. ii. p. 119.

‡ *A Treatise on Neuralgia*, by R. Rowland, p. 18. 1838.

§ *American Journal of the Med. Sciences*, Oct. 1854, p. 423.

|| *Gazette médicale de Paris*, 1845, p. 547.

¶ *Dublin Journ. of Med. Sc.* 1838, vol. xiii. p. 53.

** *Trans. of the Med.-Chirurg. Soc.* vol. viii. 1817, pp. 246 et seq.

†† Case of Maréchal, cited by Marchal de Calvi, in *Annales de Chirurgie*, 1844, vol. iv. p. 69.

†† *The Lancet*, 1846, vol. ii. pp. 266, 267.

§§ *Annales de Chirurgie*, loc. cit. p. 76.

||| *Lehrbuch der Nervenkrankheiten*, 3rd edit. vol. i. pp. 23–35.

¶¶ *Rech. sur les névralgies consécut. aux lésions des nerfs.* Paris, 1860.

*** *Surgical Observations, with Cases and Operations.* Boston, 1867, p. 471.

††† *Névralgies réflexes de l'orchite*, in *Gaz. méd.* Paris, 1869, Nos. 1 and 1870, Nos. 1 to 5.

††† Joerdaens, apud *Hufeland's Journal*, vol. iv. p. 227, cited by D. Payne, in his *Medical and Physiol. Commentaries*, vol. i. p. 425.

ient became rational, and remained so when pressure was made on the and.*

Mr. Sherwin has seen a woman, who, after having been bled, was attacked with pains in the arm, neck, and face, with spasms in those parts, and delirium. After the symptoms had continued a fortnight, a deep incision above the scapula quite cured her.†

I need not speak here of the delirium that follows amputation or other great operations. The causes are many that bring on delirium in such cases. Among the principal causes I will point out a great loss of blood, and the anxiety of the patient.

Aphasia.—An interesting case shows that this affection also can be produced by an irritation of a nerve. Dr. Guyot has seen aphasia occurring every time an attack of facial neuralgia took place; the patient was cured by quinine.‡

Inflammation.—Cases of inflammation of the eye due to a reflex action are so frequently met with, that I need not stop here to prove their existence. Any one who will read the facts published by Dr. W. Mackenzie, in his admirable work on the Diseases of the Eye, by Mr. R. Taylor,§ by Dr. Brondeau,|| and by several other more recent writers, among whom I will only quote Dr. J. Maats,¶ a pupil of Prof. Donders, will soon be convinced that an inflammation of any part of one eye (the retina, the cornea, the conjunctiva, &c.) may be caused by a wound or an inflammation of the other eye; and that if the first diseased organ is extirpated, the other is often soon cured. Cases of ophthalmia owing to a wound or a neuralgia of the infra- or supra-orbital nerves, or caused by an irritation of the dental nerves, are also not rare. Dr. Busschaert has published a curious case of ophthalmia produced by obstruction of the external auditory canal.**

Dr. Rowland relates several facts which seem to prove that an inflammation in one side of the brain may be caused by an injury to a nerve in the other side of the body.†† To the cases of

* *Course of Lect. on the Physiol. and Pathol. of the Nerv. Cent.*, p. 185. 1860.

† Duncan's *Medical Comment.* vol. iv., cited by Mr. Hamilton in *Dublin Journ. of Med. Science*, vol. xiii. p. 51, 1838.

‡ *Gazette hebdomad. de Médecine*, 1867, p. 266.

§ *Medical Times and Gazette*, 1857.

|| *Des Affections sympath. de l'un des Yeux.* Paris, 1858.

¶ *Nederlandsch Archief voor Genees- en Natuurkunde*, Deel ii. 1^o Aflevering, p. 8-52. Utrecht, 1865.

** *Gazette des Hôpitaux*, 10 octobre 1857.

†† *On the Nature and Treatment of Softening of the Brain*, p. 67 et seq. London, 1851.

inflammation of the brain mentioned by Dr. Rowland, I might add several others, among which the most significant have been recorded by Hennen,* R. Bright,† and Champsaur.‡ The well-known fact that sometimes, in traumatic tetanus, the spinal cord becomes inflamed, shows that a peripheric irritation may produce inflammation in that organ.

Inflammation of the testicle is also sometimes produced by reflex action from an irritated nerve, as in cases by Sir Benjamin Brodie,§ Barras,|| Marrotte,¶ and others. Sir Astley Cooper says that by irritation morbid actions are excited in distant organs, and adds: 'thus inflammation is produced in the testis from irritation in the urethra.'

Inflammation of the abdominal or thoracic viscera may also be produced by a reflex action. Proofs of this assertion are abundantly furnished in Lecture X. of my work *On the Central Nervous System*. A reflex inflammation may be brought on to such a degree as to cause an ulcer, which nothing can heal until the cause (viz. the irritation of a nerve) is removed. Sir Astley Cooper mentions several cases of that kind.††

As well shown by Mr. J. Hamilton, there is sometimes a deceptive appearance of inflammation with suppuration.‡‡

Coma.—This most dangerous morbid state may also be caused by a peripheric nervous irritation.

In a case of Hirsch, quoted by Dieffenbach,§§ convulsions and coma accompanied local neuralgia, caused by venesection. The patient was cured by deep incisions over the wound. An immediate cure of coma and convulsions was also obtained in a case similar to the preceding, observed by Dr. Wilsch. Another case somewhat similar, has been recorded by Mr. G. Bell.¶¶

* *Military Surgery*, p. 191.

† *Reports of Medical Cases*, vol. ii. pt. i. p. 52.

‡ *Thèse inaugurale*, p. 22. Paris, 1860.

§ *Lectures on Local Nervous Diseases*, p. 16. 1837.

|| Cited by Notta, in *Archiv. de Méd.* etc., p. 547, sept. 1854.

¶ *L'Union médicale*, p. 155, 1851.

** *Lectures on the Principles and Practice of Surgery*, by F. Tyrrell, p. 4. 1824.

†† *Loc. cit.* pp. 7, 8.

‡‡ *Dublin Journal of the Medical Sciences*, vol. xiii. pp. 50, 55. 1838.

§§ *British and Foreign Medical Review*, p. 332, vol. xxi. 1846.

||| *A Treatise on Diseases and Injuries of the Nerves*, by J. Swan, p. 1834.

¶¶ J. Swan, *loc. cit.* p. 119.

be explained by the fact that that nerve sends a branch to two muscles thumb, as not only these two muscles, but others, and especially the r and the opponens, were notably atrophied.

neuralgia very often produces atrophy in neighbouring es. My friend and pupil, Dr. Cl. Bonnefin,† has seen en cases of muscular atrophy caused by a neuralgia. A kable fact, observed in those cases, serves to explain how trophy was produced—there was a marked diminution of rature, most likely due to a spasm of blood-vessels. The it of blood was consequently diminished, and the wasting ed owing to the lack of nutritive fluid. Some of the of wasting palsy related in Dr. Roberts' excellent work,§ kely belong to the class of reflex atrophy.

a case recorded by Vallez, a wound of the infra-orbital produced a reflex atrophy and paralysis of the face on me side.||

rophy of the cellular tissue.—In some of the cases of atrophy e side of the face, which Schott and Romberg have called trophoneurose, there was an irritation of some sensitive , probably acting by a reflex influence. I saw a case of rare affection, three years ago in Boston (United States); obable cause was an irritation of the dental nerves, which oced convulsions in one side of the face (the side where the hy was afterwards observed). I will refer for arguments

† cited by Marchal de Calvi, in *Annales de Chirurgie*, vol. x. obs. 7, p. 73.

leading to the conclusion that the cellular tissue is alone atrophied in that affection, to a remarkable essay by Dr. L. Lande.*

Hypertrophy.—Notta mentions cases of hypertrophy of the face and tongue caused by neuralgia.† I have seen a case of considerable hypertrophy of the bones and of the cellular tissue, in the face, which occurred after repeated attacks of neuralgia, in a lady whose general health was excellent.

Various kinds of alterations of nutrition and secretion.—The number of facts that might come under this head has considerably increased since the time that my friend Mr. James Paget‡ first showed how great may be the direct and the reflex influences of the nervous system in disturbing nutrition. *Eruptions* of various kinds (*erythema*, *pemphigus*, *urticaria*, *acne*, and especially the different forms of *herpes*) are often produced by a reflex influence from an irritated nerve, as proved by facts observed by Rayer, G. Simon, Delioux, Notta, Romberg, Hasse, Parrot, and more recently by Charcot, Bärensprung, J. Hutchinson, H. F. Damon, Purdon, and others. I have seen several cases proving clearly the production of certain eruptions by a reflex action. Sir Astley Cooper relates a curious case of *fungoid granulations protruding through an ulcer* in the cheek of a lady, who was quickly cured after the extraction of a tooth.§ In a case of *ulcerations* and wasting, probably due to neuralgia, Dr. Hooker cured the patient by dividing the popliteal nerve.|| An *erysipelatous redness* and swelling has been seen by Sherwin in a case of wound of a nerve, and a *swelling* of the foot and leg has been observed by Dr. Watson of New York, in a case of cut of the sole of the foot by a piece of glass.¶ *Edema* is a frequent reflex effect of neuralgia. Hamilton has seen two cases of that serous effusion after injuries of nerves.** Those persons who know that even *gangrene* may be caused by an influence of the nervous system, as rendered so probable by Dr. M. Raynaud,†† will not be reluctant to admit that it may be the result of a reflex influence

* *Essai sur l'Aplasia lamineuse progressive.* Paris, 1869.

† *Archives de Médecine*, juillet 1854, pp. 311-12.

‡ *Lectures on Surgical Pathology*, edit. of 1853, vol. i. p. 44.

§ *The Lancet*, 3rd edit. 1826, vol. i. p. 27.

|| *The Lancet*, vol. ii. 1859, p. 336; and *The Brit. Med. Journ.* Dec. 1866, p. 730.

¶ The cases of Sherwin and Watson are cited by Mr. Hamilton (*Dublin Journal of the Med. Sc.*), vol. xiii. pp. 51, 54, 1838.

** *Dublin Journal of the Med. Sc.* vol. xiii. pp. 41 and 43, 1838.

†† *De l'Asphyxie locale ou gangrène symétrique des Extrémités.* Paris, 1862.

from irritated nerves. Most likely it was not to embolism or to thrombosis, but also to a reflected nervous influence, that gangrene was due in three cases related by Sir William Fergusson,* Dr. Gubler,† and Dr. Grainger Stewart.‡ The case of Dr. Stewart is especially worthy of attention. *Altered sensations* are very often due to a reflex influence in cases of neuralgia, and sometimes in cases of wounds of nerves. *Diabetes* is most likely produced by a reflex influence when it comes after a peripheric injury. It may be objected, however, that concussion of the brain is then its constant cause. I have not room enough to discuss the question here, but I think that the possibility of a reflex origin to mellituria is clearly established by cases like those reported by Dr. W. R. Hill, in which a burn was the cause of the secretion of sugar.§ Cases of alteration of hair due to neuralgia or injuries of nerves are not rare, and I could easily mention many, showing changes in colour, in thickness, in abundance, and in rapidity of growth of hair.|| *Cataract and glaucoma* have been pointed out also as evidently resulting, in some cases, from an irritation of nerves.¶

SECTION II.—GENERAL FEATURES AND RULES OF TREATMENT OF THE VARIOUS AFFECTIONS CAUSED BY AN IRRITATION OF A NERVE.

The following features usually characterise cases of neuralgia, paralysis, epilepsy, and other affections brought on by a peculiar influence, exerted upon, or through, a nervous centre, by an irritation of a nerve.

1st. Previous to the appearance of a remote affection due to such an irritation, the patient has suffered for a variable time from a neuralgia or a neuritis, from a wound or a burn, or from pressure upon a nerve, by either a tumour, a displaced bone, or a foreign body.

* *The Lancet*, vol. xiii. p. 152, 1850.

† *Comptes rendus de la Société de Biologie*, pour 1854, p. 76.

‡ *The Medical Press and Circular*, Jan. 10, 1866.

§ *Beale's Archives of Medicine*, vol. ii. p. 172.

|| I found lately that in some animals the division of the sciatic nerve is almost always followed, in two or three months, by a fall of hair in the neck on the side of the lesion.

¶ *Gazette des Hôpitaux*, 1846, p. 1; *Gazette méd. de Paris*, 1840, p. 130, and 1845, p. 546; and De Brondeau's dissertation, *Des Affect. sympath. de l'un des Yeux à la suite d'une blessure de l'autre œil*, pp. 40-46. Paris, 1858.

2nd. An increase or a decrease of the irritation of a nerve is often followed by corresponding changes in the intensity of the remote affections caused by the peripheric nervous irritation.

3rd. The various modes of treatment of nervous and other affections, produced by an influence exerted on the nervous centres by a peripheric irritation, are generally quite unsuccessful so long as this irritation persists unabated.

4th. The various affections produced by a peripheric nervous irritation are frequently cured or relieved at once, or very soon after the removal of their cause, viz. the irritation.

I may add a few other characters, more or less implied, however, in the preceding: 1st. When remote affections due to a peripheric nervous irritation occur by fits, it is not rare to see the fit suddenly produced (completely or incompletely) when the diseased nerve is irritated by pressure, or otherwise (application of galvanism, for instance). 2nd. Narcotics, applied to the diseased nerve, will very frequently diminish, at least for a time, the remote affection, even, sometimes, when it consists in, or is connected with, a notable alteration of nutrition.

The above characters may all serve for the diagnosis of remote affections caused by a peripheric nervous irritation; but the only essential one consists, of course, in the pre-existence of a lesion of a nerve. It must be remembered, that if the trunk of a nerve is inflamed, all the symptoms spontaneously mentioned by the patient may seem to him to exist only at the terminal ramifications of that nerve. There is but one way to ascertain what the starting-point of these symptoms is: it consists in the examination, by pressure, of as much as possible of the whole length of the nerve, from the periphery to the neighbourhood of the brain or spinal cord. Had this rule been applied in the following case, it would not have been published and accepted as a case of reflected influence from a disease of the nerves of the thumb upon the four limbs.

Lady — was attacked suddenly by an acute pain, soon followed by redness and swelling in the left thumb; and the other fingers gradually were also attacked, and afterwards the fore-arm. There was contracture and paralysis, with hyperæsthesia. The other arm became affected in a similar way; and when the pain was violent, there was paraplegia. No benefit was obtained from powerful narcotic applications on the left thumb and hand; but the patient was cured after the use of a counter-irritant ointment rubbed over the arm.*

* Case of Pearson, in *Med.-Chir. Trans.* vol. viii. pp. 252 et seq. Pearson does not speak of the spine. Had he examined it, he would have found great tenderness between the shoulders and a little above.

In this case there had been no injury to the hand; there was no neuralgia; and the symptoms observed in the fingers and the fore-arm were those we find in cases of local meningitis, or inflammation of the sheath of nerves, at their exit from the spine. I have seen five similar cases, four of which were cured by counter-irritants applied to the spine.

The wonderfully powerful and varied influence exerted by an irritation of a nerve is not due to pain, but to an action of peculiar incident non-sensitive nerve-fibres, as is well proved by two sets of facts: first, that there may be no pain, and even no sensation of any kind, in certain cases in which, however, a peripheric nervous irritation causes a neurose,* or another affection, as, for instance, in cases of worms in the bowels; secondly, that we every day see cases of pain from neuralgia, or other diseases of nerves, without the production of any remote affection. In only few cases seen by myself or others, was there such an agonising pain as in a patient, in whom a ball had lodged in the trunk of the radial nerve, producing for many days the most excruciating pain, depriving him of sleep, and causing a continued perspiration from his face, without any other marked reflex action than a contraction of the fore-arm upon the arm.†

It may seem quite surprising, and perhaps incredible, that the same cause, viz. an irritation of a nerve, will either produce no effect at all, or produce such a variety of affections as I attribute to such a cause. But those who will take the trouble of studying the variety of effects of a clear cause of reflex action, such as, for instance, the exposure of many people to a cold wind when they come out perspiring from a very warm room, will understand that reflex effects may be exceedingly various, although resulting from the same peripheric cause.

Treatment.—Of the various means of treatment of the reflex and other remote effects of the irritation of a nerve, the most important may be classed into two groups—the local and the general means. As regards the local means, they consist chiefly in applications of revulsives or sedatives, or in an amputation or division of a nerve; while the general means consist chiefly in the use of remedies that will diminish the reflex power, or the morbid excitability of the irritated nerve.

* See my *Researches on Epilepsy*, p. 17. Boston, 1857.

† Case of Denmark, in *Med.-Chir. Trans.* 1813, vol. iv. p. 48. The patient was cured by the amputation of the arm.

Local means of treatment.—Of these means, the best theoretically are also the best practically, according to the mass of facts I have collected. The section of the injured or irritated nerve between the brain or spinal cord and the part of the nerve which is altered, is certainly the most important local means. I hardly need to say, that if this operation is to be performed, the sooner the better, in cases of hydrophobia, epilepsy, tetanus, reflex neuralgia, paralysis, &c. Of course, if there is any reason to fear that the irritating cause will persist after the time necessary for the reunion of the parts of the divided nerve, an excision of an inch or two, which will retard reunion, must be made instead of a simple division. There is no doubt that in a number of cases (especially those of long duration) this operation will not succeed; and there are many discouraging facts, showing that the alteration of nutrition produced at a remote distance from the irritated nerve will continue after the division of the nerve, owing to causes yet undiscovered, or to an inflammation of the nerve in a great length between the place of the section and the nervous centres. It would be prudent always to excise at least a very small part of the length of the nerve, to ascertain, by a microscopical examination, if it is inflamed at the place of the operation; as, if such be the case, another division ought to be performed much higher up, and even as near the nervous centre as safely possible. In a paper recently published by M. Arloing and M. Tripier,* they give good reasons for the division of all the nerves of a limb in cases of tetanus; but I do not think that this radical proceeding is essential in a large proportion of cases of tetanus.† Still less would it be essential in most other affections due to peripheric irritation.

There are cases in which, instead of dividing a nerve, all that is necessary is to gain a few days to allow a wound to heal up. I proposed, several years ago, to make use, in those cases, of a simple means, consisting in laying bare the nerve above the wound, and in dropping sulphuric ether upon it. This operation, especially if ether is often applied, may render the nerve,

* *Archives de Physiol. normale et pathol.* p. 245, 1870.

† While correcting the proof of this article I received a number of the *Boston Medical and Surgical Journal* (March 31, 1870, p. 238), in which I find a case of severe traumatic tetanus, cured by exsection of the internal plantar nerve. Dr. G. E. Foster, who reports the case, states that 'no spasm of any kind' occurred after the operation.

irritated nerve, together with applications of emollient
astringent lotions, or poultices, on the wound itself, are among
the best local means after neurotomy. I have sometimes
obtained the cure of chorea, of irregular attacks of convulsions,
of neuralgia, and even twice of epilepsy, by subcutaneous
injections of narcotics (from half to two-thirds of a grain of
atropine, together with from one-sixtieth to one twenty-fifth
grain of atropine).

It has also been derived some benefit also in cases of epilepsy with a
peripheral aura, from applications of temporary circular
ligatures, like a ring, around a limb or a finger.

Applications of ice, or even sometimes of a freezing mixture,
to the spot where a nerve is wounded or irritated, might be
expected to produce a cessation of its influence on the nervous
system or another organ. Induction of local anæsthesia by
injections of ether spray just upon and above the wound might
be employed with benefit. Before dividing a large nerve,
or cranial nerves, one of these means should be tried.

Actual cautery, applied at white heat, may also be ex-
tremely useful. It is perhaps the best, and one of the least
counter-irritant means.

I need not say that foreign bodies, tumours (neuromatous and
others), or vicious cicatrices, &c., giving rise to reflex affections,
should be extirpated.

General means of treatment.—After anæsthetics, the most

tions, the most powerful narcotics, especially opium, and also other remedies, such, for instance, as the chloride of baryum, may be borne in very large doses without any poisonous effect. It would be impossible to say more as regards the general treatment without entering into details which I have not room enough to give,* and also because the rules must vary considerably according to the kind of reflex affection to be treated, and the special features of each case.

C. E. BROWN-SÉQUARD, M.D.

* I will refer for details to my work *On the Diagnosis and Treatment of Functional Nervous Affections*, the first part of which, treating chiefly of general therapeutics, was published in 1868.

HOUGH locomotor ataxy comes more frequently under the care of the physician than the surgeon, yet in the early part of its course the symptoms are so equivocal, and so liable to be mistaken for those which belong to certain surgical diseases, that for the sake of the differential diagnosis alone, a description of this malady should have a place in every treatise of Surgery.

Among its chief peculiarities are the progressive development of the most prominent symptoms at intervals of considerable time, the irregularity with which, in the order of time, some symptoms make their appearance in different cases, and, lastly, the different way in which they are grouped together. It is to these circumstances that we must attribute, in a great measure, the difficulty which frequently attends the differential diagnosis of this disease.

The symptoms observed in different cases on record are the following:—Strabismus, diplopia, amblyopia, amaurosis, ptosis, paralysis of both pupils or only of one; shifting pains in different parts of the body, chiefly in the extremities; cutaneous hyperæsthesia and loss of sense of temperature; ataxy, incoördination of voluntary movements; incontinence of urine and hæmaturia; loss of electro-muscular contractility in a greater or less degree; occasionally but rarely some paralysis of the fifth, seventh, eighth and ninth cerebral nerves; spermatorrhœa with loss of sexual power and desire; œdematous

examples, the symptoms are grouped in the following way, made their appearance in the order of time in which they mentioned.

Case I. Strabismus and diplopia; pains in the legs with numbness of ataxy or unsteadiness of gait; numbness of fingers, followed by pains in arms and unsteadiness of voluntary movements, or ataxy; incontinence of urine.

Case II. Great nervousness and external strabismus of right eye, was of both legs and of right arm; numbness in dorsum of each foot, in right as high as the wrist, and in joints of little and ring-fingers of left hand; numbness round the mouth; pains in upper and lower extremities, and in ataxy; incontinence of urine and dysuria; some impairment of smell and rachialgia, and griping in the bowels; loss of sexual power and desire.

Case III. Pains with numbness and heaviness of legs; pains in abdomen, chest; ataxy; pains and numbness in hands and arms followed by analgesia; incontinence of urine and dysuria, alternately; hæmorrhoids; sexual power.

Case IV. Hæmorrhoids, with pain and numbness in sacrum and perineum; heavy forcing pains in rectum, with tightness and weight in abdomen. subsequently pains in legs. Both pupils contracted to size of pin's head; loss of movement; loss of taste and smell; impaired sensation and motion of side of nose; great numbness of feet and legs, and analgesia; numb fingers; 'quivering' of muscles; exalted reflex excitability of skin of and legs.

In many cases the pains in the limbs are for an indefinite but sometimes for a very long, period the only precursors of other symptoms with which they are subsequently associated. They consist of two kinds—the one more or less dull, or gnawing, and frequently described by the patient as neuralgic; the other more acute and lancinating, like electric shocks. The former are more continuous; the latter are paroxysmal, suddenly, in paroxysms which last from a few hours to days, and as suddenly disappear for an indefinite period. during the paroxysm the pain is not continuous, but paroxysmal, although it may recur in rapid succession at very short intervals, and may either fix on some particular spot, or pass from one part to another with the rapidity of lightning. The parts which most frequently suffer are the lower and the upper extremities.

In other instances the ocular disturbances are for a long time the only symptoms that excite attention. These are either internal or external strabismus, or amblyopia followed by amaurosis. In more than one-half of the cases of locomotor ataxy, paralysis of either the third or the sixth cerebral ner-

ing, but more frequently it is limited to one eye. In other cases there is no perceptible strabismus, there is never diplopia or double vision, when the patient looks in any particular direction. The strabismus is not unfrequently accompanied by more or less ptosis and dilatation of the pupil. Sometimes one pupil is dilated while the other is contracted; sometimes both pupils are reduced to a very small size, there is no other apparent affection of the third nerve or other ocular nerves.

Nystagmus is occasionally one of the earliest symptoms of locomotor ataxia. It rarely disappears, or even remains stationary, but generally increases at a variable rate, and often terminates in blindness.

Ataxia, incoordination, or loss of power to control the voluntary movements, is an invariable and essential symptom, which makes its appearance in different cases at different periods of the disease. Occasionally it is first in the train of symptoms, but generally it is preceded for a variable length of time, either by peculiar shifting pains, or the ocular disturbances already mentioned. According to my own observations, which have been confirmed by those of other investigators, the disorderly movements occur under two different forms. First, they are usually manifested in the lower extremities, as simple unsteadiness of gait; the patient staggers or totters more or less, as if a person partially intoxicated. At the same time he usually complains of heaviness about the legs, of fatigue

ways on surrounding objects; but, at a later period, he cannot move without looking at his feet.

When the disorderly movement extends to the upper extremities, the patient is unable to dress himself, or button his clothes, to write, pick up a pin, or execute movements of a similar nature.

As the disease advances, another kind of disorderly movement supervenes. This is of a jerking character, and arises from spasm of the muscles, which the will puts in motion but is unable to control; for the patient has lost the power of regulating the degree of their contraction. Once excited by the will, the muscles contract spasmodically beyond the degree intended, and flex or extend the limb with a sudden and uncontrollable jerk.

These two kinds of incoordination are associated together in different degrees in different individuals; and, according to my own observations, the first kind is that which generally prevails in the early stages of the disease. At a later period, the second or spasmodic kind of disorder increases. All the voluntary movements are more or less hurried and precipitate. The patient seems to be walking upon springs; he proceeds with a kind of prancing gait, and brings his heels to the ground with a kind of kick. If he attempts to take hold of an object with his hands, he probably thrusts it from him by a spasmodic jerk of the arm.

The motor ataxy is usually accompanied, and occasionally preceded, by cutaneous *anæsthesia*, to a variable degree and extent. The fingers, toes, arms and legs are the parts chiefly affected. The patient commonly feels as if he were walking upon something soft like wool; unless he looks at his feet he is not certain that they have reached the ground. Sometimes he can scarcely feel that he has any feet at all, or seems to be 'walking on air,' or 'on his ankle-joints,' or 'on his hip-joints,' when the numbness extends up the thighs.

Analgesia, or loss of sensibility to pain, is frequently experienced to a variable degree and extent; or painful impressions are conveyed to the sensorium with unusual slowness. In one case that I saw, three or four minutes elapsed before the patient experienced any sensation of pain in the part that had been pricked; and, in another, it was not till after the very long interval of *twenty minutes* that the patient, without being asked, complained of smarting in the part which had been pricked with a needle.

by Dr. Charcot of Paris. The knee-joint is almost always the seat of the disease, which occurs suddenly as an oedematous swelling. The part is neither red nor hot, nor is there any constitutional disturbance or fever. In the presence of strabismus and urinary affections, it may last only a few days, or be prolonged and give rise to permanent deformity. In the latter case the bones and cartilages of the joint are often found diseased.

In the early period of the disease, before motor ataxy has appeared, and when only one or perhaps two of the symptoms are present, the differential diagnosis is exceedingly difficult. But whenever strabismus or the peculiar flying nystagmus suddenly attack an otherwise healthy person, they should suggest the possibility of their being the precursors of locomotor ataxy. In a large proportion of instances the strabismus is accompanied by amblyopia; and when it is single, the amblyopia is on the corresponding side.* Even in cases in which the symptoms have been present, and in which the diagnosis is not means difficult, I have seen operations needlessly performed on the eye by surgeons who were ignorant of the nature of locomotor ataxy. Frequently the disease has been mistaken for incomplete paraplegia. It is not long since I was employed as a surgeon to a public institution to see a case of paraplegia which proved to be a well-marked case of locomotor ataxy.

Pathological anatomy.—In this disease the spinal cord is

tion of the extremities of the posterior cornua, and so I have found the same kind of alteration in the more parts of the grey substance. In the latter case the disease is of a mixed nature, partaking of the characters of both locomotor ataxy and ordinary spinal paralysis.

Prognosis and treatment.—The prognosis, as might be inferred from the pathological anatomy, is generally very unfavorable. It is chiefly at the first invasion of the disease that any benefit is to be expected from the use of remedies. Hence the importance of an early diagnosis. One of the chief objects of treatment is to protect the patient from cold and damp, and place him in a moderate and equable temperature. A good and wholesome diet, with moderate exercise, or beer, is generally necessary. With regard to drugs, nitrate of silver appears to exercise the most direct or specific influence on the disease. Sometimes it not only alleviates the paresthesia, but it diminishes the incoordination. It should be first given in small doses, of one-eighth of a grain gradually increased to one grain three or four times a day, after meals. To prevent it from irritating the bowels or the bladder, it may be combined with opium, or with cannabis indica, or with belladonna. For relieving the neuralgic pains of the limb-pains, which so frequently disturb the patient, there is nothing so efficacious as the subcutaneous injection of morphia, beginning with one-sixth of a grain. I have always found these pains aggravated by constipation, and in this case I get the bowels to act by some means. Dry cupping along the spine sometimes affords decided relief.

DISEASES OF THE TONGUE.

THE tongue, an organ of well-known form, to which the sense of taste is chiefly referred, is composed of muscular and of gland-substance, covered by mucous membrane, continuous with that of the mouth and fauces. The more delicate appreciation of taste is limited to the upper surface, where the papillæ of the mucous membrane are remarkably developed. The proper muscular structure is striated, and arranged in two horizontal and two vertical layers; the former, immediately subjacent to the mucous membrane, are to be found on the upper and under surfaces; the latter pass vertically from one horizontal layer to the other, connecting them, but leaving intervals, which are supplied by gland-structures, supplied by ducts opening on the under surface, and secreting a fluid resembling saliva. In the anterior third, the muscular prevails over the gland-structure, which is reduced to two large lateral bodies. Abundantly supplied with nerves and blood-vessels, always in movement, and exposed to many sources of irritation or injury, this organ often becomes the seat of disease.

I know of no well-authenticated instance of atrophy of the tongue, although we have now on record many cases of progressive muscular paralysis involving for the most part not only the tongue, but also the palate and lips: such cases must be referred to diseases of the nervous centres. Congenital ankylosis of the tongue is also unknown to me. The frænum, it is true, tends in some cases to the tip of the tongue, when its division may be easily accomplished; but the defect usually produces little or no inconvenience, and the difficulty of utterance with which it is frequently associated proceeds from causes affecting the sensorium. Still the division of the frænum linguæ is an operation somewhat popular, and may be performed without detriment to the patient. Whenever the operation is performed, let it be remembered that the artery of the frænum

lingua propendula, macro-glossia, &c.) is an affection occasionally seen.

Zacchias* relates that he saw at Rome, in 1628, a male infant with the exception that the tongue projected three fingers' breadth from the mouth; the child could suck, and lived to the age of fourteen months, when it died of an obvious cause. Bertholin† mentions a case of 'linguæ portentosa' made known to him by his pupil Bagdun: a male child, born with the tongue protruded from the mouth as large as a filbert. As the child grew, the tongue increased in size of a calf's heart. Sauvages, in his *Nosologia*, speaks of enlargement of the tongue in new-born infants. Percy and Laurent‡ speak of two cases in women, in whom the enlargement was dated from birth, but rapid increase took place at the age of three. The other was that of a boy, aged sixteen, whose tongue hung three inches below the chin, was two inches and a half thick, and filled the mouth. The affection had existed from birth.

In Mirault's case, the man, aged thirty-four, had the tongue protruded from infancy; so likewise in that related by M. Mauran. In other cases the enlargement is described as commencing in the first or second years, when perhaps the better expression would have been, 'was then first noticed;' for the tongue is almost always enlarged in which hypertrophy in a slight degree might be readily looked for, especially in early life, and before attempts are made at articulation. In a few other cases the beginning of the enlargement in adults has been referred to salivation. Hypertrophy of the tongue as seen in infancy is congenital.

The measurement of the tongue in the case of a child of eleven, under the care of Dr. Humphry,|| was as follows: from the upper lip to its tip, 3 inches; from the under lip to its tip, $1\frac{1}{2}$ inch; from the angle of the mouth round the tip to the opposite angle, $6\frac{1}{4}$ inches. The circumference of the wi-

ontal direction which the inferior incisors and the alveolar process of the jaw had assumed. These were so placed as to form a wide channel, the tongue rested.

Amphry determined to remove the protruding part of the tongue by although he was aware that a fatal case had occurred elsewhere from ice.* He passed a straight bistoury from below upwards through the title to the left of the middle line, and cutting forwards and outwards left lateral flap. He then secured a vessel. Next he cut across the the tongue, divided the ranine arteries, the movements of the organ controlled by means of the prolapsing portion, which was not quite The operation was completed by the formation of an oblique lateral the right side, corresponding with that on the left. The flaps were ted so as to form a tip, and maintained in apposition by two sutures eply. The hæmorrhage was not very great, and the bleeding vessels without difficulty. A good deal of swelling of the tongue followed ion; but under frequent washing and fomentation it soon subsided, ound healed. At first the lips could not be approximated, and the ap of a tongue was always visible, though never protruding. After a however, by continued action of the orbicularis muscle, the mouth d. An apparatus was then constructed to act by pressure on the jaw, and ultimately a result was obtained highly creditable to the

effect of continued pressure on the enlarged tongue is a treatment only applicable to early cases, and is uncertain sults. In a case under the care of the late Mr. Crosse, ling, partly removed by this method of treatment, soon d, and the relief appears to have been imperfect. Some s have treated such a case by ligature; but to this ing an objection offered itself in the formation of a slough

The *écraseur* is an instrument generally unsuited to operations about the mouth, fauces, or tongue; but may be advantageously used whenever there is fear of uncontrollable hæmorrhage. There is this disadvantage attending its use: that as the chain becomes tightened, it draws nearer and nearer to the limits of the disease, so that when the operation is completed and the part removed, the line of section will often be found in dangerous proximity to the diseased structures. I think that the fear of dangerous hæmorrhage is somewhat exaggerated, except in the case of operations performed at the root of the tongue.

Acute inflammation of the tongue, attended with sudden enlargement, is a disease occasionally seen. Mr. S. Cooper remarks that it may arise spontaneously, and without any apparent cause; or else from some particular irritation, such as that of mercury or some poisonous substance.*

He mentions that, in the middle of the seventeenth century, Schlegel, who was at Paris, saw a patient in salivation, whose tongue became so enormously enlarged that the mouth could not contain it. Pimprenelle, an eminent surgeon of the time, was sent for, and amputated one half of the organ; a measure which Louis criticises as extremely violent.

The disease to which I now refer, however, proceeds from occult atmospheric causes. The swelling is sudden, perhaps occurring during the course of the night, and it produces feelings of threatened suffocation, having been preceded by loss of appetite and a general sensation of *malaise* for a few days.

The first case which came before my notice was that of a gentleman aged sixty, of remarkably abstemious habits, living chiefly on vegetable diet; the second, that of a young married lady, residing in lodgings in a healthy part of London. In this latter case the swelling of the tongue was so great that the mouth could not be closed; the teeth became coated with sordes, and a profuse watery and highly fetid discharge flowed without intermission for many days. The patient feared to close her eyes in sleep; and so difficult was the respiration, that at one time the question presented itself, whether it might not be necessary to make deep incisions into the swollen organ, or even to open the tracheæ. However, by constant attention to cleanliness and gargling, by the local use of ice, by supporting the strength with wine and quinine, and the administration of nourishment by enemata during a short time when deglutition was impossible, the swelling subsided, and the patient recovered. For some weeks, however, she remained in an enfeebled state. I saw about the same time two other cases in St. Bartholomew's Hospital. In both the disease was rapid in its course, the accession lasting but a few hours, and the swelling usually subsiding within a week.

* *Surg. Dict.*, ed. 1830, p. 1102.

But during the period of acute œdema of the tongue, the patient may die suffocated; and the surgeon should bear in mind that such an accident may be prevented by his making free and deep longitudinal incisions into the swollen organ. 'I have seen a patient,' observes Mr. Erichsen, 'who was nearly suffocated by the immense size of his tongue, relieved at once, and got nearly well in the course of a few hours, by such incisions.*' These incisions should be made along the upper, rather than the under surface of the tongue, that the ranine artery be not wounded; and there is but one caution: the œdema may so far involve only one side of the tongue as to cause the lower surface, which yields the more readily, to be turned *directly upwards*, in which case the incision made above passes in truth into the tissues normally inferior. Such a case happened in the practice of Mr. Wormald, who found that, upon the subsidence of the swelling, the incision made above gradually acquired a directly inferior position.

I have known tracheotomy deemed necessary in such a case, although the symptoms do not usually demand it. The condition of the mouth prevents the administration of medicines during the acute stage. Afterwards we must be guided in the selection of remedies by the state of the pulse. Should there be much fever, saline and diaphoretic remedies are suitable. But more commonly the indication is to support the strength, by the administration of tonics, wine, and nutritious diet. For the mouth, a gargle containing alum, or borax and honey, or tincture of myrrh, may be used; and the surgeon may give much relief by carefully wiping away the thick viscid secretion from the teeth, and by touching any ulcerated spots with a weak solution of nitrate of silver.

In the Museum of Guy's Hospital there is a preparation of a portion of a tongue, weighing two ounces and three drachms, removed by ligature by the late Sir Astley Cooper, in consequence of enlargement following salivation produced by mercurial medicines in the treatment of syphilis. And another preparation (1672) illustrates the extreme effects of mercury, namely, mortification of the organ; a result extremely rare, and due to the injudicious and long-continued administration of this powerful agent.

Abcess.—I have seen several cases of abscess of the tongue.

* *Science and Art of Surgery*, 3rd ed. p. 804.

One case was that of a lady of middle age under the care of Mr. Stanley. She had a firm tumour, the size of a large pea, embedded in the substance of the tongue. Considering the patient's age, and the symptoms attending the formation of the tumour, Mr. Stanley could not dismiss from his mind the idea of carcinoma. But its situation in the middle of the substance of the organ militated against that idea. He made an incision into the mass, and a small quantity of healthy pus escaped. The wound cicatrised, and the lady has continued in good health ever since. Mr. Erichsen * says, 'abscess of the tongue, though rare, occasionally occurs. A boy was brought to me some time ago with an elastic fluctuating tumour of slow growth and about the size of a small plum, situated deeply in the centre of the tongue. On puncturing it, about half an ounce of healthy pus was let out, after which the cyst speedily closed.' Other similar cases are met with scattered through medical literature, and they show that the gland-follicles of the tongue are liable to suppuration.

Chronic ulceration.—'The most frequent forms of disease in the tongue,' observes Sir W. Lawrence,† 'are, ulceration, generally superficial, sometimes more deeply seated; swelling and thickening of the mucous membrane; swelling and induration of the substance of the organ. Ulceration often exists in conjunction with swelling of the mucous membrane and with induration of the lingual substance. The more formidable diseases of the tongue are syphilitic or cancerous; the former being by far the most numerous. Disorder of the digestive organs is sometimes the source of the mischief, giving rise to affections which are usually superficial, but sometimes of more serious character. When maintained by imprudent indulgence in drinking, there is often a sore state of the organ, the epithelium being smooth, reddened, or white and opaque. When the surface is superficially ulcerated, the ulcerations are generally small, circular, and of greyish colour; occasionally there is thickening of the mucous membrane, and deeper ulceration.' In the same volume of the *Medical Gazette* (p. 800), Sir W. Lawrence relates some cases in illustration of these points.

A healthy-looking gentleman, between forty and fifty, who had always enjoyed excellent health, but who had lived freely, became the subject of the following condition of the tongue:—the mucous membrane over the greater part of the organ was unnaturally smooth, more or less opaque and white, here and there raw, as if it had been ulcerated. It had been in this state, generally causing great pain, for four or five years. Under a course of mild mercurial medicines, the employment of tincture of myrrh as a gargle, and abstinence from strong drink, the pain ceased in a week; and the general condition of the

* *Science and Art of Surgery*, 5th ed. vol. ii. p. 359.

† 'Clinical Lecture' reported by Mr. Coote, *Medical Gazette*, 1845, p. 703.

Some years ago a patient, about sixty years old, came from the to consult Sir W. Lawrence. The man was of sallow aspect, and had sely. On the upper surface of the tongue, in a space about an inch and ng by an inch wide, the mucous membrane was thickened, indurated, ed into irregular prominences, which were smooth and not ulcerated. mplaint had existed some months, having become so painful that he uly take food; washes and gargles had been of no benefit. There was ion of syphilis in this case; and the habit of drinking, especially spirits, ly accounted for the disease. Sir W. Lawrence prescribed the com- ecotion of sarsaparilla with compound decoction of aloes three times a r grains of the extract of hyoscyamus at night, and a strictly regulated hen again seen at the end of a fortnight, all the pain had ceased, and l masticate anything. He considered himself well, and returned to try. I saw a tall, spare-looking man, aged sixty, March 18th, 1860, in e gums had shrunk from the teeth, leaving the roots denuded to the border, and exposing them to an accumulation of tartar. The patient i freely, having been a publican. The whole of the tongue was swollen, and of glistening red hue. There was no trace of the papillæ, except at , where the papillæ circumvallatæ appeared hypertrophied. In those s where the organ rubbed against the exposed and roughened teeth, the membrane was white and thickened, and seemed to surround numerous d unhealthy-looking ulcerations. I directed this patient to have the amined and cleaned by a competent dentist; imposed a strictly regulated lered a gargle containing alum, and occasionally aloetic aperients. Con- s improvement took place; but after a time the patient declined to o the strict regimen which was considered necessary.

* smooth glazed condition of the tongue differs from that only seen in persons suffering from chronic dyspepsia, h habitually abstemious. In these latter cases, the as membrane has a red, uneven, and raw aspect; it is r fissured, ulcerated in spots; the papillæ may be univer-

an indurated condition of the tongue neither mangna syphilitic.

A hatter, aged sixty-one, of unimpaired constitution, who had no syphilis since the age of eighteen, found that whenever he had been exposed to the fumes of some secret composition used in his trade, the tongue became cracked, and dry, and remained so two or three days. For the last time it had swelled and become painful during the winter months, but returned to its natural state with the approach of warm weather. About four years before he was seen, the back part and left side of the tongue swelled in an unusual manner, and became hard. This was attended with a dull pain. The surface was uneven and tuberculated, and presented several superficial ulcerations not extending beyond the mucous membrane. The tongue resembled that of scirrhus. The absorbent glands under the jaw were affected. Iodide of potassium was tried without effect. A course of mercury was next administered, under which the gums became sore, and the tongue recovered so much of its normal character that the patient was pronounced convalescent.

Nævus of the tongue.—

On March 15th, 1852, I saw a private patient under the care of Sir W. Lawrence, suffering from congenital venous swelling involving the right half of the tongue. The part had a livid-blue colour, with rounded and elevated surface, marked by large tortuous veins. Sir W. Lawrence made an incision into it, and evacuated a quantity of fluid venous blood, by which the tumour entirely disappeared. A ligature was then applied to the pedicle, and hæmorrhage, when there was renewal of the swelling to even a greater extent than before. The venous trunks still communicated one with another. The case did well. A student at St. Bartholomew's Hospital showed me in 1853, on his own person, a large congenital venous nævus of the tongue occupying its right half, including both upper and under surface. At first it used to bleed: but it had long ceased to cause him inconvenience.

the hæmorrhage, though profuse, may be generally controlled by ice. When the ligature is to be applied, it is usually secured by a strong curved needle; the tongue being drawn out by an assistant, who holds the organ, covered at the part by a towel, between the thumb and fore-finger of the hand. No instrument can press and hold the tongue with the certainty as the human hand.

May 17, 1847, Sir W. Lawrence removed a small vascular tumour of the gum from a young lady aged nineteen. She had previously had removed a tumour from the gum of the lower jaw opposite the incisor teeth, one of which had been extracted. On microscopical examination of the extirpated mass, he found that the papillæ of the tongue, about the centre of the tumour, were hypertrophied and agglutinated into a prominent and flat-surfaced mass of white colour. It was composed of white fibrous tissue, of stringy, fibrous tissue, and of an immense quantity of mucus and of epithelial cells. It seemed, in fact, as if the follicular portion of the mucous membrane was much hypertrophied, and covered by altered and thickened papillæ.

Paget says, 'In the Museum of the Middlesex Hospital is a tumour, one and a half inches long, which was removed from beneath the tongue, where it looked like a ranula; and in the College Museum, No 190, is one taken from the substance of the tongue.' This surgeon also states that he removed an elongated tumour, about half an inch in diameter, from the gum of a young man, from the substance of which, near its base, it had been growing for three years. It was firmer than most others of a similar kind, yet succulent, and formed of an areolar filamentous tissue, abundantly nucleated.*

accuracy of this description, and Mr. Erichsen questions how so small a duct can be dilated to so large a size as is occasionally attained by these tumours. This doubt, however, disappears when one reflects upon the enormous size which the minute ducts of the mammary gland may attain in sero-cystic sarcoma.

In one case of ranula under the care of Sir W. Lawrence, the little finger could be inserted for a short distance into a cylindrical tube, pursuing the normal course towards the gland. And in another case under my care, the patient, a domestic servant, presented herself with a hard swelling near the *frænum linguæ*, which had been pronounced 'a cancer.' Upon examination, I detected in Wharton's duct one of those phosphatic concretions, about the size of a pea, common to the ducts of the salivary glands; it was removed with ease, and the patient recovered in a few days.

But cysts may form in the floor of the mouth from any of the causes which lead to the development of cysts in other parts of the body.

Many years ago Sir W. Lawrence requested me to give him assistance in the following operation. A young lady had a swelling on the floor of the mouth. It was firmer and more solid than the common ranula. Sir W. Lawrence made an incision through the mucous membrane, and endeavoured to extirpate the cyst entire; but finding it too large, he made an incision into it, and, with the handle of the scalpel, took out a quantity of thick matter, the consistence of putty. He then removed the entire cyst, which was but loosely attached to the surrounding parts. The patient recovered.

Then we meet with other cases in which a cyst, containing a fluid of watery consistence, extends from the floor of the mouth under the sterno-mastoid muscle to the middle of the neck. Such a tumour can scarcely be regarded as an enlargement of Wharton's duct. The proper conclusion seems to be, that cysts may form in this situation from four sources. First, dilatation of Wharton's duct. Secondly, dilatation of one of the sublingual ducts. Thirdly, dilatation of a mucous follicle. Fourthly, dilatation of a bursa mucosa, said to exist on the outer surface of the *genio-hyoglossus* muscle. The contents of a dilated Wharton's duct are a perfectly clear and thick albuminous fluid of the consistence of white-of-egg; occasionally we meet with phosphatic concretions. An enlarged mucous follicle is filled with a putty-like substance consisting of epithelial cells filled with granular fat. An enlarged bursa, which may extend down the neck, contains a clear serous fluid, in which blood-discs may be from time to time detected in various quantities. I have no information respecting dilatation of the sublingual ducts.

of the eye may be traversed
on, so as to excite inflammation, and its subsequent
ion. Mr. Skey employs a single silken thread passed
the floor of the mouth to the most depending part
yst; and I have seen benefit result from this treat-

gue is subject to injuries of various kinds; the most
of which are those inflicted by the teeth in the con-
its of epilepsy. There are instances known where a
of the organ has been completely severed by the spas-
osure of the mouth; but the injury is rarely of a nature
re a suture. Wounds heal very readily; and a foreign
however small, is irritating to the mouth. But a suture
necessary, be applied. In the Museum of Guy's
there is a specimen (1674) showing a piece of tobacco-
bedded in the substance of the tongue; the soft parts
ed over it, so that its presence was not readily detected
life. Frequent hæmorrhages ensued; and the case
ultimately fatal. Fatal hæmorrhage rarely occurs from
e wound of the tongue. The bleeding generally ceases
e application of ice.

ure of the lingual artery above the great cornu of the
les in the neck, for the purpose of controlling hæmor-
rom the ranine artery, is recommended, but is rarely
L.

ilitic affections of the tongue are mostly ulcerative, and

primary sore. Syphilitic induration of the tongue may occur alone, in which case it has been mistaken for cancer. The diagnosis is not always easy.

A female, aged thirty, servant in a family of respectability, presented herself at the hospital, with a foul ulcer of the tongue on a hard base, situated between the middle line and the edge of the organ, midway between the base and the apex. It had existed three months; there were no other symptoms. After due consideration, it was thought that the disease was syphilitic; and she was put upon the proper remedies, namely, two grains and a half of mercury and chalk three times a day, when the ulcer healed, and the induration disappeared. We then learned that the patient had suffered from syphilis, under peculiar circumstances, which it is unnecessary here to relate.

A gentleman, aged forty-five, consulted Sir W. Lawrence with an indurated knot the size of a sixpence on the middle of the dorsum of the tongue. It had resisted common treatment, but yielded to a slight mercurial course. No other constitutional symptom of syphilis was present, and the primary sore, which was little more than an excoriation, had happened eight months before.

From the upper surface of the tongue there may project a mass of hypertrophied epithelium, which, blended together, constitutes a mass analogous to the mucous tubercle commonly seen about the scrotum, the vulva, or anus. For the treatment of syphilitic affections the reader is referred to the proper section.

Cancer, as it affects the tongue, usually occurs in one of two forms: either interstitial deposit, when it is called scirrhus, or more superficial development, when it is called epithelial cancer, or epithelioma. Both forms are of serious nature. I have not hitherto known any well-recognised case in which, even after the most complete and early removal, the cure has been permanent. In both instances microscopic investigation detects the usual elements of cancer, namely, nucleated cells and connecting tissue.

We have first to inquire what is the cause of cancer of the tongue. Is it due to local irritation, such as is excited by an irregular or broken tooth? If such were the case, how infinitely more common should it be! how frequently should we notice its ravages in the mouth! Many experienced dental surgeons deny that irregularities in the shape of the teeth will do more than determine the exact seat of development of cancer in constitutions previously so disposed; and the same may be said of the irritation excited in other ways. Let us take, for example, the irritation produced by a clay-pipe. How many are there

to smoke unaffected by cancerous disease to the end of a long life! How often have we to remove carcinomatous or cancerous growths from the mouths of persons who have rarely or even never smoked at all!

I lately extirpated such a tumour from a man aged seventy. He had led a most abstemious life, and had rarely indulged in any luxury, however small. The wound of the tongue cicatrised; but the patient died eight months afterwards of secondary disease in the glands of the neck.

Still the irritation excited by the constant presence of a tobacco-pipe cannot be ignored. In 1848 I examined the body of Major-Gen. D., aged sixty-three, a patient under the care of Sir W. Lawrence, who gave me the following history of the case. This officer had been above thirty years in India, where he had lived solely. He was always in the habit of smoking a short clay-pipe, the end of which kept firmly wedged between the floor of the mouth and the under surface of the tongue. About July 1847, he noticed a slight impediment to swallowing; the food, as he said, was caught in its passage through the fauces. A physician applied caustic, but without avail; the disease increased, and the left side of the tongue became swelled and hard. The patient died June 19th, 1848, without peculiar suffering, and rather from exhaustion than from suffocation. On examination after death, a deep ulcer was found, the size of a shilling, with red raised edges, on the left side of the under surface of the tongue. It penetrated the root of the organ, and spread on the dorsum into an ulcer three inches broad and two inches from before backwards, in which direction it invaded the soft palate and tonsils, and had destroyed the epiglottis and the glosso-epiglottidean folds. The aperture of the glottis was nearly blocked up, and the disease spread into the interior of the larynx. There had been little or no hæmorrhage during life. There were six or seven small absorbent glands, infiltrated by masses of opaque yellowish-white matter. The morbid growth consisted of epithelial cells, and similar bodies were embedded in the tongue along the course of the penetrating ulcer. This gentleman's father had lived to a very advanced age.

Hard cancer or scirrhus of the tongue commences as a firm and incompressible knob on the edge of the organ, situate often opposite the last molar or wisdom tooth; or so far towards the tip that it is often beyond the reach of surgical interference. The patient at first complains that the tongue is sore, and he often attributes this symptom to the irritation of an unsound tooth, and the uncontrolled movements of the jaw during sleep. Soon the act of deglutition becomes painful: the patient fears either to eat or drink; a sharp pain extends along the Eustachian tube towards the ear; the saliva flows from the mouth profusely. The symptoms become aggravated during sleep; the nights are disturbed by the secretion accumulating in the throat and the resulting cough. Often the patient is roused by a painful compression of the tongue between the jaws; and as the ulceration

extends, hæmorrhages occasionally supervene. Sooner or later the submaxillary absorbent glands become enlarged by the same cancerous deposit, when first a circumscribed and then a diffused tumour forms in the neck, covered by thin and reddened integument, which, should the patient survive a sufficient length of time, ulcerates, leaving a foul open sore, discharging a thin fœtid fluid, with occasional hæmorrhages. But with the progress of the disease emaciation supervenes, and the patient dies with the usual cachectic indications of cancer.

But this is not the only form of malignant disease affecting the tongue.

In 1847, Mr. Ceely of Aylesbury removed from the tongue of a patient a soft, friable, and irregularly-lobulated mass of reddish-brown colour, with vascular red points, but without any distinct arrangement of vessels, composed of cauliflower or club-shaped cells with large nuclei, blood-discs, and a thickish matter coagulable in alcohol. The tumour, said Mr. Ceely, when examining it immediately before its removal, protruded from the mouth; the tongue was thrust to the left side, and was swollen; the lips were cedematous; the surface of the tumour was in many places black, and a sero-sanguineous fluid oozed from its upper part. By retracting the angles of the mouth, and directing the patient to thrust the tongue forward, and at the same time raising the tumour over the lower lip, it was brought so far forward as to become girt round its broad base by the whole circumference of the mouth, where it remained fixed. It then felt quite elastic, and measured longitudinally five inches, transversely three inches. While it was in the above situation, I placed my fingers, continued Mr. Ceely, under the tumour, and gently raising it from the base I peeled it off, when a large jagged surface was left, which was soon covered with coagula from bleeding vessels. The hæmorrhage was moderate, and easily repressed with the matico-leaf in coarse powder. The portion removed very much resembled a recently detached placenta, though perhaps less firm; it was a spongy, granular mass, with interstitial fluid blood; it had none of the shaggy filamentous appearance of cauliflower excrescence detached from the uterus. The weight of the mass removed was seventeen drachms and a half. On April 8th, 1850, Mr. Ceely reported that the man was still alive, although the morbid growth was reproduced within a week.

I saw near to Town, in 1848, a patient in whom there was scarcely any vestige of tongue remaining, the whole organ having been affected with a similar friable degeneration. The patient could still articulate, and suffered less than might have been expected.

In 1860, a patient under the care of Mr. Harle, surgeon, of Islington, presented himself at the hospital, in whom a portion of the upper surface of the tongue, the size of a sixpenny-piece, was occupied by an opaque white layer or mass, consisting of thickened epithelium, rising above the level of the surrounding parts. Under the microscope the substance was seen to consist of epithelial scales, dried and deformed, and containing but few nuclei. There was no tendency to indefinite propagation, as in cancer, nor to any implication of the neighbouring tissues. The papillæ under the white mass seemed enlarged.

was no swelling of the neighbouring absorbent glands. The disease had twelve years without material change; the patient was a healthy aged forty, and no surgical interference was indicated. But when this mass was raised, and the state of the papillæ seen beneath, the resemblance to epithelioma was obvious. And it was found, by microscopic examination of the softer epithelial cells carried out this resemblance further. That the case was not of cancerous nature was proved by the history; no source of irritation was discovered by the most careful examination; and the case to illustrate the fact, that in malignant disease there is something the mere increase of the epithelial elements. It was in all probability character that a tumour, called supposed epithelioma of the tongue, which was presented to the Pathological Society, December 6th, 1859.* taken from a patient aged seventy-nine; had existed sixteen years, been six years under observation. It involved the tip, the right half left side of the tongue; the pharynx and larynx were atrophied. After an examination of the body was made, when no malignant disease was seen elsewhere. In 1857, I saw a similar case under the care of Mr. Ceely of Ry. The patient was a sailor, of very intemperate habits; and about years ago he presented himself with a small epithelial growth on the about the size of an almond.

Cancerous tumour of the tongue should be extirpated, when possible, by the knife. The patient must sit in a chair, with head supported: chloroform had better not be given in these cases, because the concurrence of the patient is needed. The assistant grasps the tip of the tongue, covered with a towel, with the fore-finger and thumb of one hand, while with the other he pulls back the cheek. The surgeon then seizes the tumour completely within the blades of double-hooked forceps, and with a sharp round-pointed knife, of somewhat large size, removes away the enclosed mass in a few seconds. The whole must be grasped with the forceps, that none be left behind. The hæmorrhage renders any subsequent examination impossible. The knife should be round-pointed, that none of the series on the floor of the mouth be wounded. It may be done in different ways, according to the fancy of the operator; but a straight knife is always the best. The hæmorrhage must be controlled by ice. An artery which commonly needs a ligature is the sublingual; and this can only be wounded by the surgeon using a sharp-pointed knife. No ligatures or sutures are usually desirable to the wounded tongue; they irritate the wound, and excite the flow of saliva. The cut surfaces granulate and heal, leaving perhaps an elevation of the substance of the tongue, from its being puckered up at the seat of operation.

* *Path. Soc. Trans.* vol. xi. p. 240.

This, however, can be distinguished from the reproduction of disease by the absence of induration.

The *écraseur* has supporters as a proper instrument for the performance of these operations. When both lingual arteries may be divided, it is to be preferred to the knife, and M. Chassaignac has amputated the tongue close to its base with this instrument. As the disease threatened to involve the entire organ, its complete extirpation was resolved on. Two or three days before undertaking the operation, M. Chassaignac passed a drainage-tube from without round about the base of the tongue, immediately above the great cornu of the os hyoides. The point where the tube was introduced was a little to the right of the middle line. Having thus prepared a passage for the chain of the *écraseur*, the tube was withdrawn as soon as the chain was in position. Then commenced the separation of the organ. Two minutes were allowed to elapse between each movement of the instrument; and the transverse division of the organ was effected in exactly half an hour. The chain was then passed behind the base of the separated organ, and made to embrace the muscles and tissues attached to its under surface. In thirty minutes more this second part of the operation was completed, so that the entire time occupied was exactly one hour. The quantity of blood lost was quite insignificant, and the poor man did not evince great suffering. On the eighth day after the operation he was going on well.* A subsequent account reports the patient's recovery. A yet more serious and complicated proceeding consists in the introduction of the chain round the root of the tongue by means of an incision into the floor of the mouth from without. A second chain is required to free the severed tongue from the muscles and soft parts; this double operation takes about one hour for its completion. The jaw has been divided at the symphysis and the two halves pulled asunder in order that the surgeon might gain more room for passing the chain down to the root of the organ. But the amount of space thus obtained does not equal the surgeon's expectations.

When the cancerous degeneration is limited to the side of the tongue, no such proceeding can be necessary, for the severed arteries are but small. But when the operation involves an extension beyond the mesial line, so that one or both ranine arteries must be cut across, and that, too, towards the base of the organ,

* *Med. Times and Gazette*, May 14, 1859.

under such circumstances the *écraseur* may be used with advantage. With it may be combined the employment of the ligature, as in the following case.

In 1859 I removed by the knife a cancerous mass from the left side of the tongue of a man aged forty-two. The hæmorrhage was considerable, the wound being very deep; but it ceased after the prolonged use of ice. The parts cicatrised completely, and the patient left the hospital apparently well. In a few months the disease returned in the cicatrix, and the patient came to me with the request that I would again remove the hardened parts, in order to give him even temporary relief from pain. As the induration extended quite to the middle line, and the ranine arteries would of necessity be divided, I proceeded as follows:

Chloroform having been administered so as to produce complete insensibility, the tongue was drawn from the mouth by an assistant; the chain of a small *écraseur*, armed with a needle, was then passed through the middle of the tongue beyond the base of the tumour, and made to cut from within outwards, so as to sever the posterior connections. Next a slight incision was made with scissors into the point of the tongue, so as to guide a ligature round the remaining attachment of the partially severed mass. A strong knot was tied, the principal part cut away, and the patient was removed to bed.

In December 1861, Mr. Nunneley described, before the Royal Medical and Chirurgical Society, the particulars of an operation for the complete extirpation of the tongue, in which it was asserted the cure had been permanent for some months. A small incision was made anterior to the os hyoides through the integument, mylohyoid, and geniohyoid muscles, by which a curved needle, to which was attached the chain of the *écraseur*, was introduced so as to carry the chain completely round the base of the organ. In consequence of some imperfection in the chain, it was removed; strong whipcord was substituted, and pulled with all the force possible. During the course of treatment hæmorrhage supervened, but it was easily arrested by a solution of tannin. Ultimately the organ came away, and the wound cicatrised. Some doubt, however, was expressed by the Society as to the exact nature of the disease.*

The tongue has been completely extirpated by the knife. Gaglioli has removed large portions by making an incision into the floor of the mouth from one angle of the inferior maxilla to the other. But Mr. Syme first, and subsequently Mr. Fiddes,† have extirpated the entire organ, by making an incision

* *Med. Times and Gazette*, Dec. 21, 1861, p. 648.

† *Edin. Med. Journal*, June 1859.

through the lower lip, chin, and lower jaw, pulling aside the two halves of the bone, and dissecting the diseased mass, which is by these means fully exposed, from the os hyoides. Fiddes points out the importance of securing one lingual artery before the other is divided. He has operated thus twice, and speaks well of the results of his cases. The proceeding has, however, yet found favour generally with modern surgeons.

Now, in cancer of the tongue the greatest distress to the patient arises, first, from the pain in every movement of the organ, extending to the ear and over the side of the head; secondly, from the profuse flow of saliva, which keeps him constantly wet and unable to articulate with comfort. The object of diminishing the sensibility of the tongue, and of checking the excessive secretion by the division of the gustatory nerve, which it lies close to the ramus of the jaw, suggested itself many years ago to Mr. Hilton;* and Mr. Moore, of the Middlesex Hospital, has recently repeated the operation.† By dividing the gustatory nerve, he remarks, between the disease and the brain, all the consequences of the irritation of that nerve are necessarily suspended. No sensation from the tumour can be conveyed along it; no reflected irritation can reach the collateral branches of the fifth nerve; no stimulus to an exaggerated secretion of saliva can be given to the salivary glands. A patient on whom this operation has been performed should speak more freely and swallow with less difficulty; should be relieved of pain in the tongue and jaw, temple, and crown of the head, and of the incessant annoyance arising from the dribbling of saliva; should sleep better, and be better nourished than before.

The mode of performing the operation, as adopted by Mr. Moore, differs in some respects from that adopted by Mr. Hilton. The latter surgeon sought for the nerve in the floor of the mouth, and exposed it by making an incision along the mucous membrane close to the sublingual gland. The former divided the nerve just behind the last molar tooth, as it escapes from the cover of the pterygoideus internus muscle. It is not more than half an inch distant from the tooth, and is covered only by the mucous membrane.

The guide to the nerve is the last molar tooth. On passing the finger into the mouth, within and beyond that tooth

* *Guy's Hospital Reports*, vol. vii. p. 263.

† *Med.-Chir. Trans.* vol. xlv. p. 47.

along alveolar ridge can be felt narrowing as it ascends into the ramus. Behind, below, and parallel with the ridge, is a shallow groove in the bone. The nerve passes along that groove. Therefore an incision three-quarters of an inch in length made within the mouth and down to the lower jaw behind the last alveolar tooth, and in a direction crossing the course of the nerve, must divide it. It is advisable to operate with a curved bistoury, since the alveolar ridge might shield the nerve.

The patients have expressed themselves relieved; the tongue being rendered completely numb.

Mr. Moore has combined with this operation ligation of the corresponding lingual artery, so as to cut off as much as possible the supply of blood to the affected part. The results are sufficiently encouraging to merit further trial.

It must be remembered, however, that there are ulcers of the tongue of very unfavourable aspect, which are not cancerous.

A patient, aged thirty-seven, had an ulcer of the tongue of eight months' duration. It was situated on the right half of the tongue, and had a coarsely granulated base, with inverted edges and undermined margin exposing the muscular tissue. The patient died labouring under severe tubercular disease of the lungs, and after death there were found tuberculous deposits both in these organs and in the larynx, but no trace of cancerous disease.*

The rule, therefore, holds good that in cases of doubtful ulceration of the tongue, all sources of irritation, such as the sharp edge of a decayed tooth, &c., should be removed, and great attention should be paid to the general health before we arrive at a final diagnosis.

HOLMES COOTE.

* The preparation is preserved in the Museum of St. Bartholomew's Hospital.

DISEASES OF THE NOSE.

I. AFFECTIONS OF THE EXTERNAL PART

ACNE ROSACEA. GUTTA ROSACEA VEL ROSEA.

THE skin of the nose is especially liable to the various conditions commonly included under the term 'Acne cea.' The appearances presented vary greatly in different They depend not only upon the stage the malady has reached, but also upon the particular form or variety which be under observation. In most cases, however, if not the earliest symptoms are due to excessive injection, followed by congestion of the superficial capillaries of the skin. Soon after the sebaceous follicles usually become involved to a more or less extent. But these organs are not the parts primarily affected. Hence many authorities* object to the application of the generic term 'acne' to the affections under consideration.

In the simplest, and perhaps most common form of rosacea, the skin at the spot affected is more or less inflamed, red, and generally has a somewhat shiny, greasy appearance. There is no obvious swelling or deformity of the part. A sense of heat is often experienced by the patient. This, however, is not constant, but comes on after food, or after exposure to changes of temperature. The tip or one side of the tip of the nose is often the only part affected. Occasionally the paranasal area is also affected, as though it had been *frost bitten*.

In another and more severe form the greater part or even the whole of the countenance presents an unnatural degree of redness, due to numerous dilated and congested vessels, ramifying in the skin, which can be seen by the naked eye. The tint varies according to the circumstances from a brightish red to a dusky purple.

* See especially Erasmus Wilson, *Diseases of the Skin*, p. 208.



mes the sebaceous follicles are involved, and the surface of the skin becomes more or less marbled and irregular.

These two forms of acne rosacea are much more common among women than among men. In women they are most frequent at the age of puberty, and at the climacteric period. In men they most frequently occur at the commencement of the decline of life. They do not appear to result from habits of intemperance, but are rather due to disorders of the digestive system, to exposure to the weather, and in the female to menstrual irregularities.

There is a third form, however, especially affecting, and often confined to the nose, which is most frequently, if not invariably, met with among men who have been habitual 'spirit drinkers.'

It is rarely if ever seen in any one under fifty years of age. The 'spirit-drinker's nose' is more or less swollen and shiny and fleshy. It is of a dusky or purplish red colour, and somewhat hard to the touch. On close examination the skin is seen to be traversed by numerous distended blood-vessels, but between them its appearance is comparatively natural. The sebaceous follicles are generally enlarged and inflamed; and the aspect becomes irregular and pimpled. At a later period the swelling increases, and the nose becomes protuberant. Its surface is covered with tubercles or blotches, or studded by minute pustules, or disfigured by adhering scales and crusts. The cutaneous venules are greatly distended, and sometimes varicose; and their dark purple ramifications become characteristically manifest.

Treatment.—Attention to the general health is in the first place necessary. But very diverse treatment is required in different cases. Disorders of the digestive system, and sexual irregularities must be especially inquired into, and if possible corrected. There is usually more or less general debility. In some instances tonics and stimulants are useful from the first; in others free purgation is indicated as a preliminary measure. Any such habits or practices as may seem to have caused, or aggravate the local capillary congestion upon which the affection primarily depends must be avoided. Moderation in the use of alcoholic drinks must be insisted on; and in some cases total abstinence may be advantageously enjoined. But general treatment alone rarely suffices to restore the reddened nose to its natural condition. Local remedies are requisite, and in recent cases are often speedily efficacious. In the early

stages and when the affection is slight, frequent bathing during the day with water as hot as can be borne, and bathing every night and morning with a weak tepid solution of bichloride of mercury (gr. j. or ij. to \mathfrak{z} j.) may be recommended. Wilson advises washing with cold water and juniper tar or carbolic acid soap twice daily: the ablution is to be followed, if much irritation is produced, by the application of benzoated zinc ointment. Subsequently an ointment of hypochloride of sulphur is to be rubbed in.* According to Hebra† (and Alibert) sulphur in the form of a soap, or an ointment, or as a solution of liver of sulphur, is the best application. The part affected is to be thoroughly washed at bed-time with soap and water. The remedy is then to be well rubbed in, and not washed off until the following morning. In some obstinate cases the chloride or the iodide of sulphur in the form of ointment may be used with advantage. It must be borne in mind that the various applications mentioned are all of them more or less stimulating and irritating, and that therefore (as pointed out by Hebra) the immediate effect of their application is apparently to do harm; and 'it is only after a little time that the good which they do becomes evident.'

In cases of greater severity and longer standing, in which the cutaneous vessels are much distended, and the swelling of the part affected is considerable, Hebra recommends a number of longitudinal incisions to be made into the dilated veins, especially those of which the loops are visible. After the blood has flowed for a little time, the parts are to be touched with a brush dipped in the *Liquor Ferri Perchloridi*. Very little reaction follows. This method of treatment has been followed by very satisfactory results. In some instances it has been necessary to repeat the scarifications on several occasions. Scarification of the vessels without the application of the perchloride is not, as a rule, effectual.

INTEGUMENTARY HYPERTROPHY. LIPOMA NASI.

In cases of acne rosacea of long standing and severe form, the integumentary structures of the nose sometimes become irregularly hypertrophied, and 'fleshy excrescences' or out-

* Wilson, *op. cit.* p. 211.

† Hebra, *On Diseases of the Skin* (Sydenham Soc. Translation), vol. ii. p. 334.

growths result.* These outgrowths vary greatly in the degree of development they attain, and the appearances they present. In some instances they are scattered, sessile, and wart-like. In others they are aggregated and flattened, and extend more or less uniformly. In others again, they are pedunculated, and form irregular globular or lobulated masses, which give rise to great disfigurement. In extreme cases they hang down as far as the chin, and interfere with respiration and speech. Sometimes they project upwards to such an extent as to obstruct the vision of one eye or both, so that the patient cannot see objects in front without changing the position either of his head, or of the nasal protuberance. They are generally dusky or purplish in colour, and somewhat cold and greasy to the touch. They rarely occur in individuals under fifty years of age, and scarcely ever in women. They appear to be less frequently met with now than in former times.

When cut into, these growths present a brawny consistence; and in places they are occasionally almost as hard as cartilage. They consist of hypertrophied and infiltrated skin and connective tissue, with dilated venules, and enlarged sebaceous follicles. They never involve the cartilages or mucous membrane, nor do they encroach on the interior of the nostrils. They contain, as a rule, little or no fatty tissue. Hence their common appellation (*lipomata nasi*) is obviously inappropriate.

Treatment.—The only treatment that can be recommended is removal by the knife. In operating, the growths may either be simply cut off one by one; or an incision may be made in the median line down to the cartilage, and the skin with its outgrowths dissected off on either side. Care must be taken not to cut through the cartilages. During the operation, the surgeon or his assistant should keep a finger in the nostril to serve as a guide. The bleeding which ensues may be readily checked by pressure or by the application of styptics. There is much less danger of erysipelas and other serious results than is commonly supposed. The exposed surface usually heals favourably under cooling and soothing applications. When cicatrization is complete, it is remarkable how little disfigurement remains. In some rare instances there has been recurrence of the growth after extirpation.

* Hebra describes these 'fleshy excrescences of the nose' as characterising 'the third degree' of *acne rosacea*. Other authors include them under the term '*acne hypertrophica*.'—Hardy, *Leçons sur les Maladies de la Peau*, p. 115.

LUPUS.

Lupus exedens in a large majority of cases first attacks, and most seriously affects, the skin of the nose. So also does lupus erythematosus, but in a somewhat smaller proportion of cases. Lupus non exedens, on the other hand, most frequently commences on one or other of the cheeks or lips, but in its onward progress it often invades the nose. Hence it might appear that only the first named form, and perhaps the second, demand special description in the present Essay. But all three forms of lupus appear to be associated with similar constitutional peculiarities; and they present, in their course and results, certain characteristic phenomena which are common to all of them. Moreover in some cases the distinctive features of the form under observation are not well marked; or two forms may exist at the same time in different parts. In other cases one form may seem to pass almost imperceptibly into another. For example, what is lupus non exedens on the cheek may become lupus exedens when it invades the nose. And thus even the different described forms and varieties of this affection may be regarded as only so many different modes of expression, as it were, of the same, or of a nearly similar morbid condition. It is scarcely possible, therefore, to discuss thoroughly any one form of lupus without reference to the others.

All the forms of lupus are ordinarily classed among those affections which are considered to denote the strumous or scrofulous diathesis. All occur most commonly, if not invariably, during youth or adolescence. All present the following characteristic phenomena. First, infiltration of the skin (or mucous membrane) by material of imperfect organisation and low vitality; secondly, slow, insidious, and progressive destruction of the normal tissues; thirdly, absolute persistence of more or less depressed scars after cicatrisation;—in other words, deficient restoration of the parts destroyed.

Notwithstanding the close relationship thus indicated as existing between them, it is nevertheless of some importance to distinguish the different typical forms of lupus one from another.

The broad distinctive features generally recognised are as follows.

Lupus exedens and lupus non exedens in their earlier stages

ance. Atrophy and absorption subsequently take place, lupus non exedens without ulceration, but the destructive does not as a rule extend so deeply. It is stated that in cases the papillary layer only of the corium is destroyed.

may now proceed to discuss some of the more special in the history and treatment of each form of lupus.

Lupus Exedens. Noli me Tangere.

form of lupus usually commences on one or other of the tip, or the columnna of the nose. It may commence in the skin or in the mucous membrane; or it may attack simultaneously, in which case its progress is as a rule completely rapid.

the other forms of lupus it occurs most frequently among people, or among those who are about entering upon life.* It sometimes commences during early childhood, rarely, if ever, after middle age. It is somewhat more common among females than among males, and is especially to affect those who are of fair, or even rosy complexion, thin delicate skin, auburn hair, and light grey or blue

It is associated, as a rule, with other indications of a scrofulous constitution or tendency in a more marked degree than is constantly the case with lupus erythematous, and perhaps also than with lupus non exedens.

On first inquiry or merely superficial examination, it appears that the general health of the patient is good. On full investigation, however, evidences of more or less functional derangement of the chylipoietic viscera may generally be covered. The tongue is redder than natural, especially the margins, and towards the tip; and the middle portion is covered by cream-coloured fur, through which the redness projects. The breath is more or less offensive. The evacuations from the bowels are irregular and unhealthy. Other symptoms of 'strumous dyspepsia' often present themselves. The processes of assimilation, nutrition, and sanguification are carried on feebly; and there is usually more or less general anæmia with pallor and flabbiness of tissue.

Lupus exedens first shows itself as one or more brownish-red papules which increase very slowly in size, and often remain for a long period without manifesting any change. After a time other papules spring up around between those which originally appeared, and all being gradually blended together a tubercle is formed. The tubercle at first may be about as large as a lentil or small horse-bean, but it is liable to considerable variation in extent and thickness. Its surface may be smooth or covered with thin silvery scales. At a later period one or more minute yellowish pustules appear. These increase in number, and become more distinctly pustular, and irregularly confluent. The epidermis becomes raised, and then gives way, and, with the ill-conditioned purulent secretion and tissue debris accumulated beneath, soon dries up into a thick greyish-yellow crust. On removal of this crust an uneven ulcerated surface is exposed, which is slightly raised in some places, and in others is covered by a purulent or semi-purulent viscid exudation which speedily concretes on exposure to the air. A new crust is thus formed, and beneath the ulcerative process goes on. The crusts vary in thickness and general appearance according to the quantity and proportions of the materials entering into their composition. They vary also in the degree of intimacy with which they adhere to the subjacent surface. The ulcerating surface in some places is papillated or mammillated and red, in others is depressed, excavated, and comparatively pale, or greyish. Its edges are somewhat jagged in outline, but bevelled almost or altogether on the level of the surrounding skin. The skin is more or less reddened, but rarely swollen or indurated.



by extent. This process goes on slowly and insidiously. Papules, tubercles, and 'pustules' appear at various points round the circumference of the ulcerated patch; and the area gradually extends until a more or less considerable portion of the countenance is involved. The skin, with the hair, sebaceous follicles, and sudoriparous ducts, and also portions of the subcutaneous areolar tissue, are liable to be destroyed in turn. But the destructive process rarely goes deeper, except in and about the nose and eyelids. In the cartilages and the small muscles which move them are also destroyed; and great deformity results. This, as may be understood, most frequently occurs in cases in which the nasal membrane is simultaneously affected. Sometimes the nasal bones and the vomer may suffer to some extent. The graphic descriptions given by the older writers (as well as the modern ones) of the direful ravages of 'lupus,' seem to represent the effects of rodent cancer rather than to those of the disease now under consideration. The distinction between the two affections is sufficiently clear, as will be hereafter explained, and the distinction is one which for practical reasons should be carefully made.

When the processes of destruction and ulceration thus decrease and extend circumferentially, it very often happens that the processes of repair and healing commence centrally, and the disease spreads. In such case the ulceration may become for a time more or less 'serpiginous' in character. In this way, after months or years, cicatrization at length advancing more rapidly than ulceration, and the disease having as it were run itself out, a natural cure may result. Sometimes, however, somewhat more frequently, the destructive process does not stop short. The surface loses its irregular papillated appearance; the excavated spots become pink; the secretion is increased in quantity, becomes thicker and for a time more purulent in character; and cicatrization commences at the periphery, and gradually advances towards the centre. The cicatrices are at first covered by a thin shining epiderm, through which the red remaining portions of the corium and adjacent tissues can be seen. After a time the epidermis increases in thickness; and the cicatrices become paler, but are marked by slightly elevated lines or ridges radiating from a central point, and are sometimes marked here and there by small vessels. Still later the cicatrices become almost white,

and more or less contracted, and irregularly depressed. The contraction gives rise to various deformities, or increases the deformities already produced by the preceding destruction of tissue. The point of the nose may be drawn down; or one or other of the alæ may be turned upwards, or the eyelids or lips may be everted. The probable stability of the cure may be inferred from the character of the cicatrix. If the new skin is soft and free from tenderness, and soon presents a nearly natural colour, it may be considered sound; but if on the other hand it remains for any length of time unnaturally red and indurated, or if it continues delicate and membranous in appearance, and is traversed by numerous tortuous blood-vessels, there is considerable probability of a relapse.

A well-established tendency to relapse, or to reappearance of the disease in some part before unaffected, is a noteworthy feature in the clinical history of lupus.

The interstitial changes which take place during the progress of lupus have been made the subject of careful microscopic investigation by Wedl, Auspitz, Rindfleisch, Neumann, and others.* It would appear that in the earlier stages the cells of the rete Malpighii multiply rapidly, and become granular and fatty; the corium becomes succulent, and is infiltrated, and more or less distended with new connective tissue elements, rounded or oval cells with nuclei and nucleoli. The subcutaneous areolar tissue is similarly affected to a variable extent. The normal tissues are no longer healthily nourished, and consequently become atrophic. During the later stages the atrophied tissues and the new elements break down, and become transformed into a molecular detritus. This, mingled with free exudation, helps to make up the characteristic discharge from the ulcerating surface.

The diagnosis of lupus exedens may sometimes be difficult during the earlier stages. But when the stage of gradual advancing ulceration is reached, the disease is as a rule easily recognisable. The affections with which it is most likely to be confounded are 'common strumous ulceration,' syphilitic ulceration, and rodent cancer. It is of great practical importance to distinguish accurately between these several affections, because each requires different treatment.

* See Wedl, *Pathological Histology* (Sydenham Society Translation), p. 333; Auspitz, quoted by Bazin, *Dict. encyclop. des Sciences médicales*, art. 'Lupus'; Neumann, *Lehrbuch der Hautkrankheiten*, 1849, p. 218.

constitutional tendency ; secondly, that when once established 'the lupus process,' if uninterfered with, may go on spreading during an uncertain period, and to an indefinite extent, as if it were to propagate itself; thirdly, that sooner or later the morbid processes come to an end, tissue destruction and healing ensues.

The natural termination of lupus which thus takes place may be considered to be due either to the disease having, so to speak, worked itself out, or to constitutional changes, from advanced age or other cause, having rendered the general condition favourable to the continuance of the local affection. The facts regarding the natural history of lupus are strongly supported by the results of different methods of treatment. And if they serve to indicate the double direction the treatment should take in order to effect or hasten the cure. Experience has amply shown that, as a general rule, comparatively little permanent good is to be expected from local remedies alone. Various sorts of stimulating and soothing applications have been repeatedly tried in vain. In many cases, even after the most powerful caustics, the characteristic ulceration has again recurred when the immediate effects of the escharotic had passed away. So also the most judicious constitutional treatment has been tried over and over again, and persevered in for months, or even for years, without obviously producing any good effect, unless or until proper local treatment has also been adopted. It may, and probably does, occasionally happen that the disease is prevented from becoming established by the application of caustics, or that when the morbid process has nearly worn itself out, a cure is effected under the influence of constitutional remedies. But in the majority of cases it is to trust to neither one nor the other line of treatment alone, but to adopt both. And if this be done judiciously, it will be found that lupus exedens is by no means the unmanageable malady as was formerly supposed to be, and that it is far from deserving its common appellation of 'noli me tangere'—an appellation that by misleading may prove mischievous.

In the treatment of lupus exedens it is, in the first place, desirable to correct as far as possible all that may be wrong in the general health of the patient by improved hygiene, and by medicines as are calculated to remove or overcome any disorder of the digestive system that may be found to exist. Good food, including a liberal allowance of meat and

Vegetables, and some wine or malt liquor, are especially indicated. Sea air occasionally appears to be hurtful rather than beneficial; and in some cases alcoholic stimulants of every kind seem to disagree: but these are probably exceptional cases, though several such have come under my observation.

Various tonics, as the mineral acids, vegetable bitters, quinine, and some of the preparations of iron, especially the syrup of the chloride or the syrup of the phosphate, may be advantageously administered, and varied from time to time—due regard being always paid to the condition of the digestive organs, and to the action of the bowels. But the medicaments which are especially useful, are cod-liver oil, and the liquor arsenicalis, or Donovan's Solution. Cod-liver oil should always be administered in such doses as the stomach will bear. Its influence for good in the large majority of cases is marked and indisputable. In most cases also arsenic appears to be a very valuable remedy. Mr. Hunt, to whom is due the credit of having especially urged, not of having first suggested, its administration in this malady, considers it 'absolutely a specific.' He seems inclined to rely on it almost exclusively.* He recommends it to be given in the largest doses, ascertained by careful trial, that the patient can bear continuously without injury. In some cases very large doses are tolerated.† The results of general experience testify to the beneficial effects of arsenic, but seem scarcely to confirm all that Mr. Hunt has said in its favour. Indeed, it is worthy of note that he refers to the possible necessity of continuing the use of the drug, 'month after month and year after year,' and even 'for half a life-time.' Any 'specific' influence it may possess, therefore, may be at best very slow in becoming manifest, and can scarcely be considered worth waiting for, when a more speedily effective treatment is at command.

Due regard, then, being given to the general health, and such medicaments being administered as are likely to modify the morbid constitutional tendency, it is, in the second place, of the greatest importance to arrest locally the spread of the disease. This can be effected most speedily and most certainly by the absolute destruction of those parts and tissues in which the

* See *British Medical Journal*, 1862, vol. i. p. 8; also, *On the Treatment of Lupus*, by J. L. Milton. (Hardwicke.) London, 1866.

† See a remarkable case detailed by Dr. Andrew, in which the patient took half drachm doses of the liq. arsenicalis three times daily for a considerable period.—*British Medical Journal*, Sept. 8, 1866.

'lupus process' is going on. It matters comparatively little what method is adopted, or what particular escharotic is provided always that the application is thorough and efficient. It is worse than useless simply to irritate the surface of the sore; and, on the other hand, it is obviously undesirable to penetrate further than needful into the healthy tissues. No method may claim to be the best, which, on experience, is found to be most completely under control during application, and which causes least enduring pain afterwards to the patient. Judged on these grounds, the actual cautery, strong nitric acid followed by nitrate of silver, and chloride of zinc stick may be especially recommended. Their destructive influence is certain, speedy, and readily limited and controlled in accordance with the judgment of the surgeon. Chloroform is administered at the time of the application; and the pain subsequently experienced, as a rule, is neither very severe nor long. On the other hand, the pastes of chloride of zinc and nitric acid and sulphur, acid nitrate of mercury, arsenious acid and calomel, &c. &c.,* although effective, are open to the objection that, acting slowly, they often subject the patient to a long agony. Moreover, during their application the surgeon has no opportunity of limiting or controlling their action, and judging how far their destructive influence may have penetrated.

The actual cautery may be used in the form of the red-hot white hot iron, the galvano-caustic wire or pencil of Middeldorp or the gas cautery of Bruce. The last named is a very convenient and efficient instrument for use in cases in which the ulcers are small and superficial. The hot iron and the galvano-caustic pencil are more applicable if the ulcers are deep, or if it should be necessary to destroy much tissue.

Nitric acid may be applied by means of a glass rod, a glass brush, or, best of all, a piece of rattan cane frayed at one end, and cut to convenient size. The acid must be thoroughly applied along the ulcerating edges, and well pressed into the surface of the sore, and into all the irregularities

* The following formulæ may be quoted:—Chloride of zinc 1 part; gypsum, or compound tragacanth powder 3 or 4 parts; water q. s.; Chloride of zinc 3iv., chloride of antimony 3ij., starch 3j., glycerine q. s. (Startin). Vienna paste: Potassa fusa and unslaked lime, equal parts; of wine, q. s.; mix. Arsenious acid 1 part, calomel 99 parts (Dupuyroux). Calomel 3ij. ss., bisulphide of mercury 3ij., arsenious acid 3j.; mix (Startin).

any excavations. After a few minutes the whole surface should be wiped dry and comparatively clean, and then a fresh application of the acid should be made. This process should be repeated until it is clear that all the lupus-affected tissues must have been reached by the caustic. It is then a good plan to rub a stick of nitrate of silver all over the surface, and well into the cauterised tissues. The silver salt assists in forming a good protective scab, and certainly seems to deaden and diminish the pain produced by the nitric acid.

The solid chloride of zinc in stick is especially recommended by Mr. Hutchinson. It must be thoroughly rubbed along the edges and well pressed into the surface of the ulcer. It is needless simply to touch or wipe over the parts affected: the caustic must be made to penetrate.

In all cases before any escharotic is applied, it is necessary to clear away all crusts and scabs from the surface and edges of the ulcer. They may be softened by bathing with warm water, or by a short application of an emollient poultice, and may then be picked off. If the ulcerated surface and the surrounding skin appear much inflamed, it is better to attend to the general health, and to apply soothing ointments and lotions for a time before proceeding to the use of escharotics. Poultices and water-dressings as a rule are not suitable. In applying the escharotic, care should be taken to follow accurately every extension of the disease on the mucous membrane and into the cavity of the nose, as well as on the cutaneous surface. The escharotic must be applied to every portion of the ulcerating surface, especially to the edges along which extension of the disease is going on; and some portion of the surrounding apparently healthy skin should also be destroyed. The healing edges and the cicatrised portions of the surface need not be touched.

If the patient is under chloroform as recommended, all that seems necessary may be done at one operation. But if the disease is extensive, and chloroform is objected to, different parts may be attacked at different times, one part being destroyed after another.

After the application of the escharotic, the surface should be carefully covered from the air by a little scraped dry lint, and a light pad of cotton wool. This dressing may be kept on until the scab or slough separates, which usually happens in the course of four or five days. Benzoated zinc or lead ointment may then be applied, or such other dressing as may be indicated.

If the escharotic has been thoroughly applied, healing commences, and as a rule goes on satisfactorily, provided the general health is good, and that proper constitutional treatment is carried out. It sometimes happens that repeated applications of the escharotic may be necessary; and this is almost sure to be the case if the former applications have been too superficial, or if the escharotic employed has been too feeble in action. Some surgeons indeed prefer the repeated application of weak caustics to a single application of a strong one. The Hebra recommends the application of solid nitrate of silver once or twice a week. Such treatment has proved efficacious in very many cases. Hardy * speaks highly of the efficacy of an ointment composed of equal parts of biniodide of mercury and lard. The ointment of the red oxide of mercury, or that of the ammonio-chloride, or the ointments of lead, zinc, and nitrate of mercury mixed in equal proportions, may also be recommended in cases in which escharotics appear unnecessary, or in which it is deemed desirable to defer their application. The ointments are often very useful also after partial or incomplete cauterisation.

It must be borne in mind that although lupus is not of syphilitic origin, yet it may and sometimes does occur in those who are the subjects of syphilis. In such cases the appearances presented partake of the character of syphilitic ulceration, and the treatment must be modified accordingly. If the syphilitic taint is hereditary, mercurials prove useful. The iodide, mercury and chalk, or the bichloride in small doses may be given. If the patient is the subject of acquired syphilis, and especially if mercurials have been previously administered, iodide of potassium in full doses (twenty to thirty grains) three times a day, and iodide of iron are especially indicated. In such cases it is well to defer the use of escharotics, and to apply black lotion or one of the weaker mercurial ointments.

Lupus Non Exedens.

Lupus non exedens is slower in its progress, and less destructive than lupus exedens. It commences, according to the excellent description of Mr. Wilson,† 'as a small tubercle of

* *Leçons sur la Scrofule et les Scrofulides*, p. 89. Paris, 1864.

† *Op. cit.* p. 422.

a yellow, or pale amber colour, and has the appearance of jelly effused beneath the cuticle. . . . There is no inflammation, no redness around the tubercle, and a few minute vessels may be seen struggling through it, or over its surface.' The tubercle goes on extending, and is joined by others; but the yellow tint and the somewhat 'gelatinised' appearance is preserved. By and by absorption commences; the papillae, and more or less of the deeper parts of the corium are removed; and a permanent cicatrix remains. No ulceration takes place. A certain amount of pricking or itching is experienced, but as a rule very little pain, or none at all. Sometimes the disease spreads over the greater part of the face, and considerable deformity results.

Treatment.—The treatment generally should be similar to that commended in lupus exedens. But it is not necessary to excise the parts affected nearly so deeply nor so extensively in the more severe form of the disease. Caustic potash appears to be especially applicable. Tubercle after it may be destroyed at different times without occasioning much suffering to the patient. Among the constitutional remedies, iodide of iron and iodide of ammonium are found to be obviously beneficial than in lupus exedens, but arsenic does not appear to be so efficacious.* Cod-liver oil should be administered in such doses as can be tolerated.

Lupus Erythematosus.

This form of lupus usually appears on the nose; but other parts are sometimes attacked at the same time or subsequently.

It is a comparatively rare affection. It differs from the other forms of lupus not only in its non-tubercular character, but also in the fact that it commonly occurs in individuals who are otherwise healthy, and who manifest no indications of strumous disposition or tendency. Its general features have been already described (see page 239).

The disease begins with dull or purplish red patches, with which it commences, and as the period becomes depressed, and covered by scales or thin crusts composed of epidermis and dried sebaceous secretion. These scales and crusts separate, and the skin beneath is found to be attenuated and to present a scar-like appearance.

* Hunt, op. cit.

DISEASES OF THE NOSE.

According to Neumann, in lupus erythematosus the morbid process commences in and about the walls of the sebaceous follicles, which become condensed by accumulation of cells and connective tissue. The follicles lose their acinous structure and become converted into globular bodies with granular contents and finally disappear. During the progress of the malady other cutaneous structures suffer to a greater or less extent, the microscopical appearances more or less resemble those with in the other forms of lupus.*

Treatment.—In a large proportion of cases local treatment alone is requisite. The patches should be carefully destroyed by the application of strong solution of potash or some similar agent; or the gas cautery of Bruce or the galvanocautery wire may be used. It is better to proceed gradually, and repeat the application as often as may seem desirable, rather than to run the risk of destroying the tissues to an unnecessary depth and extent.

CANCEROUS AFFECTIONS OF THE SKIN, ETC.

Rodent cancer.—This formidable malady most frequently commences near the eyelids, or about the margin of the lip; but sooner or later, if unchecked, it usually invades the nose, and gives rise to the most hideous deformity. It occurs in persons who are otherwise healthy and free from all constitutional taint, but very rarely in those who are over fifty years of age.†

It first appears as a hard tubercle, the induration of which is characteristic. The tubercle is almost always solitary at first, but at a later period other tubercles occasionally, though rarely, appear and join the original one. The tubercle slowly but steadily increases in size, and involves the surrounding tissues. By and by a superficial fissure or excoriation appears, which for a time may be concealed by a scab. But later the fissure or excoriation becomes a deeply-eating ulcer; and the central portions of the growth and the tissues involved by it are destroyed. The process continues. The surrounding tissues

* Op. cit. p. 230.

† In his excellent monograph *On Rodent Cancer*, recently published (London, 1867), Mr. C. H. Moore mentions a case in which the growth appeared at the age of thirty-three. Mr. Hutchinson met with another in which the patient was only a year or two older.

ded by the solid growth or deposit; they become more indurated; and are then in turn regularly eaten away. Growth of the solid disease is not limited to the integuments the ulcerated cavity: it advances in depth as well as in superficial area, involving all the structures it encounters. It reaches the glandular textures and the bony, it spreads by the areolar and fibrous tissues, and after perforating the skull, it will grow into the very substance of the brain. In the progress of ulceration these structures are likewise removed, and the excavation becomes deep as well as wide.*

In the earlier stages there is comparatively little pain, but as the destructive process advances into the deeper parts, and especially when the eyelids are attacked, the patient's sufferings are often very severe. The lymphatic glands are rarely

This malady has been described under the names of 'rodent canceroid,' 'Jacob's ulcer of the eyelids,' &c. &c.

Its alliance to cancer is indicated by its clinical history, rather than by its microscopical structure. Its onward progress is, though very slow. It never undergoes spontaneous removal; by surgical aid, it lasts until death.

Treatment.—Constitutional treatment is altogether useless. Local excision, followed if necessary by cauterisation of the raw surfaces, or thorough destruction by powerful escharotics, is the only treatment likely to be followed by successful

Experience shows that by such treatment much good may be done. In many cases a cure has been effected which has lasted permanent for years; and in others the ravages of the disease have been checked, and the sufferings of the patient diminished. If the nature of the malady is recognised, and all the parts infected are at once absolutely removed, the patient may be saved from the most serious disfigurement and the direst suffering. And even in cases in which the disease has made extensive progress, the surgeon should not hesitate to attempt its removal. The results obtained in some cases have been decidedly encouraging.† In cases in which the disease has not advanced far, and in which the whole may be satisfactorily excised, the space left may be at once

* Moore, op. cit. p. 9.

† Cases by C. H. Moore, op. cit.; Hutchinson, *Medical Times and Gazette*, 'Cut Ulcer;' also *London Hospital Reports*, vol. ii. p. 135.

DISEASES OF THE NOSE.

covered in by a portion of transplanted integument. But in cases in which the greater part or whole of the nose is destroyed and the cavity after excision is very wide and deep, it is better to wait until cicatrisation is complete, and subsequently to perform if practicable some reparative operation. If the defect is too extensive to be covered in by operation, or if the health and powers of the patient cannot be relied on, a mask with an artificial nose, &c., may be worn.

Epithelioma.

This form of cancer occasionally, though very rarely, attacks the nose. It generally commences as a small 'wart'; sooner or later becomes excoriated, and covered by a crust; after a time ulceration commences. In these earlier stages epithelioma closely resembles rodent cancer, and is difficultly distinguished from it. But epithelioma usually commences just at the junction of the skin with the mucous membrane, or even in the mucous membrane. Rodent cancer as a rule appears in the skin at an appreciable distance from the mucous membrane. In their more advanced stage the distinctions between these two allied affections become

obvious. In both there is slowly progressive infiltration of surrounding structures with new material, and destruction by ulceration extending centrifugally. But 'there is much solid substance in the epithelioma, and the gaps which it makes by destroying the normal parts, though equally great, are less openly cavernous than in the rodent cancer.*' The outline of epithelioma is less irregularly indented, and the margins are less uniform in thickness than in rodent cancer. In epithelioma too the margins are often everted, and the surface is uneven and irregularly papillated and sometimes covered with a clear like juice. In epithelioma the lymphatic glands are liable to become affected sooner or later. In rodent cancer, as already stated, this is not the case. The generally recognised microscopical characters of epithelioma (see Vol. I. p. 61) are almost or altogether wanting in rodent cancer.

Treatment.—Epithelioma of the nose may be excised and destroyed by escharotics.

* C. H. Moore, op. cit. p. 24.

Ure refers to a well-marked case in which he effected a cure by means of mercurial paste. He mentions also another case in which an oval epithelioid sore as large as a florin was treated first by excision, next by the application of chloride of zinc, then by the application of arsenious acid, and finally under the influence of an ointment composed of equal parts of finely-sieved sulphate of zinc and cerate. The cicatrix was perfect a year after-

*
 remarkable case is mentioned by Mr. Moore in the first volume of this (p. 571), in which an epitheliomatous growth commenced on the nose, and spread over the centre of the face until it reached the size of half a 'egg.' 'The entire surface of the growth was made up of pedunculated and tumours no larger than peas, and covered with thin integument.' A similar case recently came under my care in Guy's Hospital. I removed the growth as completely as possible with the knife, and then cauterised the wound with the hot iron. Cicatrisation took place. But four months after the growth returned, and a second attempt to remove it was followed by an attack of erysipelas which proved fatal.

irrhus and encephaloma sometimes, though very rarely, arise upon the cutaneous aspect of the nose; they more frequently invade it from within. These forms of cancer demand special description here. They should be removed if practicable as soon as recognised; and it is well to cauterise the nose left bare by the knife.

DEFORMITIES AND DEFICIENCIES.

The nose is liable to various deformities and deficiencies. Some of them are congenital; others result from accident or disease. Some are more or less perfectly remediable by operation; others are best concealed by masks or artificial noses, which are now-a-days made and adapted with great skill. *Entire absence of the nose* as a congenital affection is extremely rare, but some instances are on record.

In the case of a female infant under the care of M. Maissonneuve,† the nose was represented by a plane surface merely pierced by two small holes. An operation was performed by which a columella was made from the lower lip. It is difficult, however, to 'see how this could have improved the child's appearance.' (See p. 128.)

When the nose has been entirely destroyed by disease, the mouth may be to some extent covered in by operation. But it

See Former Edition of this Work.

Bulletin de Thérapéutique, 1855, vol. xlix. p. 559; quoted by Holmes, *On the Surgical Treatment of Children's Diseases*, 2nd ed. p. 128.

would appear to be useless to attempt to form a new nose, natural and necessary framework of cartilage and bone together lost. Nothing short of an osteoplastic operation seem likely to be permanently successful. But the objections to such an operation in this part are sufficiently obvious; and so far as I have been able to learn, the results of the attempts hitherto made have not been encouraging.

In cases, however, in which some portion only of the nose is wanting, whether as a congenital deficiency or as the result of an accident or disease, a rhinoplastic operation may be taken, and often with fair prospect of success. In very many instances the most satisfactory results have been achieved (see PLASTIC SURGERY).

Occlusion and contraction of the nostrils.—These deformities of the nose are sometimes congenital, but much more frequently are the result of corresponding conditions of the ear, the anus, or the mouth. In some instances the nostrils are more or less completely closed by membrane,* in others by firm fibrous tissue, or by the continuity of integument. In other cases, again, one ala may be adherent to the septum, or even to the upper lip. In all these deformities interfere with respiration, and prevent the infant from sucking uninterruptedly. The earlier therefore these are remedied by operation the better. In most instances a simple incision of sufficient extent carefully made through the obstructing membrane is all that is requisite. The nostril must be kept patent by strips of lint or a short elastic catheter until the cut surfaces are completely skinned over. Sometimes it may be desirable to excise a portion of the obstructing membrane. In cases in which there is no indication of the opening of the nostril, the adherent parts must be gradually and cautiously divided until the nasal fossa is reached. If free respiration is not materially interfered with, or if one nostril only is obstructed, the operation may be deferred for a time.

In a case under the care of Mr. T. Smith in the Hospital for Sick Children, the right nostril had been closed from birth. The corresponding ala was also deformed and unsightly. The child suffered from snuffling and snoring. The adherent parts were divided by the knife, and the opening made was maintained for a period of three weeks by means of a portion of gum catheter. A good result was obtained.†

* Richerand, *Nosographie chirurgicale*, 4th edit. tome ii. p. 156.

† *Medical Times and Gazette*, March 28, 1863.

se, which becomes more or less nasal in character; in certain states of the weather, an uncomfortable dryness of the mucous membrane of the mouth. If the patient suffers from inconvenience, an attempt should be made to dilate any opening that may remain by the passage of bougies, or the introduction of sponge or laminaria tents. If such fail, incisions should be made in the direction and to the extent indicated by the particular features of the case under treatment. Mechanical dilatation must be subsequently kept up for a considerable period to prevent the contraction which is almost certain to recur.

Fissures of the nose and fissures extending from the nostril into the orbit towards the orbit, are occasionally met with. As a rule they may be successfully treated by plastic operations.

Deviations of the nose.—In some cases the nose inclines to one side or the other. Such deviation usually depends on a corresponding curvature of the cartilaginous septum. It is often associated with hypertrophy of the anterior extremity of the septum, and of the tip of the nose generally. In consequence of the deviation of the septum, one nostril is more or less constricted, and may even be almost completely occluded. Respiration is principally or entirely carried on through the other nostril, and the voice is liable to be affected. Deformity is sometimes the effect of injury, but in many cases its origin cannot be explained. Little can be done in the way of remedy. The attempts hitherto made to straighten the deviated nose by means of mechanical contrivances do not

II. AFFECTIONS OF THE INTERNAL PARTS.

The cavities of the nose may be inspected to some extent either through the nostrils, or through the naso-pharyngeal openings by means of a small mirror introduced into the pharynx.* They may likewise be explored to some extent by the finger, or by a probe, passed either through the nostrils or through the mouth, and round the posterior border of the soft palate. But in the normal condition of parts, certain considerable portions of these cavities can neither be brought into view nor reached by the finger or probe. Hence an absolute and precise diagnosis is often difficult, and sometimes in the present state of our knowledge, and with our present appliances altogether impossible.

RHINOSCOPY.

Anterior rhinoscopy, or inspection through the nostrils.—In order to inspect the nasal cavities from the front, the nostrils must be dilated, and a good light must be thrown in, either directly or by reflection from the mirror hereafter described.† The latter

FIG. 225.



Ordinary Bivalve Nasal Speculum.

method is generally preferable for minute and thorough examination, because by it the light may be concentrated and directed at will, upon different parts in succession, and no obscuration is caused by the head of the observer, as is apt to occur when direct illumination is employed. For the dilatation of the nostrils no instrument is more generally useful than the ordinary

* To Czermak especially, and after him to Semeleder, we are indebted for having first completely demonstrated the practicability and value of posterior rhinoscopy.

† For methods of illumination, &c., see 'Posterior Rhinoscopy,' p. 258, and 'Laryngoscopy' in DISEASES OF THE LARYNX.

nasal speculum (Fig. 225), or some slight modification of Hilton's modification * is probably the best. He recommends that the blades should be slightly curved, and somewhat wider and broader at the extremities, than they are in the instruments. The blades are introduced into the nostril and examined in the closed position in which they are kept by being brought between the handles. By compressing the handles the blades are separated, and dilate the nostril. Their brightly polished inner surfaces help to throw the light into the cavity. The blades should not be introduced beyond the cartilaginous part of the nostril.


Hudichum's speculum consists of two blades or valves joined together by a spring. 'When compressed the valves form an oval bore, about $\frac{3}{4}$ -inch length. The valve which is applied to the ala is from $\frac{1}{4}$ -inch to $\frac{1}{2}$ -inch longer than the valve which is placed against the septum. Each nostril requires its own speculum.' The valves are compressed and introduced into the nostril. When the compressing force is withdrawn the valves are separated, and the nostril is dilated by the action of the valves, which also serves as a speculum.

This speculum has obtained indisputable merits of

But it often occasions distress unless it accurately fits the nostril, and the extent to which the nostril is stretched is not so completely controlled as in the expanding power of the hand of the surgeon. At any rate, is the result of experience hitherto.

The speculum of Dr. Metz of Chapelle, consists of two portions. One of them is held in the right, and the

FIG. 226.



The Nasal Speculum of Dr. Metz.

* Made by Milliken, Southwark Street.


other in the left hand. They may be used together or separately. It is easy to understand how by means of them the nostril may be dilated, and light reflected into the cavity. To prevent dimming of the polished surfaces by deposition of moisture, as well as the other specula should be slightly warmed before introduction into the nostril.

In some cases, as suggested by Czermak,* a small oval mirror mounted at an angle upon a long slender stem, may be used to great advantage. When such a mirror is introduced to a certain or less depth through the nostril, and properly illuminated, various parts which could not otherwise be brought into view.

Posterior Rhinoscopy. Choanoscopy. Inspection of the Nasal Cavities from Behind.

In order to inspect the nasal cavities from behind, a strong light is especially necessary. The instruments required are similar to those used in the examination of the larynx.

FIG. 227.



The general arrangements for posterior rhinoscopy by sunlight.

In addition to the addition of a long narrow spatula turned up at the end into a blunt flat hook, with which the uvula and soft palate may be supported and drawn forwards. The rhinoscopic mirror, however, is smaller than any ordinarily used in laryngoscopy, and its stem should be a little longer, and bent somewhat downward near its junction with the handle. This bend enables

* Czermak, *On the Laryngoscope* (Sydenham Soc. Translation), p.

operator to keep his hand in such position as not to obstruct the light. Sometimes the stem may be advantageously bent into a double curve, somewhat like the letter *f* reversed.

In making the inspection the same general arrangements are necessary as in laryngoscopy (see Fig. 227). The patient is seated with his back to the light, his head is thrown well back, and his mouth opened as wide as possible, and in such manner as to display the pharyngeal cavity to the fullest practicable extent. The surgeon, seated opposite, by means of the spatula, held in his left hand, raises and draws forward the uvula and soft palate of the patient, and directs the light reflected from the frontal mirror into his pharynx. The speculum held by the right hand of the surgeon is slightly warmed, and introduced with its face looking upwards and forwards. It must be carried well into the upper and back part of the pharynx. The light is then made to shine upon it, by some slight movement, if needful, of the head of the surgeon or of the reflector. When all this is successfully accomplished some portion or other of the walls (or contents) of the naso-pharyngeal or nasal cavities may be distinctly seen imaged in the speculum. By varying the position and direction of the speculum other parts may be in turn brought into view. Under the *most* favourable circumstances, the two superior meatuses may be inspected more or less completely, and considerable portions of the mucous membrane covering all three arched bones may be seen; the septum may be examined throughout a great part of its extent, but the view obtained is necessarily very oblique: some portions of the posterior surface of the velum palati may also be seen: lastly, if the mirror is turned towards one side, the lateral wall of the naso-pharyngeal cavity, and the orifice of the Eustachian tube may be brought into view. But the difficulties of posterior rhinoscopy are great, and it is rarely that the cavities may be inspected to the extent indicated: although this is theoretically possible in most cases, and has been actually accomplished in many. Nevertheless, persevering attempts should always be made in doubtful cases. Experience has amply proved that this method of examination, even when incompletely carried out, may afford most valuable aid not only in the diagnosis, but also in the treatment of various affections of the nasal cavities. To insure success considerable practice on the part of the surgeon, and sometimes great patience and self-control on the part of the patient are requisite.

The principal difficulties arise in the first place from peculiarities in the anatomical conformation and proportions of parts. Thus the opening of the mouth may be small; or the tongue may be disproportionately large; or the posterior nares may be narrow; or the free border of the velum palati may be naturally so near to the posterior wall of the pharynx as to leave little or no available space between them. This last is a frequent source of great difficulty. Such difficulties cannot be removed though by patience they may be to a certain extent circumvented. Secondly, there are other difficulties of physiological origin. Thus the tongue may be difficult to manage; or the soft palate may be extremely sensitive; or the natural tendency of the velum to become raised, tightly stretched, and closely approximated to the posterior wall of the pharynx, when respiration is carried on through the mouth, may come into play. All such difficulties may generally be overcome by perseverance and practice. The patient may himself depress his tongue by means of a rectangular spatula, or he may learn to keep it well down in the floor of the mouth. The soft palate, though highly sensitive at first, soon becomes accustomed to the steady tone and support of the flat hook, and in some cases may be voluntarily kept in such position as to render the use of the hook unnecessary. It is not improbable, as suggested by Czermak, that some simple means may be discovered by which the natural sensitiveness of the soft palate may be temporarily diminished. The administration of bromide of potassium, recommended by Gosselin† as a preliminary to the operation of staphyloraphy, appears to me from my own experience worthy of more extensive trial than it has hitherto had, although it has not proved so efficacious as was expected. I have tried the chloral hydrate in some cases, but not as yet in a sufficient number to enable me to express any decided opinion as to its merits. The elevation and tension of the soft palate, and the approximation of the velum to the posterior wall of the pharynx, to which allusion has been made as frequent sources of difficulty, occur naturally when respiration is carried on through the widely opened mouth. But it is quite possible to breathe almost entirely through the nose, even when the mouth is open. When this is done the parts become at once relaxed, the velum falls, and the associated difficulty is at an end. I have found it

* Op. cit. p. 76.

† *Gazette médicale*, avril 14, 1860, p. 223.

st plan to direct the patient to breathe through the nose as though snoring.

The method of posterior rhinoscopy thus described is, I believe, the best yet devised for purposes of examination alone. The speculum and the hook which supports the velum may be moved as occasion may require independently one of the other. But in cases in which it is desired to direct the application of remedies by sight, this method has the obvious disadvantage of occupying both hands of the surgeon. Various instruments have been devised in which speculum and palate hook are so connected that they can be held and manipulated by one hand, the other being left free to operate. The best of such instruments yet contrived is, I believe, that of M. Duplay.

NASAL CALCULI. RHINOLITHES.

Calculus concretions are sometimes formed in the nasal cavities. Instances are rare, but a considerable number have been placed on record.

Bartholin relates the case of a young Danish female, who, after suffering for a long time from headache, passed from her nose several calculi, which in size and shape resembled date-stones. Clauder witnessed the case of a woman, sixty years of age, who expelled from her nose a very hard rounded concretion, as large as a hazel nut. Khern noticed several pisiform calculi which had escaped from the nostrils of a young person suffering from violent headache.* Such and such like cases must be taken for what they may be worth. Many others which are well authenticated, are quoted or referred to by Demarquay in his elaborate memoir on the subject.†

These concretions are for the most part met with in the inferior meatus of the nose. They may originate in the frontal or maxillary sinuses, and from thence pass into the nostrils. They vary in size, and increase by the deposition of successive layers. They may even attain such dimensions as to block up the nostrils, and to cause deviation or partial destruction of the septum. They are uneven on the surface, and may be black, grey, or whitish in colour. In many instances they are formed round extraneous bodies, which thus constitute their nuclei. The nucleus may be a bead, a cherry-stone, an incisor tooth, a small detached portion of bone or tooth, or some other body. Often, however, no definite nucleus can be discovered; and in such cases the cause and starting point of the calculus formation

* Cloquet, *Osphrésiologie*, p. 627.

† *Archives générales de Médecine*, 4^e série, vol. viii. p. 174.

remains obscure. Chronic inflammation of the mucous membrane or of the lachrymal gland may possibly give rise to alterations in the secretions leading to depositions of calculeous matter. Crust-like lumps of dry hard secretion are often formed in cases of ozæna &c., and if any of these should be retained in the recesses of the nasal cavities it is easy to understand how they may form nuclei upon which successive deposits may be made. Such deposition may also be favoured by certain abnormal anatomical conditions; as for example contraction of the nostril, or narrowing of the lower meatus to such an extent as to hinder the due expulsion of the secretions. Graefe considers that the gouty dyscrasia favours the production of nasal calculi. On the whole it would appear that the most frequent cause is the retention of some foreign body in some part or other of the nasal cavity.

The calculi themselves (according to the analyses of Boichardat) consist of the phosphates and carbonates of lime and magnesia, chloride of sodium, and mucus or some other animal matter.

The ordinary symptoms are dryness of the affected nostril accompanied by a sense of weight and fulness; frequently more or less obstruction of respiration; pain of an intermittent neuralgic character referred to the nose or forehead; and sometimes inflammation, and swelling of the adjacent parts, with copious discharge of mucus and pus from the nares. The sense of smell may be impaired or abolished. The eye may suffer sympathetically, and may be suffused or bathed with tears; or there may be epiphora from obstruction of the nasal duct.

In most cases the calculus may be detected on careful exploration of the cavity by means of a probe; or it may be seen on rhinoscopic inspection. The sensation communicated through the probe, and the dull dead sound sometimes emitted when the calculus is struck may be considered diagnostic. But should the calculus be inaccessible to touch as well as to sight, the symptoms manifested may mislead the surgeon. Thus, the presence of a polypus may be suspected on account of the obstruction to respiration that may arise; or some primary affection of the bones may be supposed to exist on account of the deformity of the nose which sometimes occurs, and the accompanying suppuration; or in consequence of the fetid and disgusting odour frequently exhaled the surgeon may be disposed to put the case down as one of 'idiopathic' ozæna.

Treatment.—The treatment obviously resolves itself into the removal of such concretions as may be discovered, and the careful and thorough washing out of the cavities by means of the nasal douche * or syringe.

There may be some difficulty in removing the calculi, especially if they are firmly impacted in the lower meatus, or if they are situated in the upper part of the nose. The most convenient instruments to use are thin-bladed well-curved dressing forceps, and small scoops. But care is requisite in seizing the calculus, otherwise the pressure of the instrument may only serve to push it on more deeply into the cavity. Forceps made in two distinct portions, which may be introduced separately one on each side of the calculus, and afterwards locked together, (like midwifery forceps) would probably prove most applicable. Forceps made on the principle indicated are invaluable for the extraction of foreign bodies through narrow passages. I have used such forceps during several years past for removing foreign bodies from the ears and nose with the greatest ease and most perfect success. Each blade can be in turn insinuated beyond the foreign body or calculus. But in using ordinary forceps one blade is often necessarily withdrawn while the other is being put into position, and as a frequent result the foreign body is pushed further on instead of being seized. Care must be exercised in extracting the calculus lest from its bulk or by the asperities of its surface the surrounding parts should be lacerated.

A case of calculus of the nose is mentioned in the *Gazette des Hôpitaux* for 1859, in which attacks of intermittent pain occurred. These were at first considered to be neuralgic. Subsequently the symptoms were attributed to necrosis of the nasal bones. Ultimately the calculus was discovered and got rid of by trephining at four sittings, at intervals of a fortnight each. The cure was followed by some slight deformity of the nose.

When the calculus is extracted soothing measures must be adopted, as already indicated, to allay the irritation set up, and to cleanse the cavities. If there is reason to suppose that a gouty diathesis or tendency has contributed to the production or increase of the calculous formation, or if any other constitutional taint is discovered, appropriate general treatment should be prescribed, and thoroughly carried out.

* See p. 276.

media of communication between the vessels within the c and those which are distributed externally. The tissues support them are in close relation with bone or cartilage are subject to injuries, abrasions, and ulcerations; the vessels themselves are liable to become distended and con from various different causes. Hence it happens that bleeding from the nose is of very frequent occurrence. In a large portion of cases such bleeding is of trifling importance; in some cases it seems to afford relief, and to be really beneficial. Sometimes, however, it is symptomatic of visceral disease or degeneration of blood. Occasionally, if unchecked and frequently recurrent, it may prove serious, dangerous, and fatal. The blood usually escapes from the nose drop by drop but sometimes it flows in a fine stream. As a rule it comes from one nostril only, rarely from both simultaneously.

Epistaxis may be 'accidental' or 'spontaneous' in origin. *Accidental epistaxis* resulting from a blow upon the nose or without fracture of the bones, is very common. In some cases it may be brought on by a very slight blow, or by unusual exertion, or even by sneezing, or blowing the nose violently. It rarely goes on to a serious extent.

Spontaneous epistaxis arises from many different causes. It may occur as capillary hæmorrhage consequent upon active or passive congestion; or it may result from ulcer extending into some one or other of the larger vessels.

in young women capillary hæmorrhage from the nose is occasionally vicarious of suppressed menstrual flow, and under such circumstances may be considered salutary.

Bleeding from the nose frequently occurs in scurvy, and fever, and in other affections associated with an impoverished state of blood. In persons of hæmorrhagic diathesis it often becomes a source of great anxiety and difficulty.

The epistaxis of declining or advanced life should never be lightly regarded. It is usually preceded by mechanical or organic congestion, which may result from over distension or rupture of the vessels conveying the blood from the brain, and may betoken the existence of some visceral disease of more or less serious character. The blood that flows is often venous in character.

Treatment.—In the treatment of epistaxis discrimination and judgment on the part of the surgeon are demanded as frequently as promptitude and skill. It is one thing to devise and apply the proper means for the immediate arrest of the hæmorrhage; it is another to decide whether or not it is better to make the attempt. In very many cases the conditions which have given rise to the bleeding require treatment, rather than the arrest of the mental and temporary flow of blood which indicates the existence of such conditions. In some instances the non-cessation of periodical or habitual epistaxis may betoken the approach of danger; and in others the sudden arrest of the bleeding by surgical interference may be followed by symptoms of the gravest import.

Accidental epistaxis may usually be arrested by the simplest means. So also as a general rule may the spontaneous capillary epistaxis of early life. Cold applications to the nose and forehead, snuffing cold water up the nose, the introduction of a piece of lint into the nostril, and such like remedies are generally efficacious. The nursery remedy of slipping a cold key down the throat, or far better of suddenly dashing cold water over the face, seems sometimes to answer almost magically;—possibly, as has been suggested, through some reflex action excited in the sympathetic nervous system. In some cases the vertical elevation of the arms as high as possible above the head is successful. The rationale may be considered doubtful, but the fact appears to be well established. This plan has been especially recommended (if not first suggested) by Dr. Negrier of Angiers,* who

states that he has never known it fail to arrest the bleeding. It sometimes answers in cases of severe injury. Malgaigne mentions the case of a young man who had sustained a severe fracture of the nose. Cold lotions, &c. failed to arrest the copious bleeding which ensued. The patient was directed to raise his arms perpendicularly. The hæmorrhage ceased almost immediately and did not recur. Dr. Keith of Aberdeen speaks most strongly of the excellence of this method. Another simple and often very effectual method consists in firmly compressing the nostril of the affected side against the septum. Observation teaches that in a large number of cases the bleeding spot is somewhere near the anterior and lower borders of the septum. In such cases it is obvious that the hæmorrhage may be arrested by pressing upon the nostril or nostril with the finger, or with the finger and thumb. The compression should be maintained for a quarter of an hour or more, and the patient gently relaxed. The patient should be warned against sneezing or blowing the nose for some time after the pressure has been removed. The efficacy of the method is established by ample experience. Dr. Chapman recommends the application of an india-rubber bag containing water of a temperature of 115° to the back of the neck, and gives some remarkable instances of the success of this mode of treatment.* Dr. Graves † speaks very highly of the good effects of dry cupping in some forms of epistaxis. In all cases the head should be maintained in its natural erect position. If the head is bent forwards the large venous trunks may be obstructed, and congestion and continuance of the hæmorrhage consequently favoured. If the head is thrown backwards or the patient is in the recumbent position the blood may flow backwards, and almost imperceptibly trickle down the pharynx long after all bleeding may be supposed to have ceased.

If none of these simple methods prove effectual, a stream of cold water containing in solution some astringent may be directed into the nostril. This may be accomplished by means of an ordinary syringe, or better still by the nasal douche (see p. 276). Or some astringent, as the persulphate or perchloride of iron, or burnt alum, or tannic acid, on a dossil of lint or a piece of sponge may be applied to the interior of the nostril; or powder

* See *Medical Mirror*, February 1, 1870, p. 25.

† *Clinical Lectures*, vol. ii. p. 316.

matico, or tannic acid or persulphate of iron may be introduced either mechanically or by insufflation. But such means, however certainly effectual, should not be recklessly adopted. I have known the introduction of matico cause the immediate arrest of the hæmorrhage with the after consequence of most alarming symptoms.

If these means fail, there remains to the surgeon the easy and certain expedient of plugging the anterior and posterior nares. When this is done it is physically impossible for the hæmorrhage to continue after the nasal cavity is once filled. For want of knowledge of this method more lives than one have been sacrificed. Gross refers to an example.* Two have come under my own cognizance.

The operation of plugging the nares is best performed by means of Belloc's sound. This consists of a silver canula (slightly curved at the distal end, and six or eight inches in length), within which freely slides a stylet some four or five inches longer than the canula. The proximal portion of the stylet is rigid to an extent corresponding in length to the straight portion of the canula. Its distal portion is made of a piece of 'mainspring' which tends to curve round in the proper direction, and is terminated by a perforated button. The operation is performed as follows. The eye of the button is threaded by a long piece of strong silk, and the stylet is drawn up until the button lies against the end of the canula. The instrument thus armed is passed along the floor of the nose into the pharynx. The stylet is then pushed along the canula, and the spring turns round the posterior border of the velum into the mouth. The silk is seized and drawn out through the mouth, and a portion of sponge of sufficient size to fill the nasopharyngeal opening somewhat tightly is attached to it. The instrument is next withdrawn, carrying with it through the nostril the other end of the silk. By means of traction upon the silk and manipulation with the finger, the piece of sponge is then fitted into the nasopharyngeal opening in such a manner as to block it up completely. Lastly, a second plug of lint or sponge is attached to the other end of the silk, and firmly fitted into the nostril. It is obvious that this operation may be accomplished as effectually, though perhaps not so easily, by various other instruments, as for example a

* *System of Surgery*, vol. ii. p. 333. Philadelphia, 1866.

long bent probe, the stripped feather portion of a quill, or a gum elastic catheter. The last named instrument properly threaded throughout its length, and carrying a loop of silk hanging through its eye answers extremely well. The plugs should not be allowed to remain more than about forty-eight hours, for they are necessarily a source of discomfort and irritation, and are liable to give rise to the formation of matter, which may speedily become offensive, and possibly poison the system. Professor Gross* says: 'I have seen several cases where from this cause the patient lost his life, being seized with a low form of fever, attended with delirium, which nothing could arrest. In one of the cases there were marked symptoms of pyæmia.' The plugs may be easily removed by cutting the silk which connects them, withdrawing the anterior plug from the nostril, and pushing the posterior one on into the pharynx, and bringing it out through the mouth by the finger, or a pair of vulsellum forceps, or by means of one end of the silk which may have been left to hang through the mouth. After the removal of the plugs the cavity of the nose should be thoroughly but gently washed out.

In all cases of frequently recurring epistaxis, and especially those which depend on constitutional causes, it is of great importance to carry out the general treatment which the circumstances of the particular case may indicate. Rest, avoidance of excitement, and general hygienic measures must be enforced. In some cases it may be desirable to administer gallic acid or acetate of lead for a time; but much more frequently, especially if the strength has been reduced by repeated bleedings, ferruginous preparations, and *par excellence* the tincture of the perchloride of iron, are indicated.

It is almost needless to add that if the epistaxis is due to the presence of a polypus, or to any specific ulceration, no permanent good can be effected until the polypus is removed, or the ulceration is partially or completely healed.

CORYZA.

The term coryza designates a symptom rather than a disease. It implies an excessive discharge of mucus, or 'running from the nose. The discharge may be watery and clear, glairy and thick, or even muco-purulent. It may depend up

* *System of Surgery*, vol. ii. p. 334. Philadelphia, 1866.

rrhal, strumous, or syphilitic affections, or upon the presence of a polypus. So long as there is no well characterised or, the term coryza is applicable. But if the discharge becomes horribly offensive, or if it has been so from the first, the term 'ozæna' is employed.

Catarrhal inflammation of the pituitary mucous membrane, or, in the popular designation 'a cold in the head,' is by far the most frequent cause of acute coryza. This common malady needs no discussion here; but it is worthy of note that many different affections of the nasal mucous membrane, and even atrophic growths are often incorrectly described as taking origin from 'severe and repeated colds.' In some instances, doubtless, such an explanation is correct. But in others the coryza and other symptoms are the effects and indications, rather than the causes of the development of the special malady that may have arisen.

Chronic catarrhal inflammation is often attended by troublesome persistent coryza, and associated with more or less inflammatory thickening of the mucous membrane, and consequent obstruction to free respiration. This condition is most liable to supervene in delicate strumous children, or youths of either sex. In some cases local treatment by means of mild astringent washes, or the insufflation of powders of similar character, combined with general measures, appears to be beneficial. A warm dry atmosphere, good nourishment, tonics, and cod-liver oil are indicated in most cases of long standing. Occasionally superficial ulceration occurs in cases of simple catarrhal inflammation, whether very acute or of long standing. But such ulceration is a rule is very slight, and, although painful for the time, very soon heals. It rarely causes the discharge to become foetid.

Hypertrophy of the Pituitary Membrane.

The pituitary membrane sometimes becomes thickened and hypertrophied throughout the whole or greater part of its extent, both nostrils being equally affected. More frequently, however, the thickening is limited to those portions of the mucous membrane which cover the inferior turbinated bones. The mucous membrane of these parts is naturally thick; and its thickness depends, according to Kölliker,* not only upon its

numerous glands, but also upon the presence of abundant, almost cavernous, venous plexuses in its substance. This affection is always associated with coryza. Sometimes it follows long-continued coryza. Sometimes it appears to arise spontaneously, from some general or constitutional, rather than local, cause. But in such case it is usually attended by more or less copious watery or mucous discharge, although the tumefaction itself occasionally appears dry on the surface. The other symptoms usually are a sense of discomfort, more or less interference with free respiration, and a variable degree of alteration of the voice. On inspection a red opaque swelling may be seen, somewhat velvety in appearance, and solid to the touch. The extent as well as the precise situation of the swelling varies in different cases, but it is usually symmetrical. Sometimes the turbinated bones are themselves affected, and become unnaturally soft, porous, and swollen. This affection, especially if confined to the mucous membrane covering the anterior and lower parts of the turbinated bones, may be mistaken by the patient, and even on casual examination by the surgeon, for polypus. But there is no difficulty in making the diagnosis. The non-pedunculated, broad attachment of the swelling, its immobility during respiratory efforts and on pressure by the probe, together with its redness and opaque (not semi-transparent, glistening) appearance, serve to distinguish hypertrophy of the mucous membrane from polypus. It is important, however, to bear in mind that this condition may be and indeed often is, concomitant with the existence of polypus in the upper part of the nasal cavity, and that its presence may conceal the polypus from view.

A case recently came to my knowledge in which a surgeon removed a small polypus from a patient who was also suffering from hypertrophy of the mucous membrane. Another surgeon seeing the case declared that there was no polypus, that a grave mistake had been made, and that no operation ought to have been performed. The consequences were not satisfactory. The polypus was allowed to increase, to the extreme discomfort of the patient, until its existence became indisputable, and its removal far more difficult than it would have been at an earlier period.

Treatment.—The hygienic measures suitable in strumous affections generally may be recommended. Cod-liver oil and the syrup of the iodide of iron often seem to do good. Locally, a single leech may be applied to the swollen membrane, or numerous fine punctures may be made with a needle. Such measures are sometimes obviously beneficial. Styptic and as

gent solutions may be injected or applied by means of a camel's-hair brush. Or the surface may be touched from time to time with solid nitrate of silver. As a general rule, however, whatever treatment be adopted the case is likely to prove tedious and troublesome.

In a case detailed by M. Nélaton,* the nostrils were obstructed to such an extent that it was found necessary to remove portions of the bones as well as of the hypertrophied mucous membrane. This was done by means of scissors. The result appears to have been satisfactory, but treatment so severe is very rarely indicated necessary.

Syphilitic Affections of the Mucous Membrane of the Nose.

Various syphilitic affections of the mucous membrane of the nose, with or without ulceration, are often attended by obstinate coryza. Such affections occur commonly during the secondary stages of the malady, and are to be distinguished from the more serious affections which arise during the tertiary stages, and which commence as a rule in the deeper structures. Sometimes, however, in the secondary stages, ulceration, at first comparatively slight and superficial, if unchecked by treatment may penetrate deeply, and involve to some extent the bones or cartilages. In such case the inoffensive coryza may give place to an almost intolerable ozaena. This is especially liable to happen in strumous subjects. Syphilitic coryza commences like a common cold in the head, with increased secretion of mucus, which for a long time remains clear, and of the consistence of thin starch. The discharge is increased by exposure to cold, and by indulgence in alcoholic drinks. If the patient should happen to take cold, the discharge becomes profuse, ropy, opaque, and yellowish or greenish in colour. More or less uneasiness and tenderness are felt in the nostrils, together with a sense of stuffiness or obstruction. The discharge abates at intervals, especially during sleep; the pituitary membrane feels drier than natural, and crusts of dried mucus are found below the inferior turbinated bone. The patient is frequently inclined to blow his nose to clear it of the excessive secretion, and sometimes traces of blood appear in the handkerchief. On rhinoscopic inspection the mucous membrane is seen to be of a darker or deeper red than natural, and here and there superficially eroded. A similar condition

* *Clinical Lectures* (American Translation by Atlee), p. 414.

may extend throughout the recesses of the nasal cavities, and into the ethmoidal cells, and the frontal and maxillary sinuses. It may be attended by headache, alteration in the character of the voice, and impairment or abolition of the sense of smell. Sometimes it appears late; and it often obstinately persists long after all other secondary symptoms have subsided.

Treatment.—The administration of mercury is requisite, or at any rate more likely to prove speedily efficacious, than any other mode of treatment. In cases of long standing, in which the syphilitic symptoms generally have disappeared, this alone persisting, small doses of the bichloride may be given twice daily in a couple of ounces of distilled water, and persevered in if necessary for two or three months, as recommended by Mr. Ure. Or a single grain of blue pill or grey powder may be given every night for a similar period. In the more recent cases the use of the calomel vapour bath* is especially to be recommended. The inhalation into the nose of the mercurial vapour produces a well marked beneficial effect. Mercurial lotions injected or applied by means of a camel's-hair brush are useful. All crusts and accumulations of secretion should be washed away by syringing or by the douche. Mild astringent or detergent solution should be used. Any ulcers or erosions that can be seen, and reached by the aid of the rhinoscope, may be advantageously touched with nitrate of silver, or sulphate of copper.

Syphilitic Coryza in the Infant.

Coryza accompanied by the characteristic snuffling and difficulty in sucking is in a large proportion of cases among the earliest indications of congenital syphilis. In addition to the ordinary constitutional treatment, careful but thorough syringing through the nostrils is especially to be recommended. Weak solutions of chlorate of potash or borax may be advantageously used as washes. When the discharge is cleaned away, it is often a good plan, if practicable, to pencil over the mucous membrane, where it may seem to be fissured or ulcerated, with a weak ointment of nitrate of mercury and oxide of zinc.

If the discharges are not carefully washed away, they not only tend to irritate and empoison the hitherto unaffected part, but they are liable to become highly offensive. When once the

* See Mr. Lee's article on SYPHILIS in the First Volume of this Work.

zænic condition is fully established, the treatment of the case becomes much more difficult and tedious, and the probable ultimate result much more serious. The bones and cartilages even if they escape actual destruction by caries or necrosis, may fail to become fully and perfectly developed, and the nose may consequently present in after life the characteristic appearances of the *nez écrasé*.

OZÆNA.

The term 'ozæna,' like 'coryza,' designates a symptom rather than a disease. It implies a foetid discharge from the nose, which communicates an almost intolerable odour to the breath. The discharge varies in character and quantity in different cases, and under different circumstances. It may be thin and sanious, or thick, tenacious, and semi-purulent; almost colourless, or yellowish, or greenish yellow, or stained with blood; it may be scanty or profuse; sometimes it quickly dries up into thick hard crusts; sometimes the crusts formed are few and thin. But in every case there is the characteristic, though somewhat variable, offensive odour, which often embitters the life of the sufferer by rendering him repulsive to all about him.

Ozæna arises from various causes, and is associated with many different conditions. The treatment requisite varies with the cause of the malady, and the constitutional and local conditions present. It is therefore of the highest importance to form, if possible, an accurate diagnosis, and not to rest satisfied with simply designating as 'ozæna' every offensive discharge from the nostrils. Weber* considers it would be well 'to reject the term altogether, since it serves, as a rule, as a cloak for incomplete diagnosis.' In all cases careful and thorough rhinoscopic examinations should be made from time to time.

Among the common causes of ozæna may be enumerated syphilitic, strumous, and lupous ulcerations of the mucous membrane, with or without caries and necrosis of the bones and cartilages; the presence of foreign bodies; and the retention of fragments of bone that may have become necrotic or carious as the result of injury. There remain, however, a considerable number of cases not assignable to any one of these causes, nor,

* In his excellent article on 'Diseases of the Nose' in the *Handbuch der allgemeinen und specialen Chirurgie*. Pitha und Billroth: Erlangen, 1866. Band iii. 187.

indeed, to any other cause at present well made out. To such cases the term *idiopathic* or *constitutional ozæna* is applied.

Ozæna must be carefully distinguished from foulness of breath depending upon carious teeth, ulcers in the mouth, affections of the tonsils, &c. The distinction may generally be made by noticing the effect produced when the mouth and nose are closed alternately. In ozæna the odour is most perceptible when the mouth is closed, and the patient breathes through the nose. The opposite is the case in affections of the mouth, &c. A source of fallacy sometimes arises from the dribbling or falling of the nasal discharge into the pharynx.

Treatment.—The treatment requisite depends obviously upon the cause to which the discharge is attributable, as well as upon the local and constitutional conditions with which it is associated. But certain general principles and methods of treatment are more or less constantly applicable in all cases. These may be advantageously described at once. The more special indications are stated in connection with the descriptions given of the several affections which are liable to be, and very commonly are, accompanied by ozæna.

In all cases, proper constitutional treatment is of very great importance; but it rarely proves of much avail, unless the necessary local treatment is fully carried out.

In the local treatment of ozæna, from whatever cause, the general indications are—first, to remove all possible sources of irritation and obstruction, whether crusts of dried mucus, sequestra of bone, or foreign bodies; secondly, to deodorise the nasal cavities, and to wash away all foul discharges so frequently and thoroughly as to prevent any accumulation taking place; thirdly, to modify the special morbid condition of the parts from which the discharge is derived.

These indications may be generally more or less completely fulfilled,—first, by any needful use of forceps or other surgical instruments; secondly, by washing out the nasal cavities; thirdly, by inhalation of vapours; fourthly, by insufflation of powders.

The nasal cavities may be washed out either by means of a syringe, or by the nasal douche. The syringe should have a long slender nozzle with a bulbous extremity, perforated by a rose of small holes. Such a nozzle, connected with a small hand ball enema or injection syringe, forms a very useful and generally applicable instrument. But in a large proportion of

ses, no amount of syringing is so effectual, not to say so easy, as the use of the nasal douche. This method of washing out the cavities of the nose is a practical application of a physiological fact discovered by Professor Weber of Halle.* It was some years ago introduced to the notice of the profession by Dr. Thudichum, to whom every acknowledgment is due.† But at present it appears to be less generally known and adopted than it deserves to be. Its value in some cases can scarcely be exaggerated. Indeed, its introduction may almost be said to have done as much to improve the treatment of certain diseases of the nose, as the posterior rhinoscopy of Czermak has done to aid their diagnosis.

The apparatus needful for the nasal douche consists simply of a cylindrical nozzle of horn, vulcanite, or glass, rounded and perforated at one end, and by the other connected with one extremity of a yard or two of elastic tube, through which a constant stream of liquid can be kept up from some appropriate vessel by hydraulic or atmospheric pressure. The nozzle should be three or four inches in length, and of such calibre at its rounded extremity as to fit accurately in the nostril of the patient who is to use it. The close application of the inner walls of the nostril to the nozzle is a matter of considerable practical importance. It is therefore necessary to have nozzles of different sizes for different cases. The rounded end should be perforated by five or six small holes. The elastic tube should be intercepted by a stopcock at a short distance from the nozzle. The far end of the tube may open into the bottom of the india-rubber bag or glass receptacle on stand, as recommended by Dr. Thudichum; or it may be connected with a piece of metal tube, bent syphon-like, and terminated by an inverted funnel-shaped metal, glass, or earthenware footpiece, which may be sunk in the liquid contained in a jug or any convenient utensil.

* Some years ago it was discovered by Professor Weber of Halle, that when the side of the nasal cavity is entirely filled through one nostril with fluid by hydrostatic pressure, while the patient is breathing through the mouth, the soft palate completely closes the choanæ, and does not permit any fluid to pass into the pharynx, while the fluid easily passes into the other cavity, mostly round over the posterior edge of the septum narium, and escapes from the other nostril, after having touched every part of the first half of the cavity of the nose, and a great part—certainly the lower and median canals—of the second half.—Thudichum, *On Polypus in the Nose and Ozæna*, p. 17. (Churchill, 1864.)

† *Lancet*, November 24, 1864.

The method of using the nasal douche is illustrated by 228. The receptacle is filled with the liquid to be used, placed on a level somewhat above that of the head of the patient. The precise height may be varied according to the degree of force with which it is desired that the stream should flow.

The tube is either filled by the simple descent of the fluid if the syphon action is adopted, by pouring a sufficient quantity of fluid into it, and then immersing the footpiece in the jug, or by drawing fluid into the tube after the footpiece has been immersed. The stopcock is then shut off; and the nozzle is inserted well into one nostril of the patient, and held by the corresponding hand. The other hand controls the stop-

FIG. 228.



Method of using the Nasal Douche.

The patient, seated before a basin, and holding his head in a natural erect position or slightly inclined forwards, opens his mouth widely, and breathes through his mouth only. The stopcock is then turned, and a stream of liquid enters one nostril, fills to a certain extent the corresponding nasal cavity, passes round the posterior border of the septum, and finally issues from the other nostril, after having more or less completely washed the nasal cavities on both sides. The force and fulness of the stream may be regulated by the stopcock, also by varying the height of the receptacle above the head of the patient. The extent to which the cavities are filled depends

some degree upon the position of the head. By momentarily compressing the open nostril, the fluid may be made to rise higher than it otherwise would.

It is sometimes almost astonishing to witness the quantity of masts and foul discharges that may be thus washed away with the greatest ease, and with no discomfort to the patient. So long as respiration is carried on through the widely-opened mouth, scarcely a drop of fluid escapes into the lower parts of the pharynx. The moment the mouth is closed, the soft palate falls, and all the disagreeable indications of downward flow of fluid disappear. The liquids to be used with the nasal douche must be changed from time to time, according to the nature and condition of the case. But in all cases some saline solution should be used, rather than plain water, which is found to irritate the mucous membrane much more than a solution of higher specific gravity. As a general rule, about a teaspoonful of common salt, or of a mixture of equal parts of common salt, and phosphate or carbonate of soda, to a pint of water forms the best ordinary wash. The slight alkalinity of the solutions of phosphate and carbonate of soda helps to soften the crusts. To such solutions as bases, various deodorising, detergent, astringent, or 'alterative' remedies may be added, according to the indications afforded in the particular case.

It is scarcely necessary to add, that this method is not applicable in cases in which the soft palate is destroyed, nor in those in which either side of the nose is absolutely, or even very partially occluded. In cases in which the septum is extensively perforated the method is useful, but it obviously loses more or less of its full efficacy.

The inhalation of vapours or of 'atomized' fluids is sometimes very useful. If such vapours should be of irritating character, it may be a good plan to direct their passage, and to defend the other parts, by inserting into each nostril little hollow cones such as may be readily extemporised by folding round a piece of card or paper.

The insufflation of powders may sometimes be accomplished by their being 'snuffed' up by the patient. But this is a comparatively ineffectual way in any case, and altogether inapplicable if the nasal cavities are obstructed. A far better plan consists in blowing the powder up through a small tube inserted into the nostril. A piece of glass tube properly curved, or even a quill, with a bit of india-rubber tube attached, forms

an efficient instrument easily extemporised. The most perfect instrument for the purpose is that which bears the name Rauchfuss's Insufflator (Fig. 229). The powder is introduced into the tube through the opening (a), which is then covered by the sliding tube (b), the extremity (c) is introduced

FIG. 229.



Rauchfuss's Instrument for blowing powders into the Nose.

the nasal cavity, and then by a sudden compression of india-rubber ball (d) the powder is driven out, and diffused in all directions.

Syphilitic Ulceration, Caries, and Necrosis.

The comparatively superficial syphilitic affections of the mucous membrane already referred to (p. 271), if unchecked treatment, may gradually go on until deep and extensive ulceration results, and the bones and cartilages are more or less seriously implicated or destroyed. In such cases the coryza of the earlier stages gives place to ozæna; and sooner or later the general conformation of the nose may be changed. The same results occur, as already stated, not only in acquired, but in congenital syphilis. All cases of long-continued syphilitic coryza, followed, in spite of treatment, by ozæna, demand the most careful watching and skilful management on the part of the surgeon. In such cases intracranial mischief of the most serious character sometimes supervenes. Numerous examples are given by Dr. Weber in his paper on this subject in the third volume of the *Medico-Chirurgical Transactions*. Occasionally pyæmia is set up.

In a case, mentioned by Dr. Weber, cessation of the sanious ozæmic discharge was soon succeeded by head symptoms; four days later the patient was seized with rigors. Other indications of pyæmia came on. Death occurred on the thirteenth day after the first rigor. Post-mortem examination revealed the thromboses of the left ophthalmic vein and cavernous sinus, together with purulent meningitis about the under surface of the left cerebral lobe. Several abscesses were found in the lungs and liver, and purulent effusion in the pleura.

The most destructive syphilitic affections of the nose, however, ~~se~~, as a rule, during the tertiary stages of the malady, and are liable to occur in strumous subjects. It would appear that these affections generally commence as gummatous periostitis.* Caries and necrosis of the bones, softening down of the osseous material, ulceration of the mucous membrane, and discharge of matter in turn ensue. The discharge is very offensive. The disease advances more or less rapidly, until the vomer, the perpendicular plate of the ethmoid, more or less of the cartilaginous septum, and the turbinated bones, as well as the floor, of the nose and other parts may be destroyed. But it is very rarely, if ever, that the experienced surgeon has the opportunity of tracing from the commencement the progress of the malady. He sees the more or less direful results of want of proper treatment, or the effects of mismanagement. He can often only guess at what may have been the early history of the case. Sometimes the bones are destroyed piecemeal, and numerous small fragments exfoliate one after another. Sometimes considerable portions become at once necrotic. Every pathological museum contains illustrative specimens, and almost every hospital surgeon of experience has met with examples.

The most serious consequences may follow either from extension of the mischief to the interior of the skull, or from some incidental or accidental complication.

Roussseau mentions a case in which a large irregular portion of the ethmoid (nearly a fourth part of the whole bone) fell through the posterior nares into the throat and almost caused suffocation. The patient died next day with cerebral symptoms, probably due to extension of inflammation through the cribriform plate.†

But short of such fatal results, the loss of portions even of the osseous and cartilaginous parietes of the nose is liable to be followed by permanent deformity, which may be more or less disfiguring. It therefore behoves the surgeon to foresee, and, if possible, to avert by vigorous and judicious treatment the impending dangers.

It was formerly the custom to attribute the ravages of syphilis, in the nasal region especially, to the abuse of mercury rather than to the disease itself. But I believe most surgeons are now agreed in considering such an opinion erroneous. Many cases

* See the article by O. Weber, in 'Pitha und Billroth,' op. cit. p. 189.

† *Clinique médicale*, tome i. p. 546. Paris, 1868.

of syphilitic caries and necrosis, attended by deformity, are on record, and others present themselves from time to time in which there is no evidence whatever that mercury has been administered, but rather the opposite.

Sir Benjamin Brodie mentions a good example. A gentleman had a chancre on the nose. No mercury was given. Two years afterwards he had extensive disease of the bones of the nose. Epilepsy, mania, and death ensued. Sir B. Brodie had no doubt but that the disease extended through the cribriform plate of the ethmoid and ultimately affected the membranes and the brain.* Dr. Graves alludes to two analogous cases in each of which it is worthy of note the strumous diathesis was pre-eminently marked.†

The diagnosis of syphilitic disease of the nose is not as difficult; but sometimes it is necessary to be very cautious in forming an opinion. It must be borne in mind that caries and necrosis of the bones of the nose, accompanied by ozæna, may arise from various other causes as well as from syphilis.

The history of the case, the aspect of the patient, the presence of other symptoms, especially of general syphilitic cachexia, and the extent of the mischief, &c., afford most important aid in diagnosis.

Ulcers, periosteal swellings, perforations of the septum, and portions of necrosed or carious bone, &c., if present, can only be more or less readily seen on rhinoscopic inspection, or felt by exploring the cavities with the fingers or probe. Sometimes, however, comparatively little can be discovered beyond ozænic discharges and crusts, and some swelling, with at most a few superficial erosions of the mucous membrane. But it must not be hastily assumed that neither deep ulceration, nor necrosis, nor caries exists. For the parts affected may be high up in the nose, or concealed from view by the swelling of other parts, or by the presence of the discharge. It is probable that in all cases of syphilitic ozæna there is deep ulceration of the mucous membrane, and in a large proportion of cases the bones are more or less seriously implicated. It is easy to understand how a small sequestrum of bone may remain hidden in one of the recesses of the nose, and yet serve to keep up the offensive discharge.

Syphilitic periostitis of the bones of the nose is usually attended by violent racking pain in the nose and forehead. In some cases in which the nasal bones, and nasal processes of

* *London Medical Gazette*, 1844.

† *Clinical Lectures*, vol. ii. p.

superior maxillary bones are affected, there may be more or less swelling visible externally, accompanied by pain, tension, and tenderness. The local pain is often associated with severe syphilitic headache, and the sufferings of the patient are extreme, especially so when he is warm in bed.

Treatment.—The importance of local treatment can scarcely be exaggerated. But while such treatment is being carried out, it is almost equally important that appropriate constitutional remedies should be administered.

The local treatment essentially consists in the removal of all sources of irritation, as dried crusts of foul mucus, loose fragments of bone, &c., and in washing away the discharges so frequently and thoroughly that they have no time to accumulate and decompose.

Careful inspection and exploration of the nasal cavities should be made from time to time; periosteal abscesses should be opened early by the knife; all loose or carious fragments of bone should be removed by the forceps or other instruments as soon as they are discovered. Ulcers may be touched with nitrate of silver, or sulphate of copper. A long probe, one extremity of which has been dipped into a little fused nitrate of silver, forms the most convenient caustic-holder for this purpose.

The nose should be frequently and thoroughly washed out, either by the nasal douche, or, if the septum is extensively destroyed, by means of a syringe with a long nozzle. Solutions of chlorinated soda, or of permanganate of potash, may be added to the simple saline basis already described (p. 277). Or chlorate and permanganate of potash together, or chlorinated soda and borax, in solution, may be advantageously used. Dr. Hudichum especially recommends the permanganate. In some cases a weak solution of the bichloride of mercury, or the common yellow lotion is of service. After the nose has been thoroughly washed, the immediate inhalation of the vapour of iodine, or the insufflation of Trousseau's mercurial powder,* often proves very beneficial. The inhalation of iodine is most useful in the tertiary stages of the malady, the mercurial insufflation in the secondary stages.

The constitutional treatment must be regulated in accordance with the stage of the malady and the condition of the

* Red or white Precipitate, 1 part; finely-powdered Sugar, 60 parts. Op. cit. 547.

patient. In the secondary stages, especially if the bones and cartilages are not implicated to any formidable extent, mercurials are indicated. They may be administered either in small doses taken internally,* or by inunction; or the calomel vapour bath may be employed. The last method has the advantage of conjoining local with general treatment, inasmuch as the patient may inspire some of the vapour into the nose. The treatment must be very carefully watched.

In the tertiary stages, in which the affection has commenced as a periostitis, and especially if the bones and cartilages are extensively destroyed, or if mercurials have already been administered at an earlier period, iodide of potassium is indicated, and mercurials are likely to prove prejudicial. Iodide of potassium should be given in full doses (g. \mathbf{xx} . to gr. \mathbf{xxx} . or even more) three or four times a day. It may be conjoined with iodide of iron and cod-liver oil. These remedies are particularly valuable in cases in which there is pronounced cachexia and great loss of strength. Other tonics, as dilute nitric acid, quinine, &c., may often be advantageously given.

Lupous and Strumous Ulceration, &c., accompanied by Ozena

Lupus, as already stated (p. 239) may first attack the mucous membrane of the nose. In this case the diagnosis is not so easy as when the skin is primarily affected.

Lupus exedens commencing in the mucous membrane corresponds very closely to the affection that has been described by Mr. Ure† and other authors as ‘*erosive ulcer of the follicles*.’ The earlier symptoms resemble those of a simple catarrhal affection of the pituitary membrane. The mucous membrane covering the cartilaginous septum usually suffers first. The patient experiences a certain degree of tenderness and sense of fullness, especially after exposure to cold air; and a constant inclination to blow the nose, or to pick away the crusts which are constantly reproduced. Small red angry-looking tubercles can be seen on inspection, and these sooner or later give place to spreading ulceration. But the ulcers preserve to some extent a tubercular character, and the apices of the tubercles are concealed by dry, hard, tenacious crusts. As the ulceration proceeds, the cartilage suffers, and the septum soon be-

* See p. 272.

† See First Edition of this Work, vol. iii. p.

forated, especially if the mucous membrane has been affected simultaneously on both sides. By further extension one ala both may be destroyed, and the bones may be implicated. At last, very serious deformity may ensue from the falling in of the bridge of the nose, or from extension of the disease to the external parts. The discharges vary in offensiveness in different cases, and at different periods.

It is of the highest importance to distinguish affections of this kind from those that are of syphilitic origin. If such distinction is not made, very unjust suspicions may attach to the character of the patient; and the surgeon may be induced to adopt a line of treatment which may prove positively injurious. In making diagnosis, attention must be directed to the history of the case, the age of the patient, the absence of other symptoms of syphilis, and especially the absence of the characteristic cachexia which almost always accompanies the more serious syphilitic affections of the nose, and also to the comparative slowness with which the malady has progressed, and the lateness of the period at which the bones have become affected. The presumptive evidence thus acquired as to the nature of the case receives the confirmation if the little papillary infiltrations of lupus can be recognised, and the tubercles present a compact granular appearance; and still further if the disease progresses very slowly, and there is a manifest disposition to healing here and there, even while the ulceration is extending.

The treatment of lupus has been already fully discussed (p. 43). It may be added, however, that the actual cautery in the form of the galvano-caustic pencil, or platinum wire, or the red-hot iron, is especially well adapted for the destruction of lupus in the nose. Caustic potash and chloride of zinc are not so easily controlled. From their deliquescence they are apt to cover the healthy parts. Solid nitrate of silver, thoroughly and repeatedly rubbed in, is sometimes efficacious. Mr. Ure recommends the application of a thin layer of a paste composed of one part of chloride of zinc with two or three parts of gypsum powdered tragacanth, with a little alcohol. Sometimes after ulcers are healed, a sense of rawness or tenderness of the internal membrane remains, depending upon chronic inflammation.

In such cases, mild astringent washes are useful; and the nostrils may be advantageously plugged with loose drosses of cotton wool, so as to obviate irritation from the passage of cold

Ozæna, associated with Ulceration, Caries, and Necrosis from other causes.

Simple catarrhal ulcerations of the mucous membrane are usually only superficial erosions, and are rarely accompanied by ozæna. In some cases, however, they precede, or are associated with, the so-called idiopathic ozæna described in the next section.

As a rule they heal speedily under simple treatment, when they are not irritated by 'picking' on the part of the patient, or undue interference on the part of the surgeon.

Ulceration of the mucous membrane, with more or less extensive destruction of the cartilages and bones, sometimes occurs among the sequelæ of severe fevers—as scarlet fever, measles, smallpox, typhus, and, more rarely perhaps, erysipelas and chronic pyæmia. The affection may begin during the height of the fever with swelling of the mucous membrane accompanied by pain in the part affected and in the forehead and headache. There is often œdematous swelling of the lower part of the nose and upper lip. Subsequently abscesses may form, and the cartilages and bones may become implicated and perforation of the septum may result. The perforation is generally more or less circular, and has but little disposition to extend.

Injuries followed by necrosis, abscess, and the retention of sequestra of bone, are not uncommon causes of ozæna.

Foreign bodies retained in the nasal cavities are very liable to cause ulceration, caries, and necrosis, and to give rise to distressing and obstinate ozæna. In such cases the presence of the foreign body is often unsuspected, and only discovered after a long period of suffering on the part of the patient.* The necessity for careful examination, and the treatment required are sufficiently obvious.

Idiopathic or Constitutional Ozæna.

There remain certain cases, and these not a few, in which, even after the most careful and repeated inquiries and examinations

* A very remarkable illustration is afforded by a case detailed by Dr. W. Hickman, in which a steel ring was removed from the posterior nares of a girl sixteen years of age, after having been lodged there for more than thirteen years, during which period it had occasioned horrible ozæna and other severe symptoms.—*British Medical Journal*, Sept. 28, 1867.

surgeon may utterly fail to discover any obvious cause the horribly offensive discharge from the nose which renders patient's life miserable, and his society almost unendurable. Neither caries nor necrosis of the bones can be made out, nor an ulceration of the mucous membrane, or at most only superficial excoriation. But the mucous membrane generally appears unnaturally and irregularly reddened, and often attended and atrophied in some parts and swollen in others. The colour of the discharge is altogether indescribable, but it cannot be mistaken.* The discharge varies in abundance, and the crusts which are formed vary in thickness and dryness. The discharge often becomes worse in young women during the menstrual periods.

It has been suggested in explanation of this form of ozæna that there may exist some anomalous condition of the secretions which renders them liable to undergo a peculiar process of deposition, and that such condition depends upon some unexplained constitutional cause. It has also been suggested that

the offensiveness of the discharge from the nose may be considered analogous to the somewhat similar offensiveness of the perspiration of the feet, axillæ, &c., not unfrequently met with in persons who are nevertheless scrupulously cleanly in their habits. It seems to me not improbable that, at any rate in some cases, the affection may really be the result of a form of lupus nonexedens implicating chiefly the mucous membrane of the upper and back parts of the nasal cavities. The age at which the affection commonly commences, the constitutional diathesis with which it is associated, the comparatively slight detriment to the general health, its obstinate persistence, the appearance of the mucous membrane, its wasting, and the destruction of the sensitive papillary layer as evidenced by the loss of smell, and lastly the beneficial effects of the same remedies that are useful in lupus nonexedens, all seem to indicate the existence of some close relationship between the two maladies.

Idiopathic ozæna sometimes commences during childhood, and in spite of treatment continues until after the middle period of life. When it has continued long, more or less alteration in the contour of the nose is sometimes observed: thus there may be more or less dilatation of the nostrils, and some flattening of

* The French name 'Punaisie' is suggestive, but it fails to convey a full idea of the character of the stench.

the bridge of the nose. The sense of smell is permanently impaired, and sometimes altogether lost.

Treatment.—In the treatment of this form of ozæna the nasal douche is especially valuable. The solution used may be varied from time to time; in some cases one solution, in others a different solution answers best. Solutions of permanganate of potash, chlorinated soda, carbolic acid, &c., may be used to deodorise and cleanse away the discharges, and solutions of nitrate of silver, or bichloride of mercury, or sulphate of copper, or the ordinary mercurial lotions, may be used as so-called 'alteratives;' and alum, sulphate of zinc, tannic acid, &c., as astringents. The strength of the solution must be varied according to the effects produced. Care must be taken that poisonous solutions are not swallowed. It is well to begin with weak solutions, and gradually increase their strength. The smarting produced should not continue for more than a minute or two after the washing has ceased. If it does, too strong a solution has been employed; or the specific gravity of the solution has not been properly regulated by the addition of the common salt or phosphate of soda. After the crusts and discharges have been washed away, iodine vapour may be advantageously inhaled for a few minutes. A little tincture of iodine may be added to hot water and the vapour inhaled; or the patient may 'smell' a portion of solid iodine, or a sponge moistened with the tincture, contained in a small glass or wooden box perforated above and below. Chlorine and bromine inhalations may also be recommended.

In the treatment of this form of ozæna, Trousseau speaks most favourably of the insufflation of various powders. Among those most generally applicable are the subnitrate of bismuth combined with an equal proportion of talc; chlorate of potash by itself, or with seven or eight parts of sugar; borax, similarly diluted; the mercurial powders already alluded to; calomel, with twenty to thirty parts of talc, &c. The effects of the insufflation of the mercurial powders should be very carefully watched, and the insufflation should not be too frequently repeated. Two or three times a day is generally often enough. But the other powders may be used much more frequently, almost ad libitum. In all cases it is important to wash out the nose thoroughly before blowing in the powders. It not unfrequently happens that the thorough cleansing of the cavities, and subsequent insufflation of the powder, is followed by

mediate abolition of the foul odour,—for the odour belongs to the discharge. But such abolition as a rule is only temporary. When the discharge collects again, the odour returns. In some cases the application of solid nitrate of silver to the various parts of the mucous membrane, so far as practicable, does good, though at the time it occasions much pain and lachrymation. Wetzler* recommends the application of creosote ointment, (i. to 3j. cerate) to the mucous membrane by means of a camel's-hair brush.

The special local treatment must be varied from time to time according to the indications that arise. Sometimes one application, sometimes another, seems to do most good. It is necessary to persevere for some time even after the discharge has ceased, or at any rate has lost to some extent its foulness. Relapses are very common; and the patient is too often apt to be encouraged by temporary improvement to discontinue the treatment, before the condition of the mucous membrane is so modified as to render probable the permanence of the cure. The greatest patience is requisite in the management of these cases; and the most careful examination should be made from time to time, for it sometimes happens that some definite local lesion may be discovered, although the case has been long considered to have depended upon constitutional causes.

Cod-liver oil, iodide of iron, iodide of ammonium in small doses, and sometimes arsenical preparations, are the internal remedies which have seemed to be most useful.

GLANDULAR HYPERTROPHY OF THE MUCOUS MEMBRANE.

Attention has recently been directed to a class of cases in which there is more or less swelling or thickening of the nasopharyngeal and naso-palatine mucous membrane, associated with enlargement of the various glandular structures belonging to it. In one group of cases † the racemose and compound follicular glands appear to be especially affected. These glands are most numerous about the posterior nares, round the openings of the Eustachian tubes, and in the roof of the pharynx under the sphenoidal bone. When they become morbidly affected, an ex-

* See Weber, in 'Pitha und Billroth,' op. cit.

† See Dr. Andrew Clark, 'On Naso-palatine Gland Disease,' *London Hospital Reports*, vol. i. p. 211.

cessive quantity of viscid mucus is secreted. At a later period pus-like fluid may be discharged. Still later the mucous or purulent secretion, if retained for some time, may become converted 'into little foetid cheesy masses, which are from time to time extruded through the nose or mouth.' Accompanying these conditions the mucous membrane generally is more or less irregularly swollen. The ordinary symptoms are, 'discomfort, aching, or pain in the neighbourhood of the soft palate and posterior nares; tingling or sense of fulness about the root of the nose; frontal headache; a thick mucous, purulent, or cheesy secretion discharged at intervals, chiefly through the mouth, by means of snorting nasal inspirations, followed by hawking; slight perversions of taste and smell; alterations of voice; sometimes temporary deafness from obstruction of one or both Eustachian tubes, and an abundant secretion of wax in the external ear.'* In all cases presenting such symptoms, rhinoscopic inspection and digital exploration should be made.

The treatment is often tedious and sometimes unsatisfactory. The nasal douche is especially applicable in these cases. Mild astringent solutions are indicated. In some cases I have seen excellent results follow the application of a mixture of tincture of the perchloride of iron and honey (one part of the former to two of the latter), by means of a camel's-hair brush to the surface of the naso-pharyngeal mucous membrane. In other cases a strong solution of nitrate of silver similarly applied, as recommended by Dr. A. Clark, has proved useful.

Attention should be directed to the general health, especially to the condition of the digestive organs, which are frequently disordered in these cases. Purgatives followed by tonics, are generally indicated, and sometimes cod-liver oil and steel.

In another group of cases, especially described by Dr. Wilhelm Meyer of Copenhagen,† exuberant growths or vegetations are met with in certain parts of the naso-pharyngeal cavity. 'These vegetations, when examined microscopically, are found to consist of the so-called 'adenoid tissue,' and are accordingly to be regarded as overgrowths or morbid growths

* Dr. A. Clark, *op. cit.*

† *Proceedings of the Royal Med.-Chir. Soc.*, October 18, 1869. 'On Adenoid Vegetations in the Naso-pharyngeal Cavity,' by Wilhelm Meyer, M.D., of Copenhagen. Communicated by John Marshall, F.R.S.

the closed glandular structures found in or beneath the mucous membrane of the pharynx, the fauces, and the base of the tongue. They vary in form and consistency: sometimes they are solid and firm, and at others soft and highly vascular, and prone to bleed.

The presence of these vegetations in any considerable quantity impedes respiration through the nose, and gives a peculiar 'nasal' character to the voice. There is often some degree of deafness. 'The nostrils are flattened so that the nose appears compressed.' There is deficient secretion from the nostrils, and sometimes blood accumulates in the mouth. Digital examination and rhinoscopic inspection demonstrate the presence of the outgrowths. They are most frequently met with in advanced young people. The treatment consists in removing the vegetations by the knife, or by the galvanic cautery, or by repeated applications of nitrate of silver. Dr. Meyer recommends the use of 'a ring-shaped knife, mounted on a long slender handle, which is passed through one or other nostril, and manœuvred so as to sweep over the mucous surface affected. The bleeding is free, but not excessive.' Two or more operations are sometimes necessary. According to Dr. Meyer, the results of this proceeding, supplemented by the use of the nasal douche, have proved very satisfactory.

AFFECTIONS OF THE SEPTUM.

Besides the deviations and deformities already alluded to (p. 255) the septum is liable to various affections, which from their practical importance merit more special description than has been given in the preceding pages.

These affections for the most part present themselves as swellings or tumours, possessing different characters; some contain blood, others pus, others a gelatinous fluid, and others again have a somewhat cartilaginous consistence.

Blood tumours of the septum are always the result of injury, and generally are formed within a few hours of the accident. The swelling usually occupies both sides of the septum, but it may be confined to one side. In some cases the mucous membrane only presents a flattened elevation, appearing as if raised by uniform effusion beneath it; in others it is more or less disorganized. There is often a good deal of tension and surrounding redness, which render the detection of fluid difficult. The

swelling may generally be seen by aid of the speculum, or by simply pressing up the tip of the nose, and so dilating the nares. It is generally of dark purple colour, and smooth and glossy in appearance. Its base is broad, with more or less abrupt limits. The symptoms are a sense of fulness and stuffing of the nose, with impediment to free respiration proportioned to the extent of the effusion.

Treatment.—The indications are to prevent if possible inflammation and suppuration, and to favour absorption. The application of cold externally, by means either of the small ice-bag, or evaporating lotions, may often prove useful. Incisions are rarely necessary. Jarjavay cites an instance of loosening of the pituitary membrane of the septum caused by effused blood after a bruise. The swelling completely misplaced the cavity of the nose on each side of the septum, and the patient could breathe only by the mouth. An incision was made into the swelling; dark liquid blood was discharged, and a breach of continuity in the septum was detected with a probe.

Abscess of the septum.—Abscess of the septum may result from injury. It is more likely to do so in the case of a scrofulous subject, or in one suffering from fever.* It often occurs when the integument is wounded near the lower edge of the nasal bones. In such case the integument of the nose becomes œdematous and tender on pressure; the pituitary membrane is inflamed, and the portion covering the septum is uniformly turgid; the secretion of mucus is diminished; and there is more or less febrile disturbance. At a later period matter is formed beneath the mucous membrane; and a prominent swelling, causing proportionate obstruction, appears in one nostril or both. The inflammation may spread to the frontal sinuses and lachrymal passages; and headache and epiphora may result. Sometimes the lower margin of the septum, and the upper lip become affected. The swelling itself is red, shining, tender on pressure, and its base is extensive; fluctuation can be distinctly detected, and sometimes is communicated from one side to the other through the septum.

In the *treatment* of such cases it is important to prevent the formation of matter if possible, and if not, to give it free exit by

* See Mr. Fleming's able Article in the *Dublin Journal of Med. Sc.* vol. ix. pp. 16-28.

incision as soon as it is formed. In making the incision the thickened state of the mucous membrane must be borne in mind, and care must be taken to prevent the too early closure of the opening made. The discharge is generally of a thin sero-purulent nature at first, but by and by assumes a glairy consistency. Soothing, and subsequently astringent washes and injections are useful in restoring the mucous membrane to its normal condition.

Chronic abscess of the septum sometimes commences without any assignable cause, and progresses very insidiously. The inflammatory stage is sometimes so faint as to pass unheeded; and the patient is not aware that anything is the matter till the abscess is fully formed; and then suffers not so much from actual pain, as from uneasiness felt in the nares. In these cases the outer surface of the nose is never involved. In this form of abscess there is less redness, tension, and tenderness on pressure than in the acute form. The abscess, besides, is more extensive, and more likely to occur singly on either side of the septum. It may communicate with a collection of fluid in the cavity outside the nose—in the upper lip, for example. The matter secreted has sometimes a highly offensive smell.

The swellings formed by such abscesses may possibly be mistaken for polypi, unless care is taken in the diagnosis. They are still more likely perhaps to be confounded with the thickened elongated condition sometimes presented by the mucous membrane of the septum.

These abscesses should be opened early; and in some cases benefit will be derived from injecting a weak solution of nitrate of silver or some detergent solution into the cavity. Every care must be taken to prevent extension of the disease, and to obtain a cure without any permanent deformity. A guarded prognosis should always be given in these, as in other affections of the septum; for there is always danger of the destruction of the bones or cartilages, or both; and this may take place long after the apparent cure of the malady. In the majority of cases, perforation, with more or less extensive destruction of the cartilaginous septum, may be expected. Mr. Fleming states that this may be effected by interstitial absorption without ulceration of the mucous membrane, in which case the mucous membrane of one nostril adheres to the other; or ulceration may arise,

and then an opening of variable extent admits of communication between them.

When a small permanent opening remains through the septum, a very disagreeable whistling is sometimes perceived when the patient speaks. In one case under my cognisance which this gave rise to great annoyance, the small opening was enlarged by incision. The result was very satisfactory.

NEW GROWTHS. POLYPI.

Abnormal growths of various kinds are not infrequently met with in the nasal cavities. Many of them are commonly called *polypi*. But it would probably be better to use this term in a somewhat more restricted sense than is at present customary. Some of these growths are benign in character, others are malignant, others again are of doubtful malignancy. They are specifically named, according to their nature and situation, for the most part may be classified as follows:—First, Papillary Growths. Secondly, Mucous, Gelatinous, or Vesicular Growths. Thirdly, Fibrous Polypi growing from the lateral wall of the nasal cavities. Fourthly, Naso-pharyngeal Polypi, which are fibrous, but growing from the base of the skull. Fifthly, Gelatinous and Osseous Growths. Sixthly, Malignant Growths.

Papillary Growths.

Cases occasionally, though rarely, occur in which the nasal cavity is obstructed, or more or less completely blocked up by papillary growths springing from the mucous membrane. These growths bear a general resemblance to the papillomata of the mucous membranes of other parts, but so far as I have observed, the individual papillæ are acuminate. They are usually not invariably met with in children or young people. As a rule, they spring from those portions only of the nasal membrane which line the cartilages. They may be present in one nostril or in both. They are accompanied by some degree of irritation or running from the nose, and give rise to a constant degree of irritation, and disposition to pick the nose. Respiration is obstructed, and the voice affected to an extent proportionate to their development. Their presence may be readily recognised on inspection.

Such growths may be easily and effectually removed by the use of scissors, forceps, the wire snare, or the galvanic cautery.

and converted into clustering formations with very numerous sacculi, and these help to make up the mass. In some cases these sacculi become cystic in appearance, and the 'vesicular' character of the polypi is pronounced. In other cases the connective tissue elements are developed in larger proportion; and the growths are consequently firmer and denser, and approach in character the fibrous or sarcomatous polypi. Occasionally the softer polypi become comparatively condensed and opaque in places; and in some rare instances concretions of fibro-cartilaginous hardness are found in them.*

These polypi are most frequently connected with those portions of the mucous membrane which cover the superior and middle turbinated bones, and line the superior and middle meatuses. Sometimes they arise in the lower meatus, or are attached to the inferior turbinated bone; sometimes they spring from the roof of the nose; occasionally they grow from the ethmoidal cells, or even from the frontal sinuses into the nasal cavities. But they very rarely, if ever, spring from the mucous membrane covering the septum.† The relation which they may bear to the so-called adenoid vegetations growing from the nasopalatine mucous membrane or floor of the nose,‡ is by no means clearly made out; but it is by no means improbable that some such relationship exists. Both forms of growth appear to depend upon, or at any rate to be associated with distension of the mucous follicles. It is not improbable that the serous infiltration, and somewhat dropsical appearance of the pendulous mucous polypi, as well as the elongation and slenderness of their pedicles, may be due to some extent to the fact that they grow from above downwards. On the other hand the broad bases, and comparative freedom from infiltration of the sessile growths from the floor of the nose, accord with their situation and the direction in which they have to grow. Thus the effects of gravitation may help to explain in some degree the more obvious differences presented by these two forms of growth. Other differences may depend upon the special characters of the mucous follicles in different parts of the nasal cavities. It is worthy of note that the 'adenoid vegetations' and sessile growths of the floor of the nose, most commonly occur in stru-

* See case referred to by Gross, *op. cit.* vol. ii. p. 338.

† I have not been able to find a single satisfactory example on record.

‡ See p. 288.

ous children, while the mucous polypi of the upper cavities are most frequently met with in adults, although they may commence at a comparatively early age.*

In cases in which there are several distinct polypi, the lowest most anterior, having the open nostril before them, make their way downwards uncompressed; but the others are kept up and hidden from sight, and become more or less compressed and hindered in their growth. The polypi, if uninterfered with, go on gradually increasing in size until they more or less completely fill up the nasal fossæ. Sometimes they have an especial tendency forwards, sometimes backwards; sometimes they hang down behind the uvula into the pharynx; sometimes they project forwards and expand the alar cartilages, and even, in rare instances, the nasal processes of the superior maxillary bones.† It is very seldom, however, that any of the bones are seriously affected, but they may sometimes be deprived of their investing membrane, and necrosis may consequently ensue. In the case of a lady, thirty-eight years of age, who had suffered from mucous polypus for many years, Mr. Keating‡ found 'the internal osseous structure quite denuded.' The patient was eventually cured, after extirpation of the growths.

The earliest symptoms of mucous polypus in the nose are a sense of fulness and weight in and about the nostril affected; frequent sneezing; and more or less mucous discharge. For a long time before the growth becomes visible, the patient seems to be always 'catching cold.' As the growth increases, other symptoms supervene: respiration through the nose becomes more and more embarrassed; the patient suffers from 'stuffedness' and general discomfort; he cannot blow through the affected nostril, or the air passes with a whistling noise; he is obliged to keep his mouth constantly open; during sleep he lies with his head thrown back, and often snores; the voice is more or less affected, and becomes thick, indistinct, and 'nasal' in character; the sense of smell is impaired, or altogether abolished; the sense of taste is affected in so far as it is dependent upon a perfect condition of the olfactory organs; deafness sometimes occurs, either from obstruction to the Eustachian

* See Weber, in 'Pitha und Billroth,' op. cit.

† Weber, op. cit.

‡ Former Edition of this Work, vol. iii. p. 212.

tubes, or from coexistent thickening of the lining membrane of the tympanum; there is more or less constant flow of thin or thick mucous discharge, and the patient is unable to 'blow his nose' satisfactorily. The discharge is rarely offensive. Occasionally there is epistaxis, but not often. It is far otherwise in the case of fibrous polypi and malignant growths. Sometimes the lower orifice of the nasal duct becomes obstructed, and there is overflow of tears. These various symptoms become aggravated in damp, and somewhat alleviated in dry weather; for the polypi are 'hygrometric' in character, and are notably affected by the state of the atmosphere. In dry, warm weather they contract; and in moist or wet weather they enlarge, and become fuller, and paler in colour. Sudden atmospheric changes are liable to give rise to severe fits of sneezing.

The patient is often conscious of the presence of some fleshy substance, and makes frequent but ineffectual efforts to dislodge it, by blowing his nose, or introducing his fingers.

The symptoms thus detailed, indicate the probable existence of mucous polypi. Inspection, conducted in the manner already described,* generally reveals their presence, and digital exploration may aid the diagnosis. When a good light is thrown into the well-dilated nostril, the growth or growths may, as a rule, be readily recognised, and a probe may be passed round the lower parts and carried upwards to some height on either side. Sometimes the growths come more into view after the patient has attempted to blow forcibly through the affected nostril; or they may be pushed into more favourable position by introducing the finger round the free margin of the soft palate into the posterior nares. *Posterior rhinoscopy* is sometimes necessary, or at any rate desirable.

When the polypi can be thus seen and felt, the diagnosis is sufficiently easy. But in the earlier stages it is often difficult and sometimes impossible to get a view of the growths, or to reach them with the fingers. Under such circumstances the case, if it should come under the observation of the surgeon, may for some time be supposed to be one of chronic catarrh, and the discharge to be a simple coryza. In the more advanced stages no such difficulty is likely to exist, if only a careful examination is made. It is well, however, to bear in mind that hypertrophy of the mucous membrane of the inferior turbinated

* See RHINOSCOPY, p. 256 et seq.

is not always easy, however, to ascertain the exact attachment of the polypus, nor to make out at first whether the growth is single or multiple. But these are points of little practical importance. It may sometimes be difficult, but it is very important, to distinguish the benign mucous polypus from the growths of more serious nature. The light colour, semi-translucency, and soft elasticity of the mucous polypus, its slow growth, the absence of frequent hæmorrhages, and the general good health of the patient, serve to distinguish this growth from the fleshy polypus, which is more or less red in colour, opaque, firm, resistant, and which usually increases rapidly, undergoes great change of volume with change of weather, and is usually accompanied by frequent epistaxis. Malignant growths may be distinguished by their colour, opacity, consistence, and fixity, as well as by their proneness to bleed when touched, and by the offensive fœtidity of the discharge to which they give rise. Mucous and cartilaginous growths are distinguished by their softness, mobility, origin, and mode of growth.

Mucous polypi are always very troublesome, but rarely lead to serious results. Now and then they subside, and a spontaneous cure is effected. But it is never worth while to await a result; for the growths may almost always be removed without much pain, and with scarcely any risk. They are, however, very liable to grow again, especially if any portions remain unremoved, or the patient is unwilling to persevere in the necessary course of treatment.

results of such attempts have not hitherto been satisfactory, although doubtless, in some instances much good has been effected. Alum, calomel, oxysulphuret of antimony with sugar, cinchona bark, and many other substances in powder, have been used as snuffs, and different astringent solutions have been injected from time immemorial. 'At one time,' according to Professor Gross, 'a good deal of reliance was placed upon the use of finely pulverised blood-root as a snuff; and in 1859, Dr. J. H. Reeder of Illinois published the particulars of two cases in which prompt cures were obtained by strong injections of the tincture of perchloride of iron, aided by the application of a piece of sponge moistened with a solution of this article to the cavity of the nose.'* Mr. Erichsen refers to a case in which 'after the assiduous use of chloride of zinc injections, a very copious discharge of large sloughy polypoid masses took place from one nostril, which had been blocked up for many months previously.'† Nélaton recommends in some cases the repeated application of solid nitrate of silver.‡ Mr. Bryant has recently advocated the insufflation of tannic acid in powder.§ The use of bichromate of potash in a similar manner has been suggested, on the ground probably of the special influence which this salt is believed to exert on the nasal mucous membrane.|| Cases illustrative of its good effects have been published, but at present it has not been extensively tried.

Various methods have been devised for the removal of nasal polypi by operation. In judging which it is best to adopt, the peculiarities of the case under treatment must be taken into consideration. It is always desirable, however, that the surgeon should obtain as good a view as possible of the parts upon which he is operating. The patient must therefore be placed in proper position opposite a good light, or arrangements must

* Op. cit. vol. ii. p. 340.

† *Science and Art of Surgery*, vol. ii. p. 287.

‡ *Pathologie chirurgicale*, tome ii. p. 679.

§ *Lancet*, Feb. 1867, p. 235.

|| According to MM. Becourt and Chevallier, workmen employed in the manufacture of bichromate of potash suffer in a remarkable manner from the effects of the small particles of the salt which they are liable to inhale. The mucous membrane of the nose is seriously affected. Excessive irritation, coryza, and sternutation, are the earliest symptoms. The mucous membrane becomes thinned; detached portions of it come away; after six or eight days the bony septum itself becomes thin, and at last is gradually destroyed.—*Annales d'Hygiène*, July, 1863, p. 83; quoted in *Sydenham Society's Year Book*, 1863, p. 457.

sion of the growths by means of forceps. The forceps are about seven or eight inches in length, and strong, but as may be consistent with strength. The ordinary are too thick. The blades should be grooved longitudinally and well serrated along their edges. They may be straight or curved. Curved forceps are required when the growth is situated far back, and have to be removed through the nostril. In most cases in which it is intended to remove growths through the nostrils, straight or nearly straight forceps are best. The patient should be seated on a rather high chair, his head should be thrown well back, and supported by an assistant, who at the same time may advantageously elevate the tip of the nose, and hold the speculum by which the nostril is dilated. The elevation of the tip of the nose is especially requisite in cases in which it is requisite to introduce a long instrument into the lower meatus, for as already the external opening of the nostril is on a somewhat level than the floor of the nasal cavity. The closed blades of the forceps are introduced into the nose, and made to touch the floor or septum until their extremities have reached the growth, and passed to some extent the visible portion of the growth. The blades are then opened in a vertical or oblique position, and turned upwards and outwards, so as to include as much as possible of the growth. The growth is then held as tightly as possible, and is finally torn from its attachment, and removed from the nose by steady traction, the blades of the forceps on their long axis. It is better as

avulsion is accomplished ; or its pedicle, if within reach, may be divided by scissors or a bistoury.

When the polypus is situated far back and hangs down into the pharynx, it may sometimes be seized and removed by means of properly curved forceps introduced through the mouth. There is often, however, great difficulty in keeping the tongue down, and in resisting the spasmodic retching which is apt to be excited by the presence of the instrument. Moreover, it is not always easy to avoid including the mucous membrane of the uvula or soft palate between the blades of the forceps. The better plan, therefore, would seem to be, as suggested by Mr. Syme,* to detach the polypus by means of forceps introduced through the nostril, and then to push it on into the throat. In some cases, advantage may be derived from the use of forceps so constructed that the blades can be introduced separately, and afterwards locked together, after the manner of midwifery forceps.†

It is scarcely necessary to add that very great care must be exercised in applying the forceps and removing the growths. Otherwise, there is considerable risk of inflicting more or less serious damage. Many instances have occurred in which one or other of the turbinated bones, as well as large portions of mucous membrane and periosteum, have been ruthlessly torn away.

But although the method of extraction by forceps thus described is recommended by most surgical writers, and very generally adopted, I believe it is far inferior, in many respects, to the method of extraction by the 'wire snare' devised by Mr. Hilton. This method has been for many years past constantly and successfully practised at Guy's Hospital, to the almost entire exclusion of the use of the forceps. But it does not appear to be so widely known as it deserves to be. In our experience it has proved more easy and effectual, and less painful and less likely to prove mischievous, than other methods commonly adopted.

The instrument employed is figured on the next page (Fig. 230). The thumb of the surgeon is passed into the ring *a* ; his middle and forefinger hold the cross-piece *b*, and slide it backwards and forwards along the quadrangular stem *c*. The nasal portion *d*

* *Principles of Surgery*, p. 476.

† Schreger, *Neuer Chiron*, vol. i. part ii. p. 197. Quoted in South's '*Chelius*,' vol. ii. p. 732.

The instrument is cylindrical, slender, and tapering. It is connected with the quadrangular portion *c*, by means of a hinge joint *e*, so that it may be bent at such an angle as to enable the hand of the operator to be kept from obstructing the view. The slightly bulbous extremity *f* is doubly perforated in a longitudinal direction for the passage of the wire. The wire is connected on either side with the sliding cross-piece *b*, and forms loop *g* beyond the bulbous extremity *f*, after having traversed

FIG. 230.



Hilton's Nasal Polypus Snare.

the perforations. In its course the wire passes through two small holes in little wing-like processes close to the joint *e*, so that it may be kept in proper position at whatever angle the instrument may be bent. The wire I have found to answer best consists of two or three filaments of very fine annealed steel wire smoothly twisted together. Silvered copper wire is commonly used, but this bends too easily, and is apt to kink and break. Fishing gimp and salmon line gut are strong enough, and do not kink; but they are too limp to be pushed beyond the growth, if it is close to the floor or septum of the nose. Whatever material is used, it should be so proportioned in thickness as to run easily through the several holes of the instrument; and the edges of these should be so rounded as not to catch or cut the wire.

The manner of using this instrument is sufficiently obvious. The cross-piece is advanced as far as possible, and a good-sized loop of wire, much larger than that represented in the figure, is projected beyond the distal extremity of the instrument. This loop is introduced into the nostril with as much of the slender portion of the instrument as may be necessary, and the growth is encircled by moving the wire in the proper direction. The cross-piece is next drawn home and firmly held. The growth thus securely noosed, is then twisted and pulled from its attachment without difficulty, and with little or no risk of damage to the subjacent bone. In cases in which the attachment of the

growth is high up, the wire loop may be directed into position by means of a very small blunt fork mounted on a long slender stem; and in cases in which the growth is situated very back, the wire may be made to encircle it by aid of the finger passed through the mouth and round the posterior border of the soft palate. Portions of the mucous membrane adjoining the base of the pedicle of the polypus are sometimes brought away, but this is of little or no moment. It is obvious that with the wire snare it is impossible to inflict such damage to the tissue as has too often resulted from the incautious use of the forceps. In the case of any broad based swelling, such as that of hypertrophied mucous membrane or septal abscess, the wire snare rarely slips over the surface, and cannot, like the forceps, be used so that which it would be undesirable or detrimental to tear away. If the pedicle of the polypi should be very thick, it may be advisable to use the wire rope *écraseur*, with screw movement, instead of the snare just described.

In some instances polypi may be successfully removed by the fingers alone. One forefinger is thrust into the affected nostril, and the other is introduced into the posterior nares through the mouth. When the growth is reached, and held between the two fingers, it is pushed backwards and forwards until no resistance has ceased, and it is then withdrawn through the nearest opening. Morand and Sabatier succeeded thus;† Gross in a similar manner 'promptly removed a large fleshy polypoid polypus from a youth of seventeen.' In this case the tumour had a very narrow footstalk attached to the posterior extremity of the inferior spongy bone, which was therefore easily torn asunder.† But this method, though apparently simple, is rarely practicable, for reasons which are sufficiently obvious.

Dr. M'Ruer, an American physician, advocates the following method. A piece of catgut is to be passed through the nostril into the mouth, and to its extremity is to be 'fastened a small piece of soft, dry sponge, corresponding in size, when firmly pressed, to the narrowest part of the nasal passage.' The sponge is then to be drawn gently forward through the posterior nares, and through the nose. Dr. M'Ruer 'thus succeeded in at least ten cases in bringing away all the adventitious

* Former Edition of this Work, vol. iii. p. 215.

† Op. cit. vol. ii. p. 342.

hs.* This method has an appearance of clumsiness scarcely recommends it to general adoption.

In the case of large polypi, situated far back, it may sometimes be advantageous to make a button-hole-like opening through the soft palate—the ‘*boutonnière palatine*,’ of M. Amneuve.† The direction of the incision should be from backwards; and it should be made in the median line. The elasticity of the parts composing the soft palate permits a polypus of large size to be drawn through a comparatively small opening; and the same elasticity aids the subsequent closure of the wound. The polypus drawn through the opening may be cut or twisted off, or ligatured with comparative ease. If this has been done, the wound may be either left to itself, or closed by one or two sutures. So far back as the year 1747, M. de Mead, a surgeon of Avignon, slit the soft palate in a difficult manner, and at different times removed portions of the growth. He then introduced into the remaining portion several threads, the ends of which he drew the polypus forwards. His fingers passed through the mouth pushed it in the same direction. The polypus ruptured, and the tumour was withdrawn through the opening, with a noise ‘which resembled the uncorking of a bottle.’ A second polypus which subsequently showed itself was similarly treated, and the cure was complete.‡

Whichever of the methods thus described be adopted, the operation is liable to be attended by more or less hæmorrhage, and sometimes may be very troublesome, or at any rate may interfere with the proceedings of the surgeon. As a general rule, however, the bleeding soon ceases spontaneously, or may be arrested by the injection of cold water or astringent solution.

In some rare instances, however, it may be necessary to operate through the nares. In every case, if there is much bleeding, and especially if the patient has suffered severely, it is better to postpone the completion of the operation to a future period. And it often becomes advisable to operate on several successive occasions, and to remove the growths piecemeal, rather than to tax the endurance of the patient by attempting too much in a single sitting.

Thudichum has recently very strongly advocated the use

former Edition of this Work, vol. iii. p. 217.

Comptes-rendus, août 1859. Ranking's *Half-Yearly Abstract*, 1859, p. 196.

Cloquet, *Ophrésiologie*, p. 668.

of the galvano-caustic wire for the removal of nasal polypi, especially with a view to avoid the risk of hæmorrhage; and because this method, in his opinion, is attended by comparatively little pain. The apparatus which he uses consists of a modification of Middeldorpf's platinum wire carrier, connected with a battery of five large Grove's cells. The growth is encircled by the loop of wire, which is tightened by screwing up the windlass. Connection is then made, and the portion of the growth included in the noose is quickly burnt off, and withdrawn through the nostril.* The advantages of this method when practicable are manifest. Its disadvantages are that it involves the use of apparatus troublesome to prepare, not commonly in the armamentarium of the surgeon, and formidable in appearance to the patient; and that by it, the growths are removed by a tedious piecemeal process. In one case, for example, detailed by Dr. Thudichum, the polypi were removed from one nostril in no less than fifty-five different portions; and in another case thirty-three introductions of the loop were necessary. It is obvious that only the portion included by the heated loop is cut off; whereas by the use of the wire snare (p. 301), or the forceps, a great deal more than that which the wire includes or the forceps grasp, is at once removed; indeed, a comparatively small hold suffices for the extraction of a very large and extensively lobulated growth. Nevertheless, it cannot be disputed that in some cases Dr. Thudichum's plan is of the greatest practical value.

After any operation for the removal of nasal polypi, whether incomplete or apparently complete, it is well to wash out the nose regularly during some considerable period with astringent or detergent solutions, by means of the nasal douche or syringe. And in some cases the insufflation of tannic acid or other powders proves of great service, either in hindering the development of fresh growths, or in aiding the destruction of such portions as may have had their vitality impaired by the operation. It often happens that, after a partially satisfactory operation, large portions of growth may thus be got rid of and washed away; and the cure may be accomplished, or the patient may be enabled to go on in comfort for a considerable period.

The extirpation of mucous polypi by means of the ligature,

* Op. cit. p. 9.

Fibrous Polypi.

ous polypi are by no means so commonly met with in the cavities as the softer growths described in the preceding but they are far more formidable. If allowed to take course, they give rise to the most serious results. Their removal is often very difficult, and sometimes impracticable; moreover, the operative measures requisite may be attended with considerable risk.

These growths spring from the periosteum, and are connected with it by continuity of structure. They are covered by an extension of the mucous membrane, sometimes comparatively thin but more frequently thinned away. As a rule, they occur singly but in some rare instances two or more distinct growths have been found.† They vary very much in shape and size, according to the degree of development they may have attained, and in the direction in which they may have grown. Sometimes they are uniform in contour, and more or less globular or pear-shaped. But as a rule they are elongated and very irregular in shape, presenting finger-like prolongations, rounded outgrowths, or broad-based lobules in those parts which have encountered least resistance. They are firm and resistant to the pressure of the finger, and are sometimes of almost cartilaginous hardness. They have nothing of the yielding 'oyster-like' softness of the mucous polypi. They are opaque. While growing they are

method of applying the ligature by means of the double canula, is described by Mr. Wm. Ferguson in his treatise on *Operative Surgery*, 3rd edition.

usually more or less red in colour—in some cases comparatively pale, in others dark or even purplish red. When removed from the body, they are generally dull white, or yellowish. But the tint varies in different specimens, and in different parts of the same specimen. They are traversed, not only superficially, but also throughout their substance, by numerous thin-walled blood vessels, which are sometimes varicose. Hence their proneness to bleed. They cut somewhat crisply, and on section present a more or less uneven, and occasionally almost mammillated surface. They are composed, for the most part, of closely-woven strands of white fibrous tissue, which spread out from the pedicle. They contain very few elastic fibres. In different specimens a variable proportion of spindle-shaped nucleated cells, similar to those of sarcomatous growths, are to be found. Sometimes these are so numerous, and the general appearances are such as to indicate that the growth must be considered 'fibroid' rather than 'fibrous,' and that it approaches in character the malignant forms of new growth.

Instances are on record in which deposits of carbonate of lime have taken place on the surface of the growths,* as well as others in which the growths themselves have undergone a process of partial calcification.†

Fibrous polypi occasionally spring from the periosteum of the lateral walls of the nasal cavities or pterygo-palatine fossa, and somewhat more frequently, perhaps, from the floor of the nose.

In some instances they appear to have originated in the antrum, and to have extended from thence into the nose. In many cases, in which they have been supposed to have taken such a course, further investigation has shown their origin to have been very different.‡

There can be no doubt, however, that polypi of the kind under discussion most frequently take origin from the periphery of those portions of the base of the skull which form the roof of the naso-pharyngeal cavity. In such case they are 'naso-pharyngeal polypi' properly so called, and as such are described in the next section. But it is often very difficult, and sometimes in

* H. Cloquet, op. cit. p. 688.

† Bourdilliat, *Gazette médicale*, 1868; 'Fibrome calcifié.'

‡ This was notably the case in the very remarkable instance narrated by Prescott Hewett, which is quoted on page 462 of this volume, from *Medical Trans.* vol. xxxiv.

sible, to determine the precise point of origin of a fibrous is until the nasal cavities have been fully exposed to view eration.* The difficulty is increased by the tendency these growths have to acquire secondary attachments. secondary attachments probably result from adhesion place between opposed surfaces rendered bare by ulceration pressure. In the case of a little child, under the of M. Marjolin, it was found on post mortem examination though the growth evidently sprang from the basilar pro- body of the sphenoid, yet it spread along the septum d numerous adhesions to different parts of the pituitary ane, and especially to the posterior surface of the soft

tever their origin, these polypi increase in size with come rapidity. At first they mould themselves somewhat to ities in which they lie, and grow forwards, or backwards wwards, in the direction of least resistance. When more ed, they exert injurious pressure from within upon the of the nasal cavities, and in a certain sense overcome the ee they encounter. They are liable to cause ulceration struction of the mucous membrane, perforation of the and expansion and displacement of the nasal bones, erior maxillæ; or gradual thinning and absorption of ones may result from the pressure to which they are

The palate is pressed downwards and forwards; and ty of the mouth is encroached upon. In all such cases ge is proportionately distorted. The cheeks become more prominent, and the nose expanded and flattened. The pear, or actually are, more widely apart than natural. eral aspect thus produced is very characteristic. It has t inappropriately termed 'frog-face.' In some cases the on is symmetrical. In others, one side is more obviously d than the other. If one side only of the face is affected, er remaining perfectly natural, the presumption is that ase, whatever it may be, is in the antrum rather than in e, although the cavities of the latter may be encroached

e earlier stages the symptoms are similar to those pro-

case described by M. Gerdy, a polypus growing from one side, per- the septum, and appeared on the other side, and thus gave rise to the hat two polypi were present.—*Nélaton*, op. cit.

Revue des Hôpitaux, May 25, 1861.

duced by the mucous polypi, and it may be difficult to make the diagnosis. But the fibrous growths are firmer and more fixed than the mucous, and they do not undergo any change of volume under different atmospheric conditions. The symptoms therefore are more severe and more constant, and when the nose is once blocked, the obstruction to respiration is more absolute and unyielding. Moreover, the fibrous polypi give rise to frequently recurring epistaxis. This is not only a very characteristic symptom, but it may prove the source of much anxiety and danger. The discharge from the nose associated with the presence of a fibrous polypus is as a rule more scanty, and thinner, more sanious, and more liable to become foul and ozænic than the mucous discharge usually associated with the presence of the softer growths.

In the later stages, when the polypus can be distinctly seen and felt by the surgeon, and still further when from its size it has caused distortion of the face, there can be little or no difficulty in making out the nature of the growth, whatever doubt there may remain as to its precise origin and connections. Careful exploration, and pressure in different directions alternately by the fingers, and the passage of a bent probe round the free parts of the polypus, first on one side, then on the other, may often afford reliable indications of the probable position of the original pedicle, as well as of such secondary attachments as may have been formed. If any question should arise as to the possibly malignant nature of the growth under examination, conclusive evidence may be obtained by microscopical examination of a small portion removed by means of a grooved needle or fine trocar or canula.

Treatment.—The treatment requisite consists in the complete extirpation of the growth by operation. Medicinal treatment is of no service whatever, except in supporting the general health and strength under the depressing influence of the loss of blood and general suffering occasioned by the growth. Local applications are worse than useless; they may stimulate and irritate, but they can neither arrest nor destroy the growth.

The method of extirpation, that it is best to adopt, depends upon the size of the growth, and the situation of its pedicle. If the growth is not large, and its point of origin is neither very far back nor very high up, and if also, its pedicle can be reached and is not too broad, it may be removed through nostril or posterior nares, like the mucous polypus, by means of

the wire snare, the *écraseur*, or the galvano-caustic wire; or if it should be deemed desirable, a ligature may be applied, and the growth may be allowed to slough off; but I cannot forbear repeating that this method of treatment is open to serious objection, and is not free from danger. It may, however, be justifiable in certain exceptional cases. One fatal instance has come to my knowledge; and in that instance, even if the patient had survived, the operation would have been only partially successful, for it was a comparatively small portion of the growth only that had been included in the ligature.

In cases in which the growth is large, or in which its point of origin cannot be determined, or is situated very high up or far back, resort must be had to one or other of the operations described in the next section, on naso-pharyngeal polypi.

Naso-pharyngeal Polypi.

The growths so named are, as already intimated, fibrous polypi growing from those portions of the base of the skull which form the roof of the naso-pharyngeal cavity, or rather from the periosteum which covers them. The periosteum of the basilar process of the occipital bone, the body of the sphenoid bone, and the immediately adjoining parts, is peculiarly thick and vascular. Its blood supply is very free, and derived from several different sources. Hence, perhaps, its liability to give origin to new growths, and the rapidity with which such growths often increase in size.

The naso-pharyngeal polypi, properly so called, invariably spring from some part or other of the limited area thus indicated. They are, however, prone to acquire secondary attachments, and thus may arise various sources of difficulty in arriving at a satisfactory conclusion as to their point of origin. Sometimes they become adherent to the walls of the nose, and consequently may be supposed to be nasal polypi. Occasionally they extend down behind the pharynx, and in such case may appear to have arisen from the anterior surface of the spinal column.*

In structure these growths differ in no respect from the

* These several points are ably discussed in the Treatise of Dr. Robin Massé, *Des Polypes naso-pharyngiens*, Paris, 1864; and also in the elaborate essay of A. d'Ornellas, *Des Polypes fibreux de la base du Crâne, dits naso-pharyngiens*; Paris, 1854. And in the essay by M. Beuf on the same subject; Paris, 1857. Also by M. Brevet, *Des Polypes naso-pharyngiens*; Paris, 1855.

fibrous polypi of the nose already described (p. 305); and it must be borne in mind that polypi, whether fibrous or mucous, which arise from the nasal parietes, may extend into the pharynx as well as along the cavities of the nose, and thus may become, in a certain sense, naso-pharyngeal. But the growths now under discussion are distinguished by their origin, by the rapidity of their growth, by the remarkable manner in which they invade the various neighbouring cavities, and by the very serious results to which they may give rise.

During the earliest stages of the growth of a naso-pharyngeal polypus, the only obvious symptom is frequent epistaxis. The blood may pass from the nostrils, or may enter the pharynx and be ejected from the mouth. Frequent epistaxis from both nostrils must always be regarded as a suspicious symptom, and one indicating the necessity for careful examination. As the growth increases in size, respiration through the nose becomes gradually more and more impeded, and the patient suffers from a sense of 'stiffness.' At this period rhinoscopic examination at once reveals the existence of the growth; and on digital exploration it may be felt as a rounded, hard, resistant tumour. If left to pursue its course, it soon presses upon the soft palate, and thus interferes with easy deglutition, and gives rise to frequent inclination to vomit. Sometimes, when attempts are made to swallow, liquid portions return through the nose. As the malady advances, thin, sanious, or purulent, and sometimes foetid discharges begin to flow more or less abundantly from the nostrils, and the epistaxis becomes more frequent and more severe. The growth generally enters one side of the nose first, and then invades the other, either by penetrating the septum or by growing round its posterior border. Still later, the spongy bones and septum become more or less extensively destroyed, in consequence of the pressure of the growth. The nasal bones and superior maxillæ also become destroyed to a variable extent, or expanded and pushed to either side; and the countenance is consequently distorted. Sometimes the growth enters the maxillary sinus, and presses upon the floor of the orbit; sometimes it seems to pass round the bone (or through it), and issues under the skin of the cheek, passing along the cutaneous aspect of the bone, and pressing together the walls of the antrum.*

* See the remarkable case published by Mr. Prescott Hewett, already referred to; *Med.-Chir. Trans.* vol. xxxiv.

sometimes the growth sends off prolongations through the sphenopalatine and sphenomaxillary openings, and even extends under the zygomatic arch into the temporal fossa. MM. Maisonneuve and Chassaignac record cases in which growths of this kind had penetrated the pterygomaxillary fissure, and passed through the pterygoidean space between the muscles towards the face.* In other cases extensions are sent into the frontal and sphenoidal sinuses. And in others, again, prolongations may pass through the sphenomaxillary fissure into the orbit, with or without destruction of bone, and so cause protrusion of the eyeball.† As a further consequence, the optic nerve may be put on the stretch, and vision may be impaired. Very frequently the orifices of the Eustachian tubes are obstructed, and the sense of hearing is dulled. Very frequently also, the nasal ducts are compressed, and more or less constant epiphora results. But the most serious consequences follow, when, as occasionally happens, the growth penetrates to the interior of the skull. In such case no treatment is of any avail. The growth may enter the skull either through the sphenoidal fissure, or, after having caused absorption of bone, from the sphenoidal cells, or through the cribriform plate of the ethmoid bone. Suppuration, compression of the brain, coma, and death, are the only results to be looked for.

A remarkable illustration is recorded by Samuel Cooper, as having come under observation at St. Bartholomew's Hospital. The patient suffered from an enormous polypus, which occupied both nasal fossæ. The eyes were four inches apart, and the left eye was absolutely blind. Paralysis supervened fifteen days before death, which followed a period of coma. On post-mortem examination, it was found that a portion of the growth almost as large as an orange, was within the skull. The anterior lobe of the left hemisphere of the brain was almost entirely destroyed.

These growths are much more frequently met with in males than in females. They may occur at almost any age, but are most common in young people. In 122 out of 164 cases which have come under my observation, or of which I have collected particulars, the growth evidently commenced in each case before the patient had reached the age of twenty. M. Marjolin records the case of a little girl only two years of age who died from the effects of a growth of this kind, which had probably commenced

* Massé, *op. cit.* p. 17.

† See case under the observation of M. Chassaignac, quoted by Massé, *op. cit.*

during early infancy.* On the other hand, Dr. Robin Massé quotes from Richard a case in which the patient was fifty-five years old. But it is quite the exception to meet with an instance at so advanced an age. Indeed, in any case of supposed fibrous polypus of the nose, if the patient has long passed the middle period of life, the growth must be looked upon with suspicion, inasmuch as it will most probably prove to be malignant in character, and not a simple fibroma.

As already indicated, these growths not only give rise to considerable disfigurement and severe suffering, but sooner or later may prove fatal in result. Death may ensue from exhaustion caused by repeated hæmorrhage, constant discharge, disturbed sleep, and inability to take sufficient food; or from slow or even rapid asphyxia; or from implication of the nervous centres.

Treatment.—Complete extirpation of the growth by operation is the only treatment upon which any reliance can be placed.

In some rare instances, it is true, naso-pharyngeal polypi have spontaneously sloughed away. In the case of a young woman in St. George's Hospital, Mr. H. C. Johnson 'proposed to divide the soft palate, and endeavour to remove the whole growth from the pharynx; when, fortunately for the patient, the morbid tissue was attacked by rapid sloughing, which so entirely removed it, that no trace could be discovered of any part remaining.'†

A somewhat similar result occurred in the case of a boy under the care of Mr. Birkett, in Guy's Hospital. In this case it was proposed to extirpate the polypus after removal of the upper jaw; but the parents would not consent. On one occasion hæmorrhage took place to such an extent as to render necessary the application of a ligature to the common carotid artery. The tumour subsequently sloughed, and the whole came away through a large opening in the cheek. Seven years afterwards the patient was quite well. The sinus in the cheek had healed up. There was no recurrence of the growth.‡

Instances are also recorded by Saviard,§ Bonnet,|| Vimont,¶ and others. But so fortunate an issue is far too rare to justify the expectation of its occurrence.

Local applications, whether of exsiccatives, astringents, caustics, or escharotics, in solution, in powder, or in substance, are worse than useless. The actual cautery, in the form of the guarded hot iron thrust into the growth, has passed into deserved

* 'Proceedings of the Société de Chirurgie;' Paris. *Gazette des Hôpitaux*, May 25, 1861, already referred to.

† *British Medical Journal*, vol. i. 1858, p. 61.

‡ *Ibid.*, p. 119.

§ *Recueil d'Observations chirurgicales*, p. 112. Paris, 1784.

|| Bonnet, tome iv. obs. 92, p. 457.

¶ Quoted by Brevet, *Des Polypes naso-pharyngiens*, p. 16. Paris, 1855.

** See the treatises by Robin Massé and D'Ornellas, already referred to.

oblivion; and in like manner the use of the ligature has been well nigh abandoned, on account of the difficulty of its application, the dangers to which it may give rise, and the uncertainty of its results. In some cases, however, it is possible that the growth may be of such moderate size, and its pedicle so situated, that the ligature may be applied, and lead to results at any rate temporarily successful.

In all cases it is necessary to remove the growth completely; for if any portion should be left, there is very great probability of recurrence. Such being the case, and the anatomical relations of these growths, and the size they often attain before coming under treatment, being duly considered, it becomes obvious that it must be very seldom that they can be removed satisfactorily without some preliminary operation by which more space for surgical manipulation is obtained than naturally exists.

Sometimes, however, if the growth has not extended far into the cavities, and more especially if its pedicle is comparatively small, it may be removed through one or other of the natural openings. Thus, I saw my colleague, Mr. Forster, tear away a growth of this kind from the naso-pharynx of a boy in Guy's Hospital by his fingers alone. The result was eminently satisfactory. In another case I successfully removed a similar growth by means of a wire-écraseur, the end of which was so curved as to pass round the posterior border of the soft palate.

Mr. Bryant records a very remarkable case in which the growth 'had originally appeared in the right nostril, but had ultimately occluded both posterior nares, and pushed forward the soft palate. The écraseur (through the nose) was employed for its removal, but only a small and weak one could be introduced, and the screw could only be tightened gradually. In five days it came away, but the polypus did not fall off; for a time it appeared to shrivel away, but in nine months the boy was sent back to the hospital as bad as ever. The tumour was again carefully examined, and was found to be attached to the posterior wall of the pharynx. On this occasion whipcord was used instead of wire, and again the écraseur came away on the fifth day, but this time the tumour came with it.'* In this case reunion may have occurred between the parts gradually severed by the wire, or the polypus may have continued to derive its nourishment through some secondary attachment previously formed. Mr. Bryant informs me, he has reason to believe that no reproduction of the growth is again taking place.

In cases in which the pedicle of the polypus can be reached and encircled, it is probable that the galvanocaustic wire may be very advantageously employed.† But satisfactory experience

* *Medical Times and Gazette*, vol. ii. 1864, p. 126. Also *Path. Trans.* vol. xviii. p. 106.

† *Bevk, Wien. med. Halle*, 1862, p. 223. This method appears to have been first employed by Middeldorpf, in 1853. See Robin Massé, op. cit. p. 73.

of this method without preliminary operation, is as yet wanting. In every case in which the growth has attained considerable size, and is so firmly wedged into the cavities that its points of attachment cannot be easily reached, it is better, as a preliminary measure, to make a full and sufficient opening through the soft parts and bones. Various methods have been from time to time described, and practised with more or less satisfactory results. These methods may be divided into—1st, those which involve complete or partial ablation of the upper jaw; 2nd, those in which the upper jaw is temporarily displaced and restored to its position after the removal of the growth; 3rd, those in which the face is untouched, and the opening is made through the palate, with or without the absolute removal of the natural structures.

In deciding on the best operation to be adopted in any particular case, the surgeon must be guided by the size, extent, and possibly ascertained attachments of the growth with which he may have to do. In every case sufficient space must be obtained.

Ablation of the upper jaw (entire or partial) is neither so difficult nor so dangerous an operation, nor is it followed by so much after disfigurement as might be expected; and some modification or other of this method, is probably most likely to prove successful in a large proportion of cases in which the growth has attained considerable size, and has extensively invaded the various cavities.

Temporary displacement of the upper jaw involves some difficulties in the after treatment; but further experience may show how these are to be overcome, and the obvious advantages of the method may be more generally secured.

Operations through the palate alone do not always permit the full exposure of the origin of the growth. This method, therefore, is liable to prove inefficient. Its value, however, is amply attested by many successful cases.

Ablation of the upper jaw to facilitate the removal of a nasopharyngeal polypus appears to have been first suggested by Whateley, and first practised (though with incomplete success) by Syme, in the year 1832.* To continental surgeons, however, is due the credit of having fully established the practicability

* *Edinburgh Medical Journal*, vol. xxxviii. p. 322. See also Verneuil, *Proceedings of the Société de Chirurgie*, mars 14, 1860.

and the value of this operation.* It is now generally recognised as a legitimate and successful surgical procedure. Mr. Atum appears to have been the first in this country to have operated with successful results.†

The various operative proceedings for the total or partial resection of the upper jaw, will be found described in the essay on DISEASES OF THE MOUTH.

There are certain obvious objections to the total removal of the jaw. Even if the result should be successful so far as the extirpation of the polypus is concerned, more or less disfigurement remains: the removal of the floor of the orbit may endanger the eye; the teeth on one side are necessarily lost, and mastication may be impaired; and lastly, a large opening remains between the mouth and the nose, and articulation may be seriously affected. The latter several objections may be, for the most part, overcome by the adaptation of an artificial obturator at some subsequent period. The extent, however, to which the hasm left after the operation gets filled up in course of time by firm fibrous cicatricial tissue is very remarkable.

It fortunately happens, that in many cases partial resection of the maxilla suffices to afford the requisite space. The floor of the orbit,‡ and even the dental arch may be left intact.§

A good example of partial resection of the maxilla for the removal of a naso-pharyngeal polypus, is recorded by Mr. Pick as having come under the care of Mr. Holmes in St. George's Hospital.|| The patient (a man), twenty-seven years of age, suffered from characteristic symptoms. 'The operation was commenced by a single incision from the inner canthus of the eye, down the side of the nose, and continued through the upper lip in the mesial line. The flap having been dissected up, the bone was first divided by a key-hole saw through the alar tuberosity: the bone was then divided into two, leaving the floor of the orbit, by the bone forceps, carried along the lower margin of the orbit. The other attachments having been divided in the usual manner, and the bone having been severed from the soft palate, it was easily removed, and the origin of the tumour from the base of the skull exposed. The polypus was now grasped, and easily extracted. . . . The flap was re-adjusted. . . . There was very little bleeding during the operation. . . . The man made a good recovery, and left the hospital with the wound quite healed, five weeks after the operation.'

* A full history of the operation, and copious details of the various modifications adopted by different surgeons, are given in the already quoted treatise of Dr. Robin Massé.

† *British Medical Journal*, January 1858, p. 119.

‡ Guérin, *Éléments de Chirurgie opératoire*, p. 228. Paris, 1858. Maison-Neuve, *Gazette des Hôpitaux*, août 21, 1860.

§ Bérard, *Dictionnaire en 30 vol.*, tome xxviii. p. 367.

|| *St. George's Hospital Reports*, vol. ii. p. 162.

Temporary displacement of the maxilla followed by replacement immediately after the removal of the growth, may be successfully accomplished in the manner illustrated by the following case, the details of which were brought before the Académie de Médecine by M. Huguier, in 1861.*

The patient, a young man twenty years of age, had suffered during a period of about six years from the presence of a naso-pharyngeal polypus. Various ineffectual attempts at extirpation had been made from time to time. On admission the left eye was unduly prominent. The left side of the nose generally appeared enlarged, and the left ala was distended. On inspection through the nostril a reddish mass could be seen, and on digital exploration through the mouth, a firm rounded growth could be felt pressing down the soft palate, and filling the left naso-pharyngeal opening, and passing round to the right. The symptoms were severe. M. Huguier operated in the following manner. The mouth being widely opened, an incision was made through the soft palate. A narrow band or skein of thread was then carried by means of Belloc's sound along the left nasal fossa, passed through the opening made in the palate, and one end of it drawn through the mouth. This was intended to be used in making downward traction on the jaw in the later stages of the operation. Next, an incision was made through the whole thickness of the cheek, from the left labial commissure to the anterior border of the masseter. A second incision was carried alongside of the nose through the lateral attachment of the ala and through the middle of the upper lip. The flap having been raised, the superior maxilla was cut through horizontally by means of a saw. The line of section extended from immediately above the maxillary tuberosity to a point above the floor of the nasal fossa. The first incisor tooth on the right side having been removed, the saw was applied from before backwards to the hard palate on the left side of the septum. The base of the pterygoid apophysis was next cut through by means of strong bone forceps. Thus the inferior portion of the maxilla was detached from the other bones of the face, and only kept in connection with them by the mucous membrane of the hard palate and alveolar process. By means of a chisel used as a lever, aided by traction upon the band of thread introduced at the beginning of the operation, this portion of the bone was turned downwards and towards the right. In this manner a wide opening was made into the pharynx, and the polypus was seen adhering by a broad base to the basilar process, the back of the pharynx, and the base of the left pterygoid process. The pedicle was cut through, and the polypus removed in four portions by means of a gouge and strong curved scissors. Frightful hæmorrhage ensued, but the application of the hot iron proved effectual. The maxilla was replaced, but it was found difficult to retain it in position. The wounds in the face were closed by five sutures. A gutta serena mould was subsequently fitted to the parts, and worn for some time, and threads were passed round the teeth. In spite of the difficulty long experienced in maintaining the parts in proper apposition, the case terminated very satisfactorily.†

Other methods, similar in principle to the one described, but more or less different in detail, have been practised and recom-

* *Gazette des Hôpitaux*, 1861, p. 337.

† *Ibid.* p. 336.

mended by Langenbeck,* Roux,† and others.‡ In some cases it suffices to cut through the alæ of the nose along their junction with the cheek; through the nasal processes of the maxillæ and the skin and mucous membrane covering them, and through the septum. When this has been done the nose may be turned forcibly upwards, and replaced after the growths thus exposed have been removed.

The method devised and advocated by M. Nélaton,§ leaves the face untouched, and the superior maxilla undisturbed in position. The operation is performed as follows. The soft palate is first divided in the middle line throughout its whole extent and thickness. A longitudinal incision is carried along the posterior half of the hard palate down to the bone. From the anterior extremity of this incision two others are carried obliquely outwards, one on each side to the root of the alveolar process. The flaps of periosteum and mucous membrane are detached from the bone, and reflected outwards on either side. The bony palate is next perforated, and cut away by means of bone forceps. The mucous membrane and periosteum of the floor of the nose are then divided in the median line, and the flaps turned aside; as much of the vomer as may be necessary is removed. In this manner a large opening is made into the nasopharyngeal cavity, and the growth that may be there is exposed. After the growth has been removed, and its pedicle completely destroyed by the application of the hot iron or some escharotic, the flaps may be brought together. But M. Nélaton recommends that this should not be done until some time after the operation, nor until the complete destruction of every portion of the pedicle has been effected. At any subsequent period staphyloscopy may be practised. Deglutition and articulation are but little impaired, or not at all.

This method has been adopted and variously modified by many continental surgeons,|| but it does not appear to have found much favour in this country, on account of the comparatively limited access it affords to the base of the skull, and its probable inefficiency in case the prolongations of the growth

* *Deutsche Klinik*, 1861, p. 281; and Schmidt's *Jahrb.* cxiii. p. 195.

† *Gazette des Hôpitaux*, 1861, p. 354.

‡ See Robin Massé, *op. cit.*

§ *Bulletin de la Société de Chirurgie*, vol. i. p. 159.

|| M. Robin Massé, *op. cit.*, quotes or records twenty-six cases, in thirteen of which the results were perfectly satisfactory.

should be extensive.* But for these objections it would certainly seem preferable to the other methods described. The operation itself is not difficult; it is attended by comparatively little hæmorrhage; no obvious deformity results; mastication, deglutition, and articulation are but little impaired, or not at all.

The naso-pharyngeal polypus, by whatever preliminary operation it may have been exposed, must be very carefully removed; no portion of it must be left. Much bleeding is likely to take place if cutting instruments are used, but this can generally be arrested by the actual cautery, or by styptics. It is well in every case to cauterise the point of origin thoroughly by means of the hot iron, the gas jet of M. Nélaton, the galvanic cautery, or some powerful escharotic.

With the view of avoiding the various difficulties and dangers of such operations as have been thus described, M. Nélaton attempted, some years ago, the destruction of a naso-pharyngeal polypus by means of what may be called the *electro-chemical* method.† Two platinum needles, connected one with each pole of a 9-celled Bunsen's battery, were thrust into the polypus, and the current was allowed to pass for some minutes. The operation was repeated six times at intervals of ten days. The result is stated to have been eminently satisfactory. The patient left the hospital completely cured.‡ This method may be practised without any serious preliminary operation.§

Cartilaginous Growths.

Cartilaginous growths are occasionally, though rarely, met with in the nasal cavities. For the most part they are attached by broad bases to the septum and adjoining parts of the floor of the nose. They cause more or less obstruction to respiration, and alteration of voice. Sometimes they are attended by troublesome coryza and frequent sneezing; these symptoms have been very marked in two cases recently under my observation. Occasionally recurrent headache is complained of. In some cases the septum is pushed towards the opposite side, and thus the appearance on first inspection may be deceptive.

* See Mr. Pick's article in the *St. George's Hospital Reports*, vol. ii. p. 150.

† This method is worthy of note on account of the interest recently excited by Dr. Althaus's Treatise on *The Electrolytic Treatment of Tumours*.

‡ *Med. Times and Gazette*, March 16, 1867.

§ Robin Massé, *op. cit.* p. 78.

records a case, the particulars of which illustrate very well the story of growths of this kind, and the treatment to be adopted. A patient, eighteen years of age, had suffered ever since he could remember from a swelling in the right nostril. The patient had the peculiar 'frog-face,' to be characteristic of polypus. No air passed through the affected nostril. On examination a tumour was found connected with the septum and the side of the nose, and blocking up the inferior meatus on the right side. Mr. Erichsen made an incision through the side of the nose, in order to get room, and removed the growth, which was exceedingly compact in structure. There was some oozing of blood, and some erysipelatous swelling. The patient healed nicely, and the patient recovered rapidly from his distressing case.

A similar case came under my own care in Guy's Hospital four years ago. It was removed with equally satisfactory result, and when I saw the patient a few weeks ago, there was no appearance of any recurrence.

Osseous Growths.

Osseous growths springing from some part or other of the nasal bone, or from some neighbouring bone, may invade the interior of the nose, and give rise to symptoms more or less similar to those produced by other growths. The degree of extension and fixity presented on careful exploration, establishes the diagnosis. Very considerable disfigurement may sooner or later result.

In some cases cartilage in variable proportion, and variously shaped, enters into the formation of growths of this kind.† The most remarkable instances on record is that of John Shiel, an Irish labourer, twenty-four years of age, who came under the care of Mr. Morgan, in Guy's Hospital, in the year 1855.‡

The patient stated that he first perceived a small tumour in the right nostril when he was fifteen years of age. This gradually increased, and, 'at the time of his admission into Guy's Hospital, occupied the whole of the right side of the right nostril being enormously expanded and closed by the entrance of the tumour, which from its size, completely concealed the eye on the right side, and extended downwards into the mouth, being there connected with the alveolar and alveolar processes of the right superior maxillary bone. . . .

Other cases are recorded by Erichsen, *Lancet*, 1864, p. 152. Bryant, *ibid.*, 1867, p. 225.

A case recorded by MM. Trelat and Dolbran, *Bulletin de la Société de Médecine*, 1862, p. 261.

Guy's Hospital Reports, series i. vol. i. p. 403. *Sequel*, series i. vol. vii.

Illustrative specimens, Guy's Hospital Museum, No. 1666³², wax 1860 and 38¹², cast 65. This case is especially quoted because, in spite of its remarkable character, it does not appear to be generally known.

The bones apparently implicated in the disease were the ossa nasi, superior maxillary bone, vomer, and the inferior turbinated and malar bones. The description can convey an adequate idea of the size and shape of the tumour and the enormous general disfigurement produced. 'The tumour through its whole extent was almost uniformly firm, and of a bony hardness.' Morgan removed the growth (or a great part of it). 'A section of the tumour proved that it was composed of an outer hard thin shell of bone, completely enclosing a morbid mass of spongy cancellated structure, devoid of all appearance of sarcomatous or fungoid disease.'

The patient made a fair recovery, and lived for nearly seven years. He died, but the particulars of the termination of the case were not ascertained. It is, however, impossible to look at the cast † taken after death without believing that life could have been prolonged to such a period. The growth appears to have been simply enormous—larger indeed than the head itself.

True exostoses, whether cancellated, compact, or eburnate structure, and mixed bony and cartilaginous growths springing from the bones, must be distinguished from those called 'osseous tumours,' which are occasionally met with in the nasal cavities and adjoining sinuses, but in no other part of the bone. The history of tumours of the class now referred to, is at present very imperfectly made out. Instances are very rare, and opportunities afforded for investigation are necessarily limited.

The so-called '*Osseous Tumours of the Nasal Fossæ and Sinuses of the Face*,' appear, according to recent researches, to originate in the membranes, mucous or periosteal, which line the cavities, rather than to spring, like true exostoses, from the bony parietes. At any rate, they comparatively early separate from the normal bone, if even they have been in the first place connected with it. It has been suggested that these growths may be somewhat analogous in their mode of development to the subsequent separation of the antlers of the *Cervidæ*. In several instances that have come under observation, they have increased until they have attained very considerable size, and have produced very great disfigurement, accompanied by more or less severe suffering. In the majority of cases, they appear to have taken their origin in the frontal sinuses, or in the nasal

* Preparation 1666³², described as 'Cartilaginous Tumour,' which was removed from the face.

† Cast 65, wax model 38¹², Guy's Hospital Museum.

‡ Dolbeau, *Bulletin de l'Académie de Médecine*, tome xxxi. p. 107. Dr. Olivier, *Sur les Tumeurs osseuses des Fosses nasales*; Paris, 1869. *Comptes de Chirurgie*, tome iii. pp. 98 et 564.

bean, and as hard as a stone. Some time subsequently, the nose formed, and four months later the growth completely filled the left and a disagreeable discharge flowed from the nostril. Severe head-quent epistaxis soon supervened, and the discharge became purulent. circumstances the aid of M. Legouest was sought, eighteen months ppearance of severe symptoms. A hard immovable tumour with rface could be felt occupying and distending the left nostril. The was depressed, and behind and above it could be felt a larger tumour hich occupied the anterior part of the nose. The patient breathed mouth only, and swallowed with difficulty. The voice was nasal. it sanious and purulent discharge flowed from the nostril. After a iminary operation, terminating in partial resection of the upper jaw, f the growth was removed without difficulty, but *no point of bony ould be discovered*. The patient made a good recovery. In size, eneral aspect, the growth resembled somewhat the os calcis. Its uneven—smooth in some parts, rough in others. Its substance was very compact, and eburnated. It was impossible to make out any pedicle.

Similar cases are quoted by Dr. Olivier in his treatise on tumours. The general conclusions indicated are, that osseous tumours differ from ordinary exostoses, not only in their development, but also in the method of treatment to be adopted. Ordinary exostoses must be cut off from their base; but these 'osseous tumours' require the full and complete exposure of the cavities in which they lie, and they may be removed without any further difficulty. If allowed to go on increasing in size until by the pressure they cut off their own supply of nutriment. They then gradually separate and slough out, after having caused more extensive destruction of the parts upon which they may

Malignant Growths.

Osteo-sarcomatous, sarcomatous, encephaloid, and much more rarely scirrhus growths, sometimes invade the nasal cavities, or spring directly from their walls. Such growths, as a rule, commence in the bones themselves, or very speedily involve them. They generally occur either in young children, or those who have passed the middle period of life.

The age of the patient, the appearance of the growth, its proneness to bleed when touched, the foulness of the discharge to which it gives rise, and the general cachexia with which it is associated, indicate the character of the growth. If any doubt should remain, the microscopical examination of a small portion of the growth removed by avulsion, or by means of a grooved needle or fine trocar and canula, readily determines the diagnosis.

In four cases that have come under my observation, in each of which the symptoms generally resembled those associated with benign polypus, it was noticed that the soft palate, though stretched and depressed, was red and thick to the touch, instead of being pale and thin, as is commonly the case when it is pressed down by a mucous or fibrous polypus.

As a general rule, it may be stated that if the malignant character of the growth is indisputably established, it is better to let matters take their course; for in this situation it is well-nigh impossible to effect such complete extirpation as can be of any real benefit. In some cases, however, if the nature of the disease is recognised early, and its progress is not rapid, it may be not only justifiable, but even advisable to attempt the removal of the growth. But if this should be attempted, all the adjoining parts likely to be infected must be removed also. In this way, life may sometimes perhaps be prolonged, or at any rate suffering may be diminished.

ARTHUR E. DURHAM.

NOTE.—In preparing this Essay, free use has been made of that by Mr Ure in the former edition of this work. But so many additions have been made, and the sections of that treatise have been so far modified, that this Essay may be considered as an original one—for which the present author is solely responsible.

ited space allotted to this Essay compels me to
to a great extent, dwelling on those diseases
commonly fall under the special care of the
to confine myself, almost entirely, to those surgical
which are associated with, and dependent upon,
conditions of the teeth.

act of 'teething,' and the ailments of childhood con-
that process; the irregularities and deformities
on the shedding of the temporary teeth, and the
their permanent successors, would alone occupy, if
h justice to their importance, more space than can
to the entire Essay.

matter which is of the last importance in the treat-
teeth, *stopping* or *plugging* carious cavities and other
surface, is a subject of such extent, and involves so
nary discussion upon purely physical questions, that
ible to enter here upon its consideration. Indeed,
nt of those most usual and ordinary abnormalities of
rregularities of the position of the permanent set in
le, and carious cavities, entail to so great an extent
mechanical knowledge in combination with surgery,
ence and art, that it has naturally, and indeed inevit-
ed that special character of practice and of surgical

my limits allow, those surgical diseases and abnormalities, with their treatment, which are more or less connected with the teeth, under the following heads :

- I. Alveolar abscess.
- II. Painful and difficult eruption of the wisdom-teeth.
- III. Tumours of the gum.
 - (a) Epulis.
 - (b) Congenital hypertrophy of the gum and alveolar borders of the maxillæ.
 - (c) Polypus of the gum.
 - (d) Vascular tumours.
 - (e) Warty tumours of the gum.
- IV. Tumours of the tooth-pulp.
 - (a) Polypus of the tooth-pulp.
 - (b) Sensitive growth of pulp after fracture.
- V. Odontomes, or Tumours of the hard tissues of the teeth.
 - (a) Enamel nodules.
 - (b) Exostosis.
 - (c) Hypertrophied fangs.
 - (d) Dentine excrescence.
 - (e) Warty teeth.
- VI. ' Abscess ' of the antrum.
- VII. Dentigerous cysts.
- VIII. Alveolar and maxillary necrosis from
 - (a) Phosphorus fumes.
 - (b) Eruptive fevers.
- IX. Hæmorrhage after extraction.
- X. The application of obturators and artificial palates.

1. *Alveolar abscess*.—One of the commonest consequences of dental caries, and occurring occasionally independent of any apparent disease in the teeth, is this troublesome, and too often ill-understood, malady. The results of this condition, when appearing on the external surface of the face (cheek, jaw, or chin), are occasionally so remote from any obvious connection with the teeth, and the patient's symptoms so fail to indicate its association with them, that the true nature and cause of the suppuration is lost sight of, and an easily-remedied disease is indefinitely protracted.

Alveolar abscess may be defined as a suppuration around the fang or fangs of a tooth, usually carious, accompanied by

absorption and expansion of the bony walls of the alveolus or follicle, and the enlargement of the little pus-sac, the matter gradually finding its way to the surface either along a canal by the side of the fang of the tooth opening at the edge of the gum, through the gum itself at a point corresponding to the end of the root (or roots) of the tooth implicated. When, however, the fangs are unusually long, or the reflection of the mucous membrane, from the gum to the cheek or lip, is very superficial, this discharge may burrow still more outwardly, and find its way upon the surface of the face.

The pathology of alveolar abscess, especially in its commonest form of 'gum-boil,' has not been generally understood. The first stage which occurs in the development of this condition consists in the deposit around the extremity of the fang or fangs of the tooth of organisable lymph, which is soon differentiated into obscure fibrous tissue. Coincident with this development, the bone around the end of the tooth-fang becomes absorbed, and a little hollow in the jaw is produced, varying in size from millet-seed to that of a horse-bean. Thus far the change is simply one of plastic inflammation: it occurs, as far as my observations go, with every carious tooth, and in those with more than one fang is most conspicuous at the extremity of that which corresponds with the side or part of the crown where caries occurs.

These changes are often accompanied by absorption, to a varying amount, of the extremity of the fang; and this appears to be part of the same action as produces the excavation in the follicle itself, which is occupied by the plastic exudation. There is generally, too, a thickening of the alveolar periosteum, which loosens, and often loosens, the affected tooth.

It is the lymph surrounding the extremity of the fang, and occupying the little cavity produced by the bone-absorption, that is the seat of suppuration in alveolar abscess. Whatever may be the first actual point of suppuration, when sufficient pus is formed for recognition, it is found in immediate contact with the fang of the tooth—the bare naked crista petrosa, and surrounded by the half-organised lymph, as by a sac. The form of the sac varies considerably; it is usually spherical or pyriform where the fang is single, or if the tooth have more than one fang, and the fangs are distinct; but where the fangs are close together, or the root is simply cleft at its extremity, as is often the case with the superior premolars, the sac is frequently

double and bilocular. The sac is sometimes of large size, and is frequently very long; sometimes it is fringed with fimbriated processes of lymph, which occupy an elongated excavation in the maxilla. When suppuration is established, the sac enlarges according to the amount of matter formed, and the rapidity with which it is developed; and this is accompanied with dilatation and absorption of the bony walls of the abscess, especially towards the region at which the matter is 'pointing.' These latter changes are often very extensive and very rapid, so that the whole of the osseous tissue of one wall of the alveolus may be expanded and thinned into a mere papery film, and then completely absorbed in comparatively few hours, the periosteum, with the organised lymph that has been the seat of the suppuration, alone remaining as the membranous sac of the abscess. The development of an alveolar abscess is generally accompanied by a deep throbbing pain at the part affected, and often by great general swelling of the face, sympathetic of the more internal mischief. The distortion of the countenance from this cause is sometimes prodigious, especially when the upper incisors or canines are the cause of the abscess, and the attack is acute. In such a case the nose will be pushed on one side, the eyelids become œdematous, closed, and ecchymosed. In other instances, where the character of the affection is less intense, a mere indurated, indolent, and hard swelling on the side of the jaw is all that is produced; the maxilla being distended at the extremity of the tooth's fang, and but little bone absorbed.

The development of an alveolar abscess is sometimes associated with febrile symptoms of extreme severity. M. Robert* describes a case of alveolar abscess, resulting from a carious wisdom tooth, which led to necrosis of the surrounding bone and diffuse purulent infiltration of the side of the neck. The case terminated fatally.

Alveolar abscess is usually confined to the tooth or one fang (when the tooth has more than one) which has been the cause of the affection, and around which the plastic exudation has formed. Occasionally, in two- or three-fanged teeth, the intervening bone between the extremities of the individual fang becomes absorbed, and one abscess may be common to all

* *Conférences de Clinique chirurgicale*, par M. A. C. Robert, p. 145. Paris, 1860.

tion that this form of alveolar abscess is dependent on constitutional taint—scrofulous or syphilitic.

'pointing' of the matter in alveolar abscess is a question of moment, both as regards diagnosis and treatment; and the circumstance that the discharge of pus may occur upon the side of the face, entailing with it very great disfigurement, is of great importance to this consideration.

An alveolar abscess affecting the one-fanged teeth may find its exit by a gutter or channel along the course of the root of the tooth, the matter being discharged at the neck of the tooth. This occurs to the many-fanged teeth, and is most commonly in the inferior incisors—the pus flowing from the front of the gum at a point corresponding with the particular tooth affected. The commonest position at which the thinning of the alveolar bone takes place is on the outer surface of the tooth, at a point corresponding, as nearly horizontally as may be, with the extremity of the fang of the affected tooth, and *perforating the gum within the mouth*. In such common cases the diagnosis is usually simple and obvious, and the abscess is easily associated with the individual tooth that produces it. There are, however, instances in which the pointing of the abscess, after having taken a lengthened and burrowing course, is remote from its source. The most striking peculiarity of this kind is where a circumscribed collection of matter appears far back in the palate, usually at the very posterior extremity of the hard palate, without any affection of one of the six front upper teeth. This condition may readily fail to indicate its true explanation, and the source of the abscess may be sought in the

logical anatomy of the affection is the same as in ordinary cases, only the canal of the abscess is lengthened out in the narrow cancellated bone between the two compact plates of the palatal process of the superior maxilla. A critical scrutiny of the front teeth (incisors and canine) will scarcely fail to show which is the offender.

The forms of alveolar abscess which we have just considered are trivial in their importance in comparison with those which, in their advancing course, involve the integument of the face. It is a curious circumstance that alveolar abscesses, when pointing externally upon the face, have been so frequently misunderstood by surgeons, having been mistaken either for idiopathic abscess in the substance of the cheek, or suppuration associated with necrosis of more or less of the maxillary bone. To the superficial observer (and, indeed, in its consequences there is a vast difference between a common gum-boil and an abscess which pierces and discharges its contents upon the surface of the cheek; but the cases are the same, barring the point where the pus is evacuated.

The circumstances which determine the pointing of an abscess upon the surface of the face appear to be either an unusual length of the fang of the tooth, or a superficial reflection of the mucous membrane from the jaw to the cheek; so that, in either case, the abscess which is forming around the extremity of the fang does not correspond horizontally with the gum within the mouth, and thus in its course outwards it passes either above or below (as it may occur in the upper or lower jaw) the line where the mucous membrane folds from one surface to another. In some cases, however, it takes an outward course irrespective of these conditions.

I am not aware that alveolar abscess, associated with the superior incisors or canines, ever points upon the surface of the face; the bicuspid, first and second molar, and all the teeth of the lower jaw, may produce this form of the disease. In the upper jaw the abscess appears upon the cheek at a point corresponding with the extremity of the tooth's fang, under the edge of the malar bone; in the lower jaw it forms along the edge of the jaw below the buccinator muscle when the molars or bicuspid are its cause; but when associated with the inferior incisors (canines also?), it points frequently beneath, and sometimes in front of, the chin.

When an alveolar abscess is about to point externally, the

egument becomes firmly glued down to the bone around the point where the matter ultimately appears; the area within this space is red, distended, and glistening; the skin becomes thin and papery, and the epidermis scales off. If the surface be kept open, the breaking of the abscess is often tardy and delayed; but it ultimately bursts through a jagged opening, which soon changes into a small fistulous orifice, surrounded by a pouting circular lip of granulations, that sink into a depression, surrounded by the adhesions which limit the pus-discharging canal. These appearances now are very characteristic, and when once recognised cannot be afterwards mistaken. Sometimes the lip of granulations becomes elongated into a papilla, and is covered by a cuticle. I have seen one more than half an inch in length. The apex of this papilla has an orifice, which is the outlet of the fistulous canal communicating with the abscess; it frequently closes for a time, but bursts again as the matter accumulates. When once this papilla forms, it returns again and again, after division by the knife, till the cause of the whole malady (the diseased tooth) is removed. Another curious modification of the external orifice of the alveolar abscess is occasionally seen when it pierces the under surface of the chin; in such cases a *pad* of granulations sometimes forms, as large as the area of one's thumb-nail, from the centre of which the discharge flows. In children, with the milk-teeth, alveolar abscess very seldom opens on the surface of the face. I have, however, seen a few such cases. On account of the extreme thinness of the alveoli of the milk-teeth, the end of the affected fang frequently appears through the orifice of the discharge within the mouth, and often projects far as to wound and ulcerate the mucous membrane of the cheek or lip.

It sometimes happens, that after the first evacuation of the contents of an alveolar abscess the secretion becomes serous. I have known some instances in which the sac of the abscess has remained as a serous cyst even after the extraction of the tooth in which it originally depended, the secretion recurring again and again after the cyst had been lanced. In one instance, where it had happened in a child in connection with an upper incisor tooth, I found it necessary to remove a portion of the wall of the cyst, when it granulated from the interior, and was obliterated.

The *diagnosis* of alveolar abscess is really very simple, though, as I have said, it is often mistaken for diseased bone in those

examples where the orifice occurs on the surface of the face. The tooth is the equivalent in these cases of a sequestrum; and it is, as far as the pulp-vitality goes, a dead organ. The fistulous canal leads to the dead tooth, as a sinus to the dead bone; thus far the two conditions are alike, and their appearances are alike; but there are obvious differences. In alveolar abscess there is less general swelling, except at first in acute cases, and there is an absence of the diffuse indurated lymph-infiltration of the cellular tissue which is present in bone-necrosis; the canal leading to the abscess is single, usually short and direct, and not burrowing and complicated as in bone-disease;* moreover, there is generally an absence of foetor in the discharge. The locality in a doubtful case, being the neighbourhood of the jaw, is in favour of the idea of a tooth being the cause; and an appeal to the state of the teeth seldom fails to settle the question definitively. Sometimes, however, a difficulty may arise from the fact that the cause of the abscess is a mere stump, overlapped by prominent gum; or, what occasions still more obscurity, the abscess may consist of a large excavated cavity in the substance of the jaw, into which the stump has fallen and become loose and free. A probe and the elevator will in either case complete the investigation.

While alveolar abscess, when connected with the integument of the face, is very apt to be mistaken by surgeons for diseased bone, the reverse has happened where a dentist has attributed to carious teeth morbid conditions which have been coincident with them, though not produced by them. I have known this occur in an instance where the patient was suffering from scrofulous caries of the malar bone near its suture with the superior maxilla—a mistake which led to the extraction of two slightly damaged teeth in no way connected with the disease. Again, I have seen one of the sub-maxillary lymphatic glands, rather adherent to the bone and in an early state of

* I once saw a very remarkable exception to the above, nearly universal rule. A lady consulted me on account of a burrowing abscess, which first burst immediately below the lower jaw on the right side. This orifice closed and another formed lower down: this also closed, and was followed by another in the neck. The burrowing continued, and when I saw the patient, many months after the first discharge of pus, there were two sinuses opening about an inch *below* the right clavicle. A carious first lower molar had caused the suppuration, and its removal was followed by the immediate healing of the sinuses.

operation, mistaken for an outward-pointing alveolar abscess. Such errors as these, leading at most to the extraction of useless, though innocent, teeth, are trivial in comparison with these more frequent mistakes in which alveolar abscess is confounded with bone-disease, and an easily cured malady is allowed to run its course unrestrained, and permanently disfigure the face.

The *cause* of alveolar abscess is either caries of a tooth or death of the organ; the latter usually the result of mechanical violence, which has separated the pulp from its continuity with the subjacent nervous and vascular structures. It is uncommon to see the affection arise in association with sound and apparently uninjured teeth, though this sometimes occurs.

The *treatment* of alveolar abscess depends upon the stage at which the case is seen. In the earliest period, when suppuration is rather impending than established, the malady may be cut short by the extraction of the affected tooth, or by the removal of the stopping in a stopped tooth. Often it is undesirable to extract a particular tooth that may be threatening causing the abscess; and in that case recourse should be had to leeching the gum freely, the administration of brisk cathartics, and abundant hot fomentation of the face at the swelling part. This mode of treatment, when it does not arrest the malady, much mitigates the attendant suffering. When matter has formed, and the wall of the abscess has sufficiently thinned, the pus should be evacuated by a puncture made through the gum with a narrow thin scalpel. This is followed by immediate and complete relief, and the general swelling associated with the advent of the attack rapidly subsides; but the disease, in the very great majority of cases, remains in the form of a continuously pus-discharging fistula. It is very rare indeed for the pus-secretion to cease: it may be so diminished that the external orifice may close for a time, but it is pretty sure to burst out again and again; though I am content (contrary to the general published opinion) that in a few exceptional instances the disease ceases altogether, the offending tooth still remaining in the mouth.

When an alveolar abscess shows symptoms indicative of external pointing, immediate and active treatment is necessary. The offending tooth should be taken out; and in case it breaks on extracting, every vestige should be sought and removed. If matter has formed, and there has been decided thinning of

the integument, showing its near approach to the surface, not only should the tooth be extracted, but a vertical incision should be made between the cheek and the jaw, so as to cut across the pus-containing canal; else the matter is apt to *pocket*, and cause progressive absorption to the surface; and though the tooth be removed, the abscess may thus still open externally. This vertical incision between the cheek and the jaw is also useful after the removal of a tooth, even when the external opening has occurred; it cuts across the bands of lymph which glue the integument down to the bone: in this case a piece of oiled lint should be introduced and kept in the wound, to prevent the adhesion and reunion of the cut surfaces. In all cases of alveolar abscess, extraction of the diseased or dead tooth is *the* cure; and I know but of two circumstances which peremptorily interdict this mode of treatment. First, where a strongly-pronounced hæmorrhagic diathesis forbids the extraction of teeth altogether; and secondly, in those cases where the abscess is associated with the upper incisor teeth of young people in whom the jaws have not yet assumed their adult form, and where the permanent dentition is, as yet, incomplete. In this latter case it is of much importance to retain the teeth, even if reduced by decay to mere fangs, till the adult form of the jaws is established, even at great cost of suffering and discomfort. The earlier removal of the teeth would be followed by such contraction of the maxillary arch as would be incompatible with a proper replacement by artificial teeth of the natural ones missing.

II. *Painful and difficult eruption of wisdom-teeth.*—The advent of the wisdom-teeth is very often accompanied by painful and distressing symptoms, that may be protracted through many months, or even years, unless relieved by surgical interference. These circumstances arise from the position occupied by the wisdom-teeth, so close to the joint of the lower jaw, where the mucous membrane is reflected from the gum to the cheek and fauces, combined with the very common condition—that the jaw is not sufficiently elongated backwards to allow the dentes sapientiæ to range in the horizontal series with the other teeth. This mechanical difficulty not only prevents the proper evolution of the wisdom-teeth, holding them back in their bony bed, but it often perverts their direction of growth and dislocates them. Annoying and very painful as are often the symptoms attendant

difficult cutting and misplacement of the upper wisdom-teeth, they are trivial in comparison with those which occur in similar conditions of the lower.

The ordinary misplacement of the upper wise teeth is either backwards or outwards, or in both directions combined. When the tooth points backwards, every time the mouth is closed its own comes in contact with the mucous membrane, passing up to the base of the coronoid process; when the direction is outwards, which is more common, the tooth projects into the cheek, and when the jaws are brought together, a portion of the mucous membrane in this region is nipped and pinched. This is a very painful affair: the surface becomes ulcerated and extremely tender; there is a partial cicatrisation, and the structure becomes stiff and hard as well as painful. Beyond this, however, the symptoms never (as I believe) extend, and the removal of the offending tooth is always followed by complete and immediate relief.

The difficulty which most commonly occurs with the inferior *sapientiae* is attributable to insufficient room in the jaw: the tooth grows normally in direction and in position as regards its neighbour in front, but, from an imperfect lengthening of the horizontal ramus of the jaw, the birth of the crown is only partial and incomplete: the tooth is upright, but only its front cusps emerge, while the hinder cusps are still covered with gum, or even the upper wall of the bony loculus in which it was formed. This produces a terrible pinching of the mucous membrane over the tooth every time the jaws are brought together. Before, however, the enamel eminences of the tooth's crown make their appearance, the soft structures behind the second molar become much inflamed, and often suppurate, the pus-secretion appearing to be within the enamel of the tooth, between the tooth-crown and the membrane covering it. The inflammation, to which this impaction and pressure give rise, extends to surrounding tissues: the cheek and the fauces suffer; the movements of the jaw become stiff and painful; and deglutition is difficult, and attended with suffering like 'sore-throat.'

Next in frequency to the foregoing malposition is that in which the wisdom-tooth is developed horizontally forwards, more or less. It is attended often with serious consequences. This direction is sometimes combined with an inward leaning; and very rarely the crown of the tooth points outwards. This

latter dislocation is very unusual; but the most severe suffering from misplacement of the lower wisdom-tooth have ever seen was of this variety.

There is one peculiar symptom frequently associated with painful cutting of a lower wisdom-tooth, which adds to the difficulty of investigating the condition of the parts, and still interferes with the treatment: this symptom is *spasmodic traction of the masseter muscle*, of a continuous and peculiar character, the result of contiguous irritation; not a spasm which varies in intensity, but a true *tonic* spasm, the jaw being permanently *set*, so as to keep the jaws nearly closed, and susceptible only of very slight separation. The jaw can be opened to a small extent, and then is definitely fixed; as if there were a mechanical obstacle to further movement, but it is not so; for when the cause of irritation is removed, the spasm rapidly ceases, and then the mouth can be fully opened. This same form of spasm sometimes occurs from caries of the molars, irrespective of crowding. The pain that accompanies and precedes the cutting of a wisdom-tooth varies very much in different individuals; but it is frequently of a dull, aching character, like rheumatism, for which it is occasionally mistaken; it is diffuse and erratic, extending up the side of the head, and down to the shoulder. There is not unfrequently a good deal of swelling of the soft parts in the neighbourhood of the tooth, and this sometimes extends to the cheek and eyelids even, and below the angle of the jaw: the lymphatic glands beneath the jaw also occasionally become enlarged and tender.

The most distressing result, however, that occurs in severe cases is the suppuration which sometimes attends the eruption of the tooth. In slight cases, even where there is no misplacement or want of room, but simply a restrained pressure from an unusually dense or cartilaginous overlying gum, abscesses may be occasionally formed; being secreted (as it has seemed to me) within the capsule of the tooth, by that which was the 'pulp.' In severer cases, the pus burrows among the soft tissue, around the periosteum of the jaw; the neighbouring soft structures become infiltrated with lymph, and the tooth is glued irregularly to the bone; pus points at different spots, often remote from its original source; and the cheek may be undermined with a series of sinuses. I have seen this on the side of the face, from the tragus of the ear and the a



the jaw behind, to the angle of the mouth and mental foramen in front, a web of pus-discharging fistulæ, and which, after a fair cure (by the mere removal of an impacted wisdom-tooth), left the integument thin, and bound down to the bone, with the glossy, tense, cicatrix-like aspect of a recently-healed burn—a great and permanent disfigurement. While such cases are infrequency, they have very much the appearance of necrosed bone. And, indeed, they do occasionally lead to necrosis of more or less of the jaw; but the severest suffering and most suspicious symptoms may occur without such a complication. What is the precise anatomical condition of these burrowing suppurations I am not prepared to say positively. They are not apparently connected with the fang of the wisdom-tooth, like an alveolar abscess. I believe they commence in the enamel, before any portion of the crown pierces the gum; and their subsequent course of burrowing is from the matter *pocketing* in a downward and onward progress. As long as any part of the crown of the tooth is covered by gum, pus will be secreted by the under surface of the overlying portion. I have seen an impacted wisdom-tooth give rise to that form of serous cyst known as a 'dentigerous cyst.' In those instances where the crown of the wisdom-tooth projects forwards, the second molar very often suffers. The posterior fang is apt to be eroded by absorption, and the whole tooth may become necrosed; the worst of these conditions is very common: I have met with several instances of it; in four specimens of lower second molars now before me, which were removed on account of forward pressure of wisdom-teeth, the posterior fangs are excavated by absorption in all; in two the pulp-cavity is closely approached; in two it is opened, and in one of the latter there was total necrosis of the whole tooth; the periosteum was completely stripped from both fangs, which were unattached to their sockets, and bathed in pus. These affections of the second molar should be taken into consideration in balancing the merits of the plans of treatment which may be contemplated in any particular case, and will, I think, favour that course of proceeding which I believe ought in many instances to be followed, but which is not now usually adopted.

Treatment.—The wisdom-tooth of the upper jaw, when misplaced, suggests but one mode of treatment; it by far the most often projects outwards; it is then useless for mastication; and if it pinches the cheek painfully on closing the mouth, it

may be removed unhesitatingly and without compunction. The same may be said when it grows backwards and bruises the mucous membrane over the base of the coronoid process; the tooth is useless, and worse than useless, and should be extracted; it is very readily done, and, as far as I know, the annoyances attendant on the misplacement of the upper wisdom-tooth are never attended with that locked-jaw complication which so frequently attends the difficulties of the lower third molar eruption. In one rare instance, in which the upper wisdom-tooth grew forwards, it was found necessary to extract the second molar, on account of damage which it had occasioned by producing absorption of the neck of the anterior tooth.

In the treatment of these cases in the lower jaw, much will depend on the degree of impaction of the tooth; if it is simply covered over by a dense and cartilaginous gum, free lancing is all that may be required; but it will often have to be repeated. In young precocious people, who arrive very early at puberty, the wisdom-teeth often make their appearance before the jaw is ready for their reception, and are embedded in the base of the coronoid process, though in a proper upright position. The age of the patient in such a condition is an important question: if young, lancing the gum freely, and waiting for further elongation of the jaw, and more horizontal accommodation, may be fairly and reasonably adopted; but the same state of things in an older patient is not so likely to find this relief; and if the suffering of the patient is great, extraction of the impacted tooth will become necessary. Where the tooth emerges tardily, and where the anterior cusps have come through the gum while the posterior have remained covered, I have found great advantage not only in lancing, but in cutting away the gum which overlies the back of the tooth, so as to lay bare the whole of the masticating surface of the crown. Merely lancing the gum affords much less relief, and the cut surfaces are apt to unite, and then present a harder and more resisting impediment. If the impaction is severe, and there is no chance, from the age of the patient, of sufficient accommodation, it will be necessary to extract a tooth, so as to relieve the crushing tension; and the question now arises as to which of the two, the second or third molar, should be extracted. *Ceteris paribus*, it is always better to sacrifice the third than the second molar: the wisdom-tooth is less useful for mastication, and is more liable to decay; it is altogether less robust: but the circumstances of

in case I should extract the second molar, though per-
sond, without hesitation. The distressing symptoms
are peculiar to the difficult eruption of the *lower* wisdom-
the heavy pain, the sense of *tension* and *pressure*—are
dependent on the antagonism of the third and second
; the former pushing forward into place, and the latter
g its progress. The removal of the second molar puts
to this antagonism as completely as that of the wisdom-
and, in result, the one operation gives as entire and
ste relief as the other. When the second molar is
d, the wisdom-tooth comes forward wonderfully; and
ne or nearly horizontal tooth will advance in position,
r in attitude, so as to become nearly, or quite, vertical,
fair opponent to the upper second molar. In this
d position, when it can be used habitually in mastica-
e dens sapientiæ is less liable to caries. But when sup-
i has become established, every effort should be made to
the wisdom-tooth itself, as the extraction of the second
nder these circumstances, generally fails to arrest the
ge of pus.

losure of the jaws is often so complete that, until this
m has been combated, it is impossible to get at either
molar teeth. By pressing apart the jaws with some
wedge, this may be accomplished gradually, and in two
e days sufficiently to allow the removal of the tooth
l for extraction. Steel instruments have been devised on
nciple of an ear speculum, composed of two shafts or

however, invented by Mr. Maunder of the London Hospital seems to possess advantages over those previously in use. It consists of a cone of box-wood or ivory, upon which is cut a spiral screw-worm, gradually increasing in diameter and thickness of the worm: it resembles the shell of a univalve mollusc, such as a whelk-shell, only the spire is more elongated and drawn out. In cases where closure of the mouth is brought about by spasm induced through the painful eruption of a wisdom-tooth, occurring as it does in early life, the teeth in front of the mouth are almost always firm, and would resist the wedging force. Upon introducing the point of Mr. Maunder's 'gag' between the incisors, or canines, or premolars, and slowly turning the instrument, it evenly and in regular progress, separates the jaws, and, as I have thought, with quicker results than any other method.

In extracting an impacted wisdom-tooth one accident occasionally arises; namely, the crushing of the inferior maxillary nerve beneath the tooth-fang. It is immediately attended by loss of feeling in the teeth and lower lip of that side. This has occurred twice to myself. In each case sensation slowly returned and was quite re-established in a few weeks.

III. *Tumours of the gum.*—(a) Epulis: (b) Congenital hypertrophy of the gum and alveolar borders of the maxilla: (c) Polypus of the gum: (d) Vascular tumours: (e) Wart-like tumours of the gum.

Epulis.—"The term "epulis" has been vaguely applied to various tumours that are found in and beneath the gums. The etymological meaning of the word (*ἐπί, upon, οὔλα, the gum*), entirely referring to position, and not to structure, is likely to have caused and to continue this confusion. Where, however, distinction has been drawn, it has been applied to those hard and densely fibrous tumours that arise from the surface of the alveolar processes, involving the periosteum, and by their expanded growth stretching the otherwise healthy gum over them. These are essentially different in origin, history, and anatomical structure, from the loose flaps, and often pedunculated masses of gum, mere hypertrophied integument, that are not unfrequently seen in the neighbourhood of decayed teeth, and to which the designations "polypus" and "fungus" of the gum have been appropriately given.*

* 'Specimens of Epulis,' by S. J. A. Salter, *Trans. Path. Soc.* vol. v. 1854.

An epulis tumour consists of a hard dense mass growing slowly and evenly from the edge of the alveolar process usually between two standing teeth, and more commonly on the labial buccal aspect than the lingual. The point at which this growth generally makes its first appearance is beneath and involving the little tongue of gum which exists between the necks of two contiguous teeth: as it progresses in size, it displaces the neighbouring teeth, one usually more than the other: it has for the most part a broad base, and increases in basal area more than in projecting growth. The situation in which an epulis commences is liable to a good deal of variety: though usually at the free edge of the alveolar process, it may form at a distance from it; but I believe it is always associated with porous vascular bone, nearly connected with the periodontal membrane, and does not spring from the compact tissue limiting the outline of the bone. The growth sometimes commences in the tooth-socket. As I have remarked, the *endosteal* membrane takes part in the genesis of an epulis tumour, and the fibrous growth appears to burrow, so to speak, into the substance of the bone, producing a general expansion of the whole structure. The surface of the tumour is like that of the surrounding gum; it is, however, sometimes mottled, and not infrequently slightly lobulated. It is as insensitive as the surrounding gum, and is not liable to bleed: when manipulated, it is tense and elastic. The tumour varies in size indefinitely—from the size of a pea to that of a walnut, or larger. It is generally stated that epulis tumours are more common in the upper than the lower jaw: this coincides with my own experience. I have happened to have met with them in the proportion exactly of two of the former to one of the latter. The epulis tumour appears to have a certain relation to the teeth in whose neighbourhood it forms. It almost always makes its appearance where there are teeth: it usually invades one in a very marked degree more than any other near which it may be situated, displacing it and pushing it out of place: it has nothing to do apparently with caries of the tooth: the removal of the particular tooth, with excision of the tumour, is almost always accompanied by immediate and complete cure of the disease.

Occasionally these tumours appear where teeth have been removed and the gum seems to be edentulous: it will, however, generally be found in these instances that a fang of one of the teeth has been left behind, and is associated with the irritation

that has caused the morbid growth. One of the most severe examples of this malady which I have seen, consisted of a bilobular mass, the size of a large walnut, extending, on the left side of the lower jaw, from the dens sapientiæ to the canine tooth, the four intermediate teeth having been removed. The excision of the tumour had been repeatedly performed, but it always returned. Its removal on this occasion disclosed the remains of one fang of the first molar tooth in its very axis; this was extracted, and the disease did not again make its appearance. I believe this will generally be found the case where the tumour forms upon an apparently edentulous region of the jaw. Epulis has its origin in the osteal membrane of the alveolus. The bone of the alveolar processes is very vascular, and almost cancellated in its loose open structure. The endosteal membrane which lines this cancellated bone, the periosteum which covers it, and which lines the tooth-socket, and the fibrous tissue of the gum, are all continuous, and all share in the development of an epulis tumour. The bulk of the tumour consists of a dense web of fibrous tissue; and from its basal attachment, and passing into its substance, are usual small growths of bone. The fibrous tissue interlaces pretty regularly—some fibres being parallel to the surface, and others radiating from the base, intersecting the former at right angles. The bone-growths may be thin needle-like spicula or little flakes: at the point whence these bony processes arise, the tumour receives its main vascular supply, and the subjacent bone is very porous. Mr. Caesar Hawkins mentions an instance of an epulis attached only by a pedicle to the gum, in which there was a nucleus of bone. This could not at any time have had bony union with the jaw; but the circumstance indicates, in a physiological sense, the osteal character of these fibroid developments. Epulis tumours are perfectly innocent, nor do they, I believe, ever pass into a malignant character. Sometimes, though rarely, they ulcerate on the surface; but this is only under the influence of external agencies, such as produce ulceration of any portion of the gum: they may then become painful.

The histology of epulis tumours is consistent with their history—locally recurrent as long as the circumstances under which they occurred are maintained, destroyed by the removal of their local nidus, and never impregnating the system. Epulis tumours are always, I believe, a form of 'fibro-plastic'; a

combination of fibrous tissue and myeloid cells, the proportion of the two constituents varying indefinitely. In general, the main bulk of the tumour consists of fibrous tissue; but sometimes the myeloid-cell element preponderates, and may form the major portion of the growth.*

The treatment of epulis is very simple; it requires early and complete extirpation with the knife, and according to the extent and inveteracy of the case it may also need the removal of contiguous teeth and subjacent bone. These latter circumstances refer to an important circumstance in the *pathological history* of the disease: it is this, that as long as the alveolar process remains upon which the tumour grew, and which must necessarily be as long as the teeth which are implanted in it are not removed, so long will the tumour be inveterate, so often will it return, however accurately and carefully it may have been removed. Such, at least, is the very general rule. But when the alveolar process is gone, it shows no tendency to return. And it is a remarkable circumstance, that the spontaneous absorption of the alveoli that follows the extraction of the teeth of itself, in many instances, where the disease is confined only to these processes, enough to prevent its recurrence. That is to say, if the tumour be removed to a level with the gum, and the contiguous teeth be extracted, the alveolar process vanishes by absorption, and the disease no more returns; though the same operation, without the removal of the teeth, may have been performed unsuccessfully any number of times previously. In a first operation when the disease is of limited extent, it will be well to try the result of its simple removal by the scalpel without the extraction of teeth or the cutting away of more bone than can be accomplished with the knife. The bone about the base and axis of the tumour being vascular and spongy, the knife readily cuts away small portions. Any subsequent sprouting of granulations in an undue or threatening degree should be checked and repressed by some caustic, and for this purpose trichloric acid has been especially recommended. If the disease recurs, as it too often does, the same operation should be repeated, and accompanied by the extraction of a tooth or teeth whose socket or sockets have been implicated. Unless the disease have a deep hold on the bony substance of the jaw, it

* 'Specimens of Epulis,' by S. J. A. Salter, loc. cit.; 'Myeloid Epulis of the Jaw,' by J. Hutchinson, *Trans. Path. Soc.* vol. viii.

will seldom be necessary to cut away any large amount of bone; sometimes, however, this is the case; or the growth may have started deep in the socket of a tooth. In such an instance it may be necessary to extirpate a considerable amount of bone. A V-shaped portion or a cubical mass may be readily cut off by means of a Hey's saw and bone-nippers, according to the form and extent of the particular tumour. It will never, I believe, be necessary to go far below the limit of the alveolar process; for it is with the alveolar bone that the disease is essentially connected. In the lower jaw I would urge the necessity of never cutting through the entire bone, as the breaking of the maxillary arch most seriously interferes with the position of the remaining portions of the bone, and thus disturbs the normal opposition of the teeth in the two jaws relatively.

Congenital hypertrophy of the gum and alveolar borders of the maxillæ.—In 1859 I had an opportunity of seeing a very remarkable example of disease, which is, perhaps, best defined by the above title, under the care of Mr. Pollock, at St George's Hospital. At that time there was, I believe, no recorded example of this curious tumour-like growth; nor had any instance of it been published, as far as I am aware, until a case was described and figured by Gross in the second edition of his *System of Surgery*.* This case was not only the same in essence, but was singularly like Mr. Pollock's in all particulars.

Two similar cases have since been recorded. One occurred in University College Hospital under the care of Mr. Erichsen, and is described by Mr. Heath.† The other is published by Mr. Waterman,‡ an American Surgeon.

Mr. Pollock's patient at the time of her admission into the hospital was eight years old. At birth nothing was noticed unusual in her mouth, but a fortnight after a tooth was cut, and by the fifth week six had appeared. It was then remarked that the gums were full and thick, and puffy. They continued to increase in bulk, and at two years of age they were cauterised, and all the temporary teeth which had appeared were extracted. At birth the child had an unusual quantity of hair on the head, and also much on the arms and legs: when taken into the hospital this peculiarity was very striking; the hair of the head was coarse and rank, and grew low on the forehead, and in front of the

* *A System of Surgery*, &c., by S. D. Gross, M.D.; Philadelphia, 1862. 2nd edition, vol. ii. p. 535, fig. 330.

† *Injuries and Diseases of the Jaws*, p. 189. London, 1862.

‡ *Boston Medical and Surgical Journal*, April 8, 1869, p. 167.

on to the cheeks. The arms and legs were covered with hair. The patient is epileptic.

Before any operation was performed, the girl presented a most extraordinary appearance. A large mass, pink and smooth, protruded from the mouth, which the lips did not and could not cover. It was slightly corrugated or indistinctly lined; the structure was very dense, inelastic, and insensitive, and appeared skin-like on the surface. The greater bulk proceeded from the upper jaw, and was most developed in the front of the mouth; but the same condition appeared along the whole edge of both jaws, that of the lower jaw being less, and covered and overlapped by the upper.

In structure, this hypertrophied mass consisted both of an expanded and elongated development of the alveolar borders of the maxillæ, and an immense thickening of the fibrous tissue of the gum, with a proportionately exuberant growth of the papillary surface. The removal of portions of the mass by surgical operation gave opportunities of examining its precise nature. In the front of the upper jaw, where the development was greatest, the fibrous mass extended in some places more than three-quarters of an inch beyond the alveolar edge, so that it thus covered in with a dense cushion. Those of the temporary teeth which had not been extracted were deeply embedded in the mass, the crowns of the second lower temporary molars being the only ones which were now visible. The crowns of all four of the six-years old teeth (first permanent molars) had appeared on account of the slightness of the hypertrophy towards the back of the mouth. A section of parts of the removed mass displayed the remaining temporary teeth completely clothed with the thick fibrous growth, the fangs bedded in sockets, but the crowns free of bone and each closely surrounded by a serous-like chamber without any communication with the surface. One of the superior central incisors was more deeply covered than any other teeth, and was nearly an inch from the surface. The fangs of the permanent teeth were developed in accordance with the age of the patient; but I observed that the crowns were still encased in the bony loculi, though from the age of the patient the distal wall of the bony capsules should, in many of them (incisors and first premolars), have been absorbed. In the socket of the first temporary incisor a small absorbed orifice existed, leading into the loculus of the permanent tooth, such as is usually found about five years old.

But the most remarkable point of structure in this growth was the papillary surface. The epithelium had changed into a very thick and hard epidermis, beneath which, and evenly covered in by it, were enormously long papillæ. The papillæ of gum vary from about $\frac{1}{16}$ th to $\frac{1}{32}$ th of an inch in length normally; but in a section, vertical to the surface, they here appear from the $\frac{1}{16}$ th to the $\frac{1}{2}$ th of an inch; and when by maceration the outer epiderm has been removed, the papillæ stand up like the pile of plush or velvet, and may be brushed from side to side by the finger. Kölliker* has noticed that in the mouths of edentulous people, whose bare gums are exposed to the rough attrition of food, the papillæ become much enlarged and elongated, attaining sometimes the $\frac{1}{16}$ th of an inch in length; and here, under exposure to still more physical violence, the same change has taken place to a vastly greater degree. But I apprehend that this monstrous development of the papillæ has also another meaning, and that it is an essential element of the disease; it is quite in keeping with the rough and

* *Mikroskopische Anatomie &c.*, von Dr. A. Kölliker, vol. ii. p. 85. Leipsic, 54.

thick skin of the patient, the abundant development of coarse hair over surface, and also the enormous teeth. The permanent teeth which were moved by the operations are excessively large, especially the superior incisors; these are larger than any I have before seen removed from a human mouth. All these circumstances imply a tendency to a general tegumentary and papillary hypertrophy.

Treatment.—The proper treatment of this disease is obvious, that which was followed in each of the recorded cases. The projections of the projecting mass were cut away with scalpels and bone-nippers till the alveolar borders were curtailed to moderate limits. The operation was not completed at once, but was performed in detail, as the patients could bear it. In Mr. Pollock's and Dr. Gross's cases there was a slight tendency to a fresh growth where the hypertrophied masses were removed.

Polypus of the gum.—The gum is liable to a simple hypertrophy, the increase of growth—a sort of mucous tubercle being confined to the gum structure alone, and not involving the osteal membrane, nor complicated with a growth of bone. The little tongues of gum between the necks of the teeth are liable to this affection, especially towards the front of the mouth; and they sometimes grow to such an extent, being confluent with those on either side, as to cover a considerable portion of the crowns of the teeth. This condition is usually associated with uncleanly habits; and may be generally prevented hereafter by simple and easy means. The gums should be thoroughly cleansed from tartar; the gums may be scarified: for the future the teeth should be abundantly brushed with a stiff brush, and the mouth washed with an astringent lotion, of which solution of permanganate of potash may be an ingredient. I have seen a modification of this condition which was evidently syphilitic; the growths of the gum were condylomatous, and were more or less covered with very painful superficial ulcers. The symptoms readily yielded to the internal administration of iodide of potassium, and the application with a camel's-hair brush of a solution of nitrate of silver, 10 gr. ℥j. aquæ. Hypertrophy of the gum of a truly polypous form not infrequently occurs in the immediate neighbourhood of teeth which are carious at their sides, and when the disease reaches the neck of the tooth: the irritation of the continuing dental disease seems to stimulate this unwonted growth, and the mucous membrane bulges into the cavity and fills it.

The cavities of two teeth carious on their contiguous surfaces are very apt to be associated with this condition, the little polypus of gum between them rising to their masticating surfaces. Such tumours are more or less pedunculated; they have a red fleshy look, and are very liable to bleed when roughly touched. The structure of these growths is in consonance with their aspect and the ease with which they bleed; they consist principally of hypertrophy of the true mucous-membrane element of the gum, especially the papillary structure: the epithelial covering is rather diminished in proportional amount, while the papillæ themselves become enormously elongated and increased in diameter, and some of the clinical papillæ develop into the compound fungiform variety. These changes are accompanied by great dilatation of the capillaries which loop into the papillæ.

Polypous growths of the gum are so often dependent on the state of the contiguous teeth, that their treatment involves that of the teeth also. If the polypus, growing into a carious tooth, is extirpated, it will most probably return till the tooth itself is removed, or the condition of the tooth which had irritated the gum has been remedied. The caries should be cut away; ragged or sharp edges of tooth-substance should be smoothed and blunted, and the remaining cavity should be filled. The gum may still show some tendency to renewed protruding, and this may be kept down for a time by the repeated application of a strong solution of nitrate of silver, phosphate of copper, or alum, when it will probably assume a healthy aspect. If the removal of the polypus is accompanied by the extraction of the offending tooth, I believe it never returns.

Vascular tumours.—The tissues about the necks of the teeth are obnoxious to the growth of vascular tumours, which vary from passive *nævus-like* swellings to those other forms of more arterial character—*aneurisms by anastomosis*. I have met with both these forms of the disease. The most common position in which it develops itself is in the front of the upper jaw, between the incisors, or canines and lateral incisors. A rather rare instance which occurred recently under my care manifested itself in the region that should have been occupied by the left lateral incisor of the upper jaw; but as the laterals were wanting in this person, the tumour formed between the line and central incisor. It had been about six months in

reaching its then size, the dimensions of a large marble flattened on the surface. It was of a purplish colour, streaked with many vessels on the surface; it was easily compressed, but was elastic, and when pressed it became pale, exsanguine, and much reduced in size; upon removing the pressure it resumed its previous aspect in a pulse or two. The surface of the growth was tolerably smooth; the base somewhat constricted, being about one-third less than the head of the tumour. The whole of the gums were very red, turgid, and swollen, and the little tongues of gum between the necks of the teeth generally were enlarged and spongy. The patient had suffered no pain, but was conscious of a constant throbbing and pulsation. The most important symptom, however, was the hæmorrhage, which had latterly become a serious source of trouble and distress; it usually occurred at night; it would ooze from the mouth and stain the pillow and sheets, and sometimes trickle into the glottis and cause momentary suffocation.

The treatment which I first adopted in this case was determined by the very *arterial* character of the tumour. I attempted to destroy it by ligature: a needle armed with a double thread was passed through its base, and each portion of the thread was tied so as to strangulate half of its attachment. The ligatures were tied tight, but did not cut through the substance of the growth. Their effect was for the time to produce complete strangulation: the tumour was tense; and the blood could not be squeezed out of it by pressure. This plan did not ultimately succeed, for the size was not permanently diminished and the circulation was re-established. I next removed the tumour by a very tight ligature, cutting it clean off; after this it returned as before, and I finally extirpated it with a scalpel, cutting freely inwards so as to remove a portion of the spongy vascular bone which seemed to form its basal axis. The bone at the base of these growths appears always to be very vascular and open in its texture. Considerable hæmorrhage followed the operation, which ceased under cold and pressure. For a few weeks the cicatrix furnished freely sprouting granulations; these were abundantly cauterised twice a week, and ultimately yielded a healthy scar. In structure this little mass when removed displayed a complicated vascular network, which, under the action of acetic acid, with the microscope, exhibited little else than an elaborate aggregation of the nuclei of blood-vessel muscle-cells. The surface was clothed with epithelium

capillæ, like the gum. I believe that no danger can arise in these cases from the use of the knife in at once extirpating the tumour; the bleeding may be profuse for a minute or two, but it soon ceases.

I have seen a vascular tumour connected apparently with the periosteum of a loose molar tooth, in which the hæmorrhage was very severe, occurring also at night to an extent that was really alarming: the tumour was a tense pedunculated mass, attached to the side of an upper molar tooth, half of the fang of which was naked. The extraction of the tooth brought away the tumour with it, which immediately shrivelled up to half its previous volume, and became soft and flabby.*

These tumours, as far as I have observed, occur in adult and middle life; they are perfectly innocent, and show no tendency to return when carefully eradicated.

Warty tumours of the gum.—The papillæ of the gum occasionally become hypertrophied into warty growths. A singular instance occurred in the practice of Sir William Fergusson, in which the papillæ were more than half an inch in length: they were soft and shreddy, and consisted mainly of loosely adherent epithelium. The tumour was removed, and recurred several times. Another example was operated on by Mr. Cock, at Guy's Hospital, in which a warty tumour, the size of a split chestnut, formed in the hard palate of a man.† This was covered with papillæ the eighth of an inch long. It consisted principally of fibrous tissue and dense coherent epithelium. I have seen a profusion of small warts on the gums associated with warts on the lips and about the face. In all cases the growths should be removed with the knife; and any recurrence should be resisted by caustic or astringent applications.

IV. *Tumours of the pulp.*—*Polypus of the pulp.*—In carious teeth the tooth-pulp is occasionally developed into a polypus-like growth, that fills more or less the diseased cavity. This formation is a dense, gristly, pink-coloured mass, consisting of a condensation of the original dentine-forming organ: it is always attached by a constricted base to the pulp in the canal of one or more of the fangs; the mass itself is usually more or less

* 'Vascular Tumours connected with the Dental Periosteum,' by S. J. A. Salter; *Trans. Path. Soc.* vol. v. 1854.

† 'Papillary Tumours of the Gum,' by S. J. A. Salter, in *Guy's Hospital Reports*, 3rd series, vol. xii. p. 358.

rounded, frequently corresponding in form exactly to the carious cavity which contains it; sometimes, by a more exuberant growth it is forced into a polyhedral shape by the contiguous structures, whose pressure curtails its further enlargement—the masticating surface of the opposing tooth in the other jaw, the distal and proximal neighbours of the tooth from which it has sprung, the tongue and the cheek; such cases as these usually occurring where the walls of the original tooth have completely broken away, leaving a distinct interval bounded by the parts so named. In such examples it may be difficult to an inexperienced observer to make out the nature of the tumour; it may be mistaken for an *epulis*, or a gum polypus; search for the remains of the decayed tooth will settle this doubt. This polypus is very callous and insensitive to pain; it is not liable to ulcerate or bleed; but it discharges pus from its surface, especially when in contact with the walls of the carious tooth in whose cavity it forms. In structure these polypi approach, in the majority of instances, very closely to the ordinary granulations of a healing sore, consisting superficially of a mass of exudation corpuscles through which are distributed multitudes of capillary loops, and more deeply of a fibrous tissue into which these cells have developed. In these instances the most superficial cells appear to be shed, assuming the form and aspect of pus. Rarely do these granulations seem to heal, to cicatrise; and I have found a few instances in which the surface of the polypus has been clothed with a dense cuticular epithelium overlying a basement membrane folded into true papillae. Polypus of the tooth-pulp 'most often occurs in young people, and in those in whom the teeth are imperfectly calcified, presenting that peculiar globular calcification in which the substance of the dentine becomes rapidly sodden with saliva, and carious without limit from the enamel to the pulp.' 'The physiological phenomena displayed by polypus of the pulp are very remarkable as regards both the pulp itself and the tooth—their oppositeness to the whole train of circumstances which accompany the ordinary inflammation of the pulps dependent on caries, odontalgia, lymph-deposit on fangs, alveolar abscess, &c. . . . The pulp never undergoes intrinsic calcification, nor have we any evidence that dentine of repair is ever formed.'* Occasionally, though very rarely, teeth with polypus of the pulp, produce alveolar abscess.

* 'Polypus of Tooth-pulp,' by S. J. A. Salter, *Guy's Hospital Reports*, 3rd series, vol. iv. 1858.

The treatment of this condition is very simple. If the polypus is cut away, it will certainly return; caustics and the like operations appear to have no influence in repressing it. If the tooth can be extracted, it is finally eradicated; and this plan should always be adopted.

Sensitive growth of pulp after fracture.—This condition only arises, as I believe, when a tooth with a healthy pulp is suddenly fractured by mechanical violence. When a pulp is exposed, it exhibits intense sensitiveness to touch, to cold, to heat, or indeed to any external influence; and it frequently grows into a small excrescence. This excrescence is usually of a semi-transparent aspect; it is often complicated on the surface, and villus-like: when removed, like the other form of sensitive growth, it certainly returns, and of the same character as before. In microscopic structure this sprouting of the pulp differs little from the insensitive polypus; but its vitality implies a more abundant nervous supply—vascular granulations appear to the observer to constitute its histological elements. This condition remains as a permanent torture to the patient, till the tooth is extracted, when it is of course completely removed. I have known this state of tooth-pulp form a very distressing complication in a case of fracture of the lower jaw, in which a bicuspid tooth was broken and the pulp exposed: the apparatus for fixing the displaced bone, and the introduction into the mouth of food, were attended with agonising pain, which continued till its source, a fractured tooth with a quick pulp, was discovered and extracted. Afterwards the treatment of this fracture was successfully and painlessly prosecuted. The possibility, indeed occasionally probability, of this complication, in treating fractures of the maxillæ, should be borne in mind by surgeons; and it should be especially remembered, that when extreme sensitiveness and pain manifest themselves in such cases, a fractured tooth should be sought for.

V. Tooth-tumours—Odontomes.—The hard tissues of teeth are sometimes developed into tumours, which may be divided as follows:—

- (a) Enamel nodules, or submerged cusps on tooth-fangs.
- (b) Exostosis.
- (c) Hypertrophied, dilated fangs.
- (d) Dentine excrescence.
- (e) Warty teeth.

Enamel nodules are sometimes seen on the fangs of teeth forming little pearl-like tumours: they are essentially *submucous*, each consisting of a little cone of dentine, covered by a thick tubercle of enamel, which is clothed by a true enamel. These tumours are not known to occasion any symptoms, and are rather of physiological than surgical interest: indeed they are alluded to here, simply to complete the list of tooth-tumours.

FIG. 231. The accompanying illustration (Fig. 231), is from a characteristic specimen.



Exostosis on the fangs of teeth is sometimes a very painful malady. An increased development of crusta may occur upon the fang of a tooth as the secondary result of other disease in the tooth—usually caries, or it may arise spontaneously—the tooth being otherwise free from other abnormal change.

In the former case the thickening of the fang is usually general, around its circumference and principally on its third, as represented in Fig. 232. The symptoms of this condition are scarcely to be distinguished or separated from those of the tooth-disease with which it is associated.

Where the disease arises spontaneously it is apt to produce the most distressing neuralgic affections. The teeth are otherwise sound, but they become sensitive to change of temperature, even to the touch: they elongate from the sockets and swell with this there is more or less neuralgia of the trigeminal nerve, one of its terminal branches. Then one particular tooth becomes more distinctly affected than the others: all the pain seems to emanate from it: and each flash of neuralgic agony seems to start from it alone. At length the tooth is extracted and the pain follows. Then usually another tooth becomes affected and requires removal: and this may, in very severe cases, go on until all the teeth in one jaw, or even in both, have been extracted before complete and permanent relief is obtained.† In some cases the exostosis usually consists of small nodules on the rounded surfaces situated near the apex of the fang. Various forms of exostosis may, however, exist without producing

* For the structure of these growths, see 'On Two Forms of Tooth-tumours' by S. J. A. Salter, in *Guy's Hospital Reports*, 3rd series, vol. xiv. 1869.


† See a striking case illustrative of this condition, published by the same author in *Guy's Hospital Reports*, 3rd series, vol. xiii. p. 86. 1867.

apparent symptoms whatever. Fig. 233 represents a magnified section of the fang of a tooth in which this nodular form of exostosis existed.

FIG. 232.



FIG. 233.



Exostosis on the fangs of teeth consists solely of an hypertrophy of the crusta petrosa, or tooth-bone, the outer layer of the fang. It is usually solid and compact in structure: though very rarely it is cancellated and vascular.*

Hypertrophied, dilated fangs.—This title best defines the nature of a very rare form of tooth-tumour, of which I believe there are only four recorded examples. One of these occurred in the practice of M. Maisonneuve and is described by M. Forget† another occurred to Mr. Hare of Limerick, and is described by Dr. Tomes;‡ another is described by Heider and Wedl;§ and a fourth is a specimen in the Museum of the College of Surgeons, England (Preparation 1022) which has been histologically examined and described by myself.|| The latter is supposed to have been in the collection of John Hunter. The first two of these examples were mistaken for exostoses.

These tumours consist of dilatations of the fangs of the teeth; the dentine-pulp being hypertrophied into a globular mass of considerable size; and, when calcified, producing a bone-like mass, generally larger than the tooth itself. In structure these tumours consist of an outer layer of tooth-bone; then a thin

* 'Vascular Exostosis,' by S. J. A. Salter, in *Path. Trans.* vol. vi. p. 168. l. viii. figs. 3 and 4. 1855.

† *Des Anomalies dentaires, et de leur influence sur la production des maladies des Os maxillaires*, par M. Forget. Paris, 1869. Obs. iii. p. 27, Pl. ii. figs. 1 and 2.

‡ *Transactions of Odontological Society of Great Britain*, vol. iii. p. 335. 1863.

§ *Atlas zur Pathologie der Zähne*, bearbeitet von weil. Prof. Dr. M. Heider und Prof. Dr. C. Wedl, p. 3, fig. 28. Leipzig, Nov. 1868.

|| *Guy's Hospital Reports*, 3rd series, vol. xiv. p. 463. 1869.

shell of true dentine, enclosing the gigantic pulp, which may may not be calcified. In the former case, this bulky nucleus the tumour consists of a mass of osteo-dentine.

In the only two cases in which the symptoms were recorded there was pain with expansion of the jaw at the alveolar port. In Mr. Hare's case there was a fistulous communication from the tumour through the jaw opening on the cheek. This discharged pus.

In each case the extraction of the tooth brought away the tumour also: the swelling subsided, and the patient recovered completely. Fig. 234 represents the specimen in the Museum of the College of Surgeons.

FIG. 234.



FIG. 235.



Dentine excrescence.—Nodules of secondary dentine growing from the wall of the pulp-chamber into its cavity, in otherwise apparently healthy, are sometimes seen, though have not generally been associated (perhaps from imperfect observation) with any definite symptoms. The accompanying illustration (Fig. 235), however, represents an example* where the little tumour evidently produced neuralgia, of a character very similar to that occasioned by exostoses on the fangs of teeth. The tooth was a superior central incisor, which was constantly painful: the slightest touch or change in temperature augmented the pain considerably, when it often flashed over the face and nerves of that side of the head with great severity. The extraction of the tooth, which was for the moment attended with a violent paroxysm of neuralgia, completely removed the subsequent pain. The dentine-excrescence was the only abnormal condition which the tooth presented. Its pressure on the nerves of the pulp would readily explain the symptoms in the case.

* 'Dentine Excrescence within the pulp-cavity of the Incisor Tooth. S. J. A. Salter, *Path. Trans.* vol. vi. p. 164, fig. 1. 1855.

This anatomical condition is not altogether uncommon, but it often causes painful symptoms.

Warty teeth.—These are among the rarest and most important of the tooth-tumours. They consist of teeth in which the tissues are hypertrophied and folded into an irregular and complicated mass. The warty condition may affect part of the crown of a tooth; or all the parts of the tooth may be involved; the irregular mass may consist of teeth blended together.

L. Broca, in a general résumé* of this subject, which he read before the Academy of Sciences of Paris (December 30, 1867), divides warty teeth into ‘circumscribed dentinal odontomes,’

‘diffuse dentinal odontomes;’ the former being those examples in which the warty mass occupies a portion of the crown only; the latter, in which the whole tooth is implicated, its anatomical form is no longer recognisable.

Warty teeth are exceedingly rare, and I am only acquainted with twelve recorded examples—six of the circumscribed variety and six of the diffuse. Of the former, four were described by myself,† one by Mr. Tomes‡ and one by Forget;§ of the latter, two were recorded by Oudet|| (in the same individual), one by L.,¶ one by Mr. Tomes,** one by Forget,†† and another by Harrison.‡‡

In the cases recorded by myself, one was a superior lateral incisor with a warty mass, the size of half a horse-bean, projecting from the front of the neck of the tooth. A section of this

Recherches sur un nouveau Groupe de Tumeurs désigné sous le nom d'odontomes, par M. P. Broca. *Comptes-rendus des Séances de l'Académie des Sciences*, tome lxxv. p. 1117. Paris, 1867. Also in *Gazette médicale de Paris*, 2, Jan. 11, 1868.

Description of a Warty Tooth, by S. J. A. Salter, in *Trans. Path. Soc.* vol. x. p. 173; London, 1855. ‘Contributions to Dental Pathology: On Warty Teeth,’ by S. J. A. Salter, in *Guy's Hospital Reports*, vol. iv. 3rd series; London, 1858. Same title, vol. v. 3rd series; London, 1859.

A System of Dental Surgery, by John Tomes, F.R.S., p. 226. London, 1859.

Des Anomalies dentaires et de leur influence sur la production des maladies des maxillaires, par M. Forget, p. 25. Paris, 1859.

Cas d'exostoses sur des dents devenues monstrueuses, par M. Oudet, in *Nouveau Journal de Médecine*, p. 245. Paris, 1821.

Grundzüge der pathologischen Histologie, von Carl Wedl, p. 626. Wien, 1861.

A further description of this specimen is published in Heider and Wedl's *Atlas zur Pathologie der Zähne*, part i. fig. 39.

* In Tomes's *System of Dental Surgery*, p. 225. London, 1859.

† Loc. cit. p. 5.

‡ *British Journal of Dental Science*, vol. v. p. 557. London, 1862.

tooth for the microscope is in my histological collection. The second case, which occurred in my own practice, was that of the right lower dens sapientiæ. The mass, about the size of a bean, sprouted from the posterior aspect of the neck of the tooth. My two remaining examples are specimens in the Museum of Guy's Hospital: one a lateral incisor, and the other a central incisor, both of the upper jaw: from the side of each tooth a warty mass projected. The accompanying illustration (Fig. 236) represents the lateral incisor referred to.

Mr. Tomes's specimen was a superior central incisor: the wart occupied the whole of the front of the crown of the tooth. In Forget's example, the first and second lower molars

FIG. 236.



FIG. 237.



blended into one: the crowns constituting a large conical wart-like body, while the fangs were normal. This is shown in Fig. 237.

The diffuse forms of warty teeth, which have produced large tumours, have all been developed in the lower jaw. They deserve more serious consideration.

In Oudet's case a man, twenty-five years of age, exhibited in his lower jaw hard bony tumours, occupying the premolar region one on each side. They were encrusted with salivary calculus, upon the removal of which, an irregular mass of enamel was displayed, folded into shapes like cusps and tooth-edges. These masses were evidently formed from the fused and hypertrophied germs of two premolar teeth on both sides of the jaw. The right was removed easily by tooth-instruments, but the patient declined to have the other interfered with.

In Wedl's case the whole of the second lower molar tooth of the right side was developed as a large complicated mass in which the several tissues of the teeth were confounded together. It was overlying and keeping down the wisdom tooth. With a pair of forceps it was removed without much difficulty.

The specimen described by Mr. Tomes is exactly the same as the foregoing. It consisted of a confused mass of dental tissues as large as a chestnut, representing the second lower molar tooth, beneath which was buried the wisdom

tooth. It occurred many years ago in the practice of Sir William Fergusson, who excised the angle of the jaw for its removal.

But M. Forget's is the most serious recorded example of this malformation. Here an enormous tumour, as large as a turkey's egg, occupied the whole of the molar region of the left side of the lower jaw. It represented two molar teeth with their tissues blended together in utter confusion and into a shapeless mass.

FIG. 238.



At birth it was a third molar tolerably well formed. The jaw began to expand in early childhood, and continued to do so till the patient attained the age of twenty, when M. Forget removed the angle of the jaw, including the tumour. The mass was covered by gum, excepting for a space on its surface the size of a sixpence. There was some pain at times: the submaxillary glands were enlarged, and several fistulous pus-discharging canals led to the base of the tumour. The accompanying illustration (Fig. 238), is from Forget's memoir, and displays a specimen of the tumour *in situ*.

Dr. Harrison's specimen represented the two bicuspids and canine of the left side on the lower jaw. The tissues were confounded together, and without resemblance to tooth-form. It loosened and came away by itself.

As regards the treatment of these cases, much depends on the degree of the malformation. If it is only partial and a matter of appearance, the tooth may be removed and replaced by a better-shaped artificial representative, according to the wishes of the patient. But, should the growth be of large size and interfere with the action of the mouth, it should certainly be extirpated; and this, in nearly all instances, might be accomplished by extraction with forceps; or, at all events, without serious damage to the jaw.

VI. '*Abscess*' of the antrum.—The term *abscess* of the antrum conveys a very wrong impression of the real nature of this disease; it is not the suppuration of inflamed parenchyma, but the occlusion in a cavity of the purulent secretion from the surface of a mucous membrane which lines that cavity. I need not dwell on the anatomical characters of the antrum maxillare further than to notice its peculiar relations to those organs, the teeth, whose affections are by far the commonest cause in the production of the malady we are considering. These relations vary extremely, both as regards the extension of the antrum over the fangs of many or few teeth, and the degree in which those fangs approach or pierce the floor of the sinus. It may extend so as to be in immediate relation to all the teeth of the true maxilla from the canine to the dens sapientiæ, or it may be contracted to such narrow limits as only to correspond with two or three of the central ones. This variation in the size of the antrum is not mentioned in text-books of anatomy, but it is pointed out by Otto.* Occasionally a root or roots of the first molar tooth (rarely any other) extend into the cavity, free of any bony covering, and merely overlaid by the mucous membrane lining the sinus; more often, however, the palatine and external roots diverge so as to leave an interval between which the more depending sulcus of the antrum is excavated. Another circumstance in the anatomy of the antrum bearing on the purulent accumulation which constitutes abscess of this sinus, is the very variable size of the orifice which opens into the middle meatus of the nose. In some instances the aperture is barely sufficient to admit the blunt point of a probe; in others it would allow the passage of the end of the little finger. It is always much smaller in the living state when the mucous membrane lines and fills up the opening; indeed, it is the pouting of the tumid mucous membrane which closes the orifice in inflammation. The lining membrane of the antrum is liable, like all other mucous membranes, especially the Schneiderian membrane, of which it is a continuation, to inflammation and altered secretion, mucus being impregnated with or replaced by pus, and accumulating in quantity. This may occur in different degrees, both as to amount and rapidity of development; but the circumstance which here gives importance to this altered and

* *Lehrbuch der pathologischen Anatomie des Menschen und der Thiere*, von Dr. A. W. Otto, p. 180. Berlin, 1830.

ist by alveolar abscess, in some stage, associated with disease. It is stated that the affection may be brought on by a blow on the cheek; and an instance has been recorded in which the malady has occurred in a new-born child, and was supposed to result from the pressure on the cheek during a hard

symptoms of antral abscess vary much in degree; but usually commence by dull aching pain in the cheek, with redness, and fulness of the soft parts externally. In the early stage there may, or may not, be a purulent discharge from the corresponding nostril; this, however, can only occur to any extent before the aperture between the middle meatus and the sinus is closed. As the case advances and the matter becomes pent up, the pain assumes a throbbing character and increases, and constitutional symptoms manifest themselves analogous to those of acute abscess; the patient has rigors and

With this the local symptoms change, and an expansion of the whole jaw shows itself; the malar bone becomes elevated, the maxilla beneath it full and prominent; the molar teeth on the affected side appear to elongate, and, in closure of the mouth, they meet their opponents too soon; the concavity of the hard palate becomes flat or even convex; the nostril of that side is more or less closed, and, in severe and protracted cases, the lower border of the orbit becomes so pushed up as to protrude the eye. Mr. Abernethy, in describing the symptoms of antral abscess, speaks of the maxilla being sometimes 'affected,' but in what manner he

tion accompanying abscess of the antrum is, however, occasionally so severe as to implicate the periosteum, not only to the destruction of some parts of the maxilla, but extending beyond to contiguous fibrous structures, so as to involve the optic and other nerves in their passage from the cranial cavity to the orbit, producing blindness and fixedness of pupil on the affected side. Such cases are very rare; but one has occurred in my own practice at Guy's Hospital;* a second was under the care of Mr. Pollock; and a third is mentioned by Dr. Brück in Casper's *Wochenschrift*.† Necrosis of part of the jaw is one of the rarer accompaniments of this malady; but when any portion of the bone is affected, it is usually either the alveolar processes or the nasal plate of the maxilla.

The manual examination of the jaw seldom fails to indicate the fluid nature of the distending material; fluctuation can be felt, and the thinned bony parietes of the abscess convey to the touch a peculiar sensation, like the handling of dry parchment: as Jourdain very characteristically expresses it, 'quand on appuyoit sur l'os, il s'affaïssoit et revenoit ensuite sur lui-même en produisant une espèce de craquement.'‡ If any doubt does arise as to the nature of the contents of an expanding antrum, I am in the habit of exploring the cavity with a minute trocar and cannula not above half the size of a wheat-straw; by this means sufficient fluid, if present, may be made to escape to ascertain its nature, and it could scarcely do injury to a solid growth if such were the character of the expansion. A grooved needle would answer the same purpose, but hardly so readily. The fluid may be too dense to pass these narrow canals; generally, however, it is not thicker than ordinary pus, and it is sometimes serous.

The ultimate issue of these cases varies very much. Occasionally the abscess bursts into the nose: sometimes through the cheek; occasionally into an alveolar cavity, burrowing along the fang, and discharging its pus at the neck of the

* 'Case of Amaurosis from Abscess of the Antrum caused by a carious Tooth,' by S. J. A. Salter, *Med.-Chir. Trans.* vol. xlv.

† *Wiederholte Entzündungen des Antrum Highmori und Amaurose, etc.*, mitgetheilt vom Dr. T. H. Brück, in Casper's *Wochenschrift*, März 1851, Berlin. A case of a similar character was described by Prof. Galenzowski, in *Archives générales de Médecine*, tome xxiii. p. 261. Paris, 1830.

‡ *Traité des Maladies, etc., de la Bouche*, par M. Jourdain, tome i. p. 120. Paris, 1778.

ens to point; for that would generally be through the skin cheek.

the disease is known early, before it has caused the de-
struction of the fore part of the bone, there are two ways of
draining the abscess: one by perforating the partition between
the gum and the nose, which may be done; and the other
by perforating the first or second grinder of that side, and perfo-
rating the partition between the roots of the alveolar process and
the gum, so that the matter may be discharged for the future
safety.

if the fore part of the bone has been destroyed, an
opening may be made on the inside of the lip, where the
pain is most probably will be felt; but this will be more apt
than any other perforation to heal, and thereby may occasion a
recumulation; which is to be avoided, if possible, by putting
in all the common methods of preventing openings
from healing or closing up; but this practice will rather prove
harmful; therefore the drawing of the tooth is to be
preferred, because it is not so liable to this objection.*

When the abscess has formed, and when, as yet, it is only
incipient, it may be arrested by removing any carious tooth
near in the neighbourhood; and the application of leeches,
bleedings, &c., with the administration of purgatives. In
cases where the pus has already accumulated, and there
is no outlet, it will be necessary to make a free opening for the
issue of the matter: and the method which universal

advantage of removing the cause in most cases, and of allowing the discharge of the matter at the most dependent part of the containing sac. The tooth whose fangs are usually most intimately associated with the antrum is the first permanent molar; and its removal, in a case of antral abscess, is especially indicated from this circumstance, and from the frail and perishable nature of the tooth itself, which gives it less of than other teeth a long tenure of usefulness. The relation of the fangs of the molar teeth to the antrum is an important consideration in the treatment of this disease: in the large proportion of cases caries of the first molar is the cause, and its removal (opening an alveolus for further perforation) is the first step in the treatment; but any other tooth, molar, bicuspide, or even canine, whose disease should be considered its removal ought in preference to be extracted, as the absorption of the fang of a carious tooth would render the perforation of the antrum easy, while, by such a selection, this preliminary operation would remove the exciting cause of the disease. Indeed, as a rule, it is well to extract all carious teeth from the side of the upper jaw affected with antral abscess. The extraction of the abscess-causing tooth is frequently followed by discharge of the contained pus, in consequence of the fang having extended into the antrum, or the floor of the sinus having been absorbed resulting from the diseased tooth. It is generally necessary to enlarge the orifice into the antrum; and this is best effected by a naked trocar pushed up the socket of the extracted tooth; the trocar should be large, so as to break down a good deal of the floor of the antrum, thus making a free orifice for the discharge of matter. The absorption of bone which occurs around the fangs of carious teeth much facilitates piercing the antrum in this manner. Perforating the antrum occasionally requires considerable force; and it is necessary to apply force with great care: the fore-finger should be extended to the shaft of the trocar as a guard, and the instrument should be pressed forwards with an even rotating motion. If these precautions are not taken, the floor of the antrum will sometimes give way suddenly, the trocar will traverse the cavity of the sinus, and strike hard against the floor of the orbit, and it may even pierce. I once saw this accident occur in the hands of a young operator, fortunately without any serious consequences. Occasionally antral abscess is associated with necrosis of some portion of the walls of the sinus; and

When the teeth have been long removed, and the alveolar processes absorbed, the floor of the antrum would be more difficult to pierce, on account of the thick compact layer of which it is constituted by the *osteal cicatrix* after the removal of the teeth. In such a case it would be more easy to enter the antrum by perforating at the base of the malar process of the maxillary bone over the region formerly occupied by the second or third molar tooth. The mucous membrane should be divided first; and then the bone, when quite exposed, should be cut by means of any instrument suitable for the purpose: a trocar; or, as Sir B. Brodie suggests, a strong pair of scissors, closed and held firmly in the hand, should be *bored* into the part chosen for perforation.

When the antrum has been perforated, the next step is to perform the complete washing out of the cavity by injections, to insure the free and continued egress of the discharge by the natural opening. Warm water should be abundantly used at first, so as to remove all the matter, which is sometimes accumulated. This will give great relief, and the swelling and inflammation will usually subside at once. Should the tendency to re-secretion continue, an injection of sulphate of zinc, or in an obstinate case I found very efficacious, a solution (from 1 to the ounce) of nitrate of silver in distilled water, may be employed. I am in the habit of using a glass syringe with a fine nozzle; fitted to the extremity of this is a bent silver

of two kinds may occur;—first, food may pass through the orifice into the maxillary sinus; and secondly, the aperture, thus artificially made, may close and cicatrise over. To prevent this, the plan usually adopted has been, to plug the orifice with a piece of wood, which effectually prevents both ill consequences. This is, however, a clumsy method: the same and further advantages may be gained by adopting a plan which I have followed in some recent cases. After the perforation through the alveolar cavity has been made, I have taken a model, and a plate has been prepared to pass over the space occupied by the extracted tooth, and fastened to the contiguous teeth by the customary metallic bands. Through this plate a hole has been bored, which corresponds to the orifice into the antrum; and to the applied surface of the plate a very short tube has been soldered, sufficiently long just to enter the antrum, to occupy the perforation, and prevent its closure. This plate has been permanently fixed during the active treatment of the case, and the orifice in the tube has been kept closed by a plug of cork in the intervals between using the injection. By the removal of the cork, the nozzle of the syringe, which fitted the tube, could be applied, and the injection used any number of times, the tube being sufficiently short to allow the complete washing out of the cavity and the escape of the fluid. This plan of treatment has the additional advantage, that when there is no further need of keeping open the artificial orifice, the plate over the gum facilitates its closure. By removing the tube, and closing the hole by a little sheet of metal soldered on, the passage of air and fluid from the mouth to the antrum is suspended, and the healing of the wound is thereby facilitated, as will be hereafter explained. (See section on the Application of Obturators, &c.)

If in operations for antral abscess any foreign body, which may have caused or be associated with it, such as the fang of a tooth, becomes loose in the sinus, it should be remembered that the cavity is occasionally divided, as shown by M. Giralès,* by partial septa of bone projecting from its walls. In such an instance, the adventitious body may be pocketed in a circumscribed region of the sinus,† and can only be removed by some

* *Des Maladies du Sinus maxillaire*, par M. Giralès. Paris, 1851.

† An interesting example of this casualty is mentioned by Mr. Cattlin, in the *Transactions of the Odontological Society*, vol. ii. p. 38. London, 1861.

gh misplaced as regards position: further, in every
red case, the tooth or teeth thus implicated in disease,
been of the successional or permanent set, though a
ary instance has happened in the practice of a friend of the
or's, in which a temporary tooth was the dental element of

he dentigerous cysts, to which I have above referred, are totally distinct
ir origin and physiological meaning from those other tooth-bearing
n which are found, some in the ovaries of females, and others variously
uted in the bodies of either sex. The dentigerous cysts occurring in the
merely the result of misplacement of a tooth, or tooth-germs belonging
jaw which contains them. The two other forms of dentigerous cyst
belong to the individual in whom they are found, nor are they adven-
growths of disease, properly so called. They obviously represent a por-
ore or less) of another individual, and indicate some curious aberration
unction of reproduction. The ovarian tumours which bear teeth, and many
ructures found in the ovary of the human female, I conceive to be the
equivalents of the virgin-produced 'zooids' of those invertebrata which
the function of *parthenogenesis* in the reproduction of their kind: that
elopment of these so-called tumours is the physiological equivalent of
ction, though the anatomical result is incomplete. (See 'Ovarian Tumour
ag Teeth,' &c., by the author, in *Guy's Hospital Reports*, 3rd series,

Those other tooth-bearing tumours which are found in various parts of
y, irrespective of sex, cannot be susceptible of the same explanation, but
the entanglement of an imperfect, or more or less perfect, ovum within
ary one. Teeth appear to be the most constant of the structures enter-
the formation of these tumours; but they are associated with other
even in more than one instance to the production of an entire fœtus.
velopments receive a probable elucidation by those curious monstrosities

one of these cysts. These tooth-bearing serous tumours are, therefore, to be looked upon only as the occasional complications of dentition, in which there is an accidental deviation in the anatomical position of some tooth or teeth. The cysts only arise when the tooth or teeth associated with them are embedded in the substance of the jaw-bone; they do not occur after the tooth has pierced the gum. The embedding of a tooth in the bone does not necessarily give rise to these serous collections, for that is by no means an uncommon occurrence, whereas dentigerous cysts are rare. There appear to be three circumstances which may either of them produce impaction of a tooth in the substance of the maxillary bones: the tooth may be originally developed too deep in the body of the jaw, and thus, though it grow in a right direction and in a right position as regards the series, it will never reach the alveolar margin; or while it may be sufficiently superficial, it takes an oblique direction of growth, so that it lies covered more or less in the axis of the bone; or again, the position of the tooth and its line of growth may be originally normal, but from an arrest of the development of the fang it may fail to reach the alveolar edge, and so remain permanently impacted in the maxilla.* This may occur to any tooth, and has been recorded as affecting most of the permanent set; but it is more common with some than with others: it has been more often met with as regards the upper permanent canines than any other teeth. In cases where a successional tooth is impacted in the jaw, its temporary predecessor is usually retained considerably after the natural period of shedding. When a tooth is thus situated, its fang is enclosed in a bony socket lined by periosteum, as in ordinary circumstances, while the crown of the tooth is free in a little bony locus lined by that which was the so-called 'enamel-pulp.' This structure is clothed with a sort of epithelium, which is apt to assume the function of secreting fluid. After the enamel is completely formed, the soft membrane which rests upon the surface of the crown of the tooth frequently separates from it, the interval being occupied by a sort of serum. This is generally the result of some irritation or difficulty in tooth-cutting; and where the irritation runs on to acute inflammation, as in some cases of tedious eruption of wisdom-

* 'On the Impaction of permanent Teeth in the substance of the Maxillary Bones,' by S. J. A. Salter, in *Guy's Hos. Rep.* vol. v, 3rd series.

It has been associated with this condition.

Jourdain* describes three cases of dentigerous cysts. The first case he records was that of a girl, seventeen years old, in whom the right upper first and second permanent molars were inverted, and a large serous cyst had expanded around them into the antrum, with great dilatation of the body of the teeth, distortion of the side of the face, and closure of the mouth; it had existed 'some months.' In a second example, a woman, about sixty years of age, suffered from a tumour the size of a pigeon's egg in her upper jaw, for many months, closing the nostril of that side; it was caused by the expansion of a cyst around the second premolar, which was impacted deep in the substance of the bone. The third instance occurred in a child, thirteen years of age, from a monstrous permanent lateral incisor (side not stated) being embedded in the intermaxillary space above and behind the root of the central incisor; 'there was a considerable tumour, which occupied anteriorly the greater part of the region of the maxillary hollow.' It had existed for a

number of years. Guyton† describes a specimen shown to him by M. Loir, in which a cyst was developed in the left superior maxilla: this was unilocular, and was dependent upon, the presence of the tooth in a reversed position.

He is informed of a case of this disease by my late colleague, Dr. Mansby Cooper, in which great swelling occurred in the soft tissue of the superior maxilla of a young man, in whose

made their appearance. A free opening was cut into the expansion, when the wanting teeth were found inverted in the axis of a serous cyst expanding into the antrum. No account of this case appears in Mr. B. Cooper's *Lectures on Surgery*, but I took memoranda of its particulars when communicated to him.

Mr. Wormald operated on a boy, about fourteen years of age, some time since, at St. Bartholomew's Hospital, in whom a serous cyst had expanded about the second bicuspid tooth of the lower jaw; the cyst was the size of a large chestnut, and was in the axis of the bone: the fang of the tooth was not fully developed. The interior of the cyst was lined with a thick vascular membrane, and it contained a glairy fluid. Mr. Wormald has favoured me with these particulars.

F. E. Glaswald published a very learned discourse on this malady, at the University of Greifswald, in 1844, containing a complete résumé of the literature of the subject.* The text, upon which this essay was elaborated, was a case that had been under the care and treatment of Prof. Baum. In this instance a cyst had dilated each antrum to an enormous extent, and with hideous disfigurement of feature. The patient was a woman, thirty-eight years of age, and the disease was said to have been in progressive existence for thirty years. From a cyst in the right antrum a canine tooth was removed, and from the left a molar. The fluid in the cysts was purulent.

Three instances of this condition have occurred in my own practice. In one a serous cyst expanded the left angle of the lower jaw in a young man twenty-two years of age, resulting from the impaction of the dens sapientiæ. The cyst was very large, and had been twice opened to allow the escape of the serum, which had in each instance rapidly re-collected. I extracted the second molar, which was loose: this ruptured the cyst, liberated the serum, and freed the impacted tooth; its crown appeared in the alveolus of the second molar, whose posterior fang was greatly absorbed by the progress of the cyst. The case was entirely cured by this means alone; and the dens sapientiæ gradually rose into the mouth in an oblique position. The second example is very interesting in a diagnostic point of view. A girl, eighteen years of age, had an elastic fluid-containing tumour in the substance of the incisive bone, extending up to the base of the nose on the left side. She had been seen

* *De Tumore quodam utriusque Antri Highmori perversa dentium formatione exorto*, auctor Franciscus Edwardus Glaswald; Gryphæ, 1844.

er three surgeons; but the nature of the malady was retained. She had the normal number of teeth in the arch, though the character of one of them was abnormal for her age. When the patient was sent to me for my opinion, I perceived that the left central incisor was a temporary tooth; and this circumstance was a key to a correct diagnosis of the case. The temporary central incisor occupied a position which its permanent successor should have held: the absence of the permanent under such circumstances, suggested the almost impossible position which it must occupy above and behind its permanent predecessor, that is in the axis of the serous cyst. The temporary tooth was removed, and the cyst explored, to make room for the succeeding tooth. The permanent central incisor lay deep in the bone, in an upright and natural direction, its crown bare within the cyst; but upon its removal it was found that the fang was aborted, and had only grown to half its natural length. This circumstance it was which prevented its extruding its temporary predecessor, and kept it in its normal position. The retention of the tooth in its epithelioid sac furnished the anatomical grounds upon which, under favouring circumstances of irritation, the eruption arose, and the bone-expansion followed. The patient was that of a gentleman about forty years of age. The size of a pigeon's egg, had expanded around the central incisor on the left side, the tooth being impacted in

The fluid in the sac was thick, very dark yellow-serum, containing a large quantity of cholesterine

logical specimens of uncommon diseases, and those too which shorten life, must necessarily be very rare. It lies to specimens illustrative of the disease we are con-

One is in the possession of Mr. Samuel Cartwright, and is valuable and instructive in many points.* The specimen consists of a right superior maxilla—an adult bone: the teeth that remain, and the alveoli from which others have been extracted, show that the normal number of permanent teeth had developed in their natural position. The turbinated bones are gone, and the antrum maxillare is open. In the middle of the palate, starting from its base, but not attached to its lateral

through the kindness of my friend Mr. Cartwright, I have been allowed to examine the specimen. See *Guy's Hosp. Rep.* 3rd ser. vol. v. p. 328.

walls anywhere, is an exceedingly thin, delicate capsule of bone, about the size of a chestnut, white with a granulated surface. The bony capsule contains nothing but one small *supernumerary* tooth, which is *loose* and *free* in the cavity. There are three important points in this specimen bearing on the anatomy and history of these cases:—the tooth is a *supernumerary* one; it is free and wholly detached in the cavity of the cyst; and further, the expansion is not that of the antral wall itself, with the tooth's crown uncovered within it, but a distension of that which was the bony *loculus* of the contained tooth, which, by its further dilatation, would have expanded the antral wall, and probably have been confounded with it.

This latter circumstance appears to me to be of much anatomical interest in reference to those serous expansions of the antrum which are associated with inverted teeth. These have hitherto been described as cysts of the antrum itself: but I cannot conceive that a tooth being 'cut' through the mucous membrane of the antrum should produce such a result. The appearance of the crown of an inverted tooth in the nostril neither causes irritation nor increased secretion—at least no such consequence followed the three examples I have seen; and it seems to me to be far more probable that these cases have commenced as cysts within the bone at the base of the maxillary sinus, expanding into it, and ultimately filling its whole cavity and dilating its walls.

Two specimens in the Museum at St. Bartholomew's Hospital throw some further light on the pathology of this condition. One preparation (numbered 1. 119) exhibits a thin bony cyst, the size of a small Tangerine orange, with a thick membranous lining: it contained an inferior permanent canine tooth, *loosely attached* to its walls. It was removed from the lower jaw in the region of the contained tooth. The other preparation (1. 119a) is the superior maxilla of a young sheep, in which the central incisor is attached to the side of a large cyst; the *fang* of the tooth being *almost wholly destitute of bony covering*. This denudation of the fang does not appear to exist in the early stages of these cases, but seems to be brought about by great expansion of the cyst, and the progressive absorption of bone which accompanies it.

In one instance only am I aware of a *dentigerous* cyst being associated with a *temporary* tooth. It occurred in the practice of my late friend Mr. Alexander Edwards of Edinburgh. The

as a young man, in whose upper jaw, just below the tumour had developed: the tumour consisted of exostosis maxilla, combined with a bony cyst, containing a tooth, as pronounced by Prof. Goodsir to be a temporary portion of the tooth, which I afterwards saw, it to be the second, the larger of the two.*

Symptoms of a dentigerous cyst are almost wholly local, of a general expansion of the jaw-bone at some part; accompanied by a corresponding disfigurement of adjoining features, and a sense of weight and tension in the affected part. Where the impacted tooth has produced pressure upon a neighbour, the symptoms of pain and local inflammation have been more considerable, and have given rise to constitutional irritation.

On manipulation the fingers readily perceive that the bone has a central expansion of bone, and that it contains bony walls yield to pressure, and then return to shape, and a peculiar kind of crepitation which Jourdain characterises as *craquement*, like the doubling of stiff parchment; the bone is usually sufficiently thin at some part to allow of fluctuation under the pressure of alternate

the most usual symptoms, and which is also an important sign, is the absence from the mouth of some teeth which should have appeared, and have never been

The presence of a fluid-containing tumour within the space of a maxillary bone at the region of a tooth which is not and known always to have been wanting, would be conclusive as to its being a dentigerous cyst, though the absence (or the past-known presence) of every mature tooth does not necessarily prove the reverse; for the dental element may be a temporary or a supernumerary tooth.

Examples of dentigerous cysts may be referred to:—‘Tumour of the jaw depending on Cysts connected with the Presence of Teeth in a Pre-eruptive Position,’ by James Syme, Esq., in *Edin. Med. and Surg. Journal*, vol. lxi. 1838. ‘Tumour formed by the Capsule of an unerupted tooth,’ by T. Wormald, Esq., *Lancet*, vol. i. p. 758, 1850. ‘Dentigerous Cysts,’ in Stanley’s *Diseases of the Bones*, pl. xviii. p. 20 of Illustrations; 2d. ‘Three cases of Dentigerous Cysts,’ in Forget’s *Des Anomalies du Développement*, Obs. x. xi. xii. pp. 41–47; Paris, 1859. ‘Case of Bony Cyst of the Lower Jaw,’ by S. W. Fearn, in *Brit. Med. Journal*, vol. i. p. 241. 1864. Heath’s *Injuries and Diseases of the Jaws*, p. 158; 1868.

THE TEETH.

A loose tooth is a very great evil, but still more uncommon than a supernumerary tooth. The diagnosis is made by exploring the cyst either by the finger or by the probe, when a serous discharge is felt; and if the cavity be laid open, the hard unyielding nature of the tooth is discovered.

In these cases is obvious, and usually the first step is evacuating the contents of the cyst, and the teeth embedded in it, and when the bone is removed, moving some of the dilated bone. The operation is performed as early as possible, so as to prevent the cutting away much bone, and thus to avoid the absorption which would follow before the tooth returns its natural form. All cutting should be done in the mouth. Generally a portion of the tooth can be removed readily enough with a saw. If the tooth is reversed, the expansion is not so great at the alveolar border. In that case the removal of contiguous teeth, or even the sawing of the bone, has been especially the case where in the alveolar border cysts in the antrum. The tooth is like a plug in the case of the cyst, further, *i.e.*, from its thickness. The difficulty may be found in getting the tooth out of the embedded tooth; but various long-handled levers may be readily devised to suit any case that may arise.

From a persistence of the serous secretion it is necessary to inject the cyst with astringent solutions.

In these cases is, I believe, always satisfactory, and that it ever led to necrosis or other bone-disease which I saw, a fibrous tumour grew from the wound some months after the first operation, and did not recur.

Alveolar and maxillary necrosis from (a) *Phosphorus* and (b) *Caustic potash*.

The action of the phosphorus poison to the disease, and its method of introduction, or rather application, is the most distinctly proved of any of the circumstances.

is employed in the making of lucifer matches, and
ie (or scarcely with exception) it is that this disease
sted. But the circumstances which connect the out-
se with the disease that follows it, is a predisposition
lividual, consisting of some exposure of the periosteum,
; tantamount to such exposure. The only manner in
s occurs, at least in which phosphorus appears to be
in causing bone-necrosis, is where caries of a tooth
he pulp to the poison-influence, the bone-necrosis
t of the jaw. It is the poisoning of the tooth-pulp
e essence of the disease; the severe combinations of
tion, which give all the importance to the malady,
contingent and secondary consequences. It is this
e essential nature of the disease that links it (as I
that other form of maxillary necrosis which occurs in
fter attacks of the eruptive fevers; only that, whereas
osphorus disease the poison is applied to the tooth
xtraneous source—from without—in the jaw-necrosis
re fevers the poison is generated within, and alights
teeth and tooth-pulps, by virtue of their being dermal
embers of the tegumentary system, upon which system
the eruptive fever-poisons spend their chief destructive

phosphorus disease.—The necrosis and exfoliation of portions
aw-bones, dependent on phosphorus fumes as its cause,
associated with the manufacture of lucifer matches



The earliest published account of the disease which we have is by Lorinser,* of the *Bezirks-Krankenhaus* Wieden; and the first case which fell under his notice occurred in 1839, about eleven years after the opening of lucifer-match manufactories in Vienna. In this country, as far as I am aware, the malady was first recorded by Dr. Wilks, in 'Surgical Reports of Guy's Hospital,' from April 1846 to March 1847, where he remarks: 'Of the other diseases of the lower jaw, one occurred in a lucifer-match maker, with suppuration and exfoliation of bone.†' It is, however, to the continental surgeons, and those principally of Germany, that we are indebted for the complete and early account of this malady, from which all subsequent notices have been mainly derived. Besides the original memoir by Lorinser, before referred to, important contributions have been added by Strohl,‡ Heyfelder,§ Roussel and Gendrin,|| Sédillot,¶ Ebel,** and, above all, by Von Bibra and Geist,†† whose exhaustive treatise, their joint production, has given the clearest elucidation of this new disease.

The most important article on this subject that has appeared in this country has been written by Dr. Bristowe, in the form

* I have had no opportunity of consulting Lorinser's original writings: they are referred to by Geist as—'in den medicinischen Jahrbüchern des k. k. Oesterreichischen Staates, Jahrgang 1845, Märzheft.'

† *Guy's Hospital Reports*, 2nd series, vol. xii. p. 163. London, 1847. This case occurred in the hospital practice of the late Mr. Aston Key, who, as Dr. Wilks informs the author, was already aware, from his own observation, of the essential nature of the disease and its relation to its peculiar cause.

‡ *Gazette médicale de Strasbourg*, cinquième année, No. 11, 20 novembre 1845.

§ *Vierteljahrsschrift von Roser und Wunderlich*, Jahrgang 1845, Heft 3; and *Medicinische Zeitung des Vereins für Heilkunde in Preussen*, Jahrgang 1845, No. 45.

|| *Recherches sur les Maladies des Ouvriers employés à la fabrication des Allumettes chimiques, &c.* Mémoire présenté à l'Académie des Sciences, le 16 février 1846.

¶ *Comptes-rendus des séances de l'Académie royale des Sciences de Paris*, mai 1846.

** *Ueber den Einfluss der Phosphorzündholzfabrication auf die Gesundheit der Arbeiter*. Mitgetheilt von Dr. Ebel. In *Casper's Wochenschrift*, 16 März, 1851.

†† *Die Krankheiten der Arbeiter in den Phosphorzündholzfabriken* &c. Erlangen, 1847; also *Die Regeneration des Unterkiefers nach totaler Necrose durch Phosphordämpfe*, von L. Geist, Erlangen, 1852. An admirable digest of this subject has been published, in the shape of a review of Von Bibra and Geist's work, in the *British and Foreign Medico-Chirurgical Review* for April 1848.

to the Privy Council 'On the Relation of Phosphorus manufactures to the Question of Public Health.'*

cause of the maxillary necrosis and exfoliation occurring; the artisans employed in making lucifer-matches is of the phosphorus, need not be discussed; the question already settled; and the reader is referred to the authors above named, especially Von Bibra and the consideration of this point. The only question which deserved serious inquiry arose from the fact, phosphorus employed was often impure, and contained quantity of arsenic; and this Martius and Dupasquier might be the essential cause of the malady: but this was not found to hold good under more critical and examination.† It is only when the cause has been

Manufactories in which Phosphorus is Produced or Employed,' in *Report of the Medical Officer of the Privy Council*, p. 162. London, 1863. I state that my article on this subject in the first edition of this work appeared before the publication of Dr. Bristowe's Report, though from the issue of the fourth volume it did not appear till a year afterwards. I was furnished with a copy of my paper, from which indeed he has drawn his conclusions.

an agent in producing the phosphorus disease is doubtless oxigen, probably in the form of phosphorous or phosphoric acid, dissolved in saliva. The precise nature of the fumes is not exactly known; they consist mainly of phosphorous acid (H_3PO_3), which by mixture with phosphoric acid (H_3PO_4); and probably minute quantities of phosphuretted hydrogen (H_3P), and hypophosphorous are also present. Any vapour of phosphorus and hypophosphorous is speedily converted by the air into phosphorous and phosphoric acid. Phosphuretted hydrogen would be more slowly oxidised into the same at any rate, nearly if not quite the whole of the fumes when inhaled by the people would be in a state of some acid of phosphorus capable of being neutralised by an alkali. Such being the case, it is greatly to be regretted that efficient sanitary measures are not adopted to prevent the disease, surely might be done. The recommendations of Geist and others are simply to cleanliness and ventilation, and are not sufficient. It is suggested, that in all lucifer-match manufactories there should be a rigid scrutiny of the mouths of all the work-people employed. Faulty teeth should be excluded from the rooms (the *dipping* and *dipping* where the obnoxious fumes are being developed. All carious teeth should be extracted or plugged. What is more important—a very simple respirator for the mouth might be worn by the employés; it would cover the nostrils. It should be constructed on the ordinary plan but its centre consisting of a porous diaphragm, such as sponge or cloth, linen or cotton, which should be daily dipped in a solution of fixed alkalies or their carbonates. This would arrest nearly, if not the acid fumes of the phosphorus. Or the respirator devised by

long in operation, and under circumstances of great intensity, that the disease is developed; it is confined almost entirely to those who are employed in the process of *dipping* the matches into the fused and reeking phosphorus compound, and those who dry them in the same apartment. Moreover, a long exposure to the influence is necessary for the production of the disease. Lucifer-manufactories had existed eleven years in Vienna, when in 1839 Lorinser observed the first case which attracted attention; and the shortest known period in which the malady has developed itself was in an instance which occurred in Paris, and is recorded by Gendrin, in which the patient had been employed in the occupation two years before the first symptoms appeared. Another circumstance of curious import, though not invalidating the generally-received doctrine of local poisoning, is, that the sufferer may have been removed from the baneful influence for a considerable period, and nevertheless be subsequently attacked by the disease.

Strohl gives an instance of a girl who had worked as a *dipper* at a lucifer-match manufactory for five years; she left that employment, and adopted a totally different and healthy occupation, when, after three months, she was, for the first time, attacked with unmistakable symptoms of the phosphorus disease, ending after fifteen months, in exfoliation of portions of the superior maxilla.

It is very seldom that others than those employed in making lucifer-matches suffer from this malady; but that this may happen should not be lost sight of.

Pluskal* mentions an example of a little girl, seven years of age, who was in

Mr. Graham for persons exposed to carbonic acid vapour would probably be as efficacious. It consists of the mixture in equal bulk of fresh-slacked lime and sulphate of soda, through a cushion of which it is easy to breathe. The wearing of some such respirator should be compulsory with the dippers and dryers. Again, the acid vapour might be neutralised and rendered innocuous by keeping the atmosphere of the apartment ammonuretted. I believe, if these precautions were adopted, the disease would seldom, if ever, manifest itself.

There is a curious modification of phosphorus, known as 'amorphous' phosphorus, which does not emit noxious fumes when heated sufficiently for its employment in match-making. This kind, however, is not mixed with chlorate of potass (which furnishes the oxygen) for tipping the matches, but is employed to coat the rubber on the surface of the box against which, by friction, the chlorate of potass match is ignited. These matches deserve universal adoption; for not only are they made without possible injury to the work-people; but they are quite safe, being incapable of accidental ignition.

* *Oesterreichische medicinische Wochenschrift*, No. 30. Wien, den 25 Juli 1846.

the constant habit of playing with matches, standing before a wall and discharging them in the dark for amusement, so that her face was bathed in their fumes; in time she was attacked with necrosis and exfoliation of small portions of the front of the lower jaw, with the ordinary attendant symptoms. Simon narrates a case in which the disease appears to have been brought on by a person chewing pieces of ginger, which he kept in his pocket with some lucifer-matches. And Mr. Paget describes* a case in which the malady was induced by the manufacture of medicines in which phosphorus was employed.

The influence of the phosphorus fumes upon the jaw is undoubtedly local. Lorinser, who has the merit of discovering this disease, held a different opinion; but Roussel, Geist, and indeed subsequent writers generally, have adopted the opinion, which all the evidence upon this point seems clearly to establish.† I have not space here to enter upon the argument of the question; I may, however, mention one telling fact; it is, the necessity of dental caries in the individual before the disease can be produced. It has never been known to occur, excepting where the sufferer has had carious teeth; and many persons have worked in the manufactories for a long series of years with perfect impunity, who, upon the supervention of dental caries, have been attacked with the malady. Dental caries, by opening the central chamber of the tooth and exposing the pulp, seems to offer a tolerably direct channel for the poison to be communicated to the subjacent periosteum, and this, no doubt, is why tooth-destruction is a necessary pre-existing condition. What the precise nature of the action of phosphorus-oxide thus absorbed may be upon the bone, is a matter of speculation; but the particular nature of the poison, entering as it does so largely into the composition of the skeleton, is a suggestive circumstance: perhaps, if accumulated by the periosteum, it may generate on the bone's surface a condition of chemical *superphosphate*, inconsistent with osteal vitality.

The *symptoms* of phosphorus-necrosis do not differ essentially from other forms of necrosis in the same parts: they are, however, not infrequently accompanied by bronchial and pulmonary irritation from inhalation of the fumes: this has been especially pointed out by Sédillot, Gendrin, and Dupasquier.‡ And one

* *Medical Times and Gazette*, vol. i. p. 41, 1862.

† The arguments bearing upon this question, and the conclusive inference to which they lead, are stated with cogent force in an admirable clinical lecture on this subject by Mr. Simon, in the *Lancet* for 1850, p. 41.

‡ *Gaz. méd. de Paris*, 1846, No. 49.

of the patients whom I have seen, affected with this malady, has detailed to me symptoms of *spasmodic asthma* which occasionally supervened when he was employed for many continuous hours 'dipping' while suffering from a common 'cold' in winter weather. Barring this occasional manifestation of pulmonary irritation, the general health of these work-people is remarkably good.

The symptoms of the jaw disease usually commence with what is supposed to be toothache, the pain being at first pretty much localised to some one tooth that is carious, and which is probably the channel by which the poison is introduced. The advance of the disease is generally slow at first, and, as it were, undecided—indeed, indefinitely chronic; the pain is inconstant, and not early attended with more serious symptoms; presently, however, it becomes more severe and erratic, extending vaguely about the side of the head and down towards the shoulder, and with this severer pain swelling and extreme tenderness occur; the integument near the affected region becomes red, tense, and distended, while the teeth feel elongated and intensely painful when brought in contact with their fellows of the opposite jaw, and they become very loose. The gums are swollen and livid, and this condition extends to the mucous membrane of the cheek. All these symptoms increase till suppuration is established, and with them, more or less, symptomatic fever is developed in proportion to the severity and extent of the disease: the patient has rigors and pyrexia, and is often thoroughly ill. The point at which the pus finds its discharge varies a good deal: the soft parts first become very boggy, especially the gums, and matter often escapes early around the necks of the loose dead teeth; when pointing externally, its approach is accompanied by intense glistening erysipelas-like redness of the integument. The discharge of the pus is attended with great mitigation of the patient's suffering. The pus itself is often sanious at first, and very fetid, having the odour characteristic of the presence of necrosed bone. The orifice of discharge frequently leads to long burrowing sinuses, especially where the lower jaw is affected; and through these the dead bone may be detected by a probe. The swelling which attends the disease is often very great, particularly when the lower jaw is necrosed: it is diffuse and wide-spread, encasing the external and under surface of the bone in a prodigious, dense, plastic exudation. In very severe cases, previous to the discharge of pus, whi

the inflammatory symptoms are at their extreme height, the whole head, except the summit of the scalp, is involved—the eyes are closed; the nose, and even the forehead, swollen; the cheeks, lips, neck, and throat, are one continuous area of florid intumescence. It is a curious circumstance that in the lower jaw the necrosis is attended with a very large and complete development of ossifying callus, whereas none is formed when the upper jaw is affected. The large plastic exudation which surrounds the base of the lower jaw becomes converted into a mass of supplemental bone, supporting the sequestrum, which is, for the most part, naked and bare within and behind, and connecting sound portions of bone at its extremities when the whole of the maxillary arch is not involved. Geist assumes that the particular region occupied by the osteophytic incrustation about the lower jaw is the result of gravitation—an idea altogether too mechanical: it is rather to be looked upon as a physiological manifestation, and one element in that marvellous exhibition of the *vis medicatrix nature* which this repair of the lower jaw displays: the supplemental bone thus placed forms the best support for the sequestrum, and the least interferes with the functions of the mouth; while the absence of an ossifying callus at the upper and inner region of the maxillary arch offers the readiest escape of the dead bone through the thin mucous membrane which there alone covers it, and this without damaging the integument of the face.

As the bone becomes laid bare by the ulceration of the soft parts, it is observed bathed in ichorous pus, ragged and irregular in its surface, and of a dirty blackish grey colour: this latter condition is said to be invariable, as though characteristic; but it is not so, for although very general, I have seen specimens white and clean, and with the compact external layer of bone intact.

To return to the symptoms of the disease. When the extent of necrosis is very great, the constitutional disturbance is correspondingly severe; and in the early stages of the malady the patient may have intense fever, with delirium and agonising local suffering, the more distressing from the region which the necrosis occupies—interfering with or altogether suspending action of the mouth, and, by the secretion of foul and fetid matter, producing nausea, ructus, vomiting. The looseness and separation of the dead teeth is another source of annoyance and distress: as the sequestra containing them emerge from the

surrounding parts, their elongation and angularity much irritate the gums and cheek in contact with them. In the severest cases, general sphacelus of the soft parts about the jaw, with œdema of the face and neck, may supervene, accompanied, or not, by erysipelas; and death may then close a scene of terrible suffering. In other instances with a fatal issue, life is drawn out through many months of tedious illness, varied by different degrees of local irritation; till at length the patient, with vital endurance inadequate to the requirements of the disease, sinks tabid and exhausted, under continuous hectic, and not infrequently with tubercular complications.

But the pathological changes may be more terrible than those already mentioned.

Dr. Ebel describes a case which occurred in the practice of Dr. Hervier at the Hospital *Necker*, in which the patient, a lucifer-match maker, first had necrosis of the lower jaw, then of the upper, afterwards of the palate bones and the orbits, and lastly of the os frontis. He died with brain symptoms. Pus was found between the dura mater and the brain.

When the malady progresses to a favourable issue, which does in the majority of cases, the dead bone gradually loosens and becomes detached; and this is generally anticipated by the falling out of some of the necrosed teeth. In the upper jaw the sequestra are usually more broken up and smaller than in the lower, and they are shed easier, not being held in and detained by ossifying callus, which is always wanting in the upper jaw. In the lower jaw the sequestra are usually more extensive often including large portions of the body of the bone, not infrequently the ascending rami, and sometimes the coronoid processes, and even the articular condyles. The shedding of these sequestra is often hindered by the large surrounding ossifying callus which always forms when the lower jaw is the subject of this disease.

I may here mention that some stress has been laid upon the circumstance that the lower jaw has been more frequently observed to be affected than the upper. The learned reviewer of Geist and Von Bibra's work has given much detail bearing on this point; he enumerates twenty-five examples in which the lower jaw was affected, to twenty-one in which it was confined to the upper jaw. In five cases which I have seen, the lower jaw was diseased in four, and the upper in one; where

four, which occurred in the practice of a surgical friend, were confined to the upper jaw. In seventeen instances of which I have obtained particulars or seen specimens, nine were connected with the superior, and eight with the inferior maxilla. Dr. Bristowe in his investigations, met with evidence of thirty-nine cases in the lower jaw, twelve in the upper, and five in both.

The results of phosphorus-necrosis, in cases which get well, are very various. As regards general health, the patient usually recovers with an elastic convalescence after the dead bone has been removed; but the physical condition of his maxillary apparatus is often terribly mutilated. When the front part of the upper jaw is affected, it is an absolute and unrepaired loss, miserably disfiguring the patient, altering the physiognomy most painfully; when it affects more hidden parts of the jaw, the loss is not less complete, though less conspicuous.

If the affection is confined to the inferior maxilla, the removal of the sequestrum leaves a supplemental bony representative, which, for a time, more than makes up for the loss of dead bone, and for a long subsequent period efficiently performs the office of edentulous mastication, and supplies the wants of personal comeliness.

In some instances the whole of the lower jaw has been lost, excepting the articular condyles; and these, apparently connected with the new bone, have established the joint requirements of the supplemental jaw; but in other instances (cases in St. Bartholomew's Hospital, and those mentioned in Geist's later work*) even the articular ends have been shed with the rest of the sequestrum; still there has been joint movement, doubtless from ligamentous attachment of the new bone.

This repair of the lower jaw is, however, generally but temporary: for after a time—often a considerable time—the new bone diminishes, by absorption, to a mere narrow arch, and ultimately, there is scarcely enough bone to keep out the lower lip, and the chin is utterly lost. I have had an opportunity of examining this state of parts after the lower jaw had been removed ten years. Dr. Bristowe, however, mentions two instances, one after six, and one after ten years, in which the supplementary bone remained large and well formed: he describes another in which, after eight years, the new was scarcely bigger than the hyoid bone. How far the

* *Die Regeneration des Unterkiefers nach totaler Necrose*, v. L. Geist, 1852.

loss by absorption of supplemental bone, may be prevented by supplying it with a function, through the means of artificial teeth, is a question of theoretical interest and of practical importance.

The *treatment* of phosphorus-necrosis of the maxilla divides itself into that which is common to all bone-necrosis, and that which specially applies to the particular local affection in question. Upon the latter head alone I would remark.

In the early stages of the disease, when as yet it is not established, but its supervention feared, it would be of the last importance to remove the patient from the cause of the malady: pure air should be sought, abundant cleanliness, with urinary and alvine excretants, resorted to, and all suspicious teeth extracted. When, however, the local symptoms—extreme pain, swelling, and indurated infiltration of the soft parts—develop themselves, and the disease has already passed its premonitory stage, it will be advisable to have recourse to more active measures. Upon this point I cannot do better than quote Mr. Simon's remarks in his admirable Clinical Lecture, already referred to, and in the justice and propriety of which I entirely concur: "So soon as the second or inflammatory stage of the disease has thoroughly set in, the bone seems in every case to be irrevocably doomed to necrosis; and I would therefore recommend you, in the event of your being called to a case at the transition period between the two stages, when hypertrophy is passing into inflammation, to adopt without hesitation the most active measures for the relief of the periosteum and bone. Leeches and general antiphlogistic treatment *may* do good; but the consideration of the pathology of this disease, together with the analogy of other periosteal affections, leads me to believe that the only real chance of doing good would lie in still more energetic measures; and I would recommend you in any such instance, to make with your scalpel free vertical incisions through the gum wherever tenderness and swelling exist; extending your line of cut upwards in the upper jaw, or downwards in the lower, as far as the structure of the parts will allow, bringing your incisions as near together as circumstances may require, and in every point carrying them clearly down to the bone, so as to afford the utmost relief and relaxation to the overloaded and tense periosteum. I believe that this method of procedure would be the nearest approach to an effective one for checking the inflammatory stage of the disease before it

has reached an intensity which must inevitably destroy the jaw.*

When the stage of threatening has past, the extent of the inflammatory mischief appears to assert itself at once—the whole of that portion of bone which is subsequently the sequestrum appears to be stricken from the first. Prevention is at an end, and palliation rather than cure is indicated as the surgeon's mission. When pus has formed, it should be early conducted to the surface, and by judicious interference external scars may often be prevented by means of well-directed punctures within the mouth. The whole of the teeth implanted in the dead bone become loose, and apparently elongated; their crown is already sealed, and they should be extracted without delay, as they cause much irritation to the tongue and cheeks.

When the discharge has established itself, and the fetid pus is pouring from the sinuses that communicate with the dead bone, abundant rinsing of the mouth should be had recourse to, not only in the form of mere washes with water to remove the filthy secretions, but astringent and deodorising lotions, such as decoction of cinchona, and solution of alum, solution of chloride of lime, and, still better, solution of permanganate of potass. It has been pointed out by Mr. Simon, the unhappy patients are sometimes so disabled as to be incapable of gargling out their mouths, and then little pieces of sponge may be used by them to wipe away the stinking discharges.

As regards the sequestra little can be done. Sharp projecting pieces of bone may be cut off by nippers; but the bulk of the dead bone must take its own time to be shed—often a very long time. It is a period of miserable suffering to the patient, and terrible constitutional exhaustion; but it cannot be shortened or its results anticipated.

In March 1862, I saw in the London Hospital, under the care of Mr. Adams, a patient who had been suffering from the disease, affecting the entire inferior maxilla; it had been going on two years and a half; the latter four-fifths of the time with the lower jaw, except probably the articular ends, dead and lying in the mouth, bare and ghastly, bathed in ichorous pus within its huge supplemental successor.*

* The necrosed jaw was afterwards removed. The operation and its sequence were both unusual. The patient was placed in a kneeling posture, with the chin on a table, and, the mouth being open, the symphysis of the jaw was split down by a chisel and mallet. The right half of the bone was then seized by large strong forceps, and wrenched from the mouth with great force. After a

The sequestra of the superior maxilla are apt to be more detached than those of the inferior—divisible and smaller, and they can be brought away piecemeal; moreover, as the upper jaw is an attached, and not like the lower, a floating bone (with muscles ready to displace its parts when the integrity of its arch is broken), its dead portions can be removed without jeopardising the relations of the rest of the bone. Therefore a little and early surgical interference may be employed without injury, and parts brought away which if belonging to the lower jaw, had better not be interfered with. In the lower jaw the disease is usually so much more comprehensive, and its resultant sequestrum is so solid, that patient watching and palliative treatment are all that can be adopted generally till the very close of the case. Earlier interference might disturb the relations of the supplemental bone (the *natural splint*) and the portions of jaw still living, and by allowing the strong antagonistic action of the muscles attached to the two halves of the latter to act separately, lead to the permanent displacement of the elements of the maxillary arch. The articular ends of the bone often—indeed, in the majority of cases, though not always—escape the necrosis, when the whole of the rest of the bone may be destroyed; and this circumstance suggests an element of treatment which may much shorten the patient's suffering. When the necrosis has evidently passed up to the neck of the condyles, the bone may be removed by sawing or nipping across the neck as high up as possible (on either side, if both sides are affected), and then dividing the body of the jaw at its symphysis; the two halves of the bone may be withdrawn, one by one, without any external wound. Should the condyles be necrosed and follow, no harm will have been done by the previous operation, and the patient's suffering will have been much diminished and curtailed. The division of the bone may be accomplished by a Hey's saw, or some modification of it improvised to suit the particular position of the required line of section. Various curved bone-nippers may also be devised to

week the same course was adopted with the left half; but this required still greater traction. The entire bone was thus brought away, including the condyles. Ten days after the first operation the patient was seized with secondary hæmorrhage, profuse and arterial, on the *right* side from the wound, where the first portion of jaw was removed. This necessitated ligature of the right common carotid artery. The latter operation was performed by Mr. Maunder, and the patient recovered perfectly. See *Med. Times and Gazette*, July 5, 1882.

accomplish the same objects. The extreme and firm closure of the mouth sometimes renders this operation very difficult.

I need not dwell upon the general treatment of these cases as far as they are in common with those of bone-necrosis generally. The particular region affected, however, involving and disabling, as it does, the masticatory function, entails some modifications. In the long tedium of their course every effort should be made to supply, by suitable food, the deficiencies which suspended mastication entails—mashed meat, eggs, cod-liver oil, and such sustainants, should be abundantly employed; and iron salts, in moderate proportion, should, if bearable by the patient, be almost a matter of daily diet rather than an occasional medicine.

There is one point which I would allude to—more a matter of pathological curiosity, than having any other bearing; it is the enormous amount of pus which these patients daily swallow and (apparently) digest. It must be many ounces, and this without any obvious detriment to health; the patient's condition being no worse than may be accounted for by the force of the local symptoms.

Exanthematous jaw-necrosis.—I believe that the necrosis and exfoliation of the alveolar processes and portions of the jaws in children, consequent upon the eruptive fevers, is essentially the same as the maxillary necrosis in the victims of phosphorus fumes, and that it is the result of the local application of a specific poison to the vascular parts of teeth. There is this difference, however, that in the cases we are now considering the poison is generated within the individual, but with affinities for certain structures and tendencies to action upon certain organs which give its morbid consequences an equally local character. Whatever opinions may be entertained as to the homological relations of the several tissues of the teeth to those of the general integument, there can be no question as to their being members of the dermal system, and as such we should *à priori* expect that they would share the consequences which attend those particular diseases which spend their chief force on the skin. There is one circumstance, however, that modifies such an anticipation; it is the low state of vitality of the teeth, and the extremely slight nutritional changes which occur in them *when once they are formed*, and which must consequently remove them to a great extent from those transient though potent influences which would destroy or morbidly

affect vascular or rapid-growing tissues. But such a qualification does not apply to the conditions of the teeth during their development. From the time of birth till the eighth or ninth year, the jaw-bones are the seat of intense developmental nutrition in the formation of the teeth, and are among the most vascular parts of the body: about the middle of the period named, five years of age, the maxillæ contain no less than forty-eight developed teeth and developing tooth-germs. It is about this time that the poison of the exanthematous fevers appears to exert its most deadly influence on the dental system.

The form of necrosis affecting the alveolar edges of the maxillæ, and accompanied by the shedding of the teeth, which we are now considering, was, I believe, first recognised by myself* as one of the sequelæ of the exanthemata, and dependent necessarily on their previous occurrence. Many isolated cases of the affection have been described, and our museums contain specimens of the sequestra; but the supposed relation of cause and effect had not, so far as I know, been expressed before.†

This affection is by no means common, considering the almost universal occurrence of the eruptive fevers. The very large population tributary to Guy's Hospital has only furnished me with twenty-three or twenty-four cases during the last nine years; and I have reason to think that even in this I have been disproportionately favoured.

The cases are all singularly alike. A little child has just recovered from one of the eruptive fevers, most probably scarlatina; the case has been in no way unusual as to its severity or its course: within six weeks or two months of the passing off of the acute symptoms, tenderness of the mouth is complained of, and the mother notices foetor of the breath. Upon inspecting the mouth, the gum is seen to be peeling from the edge of the jaw around the neck or necks of some temporary tooth or teeth; pus is discharging, and more or less dead bone is exposed. The

* 'On the Shedding of the Teeth, and Exfoliation of the Alveolar Processes, consequent upon the Eruptive Fevers,' by S. J. A. Salter, *Guy's Hospital Reports*, 3rd series, vol. iv. Several specimens of sequestra from these cases have been figured by the author in the *Transactions of the Pathological Society*, vol. xi. pp. 209-215.


† The earliest recorded cases are mentioned by Fox in his *History and Treatment of the Diseases of the Teeth*, p. 112. London, 1806. These cases, two in number, occurred after small-pox.

modation of the bone progresses rather quickly in depth, but usually not after the first in lateral extent; the temporary teeth at the affected part become loose, and often fall out. There is no swelling; and no ossifying callus is formed in the region of the necrosed bone. In a few weeks from the first of these symptoms, the sequestrum itself becomes loose, and is easily removed, leaving a large gap, and a raw granulating surface, which rapidly heals. The necrosis almost always includes the bone which constitutes the loculi containing the developing permanent teeth, as well as the alveoli of the temporary; but it does not go further, and in the lower jaw the base of the bone is very rarely affected. I have never seen such an occurrence. The disease is frequently symmetrical: when attacking the bone about the temporary molars on one side of the jaw (its most common situation) it often manifests itself immediately after on the opposite side, and sometimes in the same regions of the other jaw. The same symmetry is observed at the front of the mouth.

As far as I am aware, this affection only occurs after the eruptive fevers,* and scarlatina is its most potent cause. In the instances which have been under my own care, two were after small-pox, five or six after measles, and fifteen or sixteen after scarlet fever. There is, however, nothing in the condition of the mouth to indicate which has been the precursor. The age at which this affection occurs is usually about five years: from three to eight are the extreme limits I have known.

The issue of these cases is simply comprised in the loss of a certain amount of bone with the contained teeth, and the consequent disfigurement. As the permanent teeth are lost with the temporary, the disfigurement is very great when it affects the incisors; but when the temporary molars and their

* I have heard of one doubtful case after continued fever (typhus or phoid); but this would form no real exception to these cases or their pathological interpretation. The continued fevers have their specific cutaneous eruptions; and their damaging influence on the tegumentary system is sufficiently attested by the falling of the hair in convalescents from them. I would not, however, be supposed to deny that jaw-necrosis, associated or not with death and exfoliation of other bones, may occur in children as the result only of extreme cachexia or depressed vital nutrition. Such instances are referred to in vol. iii. p. 762. I merely wish to assert my belief that the particular form of alveolar necrosis above described is one of the *specific sequelae* of the eruptive fevers, and is related to its cause in the manner indicated in the previous pages.



the time when the bone and temporary tooth were in effect equally destroyed, for the tooth was but not grow after that time; no fang was formed, it remained a mere tooth-crown, loosely attached of the gum.

The *treatment* of these cases involves little but and cleanliness. No operative interference is : extent of the necrosis asserts itself from the first be curtailed. Soon the sequestrum becomes loose then be readily removed by dressing-forceps. I use a weak solution of *permanganate of potass*, as a deodorising mouth-wash. Any general symptoms of health should be met on ordinary principles. If suppuration is profuse, and the bone-necrosis extensive, stimulants and tonics may be required; and in the severe cases steel would be a useful adjunct; but I would observe that in the cases I have seen, the children were, for the most part, remarkably good health, and had, almost without exception, escaped the other ill consequences of the eruptive

IX. *Hæmorrhage after extraction of teeth.*—I have seen and obstinate bleeding after the extraction of teeth is a serious and troublesome complication of this operation, not a common consequence: indeed it is very rare in the enormous numbers of the operations, and in the proportion continued bleeding follows them. The hæmorrhage from the wound of an extracted

se patients have themselves been aware of the tendency, having experienced similar consequences attendant on the extraction of other teeth. Indeed, some persons (I know some such patients of my own) refuse to have any more removed, preferring rather to suffer any amount of tooth-ache on account of the alarming hæmorrhages which have attended the operations before.

In one case which came under my treatment, there was no history of a hæmorrhagic tendency; but the patient was a young lady, and the circumstance was just as menstruation was imminent. The bleeding was very considerable and continued for two days, when under the influence of emmenagogues (warm baths and aloetic purgatives), the catamenia appeared, and the hæmorrhage from the alveolus immediately ceased. In this case, doubtless, the bleeding was a vicarious manifestation of the periodic function. A very interesting case of this distressing consequence of tooth-drawing, showing its constitutional nature, was under my care some years since. A clergyman from one of the inland counties came to town to me on account of the hæmorrhage which followed the extraction of a lower molar tooth three or four days before it was still continuing. He was reduced to a pitiable condition of exhaustion and weakness. All the local means usually adopted had been tried, but without success: the bleeding, however, speedily ceased under the use of large and frequently-repeated doses of *tannin*. The history of this gentleman and his case is curious and instructive. As a boy, from the most trivial accidents, such as a cut finger, he sustained prolonged and almost irrepressible bleeding: for his whole life, the slightest blow or bruise would cause a large and very profuse ecchymosis; a few years before, he had sustained a week of hæmorrhage, after the extraction of a tooth. This gentleman has three married children, all of whom suffer from terrible floodings at the birth of each child. He

adopted, is recorded in an early number of the *Medico-Chirurgical Transactions*. The previous history of this patient is singularly like that of the clergyman whose case I have just mentioned, in the hæmorrhagic diathesis manifested itself in childhood, and his having sustained a terrible loss of blood—the bleeding lasting for twenty-one days—when another tooth was extracted. In the treatment of this fatal hæmorrhage, the actual cautery produced only temporary arrest of bleeding; it was then determined to tie the carotid artery. The operation was performed by Sir B. Brodie, without any relief to the original symptoms; the wound in the neck only furnished another bleeding artery, and evidently hurried on the fatal catastrophe. In another recorded case where the cautery was used, the edge of the lip was accidentally burnt, and the surface thus cauterised soon commenced bleeding profusely, and continued to do so till the fatal termination of the case, which was hastened by it.

The *treatment* of this form of hæmorrhage divides itself into local and general; the local treatment consisting in the application of styptics, with continuous pressure upon the bleeding surface; the general treatment, in the rapid and abundant administration of internal astringents. The local plan of plugging with a compress, recommended by Hunter, is that universally adopted: 'In general it will be sufficient to stuff the socket with lint, or lint dipped in oil of turpentine, and to apply a compress of lint, or a piece of cork thicker than the bodies of the adjacent teeth, so that the teeth in the opposite jaw may be brought up a pressure. It has been advised to stuff into the socket some soft wax, on a supposition that it would mould itself to the cavity, and so stop the bleeding: this, perhaps, may sometimes answer better than the other method, and therefore should be tried when that fails.'† The restoration of the extracted tooth to the socket has also been tried with success. For the purpose of forming a plug of lint, a strip should be cut of an elongated wedge-shape, and this should be introduced, the pointed end first; the extremity should be introduced into the bleeding socket, and driven well home to the bottom; the strip of lint should then be folded and re-folded upon itself, so as to make the plug very solid and pressing on the entire superficies of the socket. When the compress of lint or cork has been applied, the mouth should be closed and the jaws kept permanently together, so as to exert considerable and persistent pressure by means of a broad bandage tied under the chin and over the head. Where the opposite

* 'A Case of Fatal Hæmorrhage from Extraction of a Tooth,' by Richard Blagden, *Med.-Chir. Trans.* vol. viii. 1820.

† *A Practical Treatise on the Diseases of the Teeth*, by John Hunter, p. 40. London, 1778.

less in the mouth, and the other to the exterior: where the
ing is in the lower jaw, the blades are nearly equal, the
ior one simply passing under the base of the horizontal
s: for the upper jaw the instrument is less applicable, and
uter blade requires a long curved sweep, so as to grasp the
f the head. Various local astringents have been recom-
ed in conjunction with the lint-plug: matico has been
y spoken of: I have found an alcoholic saturated solution
anin answer the purpose very well; but probably the old
dy, turpentine, has never really been superseded.

m convinced, however, that it is a great mistake only to
these cases topically: they are most obviously manifesta-
of constitutional vice, and require a corresponding general
ment. Astringents should be had recourse to at the earliest
; they can do no harm, and may be suspended at any time,
in the majority of cases they will be most beneficial. These
often extend over so long a period that constitutional
nent has abundant time to be brought into full operation.
in and turpentine have been the most successful of internal
lies in recorded cases; with the latter drug, steel might
lvantageously combined, especially in the form of the
te tincture. In the irritability of sanguineous exhaustion,
as may become necessary in large doses.

er circumstances require attention: the patient should
y a cool apartment, and local icy applications may be
his might nature is to be performed, or if the

suppressed menstruation, and constitute a vicarious discharge, and remedies should be adopted to meet this contingency.*

X. *The application of obturators and false palates in palate, &c.*—The title of this section is scarcely in keeping with the heading of my essay—‘Surgical Diseases connected with the Teeth,’ &c. The maladies which give rise to perforate and clefts of the palate are, practically speaking, very dependent on diseases of the teeth; but the particular requirements of their resultant defects, and the nature of the mechanical dentist’s art, as remedying those defects, bring this fairly within my province.

The two forms of cleft or perforate palate, congenitally induced, are very unequally suited to this particular treatment: a large number of the former may be remedied by means of surgical operation, which is always practicable where practicable; in the latter cases, the mechanical assistance of obturators is of more avail, and, in many instances, not only removes an existing defect for the time being, but is to a great extent curative.

In congenital cases, when *very* severe, this is the only operation applicable for the treatment of the great physical defect, the operation being capable of bringing into apposition the soft masses on either side of the cleft, or of remedying the defect of soft parts, which consequently require to be supplied with and foreign material.

When the cleft or perforation consists of an aperture in the hard palate only, as in most syphilitic and scrofulous cases, some congenital, an arch of metal or other hard substance spanning over the orifice, and firmly applied to the contiguous palatal surface, is all that is required. When, however, the cleft extends backwards and involves a destruction of the soft palate, and that to an amount which cannot be remedied by surgical operation, it becomes necessary to add to the metal arch an elastic flap (such as vellum or india-rubber), which supplies the mobile functions of the soft parts as they are in a normal condition. The former apparatus may be called *obturators*, the latter *false palates*. In not a few of these cases other portions of the upper jaw become implicated; the alveolar processes may be necrosed, involving the loss of the con-

* On the Hæmorrhagic Diathesis, see vol. i. pp. 718 et seq.

th and sometimes considerable masses of bone. To replace the losses various combinations of obturators, false palates, and artificial teeth have to be constructed, and they may be made in every variety, the different parts being associated together to supply lost parts and to fit by close adjustment those that remain. The materials of which obturators are made may vary; carved ivory, vulcanite (indurated sulphuret of caoutchouc), or flattened metal—gold, platinum, dental alloy, or silver—are all employed. The first, however, is now generally discarded, from its perishable nature. Vulcanite has been extensively adopted; but its thickness, and fragile character when very thin, militate against it in many cases. I much prefer sheet metal, on account of its strength, thinness, and durability. Silver, platinum, or dental alloy may be employed for hospital purposes; but gold is far better, and I am in the habit of using it, except in hospital cases. It is an object to have the sheet of metal as thin as possible without weakening it; and I employ the gold plate reduced to No. 6 of the gold-flatter's gauge, being somewhat thinner than is usual for the frames on which artificial teeth are mounted. This degree of tenuity is quite compatible with tire firmness, and it receives better all the inequalities of the gous palate: when a plate of this thickness is perfectly adapted to the palatal arch, it is scarcely appreciable to the wearer. As long as sound teeth remain in the upper jaw to which the plate may be attached, it should be supported by means of clasps or collars embracing them; when these are lost, the same object may be achieved by connecting the obturators by means of springs with a frame resting on the lower jaw. Under no circumstances whatever ought support to be obtained either by dilating sponge, or by hooks, or other processes pressing into the nasal cavity—the effect of which is to increase the size of the orifice, or prevent its natural tendency to contract. The * old and still very common method of treatment,

* Among the earliest surgical writings we find notices of mechanical appliances to repair lesions of the palate. The still common obturator, consisting of a metal disc supported by a sponge passing through the perforation into the nose, was invented by Ambrose Paré, an eminent French surgeon (*Les Œuvres d'Ambroise Paré*; à Lyon, 1541). These and analogous apparatus on the same objectionable plan, were alone employed until Bourdet devised others on correct principles (*Recherches et Observations sur toutes les parties de l'Art du Médecin*, par M. Bourdet; Paris, 1767). Bourdet employed thin sheet metal in the form of a vault wholly within the mouth, attached to the teeth at the sides, and free from any projection into the nose; and he was aware that

consisting of a disc of metal with a piece of sponge attached to its upper surface passing into the nose, cannot be sufficiently reprehended: it is filthy in its nature, and most injurious to the unfortunate patient's future condition, as the pressure of the expanding sponge causes progressive absorption of the margin of the orifice to an unlimited extent. Any plug kept in position by tight-fitting acts in the same manner. There is a preparation in the Museum of St. Bartholomew's Hospital (ser. L. no. 232), of the skull of a person who lost a portion of the palate probably by syphilis, and who obtained temporary relief by stuffing the oro-nasal aperture with a large cork, gradually adding to the size of the plug to meet the requirements of the ever-increasing orifice, till the whole of the hard palate had disappeared, the palatal processes of the ossa palati as well as those of the maxillæ being lost. The progressive absorption under this continually-advancing pressure had removed the vomer, the inferior turbinated bones, and the nasal wall of the maxillary sinuses. Such a specimen is an eloquent commentary on this reprehensible plan of treatment. I believe in no instance is it necessary to gain support by the passage of processes of any kind into the nasal cavity, either to wedge against the

when thus treated the orifice would contract and frequently close. M. F. Fauchard appears to have been the first person who combined artificial teeth with an obturator (*Le Chirurgien-Dentiste*; Paris, 1786). M. Touchard published an account of an apparatus not only to supply teeth and palate, but also a large mass of upper jaw which had exfoliated. This was effected with ivory (*Description d'un obturateur dentier présenté à la Société de Médecine de Paris*; Paris, 1814). M. Delabarre was the first person who succeeded in supplying soft as well as a hard palate. He accomplished the former with sheet caoutchouc, and have nearly all his successors (*Traité de la partie mécanique de l'art du chirurgien-dentiste*, par C. F. Delabarre; Paris, 1820). But the greatest advance in the construction of false palates was made by Mr. Snell, whose apparatus was essentially the same as those employed in the present day (*Observations on the History, Use, and Construction of Obturators, or Artificial Palates*, by James Snell; London, 1824). The same principles have been carried out with slight modifications by Mr. Stearns (*Lancet*, vol. ii. 1845, pp. 7, 260, 284, 310); and by Mr. Scrovide ('Cleft Palate: its Surgical and Mechanical Treatment,' *Trans. Odont. Soc.* vol. i. London, 1858). Dr. Kingsley of New York, has devised an ingenious arrangement which very firmly and effectually closes the cleft of the palate. It consists of a wedge-shaped plate of hard vulcanite with a soft velum, forced forward into the fissure, and resting by a ledge upon the nasal surface of the hard palate, thus reverting to the objectionable principle adopted previous to the time of Bourdet ('On the Treatment of Congenital Cleft Palate,' by Dr. N. W. Kingsley, in *Trans. Odont. Soc.* vol. v. p. 195; London, 1865).

rest of the orifice, or to rest upon the nasal surface of the plate.

But an obturator passing over the perforation is not negative in its effect. An orifice which, allowed to remain open, may continue *in statu quo*, or contract in a degree scarcely appreciable, will often rapidly diminish in size when the passage of air and fluid between the oral and nasal cavities is prevented. This is not generally known or understood; but it is of great importance in treating these cases. The obturator should be applied as early as possible—as soon as the dead bone has come away, and while the wound is in a state of granulating activity. It is surprising how the granulations will stretch across the upper surface of the obturator, and close up the orifice; and, in more advanced and neglected cases, the same condition may be imitated to some extent by frequently scarring the edges of the perforation, always taking care immediately to restore and keep in place a well-fitting obturator. Why the removal of the ill-effects of a perforate palate (passage of air and fluid from nose to mouth, and damaged voice), by means of mechanical appliance, should lessen or remove its cause, is not very apparent: it seems a curious transposition of events—an inverted sequence: perhaps the protection afforded by the plate to the granulating edge may to some extent account for it: it is, however, true, and very important practically.

Occasionally a portion of the floor of the antrum is lost by necrosis in cases of severe 'abscess' of that sinus, and its cavity becomes common with that of the mouth, by an aperture of variable size. This orifice should be immediately closed, or rather spanned over, by a plate (upon which substitutes for the lost teeth may, or may not, be fixed), and the result will be, in many instances, the development of a membranous growth supplying the lost bone, completely scarring over the wound, and forming a floor to the sinus.

The exact nature and arrangement of the artificial appliances we are considering will perhaps be best elucidated by illustrative cases.

The accompanying figure (Fig. 239) represents the upper jaw of a girl who has recently lost a portion of the hard palate from the effects of inherited syphilis. This cause is, I believe, unusual. I have not seen or heard of another case in which congenital syphilis has had such a result. The patient has suffered from characteristic keratitis, as well as from other

symptoms of the disease. The loss of the bone was attended with all the customary conditions of suppuration and fœtor, and after it came away a hole into the nose was left, as depicted in the woodcut. The patient's power of speech was seriously affected, and food and drink passed into the nose. A few days after the bone came away, I took a model of the

FIG. 239.



mouth, and an obturator plate was made as represented in Fig. 240. The girl being an hospital patient, the metal employed was 'dental alloy,' an alloy of silver and platinum, being less costly than gold. Support for the plate was obtained by collars passing between the bicuspid and first molars and the bicuspid and canines.

FIG. 240.



The application and fixing of the obturator *immediately* restored all the functions of the mouth: food was occluded from the nose, and articulation and enunciation were perfect.

This immediate restoration of the voice nearly always, I believe, occurs when *accidental* perforations of the palate are thus treated, and very soon, even where the soft palate is

involved in a cleft and an artificial velum has to be supplied. But with congenital cases it is very different; and the patient has in them to commence a new oral education.

I saw this case again about three months after the introduction of the obturator: the hole had contracted to less than half its original size, and will probably close altogether.

The apparatus needed to supply the defects of cleft palate is more complicated, but is scarcely less satisfactory in its ultimate results. The accompanying figure (Fig. 241) represents the upper jaw of a young gentleman, now about seventeen years of age. He was born with cleft palate and hare-lip. The latter was operated on in early infancy with success; and it was intended

FIG. 241.



that the cleft in the palate should ultimately have been closed by surgical operation; but the patient refused to submit to the pain. The cleft is complete from before backwards, passing to the left in the front of the mouth—an incisor being there wanting, as is frequently the case.

No difficulties were presented in adapting a hard and soft palate to occlude the cleft. A thin rigid gold plate was fitted to the hard palate, and support was obtained by clasps between the bicusps and the teeth on either side, and to prevent the dropping of the hinder part, which is frequently troublesome, fine thin clasps extended round the back molars. The velum palatæ was replaced by thin soft vulcanised caoutchouc, which I

always employ for this purpose. The accompanying illustrations (Figs. 242, 243) showing the artificial palate as seen in face on the oral surface and in profile, render any lengthened description unnecessary. In adapting a soft palate it is necessary that

FIG. 242.



the movable flap should be kept constantly in apposition with the pillars of the fauces during their movements in deglutition &c., and this is effected by means of an elastic spiral cord of gold wire, as seen in the profile illustration. This must be

FIG. 243.



of the lightest possible character, as the power of the muscles to move the flap is very feeble. The same object is sometimes effected by doubling the thickness of the caoutchouc near the hard palate.

When a false soft palate is first introduced, it often produces tickling of the fauces and retching; and it answers well to employ a very small flap at first, and gradually increase its size till it amply replaces the deficient velum.

When the flap is made of soft caoutchouc it becomes sodden

time and loses its elasticity; but it is easily replaced, and intelligent patient may do this for himself. To obviate this objection, Mr. Parkinson,* who has had much experience in treating these cases, employs a thin flap of vulcanite; but this being stiff can hardly apply itself so closely to the pillars of the sockets; still, in effect, it is said to answer well.

In cases where large portions of jaw are lost from disease or accident, admirable restorations may be made, especially by means of vulcanite, and there is scarcely any part of the mechanist's art which tells with more effect on the patient's comfort. It is scarcely possible to put a limit to the variety and extent which these useful apparatus may reach.

S. JAMES A. SALTER.

* 'On the Adaptation of Artificial Palates,' *Lancet*, vol. i. 1867, p. 41.

DISEASES OF THE MOUTH, PHARYNX, AND ŒSOPHAGUS.

TONSILLITIS. *Quinsy.* *In the tonsillaris.*—The most common morbid affection of the tonsils may be thus classed:

- 1st. Acute inflammation and abscess.
- 2nd. Chronic inflammation and enlargement.
- 3rd. Ulceration and discharging.

1. *Acute inflammation* of the tonsils is generally preceded or attended by a distinct rigor or chilliness, and general feverish disturbance. Accompanying or following such constitutional irritation, will be found pain in the fauces, and tenderness of the upper part of the throat, in every effort at deglutition. The pain at first is not acute, but rather as if the parts had been bruised, and is diffused over the back part of the mouth and the pharynx. As the case progresses, the local mischief becomes more evident. At first a mere blush, the redness of the part becomes more marked. With the altered colour of the mucous membrane, tumidity of the tonsil will be observed, and before long oedema of the surrounding soft structures. With the increase in the severity of these local changes, it usually becomes evident that the mischief is confined to one side. But the evils of tonsillitis are not restricted to these symptoms alone, nor do the sufferings usually stop short here. Pains in the jaws, and headache; fever and loss of appetite; often severe suffering in the part affected; a loaded, creamy, and swollen tongue, with very foul breath; muffled articulation; great difficulty in swallowing, and fluids attempted to be swallowed escaping through the nostrils; great fear of strangulation; the patient frequently obliged to sit forward, or lie on one side, to allow the abundant saliva to drip into a basin, or drain away on a sponge:—these are the symptoms which

mark the progress of the complaint, and comprise the chief characteristics of the attack. As the attack advances, the local examination of the mouth detects an increase of swelling, which usually runs forward in the substance of the soft palate, and on the side affected; in aggravated cases even implicating the tissues which cover the hard palate as far forwards as the incisor teeth. With the increase of swelling, a central softening spot will sooner or later be detected, either with the eye, or more frequently at a comparatively earlier period, by an examination with the finger.

Much difference will be found, in different cases, as to the period at which pus is formed in an attack of acute inflammation of the tonsils, *i.e.* as to the period when the abscess may be relieved by incision. In some instances from forty-eight to sixty hours will suffice to insure such a desirable condition. In other cases, many days will elapse before matter can be detected by the point of the finger, or upon the thrust of a scalpel. During all this time the patient passes his days and nights in misery; is not able to take any solid food, and but little fluid nourishment; obtains very little sleep, and that little of a disturbed character; has difficulty in respiration, caused by some oedema of the glosso-epiglottidean and aryteno-epiglottidean folds, and often greatly aggravated by fear, and a nervous horror of approaching suffocation. Inflammation in the acute form attacking the tonsil seldom terminates in resolution: as a general rule, it runs into abscess. This abscess will burst sooner or later if left to itself; and when burst, the urgency of the symptoms will at once commence to subside. But if the abscess can be opened by the surgeon a day or two previous to what would otherwise be the spontaneous period of perforation, by so much will the patient's suffering be alleviated.

The causes of this form of inflammation appear to be, more or less, associated with some disordered state of health, which renders the tonsils and throat especially susceptible to the influence of cold; for to direct exposure to cold may usually be attributed the immediate cause of an attack. But the patient is generally be found to have suffered from some constitutional disturbance, previous to the commencement of the sore-throat. When he is of a gouty habit. Rigors and flushes commonly herald the attack, before the gland itself becomes the seat of inflammation. Abscess, the result of simple inflammation of

DISEASES OF THE MOUTH.

the tonsil, irrespective of those cases which are complicated with **scarlet fever**, is a rare affection in childhood. It is seldom met with before puberty. It is not a frequent attendant on old age. It appears to be a disorder of youth and middle life.

It rarely occurs that both tonsils are simultaneously affected. But it does sometimes happen that after one tonsil has been the seat of suppuration, the opposite one becomes inflamed and runs into abscess. This, however, is infrequent. We occasionally meet with individuals who are subject to recurring attacks of inflammation of the tonsils, and quinsy. These attacks occur every one or two years for some time, often at shorter intervals.

Such conditions indicate especially, and prominently urge, the importance of constitutional treatment rather than simple attention to the local evil. It has constantly been found most serviceable, directly a patient has anticipated the recurrence of such an attack, to administer a stimulating emetic. The action of vomiting not only relieves, by emptying the stomach, but appears also to have some beneficial action locally. In fact, individuals subject to such attacks, and who have adopted such a course of treatment, have expressed themselves satisfied, that suppurative mischief has been warded off by recourse to an emetic in the very earliest stage of a sore throat.

As the attack is usually of a very debilitating character, an emetic of mustard is preferable, on every account, to one of antimony or ipecacuanha. Fomentations to the throat externally, inhalations of the steam of hot water pretty constantly, and a free purgation after the action of the emetic, should be the chief points of treatment. We strongly condemn as dangerous, in every sense of the word, the abstraction of blood, either locally or generally, either by leeches or by any other means. Life may be sacrificed by recourse to such treatment. Blisters appear equally unnecessary, though perhaps not equally prejudicial in their consequences. Mustard poultices appear to afford much relief in many cases; their application is often grateful to the patient's feelings; and by repeated application they can be made to keep up sufficient counter-irritation, without the inconvenience of producing a sore place; whereas a blister is a worrying annoyance to one who has already very great discomforts to put up with; and the subsequent discharge is a considerable nuisance, as it runs down and soils the linen, and stiffens the bed-clothes into which it soaks. Moreover, a blister often abstracts a large quantity of serum, and occasionally leaves a very red, irritable, and painful sore on the neck,

ded to the exhaustion following an attack of quinsy, means advantageous.

order is one that calls much upon the strength and powers of the patient. During the greater part of the food is taken but sparingly, and that in a fluid form. It behoves us to husband the forces rather than diminish heroic treatment. As there is much difficulty in eating, the less a patient is pestered with physic the less comfortable will be his day. At night it may be desirable to keep, if he be restless; and for such a purpose a few morphia will in all probability suffice.

It is the patient reduced by such an attack, that when he is evacuated by puncture, or escapes by an ulcerated tonsil, it will generally be found that some days of nutritious food are requisite, before he regains the strength he possessed in substance he carried previous to his illness.

Local treatment is more open to discussion. Some advocate the free application of the solid nitrate of silver to the surface of the inflamed membrane; but it is somewhat doubtful what amount of benefit is secured by this method of treatment. It is certainly attended by one disadvantage; it causes a good deal of local discomfort, leaves a most disagreeable taste for some time in the mouth, and gives rise to hawking and spitting, until the superficial epithelial cells occasioned by its application, are released or removed from the inflamed membrane. We are not aware that the application under the acute conditions we have described, is followed by satisfactory or beneficial results.

When the swelling of the tonsil be considerable, the surface very red, the part very tender, and yet no very distinct indication of suppuration present, it may appear sometimes advisable to make a small incision into the part to relieve congestion; or one or two incisions less deep. But as a rule, if the urgency of the case be not great, it is far better to wait the formation of pus rather than experimentally adopt an exploratory use of the lancet. Every sensible practitioner studies the feelings of the patient; every good surgeon avoids the uncertain or unnecessary use of the knife. Incisions, if made, often occasion suffering—are especially dreaded by some who have previously submitted to such treatment—and seldom appear to be necessary for the escape of matter. When, however, pus is detected

or suspected, the part should without loss of time be punctured with a knife.

There are some few precautions requisite, when any pointed or cutting instrument is used to open an abscess at the back or side of the mouth. The knife itself need only have, from its extremity, a cutting surface of about half an inch. Whatever knife be employed, it is best to protect the blade so as to cover its cutting edge up to this extent, by wrapping round it, from the handle towards the point, a piece of rag or plaster. A sharp-pointed straight bistoury is of all knives the most convenient for the surgeon's purpose.

In passing the instrument, when protected as described, into, or on withdrawing it from the mouth, the risk of wounding the lips or tongue is necessarily lessened—we may say is avoided; for no such accident should occur in proper hands. On puncturing the tonsil or abscess, an important precaution must be observed. The point of the knife must be kept turned towards the median line, especially after its point is buried in the soft tissues, and out of sight; on no account is the direction of the wound or thrust of the knife to be outwards. By due attention to this simple caution, all risk of wounding the larger vessels, lying on the outer side of the neck, is certainly avoided.

The patient's head should be fixed, either against the back of a chair, or by the hands of a third person. A sudden movement while the knife is passing into the tonsil, might lead to grave mischief, as the instrument might accidentally and unavoidably be thrust in the wrong direction.

The mouth is occasionally opened with some difficulty; and the tongue may be so much implicated in the inflammation that the patient cannot, by his own efforts, sufficiently aid the surgeon to obtain a good view of the swollen tonsil. Under such circumstances, a broad spatula gently applied to the surface of the tongue, will expose the part sufficiently to permit the abscess to be punctured in a satisfactory manner.*

* A practical hint in passing may here be thrown out, on the best method of examining the throat generally. With many patients it is often a matter of some difficulty to gain a sight of the fauces. The tongue is pushed upwards directly the mouth is opened; and when the spatula is applied with the intention of pushing down the tongue, the patient immediately appears to resist its application with a steady and firm effort. *If, however, the spatula is, without the slightest weight, laid on the tongue, it gradually appears to contract and is drawn backwards, and the fauces are exposed to full view.* No force or pressure is

on as the abscess has been punctured, the pus generally
eely on the withdrawal of the knife ; sometimes so abund-
and so suddenly that it runs into the gullet ; and from
umstance, or from the fact that it is often very fetid, it
nauseates the patient, sometimes to vomiting. Shortly
e escape of the pus, the patient usually becomes comfort-
presses himself wonderfully relieved, and soon desires to
of food. Convalescence is usually rapid and satisfactory.
; have already said, occasionally, but rarely, the opposite
comes inflamed and suppurates. Under such circum-
an early tonic and stimulating course of treatment may
site ; and the medical attendant must look, for his guide,
; symptoms which would indicate constitutional deterio-
and physical exhaustion, to be met by generous diet and
r other stimulants.

ie treatment of all cases of cynanche tonsillaris, there
; a certain amount of anxiety ; but the anxiety as to the
ms which precede the formation of the abscess, is as
; compared to the anxiety we may have to encounter when
rable symptoms are set up *after* the evacuation of the pus.
unhealthy constitution ulceration may supervene upon the
1, and be followed by sloughing sore-throat ; or œdema
surrounding soft tissues may arise, and travelling down-
round the larynx, pharynx, and œsophagus, terminate the
's life in a few days. But fortunately these are the rarer
llowing upon the evacuation of pus, the result of simple
nflammation of a tonsil, or true quinsy ; the usual history
h complaint is, that directly the abscess has burst or is
, convalescence is perfect and speedy.

sore-throat of childhood is rarely acute inflammation of
nsil ending in abscess. It is usually the symptom or
animent of scarlet fever, or partakes of that more fright-
l fatal character known as ' putrid ' or ' sloughing sore-
' of which we shall hereafter speak. The latter is usually
panied by enlargement of some of the glands of the throat
otid. In quinsy the throat-glands are seldom affected,
affected, not to any extent.

ral of the symptoms referred to above, as indicative of

e on the part of the surgeon to effect this ; the weight alone of the
appears sufficient to insure the object desired. This experiment has
stantly illustrated, on patients suffering from throat-affections, to our

the formation of pus in an inflamed tonsil, are present, to some extent, under conditions which are apt to be mistaken for quinsy.

A decayed tooth, or a broken stump, will often set up a considerable degree of alveolar inflammation, and be followed by an abscess of the gum or corresponding maxillary region. This condition occasions much difficulty in opening the mouth; the breath is fetid; the tongue becomes very foul and loaded; there is great difficulty in deglutition, and pain of face on the side affected. But generally one marked difference exists between the symptoms of quinsy, and an abscess originating in alveolar inflammation; when a tooth or a stump is the cause of the mischief, there is always more or less swelling of the face, and tenderness and swelling, especially over the maxilla affected. In quinsy, if there be external swelling present, it is not over the face or cheek; it is confined to the submaxillary region, and the neighbourhood of the parotid.

In all cases of inflammation, either within or external to the mouth or throat, the medical attendant should look with suspicious care to the condition of the mouth. There are few circumstances in practice more apt to be overlooked, than the evils which arise from the irritation of decayed teeth.

2. *Chronic inflammation*, terminating in enlargement of the tonsil and thickening of its mucous surface, is a very common condition; more especially to be observed in children, and in young persons of both sexes under the ages of from twenty to thirty. The tendency to this form of diseased action and enlargement, is mostly observed in individuals of a strumous or lymphatic habit. The enlargement frequently commences as early as the second year of life, and has been observed to occur more especially in children who have been weaned when very young, or brought up by hand-feeding. Less frequently, it does not show itself until towards the period of approaching puberty. It has a tendency to affect females in a greater proportion than males, though it is by no means an infrequent complaint in the latter sex.

The commencement of chronic enlargement of the tonsils appears to be unattended by any marked symptoms, local or general. Rarely is attention drawn to the condition of the part until either the thickness of the articulation, or loud snoring in sleep, excites the mother's curiosity, or gives rise to anxiety, when an examination of the throat is instituted. Pain

never a symptom attendant on these swellings. The only local symptoms are those which necessarily arise from the mechanical encroachment of the tonsils, in the median line, betwixt the external oral and nasal apertures, and the internal apertures of the breath- and food-passages. The intensity of these symptoms will depend on the size of the masses on either side of the uvula.

The causes which produce this chronic enlargement of the tonsils are not satisfactorily established. A careful inquiry into the history of a large number of cases has failed to detect any one specific reason to account for this diseased action. It does not appear to depend on cold, or insufficient nourishment, nor on want of cleanliness; for it will be met with in the children of the upper, as often as in those of the lower classes, and is often in those who are well provided and tenderly cared for, as in those who are the children 'of sorrow and acquainted with grief.'

This diseased action does, however, appear to affect in a larger proportion those children who have been brought up by hand or early weaned, than those who have been supplied for a sufficient time with the mother's breast-milk. The enlargement of the tonsils will also frequently commence, on the convalescence from severe attacks of the various eruptive or other forms of fever. Most frequently, however, their growth is grafted on a scrofulous diathesis.

The forms under which enlarged tonsils occur deserve some attention. One of the most frequent appears to be a uniform globular projection on either side, to a greater or less extent. In this condition the swelling may cease to grow, and remain stationary for years, without inconvenience or injurious consequences. Or the glands may slowly and steadily increase in size, until their surfaces touch below in the median space and above either side of the uvula. In other instances, the tonsils appear to enlarge towards the soft palate, and often upwards past its free border, as well as downwards so low that the lower end is hid from view by the base of the tongue, and is only with care detected when the latter is well depressed; or the tonsil on one side only may be seen to project, a perfectly rounded mass, which is attached to the fauces by a thin pedicle of mucous membrane: occasionally an outgrowth has been seen, a pendulous tumour hanging from the natural position of the tonsil.

The surface of an enlarged tonsil is generally somewhat uneven, often very much pitted; the mucous membrane thick and velvety, and seldom without an increased vascularity. In many instances all these conditions may be observed in an aggravated degree. Often, superficially ulcerated spots are seen dotting the surface of the swelling; or, what is more common, small points of thick sticky sebaceous secretion will be observed, marking the orifices of some of the ducts which lead into the substance of the gland-tissue.

The fauces, especially when the enlargement of the tonsils is considerable, may often be seen covered with a viscid yellowish semi-purulent secretion; this clings to the posterior surface of the glands, or lines the walls of the upper portion of the pharynx.

When the enlarged gland projects to any extent above the soft palate, some degree of deafness is occasionally complained of. This concomitant symptom appears to depend rather upon the thickened condition of the mucous membrane, which, extending from the tonsils, runs up to, and may line, the Eustachian tube, than upon any actual encroachment of the gland on the orifice of the tube.

Enlarged tonsils appear to be the result, not the cause, of constitutional derangement, provided they do not become large, or encroach much on the isthmus of the fauces. But if they increase in size, and much lessen the passage from the mouth to the pharynx, many unpleasant effects may be detected. The voice is generally somewhat unnatural and muffled, and the pronunciation thick. So that enlarged tonsils appear to be a perfect impediment to the production of fine sweet notes in song, or clear sonorous tones in speech. The breath is often offensive; partly rendered so by the decomposing sebaceous secretion sticking in the orifices of the glands, partly by the thick semi-purulent mucus which clings to the fauces.

The enlarged glands are constantly liable to attacks of ulceration, and sometimes to acute inflammation attended by fever. If large in children, they produce more general effects on the system than in adults. They now interfere with perfect and free respiration. At night the child snores loudly; often awakes in his sleep, and as often in a state of alarm. In some instances the natural and requisite amount of rest is so interfered with that the child's health suffers. There is also frequently some little inconvenience in swallowing, and care is observed to be

arged mass consists of condensed areolar tissue, which
uns like bands between and around the depressions above
l to. The naturally soft structure of the gland is so con-
and firm, that it assumes the character of fibrous tissue
cting the substance of the growth in every direction.

accumulation of epithelial secretion within the ducts of
nds, occasioned by the obstruction of their orifices, is the
ation of the formation of the cavities seen on a section of
and : for as the secretion increases, and cannot escape
h the obstructed orifices of the ducts, it collects in the
dilates them, and sets up chronic inflammation of the
re of the gland itself, which terminates in its permanent
ement.* This secretion will frequently be observed to
from the surface of the enlarged tonsil when it is firmly
d between the blades of the vulsellum prior to removal.

treatment of enlarged tonsils may be summed up in a
ses. Unless the voice be affected to such an extent that
comes desirable to lessen or remove them ; unless the
bance to the sleep of the child be manifestly interfering
health ; unless the breath be rendered fetid or offensive
secretion from the ducts ; or unless some other cogent
render it desirable to interfere with the mass—it is
to leave the tonsil in its slightly enlarged condition,
at the application of any local or constitutional measures.
hild so affected be delicate, and the tonsils appear increas-

on the importance of attention to diet; and with children, sea-bathing is of the utmost advantage.

Local applications are frequently recommended and are constantly adopted. Nitrate of silver rubbed over the points of the gland runs into the substance of the gland; sometimes nitric acid carefully applied to portions of the gland; and a variety of troublesome applications, have had their advocates for the arrest of these masses. But an examination of a portion removed by excision will show such structural changes, fibrous thickening of the finer areolar tissue, that the author considers a useless waste of time and material to procure absorption of such dense tissues. Not only actual slough can do good, if caustic be used; but application can offer much hope of acting on the mass; it is therefore best not to waste time or inflict pain by recourse to such useless local measures. If the treatment does not arrest, and local treatment fails, removal by the knife of a portion of the gland is necessary, and, in our opinion, the only efficient remedy. With a knife can remove all that is necessary in the gland, and the patient will be entirely rid of the complaint a week after.

When the glands are so large that some must be removed, to relieve the symptoms of discomfort which the patient suffers, the operation is successful in adults; but may be somewhat troublesome when

to expose the tonsil to view, the mouth dilator, invented by Mr. T. Smith, will effectually answer the purpose.

In the adult, there is no difficulty in the operation such as may be encountered in children. The gland can be at once seized with a vulsellum, and a portion rapidly cut off with the knife. If both tonsils be enlarged, and this is usually the case, there is no objection to a portion of each being removed on the same day. With children it is preferable to do so, as the fear of an operation is much greater when a child has had any experience of a previous one. But in an adult it may be left to the discretion or wishes of the patient, whether the operation be completed at once, or at a subsequent period. If the tonsils are very large, it is always advisable to take away a considerable portion of each. But it is by no means requisite to remove the whole of the projecting mass. If half of the mass be removed, and the cells thus laid well open, what with their becoming emptied, the relief occasioned by the incision through parts chronically inflamed, and the subsequent contraction of the cut surface by cicatrisation, the portion that is left shrinks and sinks into the side of the fauces, to give no further trouble to the patient.

After the removal of the mass, the patient generally suffers but little; or rather as one affected with slight sore-throat. The precaution to be given is, to use soft food, and to avoid exposure to cold for a few days, when the cut surface will be about healed. Hæmorrhage rarely follows to any troublesome extent on this operation; but it may, and sometimes does occur. The surgeon, always anxious to avoid such a contingency, should take the precaution, when operating, to draw as much of the gland as he conveniently can well into the cavity of the mouth inwards and forwards, so that a large portion may be seen projecting from the attached base; not more than half of the mass need be removed, and even as much as this only when the tumour is of considerable size. Such a portion will always be found sufficient for the ultimate relief of the patient, and the removal of all the previous disagreeable symptoms. In the practice of the author, such a proceeding has never been followed by troublesome hæmorrhage. Should hæmorrhage occur, iced water as a drink, or the mouth kept open for a time, may control it; or digital pressure may be requisite.

In children, hæmorrhage may be suspected only after it has continued some time, from the cheeks becoming pallid, and the

little patient showing signs of faintness. If, on examination, any distinct vessel is seen to bleed, a ligature can be applied with a little care; or recourse must be had to pressure, if the bleeding is from the general surface.

We need only refer to the proposal to remove enlarged tonsils by the application of ligatures, to condemn the plan of treatment as tedious, painful, and productive of much offensive discharge, until the slough separates.

3. *Tonsillitis maligna*.—Ulceration of an acute form attacking the tonsils, and rapidly running into sloughing of the fauces, is a very grave—often a most intractable—affection; and as rapid as it is unmanageable.

Sore-throat is the first symptom complained of, and with it the patient probably expresses himself as feeling ill. Rigor, followed by feverish disturbance, will generally supervene. The tonsils will soon be observed to be of a dusky-red colour, and swollen; and the surrounding soft tissues and uvula are already cedematous. There will generally be much pain, and some difficulty in deglutition. Patchy shreds of grey or yellowish membrane appear embedded in the surface of the tonsils. These, after a time, spread, unite, and become deeper, until a considerable portion, if not the whole, of the soft palate may become implicated in the spreading mischief. The discharge which accompanies this form of slough escapes freely from the nose; for as the havoc spreads, the passage of even fluids to the throat is rendered difficult, and is often entirely intercepted by the cedematous state of the cellular tissue below the root of the tongue—so that liquids taken by the mouth are rejected through the nostrils. The character of the discharge is usually most offensive and intolerable. The room has to be kept well aired, and a constant and liberal use made of solutions of chloride of soda or lime, to moderate the fetor occasioned by the discharge.

If the patient survives, large sloughs separate from the parts attacked, and a considerable granulating surface may soon be observed, marking the havoc of the disease which has passed. In this havoc the whole of the uvula, and a considerable portion of the soft palate, have been seen to disappear, the patient's health being subsequently perfectly restored; though there remain for life the permanent defect, the alteration of the voice, and the inability to articulate distinctly—the results of the local injury to the soft palate.

In the greater portion of severe cases of malignant sore-throat, the probability is, that they are specific, or only a type of the severer forms of scarlet fever. But still the surgeon's experience may be sometimes advantageously referred to in some of the stages of such cases; and for this reason we have taken a brief notice of them.

As in sloughing of other tissues, so in that of the throat, under all circumstances a stimulating and tonic treatment must be pursued; wine must be freely given; and with children, the muriated tincture of iron in full doses may be most advantageously administered. It is most efficacious as a tonic in such cases; can be given in small bulk; and, with the addition of a little syrup, forms a mixture children do not object to—a desideratum of no small importance in their management when sick.

A superficial and less severe form of ulceration of the tonsils is a very common occurrence. 'Ulcerated sore-throat' is a term so familiar, and a condition so well understood among the more industrious and often over-worked students of our hospitals and dissecting-rooms, that it requires but little description here. Pain and some difficulty in swallowing; a creamy tongue; soft, weak pulse; pallid face; loss of appetite, and often loss of sleep; much languor and pains in the limbs; tonsils congested, and somewhat swollen; small, irritable, superficial, and scattered ulcers on the fauces and tonsils: such are the chief symptoms found to accompany one of these attacks.

The treatment is well understood. Change of air; a stimulating gargle; a slight aperient if requisite (mercurials, however, to be generally avoided); the internal administration of bark and ammonia, or chlorate of potash; and a nutritious fluid diet, with a few glasses of wine daily. Convalescence is generally satisfactory under such treatment. In all of the more acute inflammatory and ulcerative actions affecting the throat and tonsils, the internal administration of chlorate of potash, as well as its free use as a wash or gargle to the part, appears to be attended by marked benefit. A lotion of the permanganate of potash is also most agreeable in such cases; for, as a gargle, it leaves a very pleasant taste in the mouth, and rapidly renders odorous offensive discharges. It may with safety be used solely as a gargle or wash in all throat affections. So also a weak solution of carbolic acid.

'Relaxed uvula' is by no means an uncommon complaint.

The term does not quite describe the actual state of the parts in their altered conditions; for the uvula may be thickened and not increased in length. Under such circumstances it occasions but little annoyance, and seldom requires attention. Or it may be lengthened without being thickened, and this is the most common evil; and then surgical interference is necessary.

The increase of size, or the elongation, appear to be as independent of the condition spoken of under the name of enlarged tonsil, as the two conditions are of each other.

The increase of size is usually accompanied by a thickened state of the mucous membrane, and often a slight tenderness of the part. Though the thickening may remain, the tenderness after a time will subside; and then the part is left in a permanently enlarged, though not in a persistently enlarging, condition. It seldom increases to an inconvenient size.

Should the symptoms which attend the first increase of size be troublesome, the simplest local remedy is to smear the surface once or twice with the solid nitrate of silver; and constitutionally use such measures as the general condition of the patient will indicate to the observant practitioner. Usually the condition is one which is benefited by tonics. Elongation of the uvula appears to be generally dependent on an excess of mucous membrane: for the azygos uvulae muscle does not often form any part of the increased growth. The mucous membrane may extend an inch beyond the extremity of the muscle, and hang down like a thin narrow slip of tissue, ending often in a point. When of this length, it dips into the pharynx, and the point rests over the aperture of the larynx. This contact with the larynx keeps up a constant source of irritation, and occasions a troublesome cough. Frequently the tickling of the fauces and pharynx produces nausea and a tendency to vomit. Patients sometimes, after food, on coughing, reject a portion of the meal taken. These symptoms are often attributed to other causes, until the throat is examined and the condition of the part detected.

This elongation of the uvula is not often seen in children, but most frequently in the period of middle-life. It is seldom observed in old age.

The treatment is as simple as the relief is certain. Removal by knife or scissors is the only treatment to be entertained for a moment. The point of the uvula is to be seized with a long

of forceps, and cut off so near to the base as to leave the rest of its natural length.

The patient may feel the throat a little sore for a day or two, but no treatment beyond ordinary care will be requisite. The palate is very apt to be implicated in inflammation about its neighbourhood; and constantly, under such circumstances, becomes oedematous. This condition, however, is seldom confined to the uvula alone, but implicates usually the soft palate and adjacent tissues. It depends on erysipelatous or diffuse inflammation of the cellular tissue—a dangerous and often fatal affection, which will be found treated of in the essay on DISEASES OF THE LARYNX.

Tumours of the soft palate and fauces are sometimes met with of a non-malignant character, and independent of inflammation. These usually consist of, 1. Fibro-cellular tissue; 2. Cysts; 3. Warty growths of a non-malignant character, occasionally met with attached to the soft palate, and growing apparently from the mucous membrane.

It has occurred to the author to have to remove such growths in two instances in young women; in both, the growths commenced without being noticed, and were accidentally discovered; they were readily removed, and were found to consist of simple epithelial structure.

1. The small fibro-cellular tumours usually assume a pedunculated character. They are painless; usually attached to the free border or upper surface of the soft palate; and are generally only detected when they become locally inconvenient by dropping down below and behind the root of the tongue. They are usually somewhat slow in their growth, and as they increase in size become more pendulous, often supported by a thin pedicle. They can be readily removed with scissors and forceps.

2. *Cysts* of these parts commonly contain thin glairy fluid, and are generally the result of obstructed muciparous ducts. They are readily treated by a simple free incision, and a subsequent application of solid nitrate of silver, or a drop or two of nitric acid, applied on a probe to the interior of the cyst.

Sebaceous cysts are occasionally found in the substance of the soft palate. They may be recognised by their yellowish white colour showing through the mucous membrane. They only require to be laid open, and the interior of the cyst touched for a day or two with a probe dipped in nitric acid.

Abscess of the soft palate is rare; but it should be opened as soon as detected. It generally gives no subsequent trouble.

CONGENITAL FISSURE—‘CLEFT PALATE’—AND DEFECTS OF THE PALATE FROM DISEASE OR ACCIDENT.

The occurrence of congenital fissure of the palate is so frequent; the evils which result from it are so marked; and the life of an infant is so often endangered and not unfrequently cut short, when the fissure is extensive—that a consideration of the treatment which proposes to overcome or to modify the defect, may truly be said to form a subject worthy of notice in a work on Practical Surgery.

The investigation of the early development of the mouth, explains the various degrees of extent which congenital fissure of the palate *may* assume, and such as *are* constantly found to exist in the new-born infant.

To Mr. Goodsir * the profession is indebted for an accurate account of the development of the mouth; and although, in a practical work of this character, it would be intrusive to enter very fully into such a subject, we are compelled to take a cursory view of the changes which occur in the upper jaw *in utero*, in order to clear up to a certain extent the causes of the various deformities which affect it.

In a fetus of about the sixth week, examined by Mr. Goodsir, the cavity of the mouth, nares, and pharynx formed one undivided space; the palate was entirely wanting; the outline of the future alveolar ridge was evident. With the particular description of this latter process we need not trouble the reader: but from the inner margin ‘a thin semi-transparent membranous fold passed backwards on each side, attached externally to the sides of the capacious bucco-pharyngeal cavity, bounded internally by a free edge, opposed to its fellow of the opposite side, and terminating posteriorly on the lateral walls of the pharynx.

• These folds constituted at this period, a partial division of the large common nasal, buccal, and pharyngeal cavity into a superior and inferior compartment.

• The upper wall of this common cavity was smooth and flat posteriorly; but anteriorly it was contracted, and terminated

* *Edinburgh Med. and Surg. Journal*, vol. li. p. 1.

longitudinal bar (the future septum nasi), which ran forward to be attached to the superior surface of a horse-shoe (described) at the median line (in front) and to the other in that neighbourhood.

Under the bar a deep cavity was seen, which communicated with the exterior of the face by two small foramina (the orifices of the nostrils), which constituted at this period the whole of the nasal organ.*

In an embryo of about the seventh week, next examined, the cleft had slightly diminished, but was still of sufficient extent to display the whole of the undivided nasal cavity.

In an embryo of the second month 'the cleft of the palate was diminished, existing only as a small angular deficiency in the fleshy portion, or soft palate.'

In an embryo of the tenth week, the anterior portion of the cleft 'had thrown itself into folds; and there was an indistinct depression.'

The uvula 'was well marked between the eleventh and twelfth weeks.'

At an examination made of the bones of the palate previous to, or not later than, the sixth month of foetal life, in addition to the median suture between the palate-processes of the superior maxillary bones, a distinct oblique suture will be observed, which starts from the median suture a little behind the alveolar ridge, runs outwards and forwards through the maxilla, and terminates on its anterior aspect, at the point subsequently marked by the interval between the lateral incisors and the canine teeth. So that in fact the under or palate-surface of the superior maxilla is at this period of life (third to fifth month) marked by this suture, as if the bone had been at an earlier period divided into two portions: one, an anterior small 'intermaxillary bone,' or 'os incisivum,' as it is termed, and which carries the incisor teeth; and a posterior, the larger portion, in the alveolar process of which are developed the canine, bicuspid, and molars. This suture disappears, under the healthy natural development, soon after, or even before, birth.


The variations met with in the deformities of the palate appear to be distinctly due to an arrest in the development, already described, of the upper jaw and soft palate—an arrest which occurs from some unexplained cause, at an earlier or

* This may be taken as the future 'os incisivum' or intermaxillary bone.

later period, but probably within the first twelve weeks of foetal life, in accordance with the extent of the deformity. The variations may be thus classed :—

1. The most extensive; a double cleft of the alveolar ridge with a fissure from each cleft running backwards and inward and joining together immediately behind the intermaxillary bone, and being continuous with a median fissure through the entire hard and soft palate.

FIG. 244.



Complete cleft of the palate with double hare-lip. (From a preparation in the Museum of St. George's Hospital.) *a.* The intermaxillary bone. *b.* The septum ~~nasal~~. *c.* The tongue. *d.* The cleft into the nasal cavity. *ee.* The separated halves of the upper jaw.

2. A single cleft of the alveolar ridge on one side or the other of the median line; running back into an entire cleft of the palate.†

3. The alveolar ridge entire; the cleft of the palate commencing immediately behind it and complete backwards.

4. The cleft running as far forwards as the middle of the palate-processes of the superior maxillæ, or through the palatine bones only.

* The cleft of the alveolus under such circumstances is often complete throughout on each side from front to back, and from the mouth into the nose.

† Instead of the alveolar ridge being cleft, as described, in this variety, there may be only a notch, or a large or small foramen through its base. The foramen has been partially closed by growth, or by pressure of the lip.

ure of the whole of the soft palate, or only implicating
 in the extreme of one and the other every gradation of
 ay occur.

onally, but rarely, a congenital defect is met with in
 an line, either as a circular or oblong opening: in
 l behind it the palate is closed. These openings most
 y occur in the palate-bone; and usually, when such
 ng exists, the median line of union is very plainly
 n the under surface of the soft palate and the uvula, by
 t straight thin cicatrix, showing the union of the sides
 ft palate completed, although the palate-bones, in con-
 of their defective development, were unable to perfect
 -oral septum in their immediate neighbourhood.

be observed in some cases that the gaps through the
 ridge are excessive, *i.e.* complete from mouth to nostril;
 n other cases they are but partial. When excessive,
 n the fissure is complete through the hard and soft
 the intermaxillary bone, or 'os incisivum,' becomes
 dely separated from the superior maxillæ, and forms a
 more or less round (the horseshoe lobe of foetal life
 described), which is supported by and attached to the
 inferior termination of the septum nasi. This nodule

FIG. 245.



hare-lip, showing the intermaxillary bone attached to the septum nasi.

more or less prominent, and is occasionally so much so
 appears simply to be stuck on to the tip of the nose.
 these circumstances it is but little covered with skin,
 is merely a prolongation from the integument on the
 the nose, and unconnected with the integument of the
 either side.

he nodule be dissected between the sixth and eighth
 after birth, in it will be found *not only the temporary*

incisors, but also the germs of the permanent ones; not always the lateral, but invariably those of the central incisors. Occasionally those of the lateral will be found, though they frequently are sacrificed by the freak played by nature in permitting these gaps to occur.* Whenever a single fissure occurs through the alveolus, the gap will not be found in the median line, but, as a rule, on one side or other of it. We cannot satisfactorily explain this. It may depend on the peculiar conformation of the upper jaw and its early development. We would refer those interested in the subject to the further consideration of Mr. Goodsir's observations.

It is rare indeed that the infant born with extensive cleft palate and alveolus has not also the upper lip notched or fissured opposite to the defect in the bone. Usually, if not invariably, when the cleft of the alveolus is double, the cleft of the lip will correspond; when single, the lip is usually similarly affected. But when the cleft of the alveolus without cleft of the palate occurs, hare-lip is not always present.


In complete cleft of the roof the attachment or disposition of the septum nasi varies in different cases. As a rule, the more extensive the gap, the greater the deformity. The septum will often be seen, in complete cleft of the palate, to be continuous with its lower border with the margin of one or other of the edges of the cleft; in which case one of the nasal cavities is so far shut off from the mouth, while the other is open to it. In other instances the septum dips down between the nasal fossæ almost as low as the edges of the fissure, and terminates in a well defined border, unconnected with the superior maxillæ; or will be sometimes seen bent or folded on itself, its edge turned up to one side, with one of its lateral surfaces facing downwards.

These different conditions of the septum have no practical

* When this nodule is very prominent, and appears likely to interfere with the success of an operation for the closure of hare-lip, we would strongly recommend the importance of depressing it forcibly rather than removing any portion of the whole of it. It may be forcibly broken and depressed, and made to fill the gap between the superior maxillæ, where it will almost invariably become fixed, and subsequently have shooting out from it some or all of the incisors. If these be defective, the surface of the bone forms an excellent rest for a plate to carry artificial teeth. If the os incisivum be removed, the superior maxillæ are certain to be more or less drawn together by the action of the upper lip, the latter become much flattened, the upper portion of the mouth much narrowed, and the gap in the alveolar ridge never entirely closed.

ing on the subsequent treatment of the case. Rokitsansky refers to fissures of the palate caused by the absence of the os incisivum, and attended by fissure of the upper lip *in the median line* * but he does not state that he had met with an example of this rare deformity. Sir Wm. Fergusson,† in alluding to the occurrence of cleft in the lip, says: 'Doubtless it has been called hare-lip from a certain resemblance to the fissure in the upper lip of the hare; but in the human subject it differs in this peculiar feature, that it never is in the mesial line, as it always is in the hare.'

FIG. 246.



of the palate in the median line. (From a preparation in the Museum of the Royal College of Surgeons.) *a a*. The tongue divided in two in order to expose the cleft. *b b*. The larynx similarly divided. *c c*. The two halves of the upper lip and jaw, between which is the cleft opening freely into the nasal cavity, the intermaxillary bone being absent.

Rokitsansky only alludes to the occurrence of the fissure in the median line, gives no details of any peculiarities which might be supposed to accompany a defect. Sir W. Fergusson had apparently not met with a case; and the deformity is no doubt most rare. But a specimen such as Rokitsansky refers to (result of absence of the os incisivum and fissure in the median line of the lip)—may be seen in the Museum of the Royal College of Surgeons in London. In this preparation (of which no history is recorded), the fissure of the upper lip is in the mesial line, the result of a great gap, as if a large portion of the middle of the lip were destroyed; the os incisivum is altogether absent, no vestige of it being present; and the cleft of the hard and soft palate is complete.

* Rokitsansky, *Pathological Anatomy*, vol. ii. p. 3.

† *Practical Surgery*, p. 584.

We do not attempt to enter into an explanation of cleft palate. We possess no data to justify observation has thrown any light on what is considered an arrest of development at some time prior to the fourth month. But the cause is shrouded in mystery, and is one of those a function of nutrition and growth, to account entirely beyond our present knowledge.

Thus much, however, we may state, that in the os incisivum from the superior maxillæ, we find to a somewhat similar arrangement in the jaws of certain animals, in whom the incisors occupy a distinct interval or gap, from the bicuspid and molar teeth.

Cleft palate and cleft lip are frequently found in more than one member in a family; but we cannot say that a cleft of the mouth or lip is a common result of hereditary transmission. It is rare to find a parent and offspring affected on either side, and sisters of the same family constantly present conditions of the deformity.*

Practically, the more extensive the cleft, the wider it will generally be found, and the more the surfaces forming the sides of the gap. Usually the palate is narrower in front, and widens towards the back. But much variety will be found if a large number of cases are examined.

the extent of the cleft be restricted to a small portion of the palate-bone and the soft palate, it by no means follows that the cleft is equally narrow in proportion; on the contrary, it is frequently found that these partial clefts are very irregular, and so much so in some instances, that the palate processes and the sides of the bifid soft palate appear most wanting. These partial clefts have been found to be, in some few instances, the greatest breadth met with; but such an extent, in one or two instances, that operative interference was considered unjustifiable, as there did not appear to be sufficient soft palate, sufficient tissue to allow of the edges being brought together by suture.

The primary effects of cleft palate, when extensive, are sufficient to create much necessary alarm for the safety of the infant's life. If hare-lip exist, with double or single cleft of the lip, and a broad cleft of the roof, unless the infant be supplied plentifully with its mother's milk from the first, it will very soon sink from exhaustion. It is very difficult in such cases to introduce a sufficient quantity of milk into the stomach, to obtain an amount of nutrition equal to the maintenance of the infant. The mother's milk is the only food that should be given during the first six or eight weeks. In all cases of cleft palate, the infant is unable to suck the nipple: for as the naso-buccal septum is wanting, the child cannot create with its tongue a vacuum in the mouth, and is therefore unable to draw the ducts of the

suction is not requisite, but at the same time gentle, and easily regulated by the inclination of the nurse's hand ; or a small piece of sponge cut into the shape of a nipple, and then placed in the mouth, will often be equally well, but of course the process requires as often as the milk is sucked out of the spoon. regards the early nursing of a child born with a cleft palate. The question now to be considered is, whether it may be desirable to alter, or improve, by surgery the condition of the mouth ; and if so, when it should be, and at what age such treatment may be successfully put in practice. In the first edition we stated our opinion that it was not desirable to close the fissure of the palate, by operation, and that such treatment was better deferred until when the patient was old enough to judge for himself, as he would prefer to wear an artificial palate for a cleft fissure permanently closed, and was moreover unwilling to submit to an operation without the aid of chloroform.

Since that time, however, the experience of our own, has induced us to materially alter our practice in this respect.

Mr. Thomas Smith, of St. Bartholomew's Hospital, I believe, the first to introduce the use of chloroform on the palate in children ; and he has entirely changed its administration. clefts of the palate may be successfully operated on in childhood at a very early age.

erty ; but we are of opinion that the surgeon should still be somewhat guided by the age of the child, as to when the operation should be performed.

The real object of the operation of closing the cleft in the palate is to enable the patient to articulate hereafter, plainly and intelligibly—not to enable the child to take food. An infant with extensive cleft, when first born, has often some difficulty in taking sufficient food to satisfy its wants, or to assimilate it, as it would be nourished were its palate perfect ; but this difficulty is overcome in a few days, and if proper food is taken the infant usually thrives. An operation at this age is therefore not requisite, nor is it desirable : the loss of blood the infant would sustain in such an operation would be attended by great risk to life ; the tumefaction and tenderness of the mouth subsequent to the operation would be sufficient to interfere greatly with the infant taking a proper amount of aliment ; and both the loss of blood and the insufficient quantity of food might materially interfere with the union of the parts, even supposing the infant does not sink under these circumstances. A very slight loss of blood may prove fatal in a young infant, especially in one of a puny delicate nature, and children with cleft palates often are ; but the loss of blood in an operation for cleft palate is often severe, more especially when the hard palate is dealt with ; great risk would therefore be run by the very early application of this operation,

ment of the articulation, to defer all operation until the child is two or three years of age. In the case of Mr. Smith's gag and the administration of chloroform, the operation may now be undertaken without difficulty, and readily performed, and in a large majority of cases successful. In children with entire cleft of palate, we recommend the closure of the soft palate as the first step in the operation. If this be successful, the hard palate will in most cases diminish in size in the course of some few weeks. Dr. Ehrman mentions that this occurred in cases under his observation, and we have confirmed the experience by our own. It is, in fact, one of the advantages of an early operation on the soft palate. If successful the continued action of the muscles of the soft palate cleft tends to approximate the edges of the soft palate, and what is observed to take place in any case of cleft of the alveolar ridge after the fissure in the upper lip is closed.

As the most simple form of cleft is that of the soft palate or soft portion, we propose to consider, first, the operation adopted to close it, irrespective of any extension into the hard palate.

Should the uvula alone be bifid, very little operation is required to secure union of its opposed edges; and indeed the uvula is not affected, nor the cleft detected by any defect in articulation, it is better not to meddle with it.

other. The surgeon should first seize one point of the bifid uvula with a pair of long spring forceps, and draw it forwards, transfix it near its inner border with a narrow sharp knife on a long handle, and freely cut upwards and downwards, in order to remove the mucous membrane along the whole of its inner margin. A similar proceeding on the opposite side completes the use of the knife. A couple of sutures will be sufficient to approximate the edges and keep them in place.

Mr. T. Smith observes with respect to the material to be used for sutures, that he has found fine fishing-gut best adapted for the soft palate in the young; or horsehair for the lower part of the velum and uvula; both these substances, if soaked for a few minutes in water, will become sufficiently supple for use.*

In the experience of the author, fine silk is as efficient for the purpose of sutures as any other material, and is preferable to most. Such sutures are readily introduced, and very readily removed. The latter is not a point of trivial importance, and is an advantage when the soft palate is concerned, which does not equally apply to metallic sutures. The author has invariably used silk sutures in all operations for cleft palate, and with the most satisfactory results; nor has he, from the experience and experiments of others, seen any good reason to prefer any other material.

When a cleft extends forwards through the whole of the velum, or even to a slight extent into the palate-bones, the operation is rather more complicated than that above described; and though not beset with any difficulties, requires a careful examination into the anatomy of the soft palate, and some little familiarity with the operation, before the surgeon can be confident of obtaining that success, which should usually follow attempts to close a cleft palate by surgical interference.

That the edges of a wound should meet without any traction on its margin by sutures, is a fundamental rule which applies generally to all plastic operations; but especially is it absolute in any operation to close a fissure of the palate, whether in the hard or soft portion.

If a fissure of the soft palate be carefully examined, and with the mouth open the patient make an attempt to swallow, the action of the muscles of the pharynx and tongue is such, that the edges of the fissure are approximated by the action of the

* *Trans. Med.-Chir. Soc.* vol. li. p. 85.

superior constrictors; the extremities of the bifid uvula touch, and often the gap is for an instant closed to one-third or one-half of its extent; but immediately afterwards the sections of the soft palate are drawn back to the walls of the fauces; and often the whole of the soft palate itself is so closely contracted into the side of the fauces, that a superficial observer might conclude, in some instances, that no soft palate existed. This latter effect is the result of the combined actions of the levator and tensor palati muscles contracting towards their attached extremities, and thus drawing up the sections of the velum. So that every attempt to bring the edges of the fissure together, would be opposed by these muscles, on either side, pulling away from the median line, and consequently drawing directly away from the line at which the edges of the fissure should meet to insure union.

Having experienced this evil, and at once seizing on the cause of the failure, which then so constantly attended attempts to close the fissures of the soft palate, Sir W. Fergusson proposed the division of these muscles prior to the edges being brought together. In a paper published in the *Transactions of the Royal Medical and Chirurgical Society* he fully describes the method he adopted to insure this result. Since Sir W. Fergusson promulgated his views, this operation has been commonly undertaken, and, thanks to his suggestion, has been followed by a large share of success. Sir W. Fergusson recommends the division of the levator palati on either side, by cutting, with a knife bent at a right angle, behind the curtain of the soft palate. The division of the muscles is thus most effectually secured. He also recommends that the palato-pharyngeus be divided on either side. In able and experienced hands, to cut with a right-angled knife, behind the palate, where the point of the knife is out of sight, is a simple matter; but there are few surgeons unaccustomed to these operations, who feel at ease in using such a knife in such a situation. Having observed this, it occurred to the author, some few years back, that the levator muscles could be as readily divided by cutting through the palate. The experiment made on the dead body at once proved the extreme feasibility and facility of thus being able to divide these muscles; since then every case which has come under his care has been thus treated, with the most satisfactory results. The method of procedure is very simple. First a suture is passed through one section of the soft palate at the

ts inner side. This process can be distinctly felt in the
ce of the soft palate, internal, and a very little posterior
st molar tooth. Running the knife upwards and back-
and somewhat inwards, the point may at last be seen in
, having passed through the entire thickness of the soft
and having cut, if not wholly, at any rate partially,
the tendon of the tensor palati: the knife should now
e most of the fibres of the levator. If the handle of the
: next raised, the point becomes depressed; and if the
e drawn forward, while it is at the same time made to
nwards, it travels through a considerable section of a
n the posterior surface of the palate, and insures the
of the greater portion of the levator palati. As the
ade travels downwards, the tension of the palate gives
d often the division of the muscle is felt to be suddenly
; the ligature being no longer pulled upon by it, though
s to division it will be felt sensibly and spasmodically
ting. As the knife is withdrawn through the wound,
ision of the levator muscle should be thoroughly effected.
ound, in the front of the palate, need be no more than
lth of the knife; whereas the wound behind is neces-
much longer, for the fibres of the levator have there to be
by the sweep of the knife. Provided the muscle be
ully divided, as soon as the knife is withdrawn it will be
that all voluntary and involuntary movements of the



already used will answer for this purpose; and there appears this advantage in the above method of operating, that *one knife suffices for the entire operation.*

Great care is requisite to free the edges sufficiently of their mucous covering: it is better to remove a little more than requisite, than an insufficient quantity; for the success of the operation is not interfered with in the former case, whereas the operation will most certainly fail to some extent in the latter condition.

Though Sir W. Fergusson recommends the division of the palato-pharyngeus muscles, the author has of late satisfactorily operated on several clefts of the soft palate, without having divided those muscles or touched them with the knife.

The completing stages of the operation consist in passing the sutures through the flaps, and securing their edges in contact with each other, until union has been efficiently established. Two or three fine curved needles, on rather long handles, should be ready-armed with fine white silk, slightly waxed; as soon as one needle has been used, it can be again armed, while another is in the hands of the surgeon. It is best to decide at first how many sutures may be requisite; and also to observe carefully the points at which they should be introduced; that when all are passed, their positions in each half of the palate may correspond, in distance, from each other, as well as in their distance from the margin of the wound; and also that those on one side should be directly in line with those opposite. The sutures, in each needle handed to the operator, should be at least a yard in length; and each suture should be doubled, for its whole length, before being passed. The surgeon, with the needle in the right hand, and a pair of long spring-forceps in the left, should now push the point of the needle through the soft palate (on the patient's left side) as near to its anterior margin as practicable; for although this, the highest suture, is not intended to approximate the edges of the cleft so that they should touch, still, by slightly assisting to narrow the cleft, it adds much to the utility of the one below, when the operation is completed. For the reason stated the upper suture should be passed rather near to the palate-bone.

The point of the needle having been passed through the palate, should now be pushed inwards, until it becomes visible through the cleft: one thread of the suture is then to be seized with the forceps, and gently drawn forwards, *through* the fissure and out

s on the opposite side. The loop must be drawn for-
mly, until secured by the fore-finger of the operator,
e needle may be disengaged from the rest of the suture.
ould then be on one side through the palate a single
on the opposite a double suture, the loop of which
nds to the inner side of the fissure. The *inner* end of
le suture is to be passed through the loop of the double
short distance, and the ends of the latter gently pulled,
; whole double suture is withdrawn from the palate, and
as pulled into the corresponding portion of the velum
the single suture; and this now should lie across the
with each end passed through a section of the soft palate.
nds of the suture should be secured outside the mouth,
the succeeding ones, as many as may be required, are
nd secured in a similar manner. One is usually required
lower portion of the uvula.

on as the requisite number have been introduced, each
be separately tied; carefully, so as to adapt the edges
to each other; and not too tightly, that room be allowed
sequent swelling of the soft tissues. If the sutures be
drawn, they most certainly will cut themselves out by
ing ulceration of the part through which they pass. A
ot to bring the edges together, and a second knot over
re sufficient to secure the suture, and maintain apposition
nion is safe. The ends of the sutures should not be cut

sutures on a subsequent one, the second operation is much simplified by the freedom from that bleeding which takes place whenever the muscles are divided at the same time that the cleft is pared and the sutures passed.

Mr. T. Smith's method of operating is, first to pare the edges of the fissure from below upwards, then to pass the sutures, and lastly 'to draw up the palate with all the sutures in one hand, and with the palate on the stretch to divide, if necessary, the palato-pharyngeus, and subsequently the levator palati; if the palate will not come easily together, two lateral oblique cuts may be made, one on either side above the highest suture, separating the soft from the margin of the hard palate to a small extent.'*

The operation for closing the hard palate is a very different proceeding to that described as requisite for the relief of fissure of the soft portion. The result of some experience confirms the opinion long entertained, that, when fissure of the hard as well as of the soft palate co-exists, the attempt to close the entire fissure in one operation is not advisable. Separate operations, at some short intervals, should be had recourse to, so as to close either the hard or soft portions first, and the remaining portion subsequently. There are several cogent reasons to render such practice generally desirable. In the first place, an attempt to close a fissure of the hard and soft palate at once entails a long and painful operation on the patient; and as he has to keep his mouth open while the operator is at work, it becomes a most tiring and irksome effort, if the period is a prolonged one. In the second place, the loss of blood which will often occur, in an operation on the hard palate alone, is sometimes considerable; often apt to make the patient faint, and so prolong the completion of the operation. But should the attempt be made to close the hard and soft palate at once, the loss of blood is usually much greater, and often too great to allow of perfect success in the attempt. In the third place, when an attempt is made to close the cleft in one operation, the edges of the soft tissues both in front and behind are apt to ulcerate, and often to slough.

If the soft palate be alone operated on, some ulceration may and does often supervene; but it is rarely of such consequence as to be ultimately of any importance, provided the palate has

* *Trans. Med.-Chir. Soc.* vol. lii. p. 84.

union and reparation are becoming established. If, the soft palate be first selected for closure, and the be successful, the union of the flaps converts the portion of the fissure into a foramen; and *pro tanto* the nutrition of the soft tissues of the hard palate, when detached from the bone, at some subsequent operation; hard palate be operated on first successfully, so much in front, to assist in the nutrition of the soft palate, comes under the knife of the surgeon. As before stated, operation on the soft palate aids to contract the gap left in the hard palate, even without further operation; and this will be effected in the adult, but especially in the child.

The fissure of the hard palate, or one extending to the alveolus, is generally closed without much difficulty. A simple method of detaching the soft tissues from the bone of forming the flaps which are to close the gap, is to make the lateral incision, on either side of the cleft. The incision should be close to and parallel with the alveolar margin and extend from a point opposite to the last molar forwards to the canine tooth. The flaps may now be separated from the bone with an instrument with a somewhat broad and rather bluntly curved inwards. The separation should be commenced from the incisors and proceed inwards, terminating when the edges of the gap is attained. The separation can be quickly effected without much risk of bruising the flaps; or the soft



is most apt to supervene and materially interfere with the union requisite to the success of the operation.

The flap detached from the bone should consist of all the soft tissues covering the latter—mucous membrane, areolar tissue, and periosteum. If the soft tissues are separated from the bone in the manner recommended, on the dead body, it will be found difficult (and we conjecture equally difficult in the living subject) to peel off the thick mucous membrane without the subjacent periosteum. The operation which Professor Langenbeck proposed, and to which he gave the name of ‘the operation by muco-periosteal flaps,’ appears to be identical with the method of operating introduced by Mr. Avery.*

Mr. Annandale † has succeeded in closing fissures of the hard and soft palate by the adoption of lateral incisions merely, and without the division of the muscles of the velum, either in the manner adopted by the author or that which has long been recommended and practised by Sir W. Fergusson. But in the cases related by Mr. Annandale he carried the lateral incisions back some way into the substance of the soft palate. His results were perfectly satisfactory—so much so, that his method of operating in extreme cases of fissure may probably, upon more extensive trial, supersede those already described and at present most commonly adopted.

When the soft tissues have been effectually separated from the bone, the flaps should fall inwards and downwards and meet in the median line, without the slightest traction being requisite. If the edges do not readily touch, the flaps have not been sufficiently detached from the bone on one side or the other; and care must be taken to ascertain the point which prevents their sufficient descent, and this should be freely liberated; nothing must be allowed to interfere with the free approximation of the edges.

When the fissure is extensive, and more or less complete, the free separation of the soft tissues from the bone is amply sufficient to allow the margins of the flap to meet easily in the median line; occasionally they will overlap; in which case, the edges require to be carefully adjusted. They should be carefully pared with a sharp knife, so that if a little ragged, they

* See Prof. Langenbeck's treatise entitled *Weitere Erfahrungen im Gebiete der Uranoplastik mittelst Ablösung des mucös-periostalen Gaumenüberzugs*. Berlin, 1863.

† *Edin. Med. Journ.* 1865, vol. x. p. 621.

made even; and the possibility of any mucous surface being insinuated between them be avoided, and the contact


FIG. 247.



owing from a model of a complete cleft of the palate (St. George's Hospital Museum).

entire and fresh raw surfaces be insured. The sutures to be passed as already described. There is often some difficulty experienced in passing them through the anterior part

FIG. 248.



the same model cut in two to show the perpendicular direction of the sides of the gap.


the palate; but care and patience will usually overcome any difficulties which may present themselves.

When a fissure of the hard palate is *complete*, or nearly so,

F F

the sides of it will be found to approach more or less to the perpendicular in their direction, as is shown in Figs. 247, 248. The result is, fortunately, that when the soft tissues are sufficient

FIG. 249.



Drawing from a model of a partial fissure of the palate (St. George's Hospital Museum).

detached from the bone, the flaps fall inwards and downwards and readily meet in the median line; there is always a

FIG. 250.



The same model divided, showing the much more horizontal direction of the sides of the gap.

material to permit of their arching over the defective roof, when they are once liberated from the bone.

But when the fissure in the bone is confined to the posterior

half, the portion of the palate in front of the fissure, and whatever portions of the palate-processes exist as its sides, will be observed to approach, in curve, the arch of a perfect palate as seen in Figs. 249, 250 taken from the model of a cleft of this kind; the bony fissure will often be broad, and rounded in front; and the membrane covering the bones very thin. In such cases, if the soft tissues are merely detached from the bone at the sides and front of the fissure, there is not much spare surface to close the gap; consequently the edges of the flaps do not meet in the median line, so readily as occurs when the tissues are detached from the more upright sides of more complete fissures. Often, however, a fissure of the posterior half is narrow, pointed in front, and its sides covered with thick velvety mucous membrane. Under such circumstances little difficulty is experienced in getting the edges together.

When cases with broad fissures of the posterior half of the hard palate are operated on, it is generally necessary to make lateral incisions through the soft tissues, parallel with the edges of the fissure, and close to the alveolar edge. The soft tissues between the incisions and fissure must then be separated from the bone; but great care should be taken not to injure the anterior or posterior extremities of these flaps, or in any way damage their connection in front and behind; the flaps must then be fastened together with sutures. Sometimes it will be found desirable to plug the incisions with a little soft cotton: this helps to push the flaps towards the median line, and takes off some of the traction of the sutures; this traction being almost unavoidable in instances in which the gap is an unusually broad one.

Such cases as the latter are the most unsatisfactory to deal with. In some instances it will be found best to close the soft before attempting the hard palate; for experience has taught us that the continued action of the muscles of the soft palate, closed by operation, has always a tendency to contract any opening left in the hard palate.

If the gap left should appear too large to be benefited by subsequent operation, it would be better that the patient should use an artificial palate; at any rate, until from lapse of time the ultimate benefit of the first operation can be ascertained.

Frequently happens that, under the most favourable circumstances, and with the greatest care on the *part of the operator*, a

in diameter ; in time the plug will become useless replaced by a larger one, and this will be followed increase in the size of the aperture.*

If it should happen that a patient, with hard alveolus and palate, should not have had the lip childhood, it will, in all probability, be found that the alveolus is considerable,—sometimes sufficient point of the finger to be passed through it from nose. Under such circumstances the first care should be to the lip. It will be found, subsequent to the healing that the action of its muscles, in a very short time approximated the edges of the separated alveolus so as to make them touch ; though, in consequence of position of the mucous surfaces, the chink between appears to be obliterated, unless the mucous membrane removed.

Occasionally small orifices are met with in the palate congenital, or more frequently the result of some syphilitic condition, terminating in necrosis and a portion of the bony palate. In either condition difficulty will usually be encountered in attempting

* The author would not be doing justice to the memory of Dr. Avery, in omitting to mention that he was the first surgeon to close entirely a complete cleft of the palate. Since then the operation has frequently and successfully performed in England and on the continent. Dr. Warren, of Boston in North America, the profession is credit to him having advocated the closure of the hard palate by opera-

In all instances, lateral incisions are requisite; and division of the operation may be necessary, even more so, before entire closure be effected. But, under any circumstances, should the opening be not ultimately filled up, it should generally be greatly reduced; so that the patient will be enabled, with the aid of an artificial palate, to secure himself from the discomforts attendant on a perforation of the

soft palate are usually, if not always, the result of a ulcerative action. Generally they close after a time, without the assistance of an operation. An opening, the result of a chronic ulceration of the soft palate, and of the size of a half-piece, in a patient under the care of the author, continued closed entirely, after cicatrisation of its margins occurred, without operation or any surgical treatment. However, such an opening ceases to contract, and holds its ground in respect of natural obliteration, but promises to be perpetually patent, its edges should be pared, lateral incisions made, and the margins brought together and retained in apposition by sutures.

The amount of blood lost during the performance of an operation on the palate will vary to a very great extent in different cases. But usually much more is lost in the process of detaching the palate from the bone, than in the operation on the soft palate. The anterior and posterior palatine vessels sometimes bleed profusely after division; so much so, that occasionally it is necessary to make pressure for a few moments or more, with the finger; especially on the anterior, which generally bleeds more profusely than the posterior vessels. Iced or cold water should be used frequently, to wash out the mouth during the operation, when hæmorrhage is free. The loss of blood in the operation, under the experience of the author, has been attended by evil consequences. Women are often apt to faint from a small loss of blood, when added to the fatigue of the operation. In one instance which occurred to the author, the operation was obliged to be completed while the patient was lying on the floor.

It is advisable and often necessary, to administer stimulants generally to the patient during the operation, not only to counteract or obviate the effects of the loss of blood; but also to encourage one who flags in determination. The patient should always be provided with his usual meal

previous to the operation. For some days after, *he must be plentifully supplied with nutritious fluid and soft food*: strong beef-tea, bread and milk, tea with half milk or cream, strong soup, and beer or wine. From the first hour after the operation, strict attention to this rule forms an important element in the satisfactory union of the flaps.

It might be supposed by some, who have not experience in this class of cases, that, after an operation on the soft palate, the effort of swallowing would be an effective impediment to union, in consequence of the action of the muscles of the palate; but it must be borne in mind, in the first place, that, prior to the operation, the palate-muscles had no power in themselves to assist in the act of deglutition; they could not close, between the ori-nasal cavities, on the passing food; in fact, that the process of deglutition is initiated in the back part of this or-nasal and pharyngeal cavity, by the extremely educated action of the superior constrictors of the pharynx. In the second place, the divided, and to an extent crippled, muscles of the palate are naturally passive in deglutition until reparation is established and complete. So, in practice, it is found that, as long as a patient is restricted to soft or fluid articles of nutriment, union is not retarded by the mechanical action of deglutition, however often he may be fed in the day. But if all diet be withheld, the patient's strength, already lessened by loss of blood, becomes lowered to such an extent, that reparation is rendered doubtful, and union of the parts is retarded, if not ultimately prevented. All hard food should be avoided for a week or ten days; such as solid meat, potatoes, dry bread, &c.

The sutures should not be permitted to remain in the flaps many days. Usually, in the soft palate, the two lower ones should be removed on the second day; and if there be four sutures, one of the two latter should be taken away each succeeding day: that which is highest, to be last removed. It has been supposed desirable to retain the sutures in the flaps for a longer period; but in each fresh case that comes under our notice, it becomes more evident, that if union is satisfactorily established, the retention of the sutures beyond the third or fourth day only tends to render it less effective: and if the condition of the edges of the approximated flaps is not satisfactory, and their union is not firm, the presence of the sutures will aggravate rather than diminish the evil.

The patient should not be allowed to speak until union is complete, and the sutures are removed. He should be provided with materials for writing all his wants; a small slate and pencil will be found most convenient for such a purpose. It is not requisite to confine the patient to bed after the operation; but in some instances individuals prefer to lie still for a few days, especially if they feel weak from the loss of blood. The condition of the tongue, before or after the operation, must not be taken alone as indicative of the patient's condition. All persons afflicted with cleft palate are liable to a dry, rough, and often coated tongue—the result of the constant passage of air in respiration over the surface opposed to the fissure. The injury, from the operation, to the soft tissues of the roof, generally sets up a slight degree of feverish excitement, with a very coated tongue for a few days. This condition of tongue must not be taken to indicate the want of purgative medicine. In an instance under the author's notice, this state of the tongue was mistaken for a symptom of some great general disturbance of system, and to remove it, two consecutive doses of calomel were administered within four days of the operation: the patient became salivated a week after the operation, and the wound, which had been most satisfactorily united, gave way entirely under the salivation.

We cannot too strongly urge the point that, previous to any operation for the closure of clefts of the palate, the surgeon should satisfy himself, that the patient is in a condition of health to justify the expectation, that '*union by first intention*' will follow the adaptation of the raw edges of the flaps to each other. The success of the operation depends on immediate union of the edges: if they do not unite at once, there is no hope that they will do so secondarily, when in the stage of granulation. We ought to be most particular not to recommend an operation, should the patient show any sign of disordered health or defective power—pustules, herpes, excoriated lips or nostrils; in fact any indications, trivial in themselves, and perhaps of no import in the balance which influences other operations, should in cases of cleft palate be decisive against operative interference until a clean bill of health can be written for the patient.

particulars of the following case illustrate the importance of such a pre-
- The author operated on a young lady ten years of age and apparently
-et health, for fissure of the soft palate. The operation was most satis-
-f in its immediate results, the *gap being entirely closed*. The parts

and speech is effected by the operation, in patients fully undergo it. Dr. Warren states: 'There is great facility of speech, which, so far as it has been in watch patients at a distance, is constantly improving. A man was present at a meeting of the Boston Society for the Improvement of the Deaf, about two years after the operation, who was difficult to discover the least imperfection in his speech, which it had previously excluded him from society.'*

The author has found that in many cases the improvement has been most marked and satisfactory; and, in some, sufficient to render what was unintelligible readily understood: in one case, a stranger could not have detected a defect in articulation, on the delivery of a long sentence, years after the closure of the entire palate. In all cases, however, the requisite, and much pains must be taken by the patient to acquire the power to articulate clearly such letters which, without a roof to the mouth, the tongue cannot properly articulate. It becomes, in fact, a task to the patient to learn how to articulate correctly and distinctly a new dialect. The position of the upper incisors is often very defective in these cases: always more or less so when the cleft is closed through the alveolar ridge. Such cases should be referred to the hands of an experienced dentist, when the surgeon has completed his work; for this irregularity of the teeth is a complete impediment to the improvement in speech, although the palate may be most satisfactorily

ve devoted attention to the subject it refers to, were he, clusion, not to mention the names of Roux, Cloquet, er, Mütter, Pancoast, and Dieffenbach, with those of i, Avery, and Fergusson. Each has so far aided in turn, rove this once apparently difficult and almost hopeless on, that it can now be confidently looked upon as certain t closure of the most extensive cleft, and almost certain ove thereby the most defective articulation.

DISEASES OF THE LIPS.

congenital defects of the lips and of the mouth, and the ent applicable to each, are described elsewhere in this as also the diseases of the tongue, and their treatment. following section will be described the diseases of the ws, and floor of the mouth.

diseases of the lips are usually so marked, that they affect the appearance of the individual; and they are o serious, that they demand the most careful attention of geon.

le '*cracked lip*,' when superficial, may be the result of ntinued cold weather, acting upon a constitution some- at of order. The crack is usually near to, or at the middle ip; is not deeper than through the mucous covering; is xcessively painful when stretched, and readily bleeds if es be accidentally and suddenly separated. It may gene- relieved by slight attention to constitutional measures; rative, with aperients, if requisite, or such other remedies ondition of the patient may indicate. The application tic will often relieve the pain at once; and some simple nstantly applied keeps the surface soft and supple, and s its edges from getting dry and tender.

ld such a slight but painful crack be neglected in early d no measures taken to relieve the patient of this trouble- ccurring sore, it will often happen that the crack deepens, comes more permanent in character: it becomes located part; and appears sometimes so deep, that it might be ed that the middle line of the lip had been grooved out eration; a very unsightly furrow, occasionally of con- le depth, is thus formed, to the great deterioration of the al appearance of the individual. Under such circum- s, the relief, which in *former years would have been readily*

obtained, is now no longer so available ; for though the ulcerated base of the furrow may heal under judicious treatment, it will leave a chink or gap in the middle of the lip ; and nothing short of paring its edges will remedy it. Under these circumstances, an operation may be recommended and undertaken, should the patient be in any degree desirous to have the disfigurement remedied. The edges when pared should be brought together with one or two fine sutures.

Slight superficial fissures, indicated by their white lines, are occasionally to be observed on the upper and lower lips, in children as well as in adults ; presenting an appearance as if at some former period slight ulceration had existed, though now healed and sound. If the teeth in the child be examined, they will be usually found to mark the characteristics of congenital syphilis ; and occasionally other symptoms of hereditary taint will present themselves, or may be detected as having existed. These marked fissures of the lips, have been constantly observed to be associated with an inherited stain.

Cracks of a severe character are generally observed in the under lip ; but fissures are not at all uncommon in the upper one. These usually occur in children, and are mostly associated with, if not always indicative of, a scrofulous constitution ; enlarged cervical lymphatic glands are constantly present in such children. Such a condition of lip is often obstinately persistent, and can only be combated by such measures as are advisable for the improvement of the general health. If very painful, an occasional application of a strong solution of nitrate of silver will relieve the extreme sensibility, for this will be sometimes very distressing in the movements of the lips : indeed so much so occasionally, that it interferes with the ready and comfortable use of the lip in conversation or during eating.

Fissures or ulcers of the commissures of the lips, or of their inner surfaces, should be very carefully looked to. In children they are usually transient, innocent, and associated with some passing constitutional disturbance ; or more obstinate, and grafted on a strumous habit. In adults, or after the period of childhood, such ulcerations or cracks about the commissures must be regarded with extreme care and some suspicion, under apparently the most innocent and unsuspecting circumstances : as a rule, they are the results of former syphilitic taint, and form one of the numerous varieties of secondary syphilitic affections ; other indications, confirmatory of such contamination,

will seldom be found wanting; and the condition of the part will be subservient to the treatment adopted for the general condition of the patient.

A troublesome, and often a recurring evil, which the practitioner has to combat, is the tendency, in some persons, of inflammation of the mucous membrane of the mouth to run into aphthous ulcers. Such ulcers sometimes become deep, and may be some days before they assume any tendency to heal. They occur on the side of the tongue, the lips, and frequently on the frenum of the tongue. They are always painful, sometimes exquisitely so; and their extreme tenderness may last many days, though the ulcer itself be not more than the flat surface of a split pea. The ulcer is generally ashy on its surface, as if covered with a superficial slough of mucous membrane; the surrounding membrane is red, and slightly swollen.

These ulcers, whether aphthous or more extensive, are generally attendant on some constitutional disturbance. In some individuals they constantly recur; and in such persons, either some peculiar defect in health, or perhaps some local climatic or other influences, may be their exciting cause. The pain of the ulcer can be at once relieved by touching the surface gently with a point of nitrate of silver. It is as satisfactory as it is unaccountable, to observe how immediately the application of this caustic removes the often exquisite pain—pain which may render the patient almost unable to eat or talk. The ulcer usually ceases to be felt after the caustic is once applied, and heals without further trouble; a second application is rarely requisite.

The constitutional measures must be guided by the circumstances of the case, as the recurring form of ulcer is usually associated with a delicate state of health. Every attention must be given to the improvement and invigoration of the system. But medicine will frequently fail to produce that which change of air, soil, scene, and circumstances effect in a short time, and as frequently in an unexpected manner.

Vascular growths of the lips, &c.—The tissues of the lips and cheeks are favourite localities for the commencement of nævi, or vascular growths, whether of arterial or venous constituents. These masses are unsightly at best, and are usually brought to notice of the surgeon at an early period for treatment. The mode of the treatment is necessarily various; and the variety

in treatment is dependent, in a measure, on the amount of the diseased structure. The *nævus* may be but a small red superficial patchwork of vessels; or it may be a large soft spongy cluster of veins, spreading over a very large section of one side of the face. The small arterial bright vascular patches are most common; though often a very large spongy venous *nævus* on the cheek may be observed in very early life.

When a small arterial *nævus* occurs on the margin of the inner surface of either lip, and is entirely superficial, it may be disposed of in several ways; but the application of the ligature is the most rapid, the most certain, and not more painful than any other. A needle passed through the base, and a fine ligature tied under the needle tightly round the mass, rapidly and surely effects its removal. In the course of a few days the slough will have separated; and in a few more, the wound, which at first may appear formidable after the removal of the slough, quickly contracts and cicatrises.

If the diseased mass of vessels is larger, and dips into the substance of the lip, and has not spread laterally, but involves more in depth than in breadth, it may become a question whether it would be better to cut out the diseased mass, as in an operation for cancer of the lip, or to destroy it by the repeated introduction of ligatures through its substance, tied tightly, so that the mass be cut up into several pieces. If the mass involves the whole thickness of the lip and the corresponding surfaces of mucous membrane and skin, and, though extending towards the root of the lip, does not spread laterally, and can therefore be readily removed by incision, there are reasons for such treatment being preferred. The disease is at once got rid of; the relief is effectual; the operation is simple; the recovery is quick; and the scar left after union is but slight. The only precaution requisite in the performance of such an operation is to take care that the incisions are made through healthy tissue, and beyond the diseased vessels. The margins of the wound are to be brought together, as in the operation for hare-lip.

But rather than sacrifice skin or mucous membrane, if either or both are much implicated, it is far preferable to have recourse to ligature: if the disease spreads laterally, or extends in any degree into the cheek, ligature alone must be used.

Under the circumstances of ligatures being applied to such *nævi* of the lips and cheeks, it should always be borne in mind that scar must be avoided as much as possible, and mucous

membrane saved as much as can be. The one is an eye-sore; the loss of the other is apt to produce distortion of mouth and a contracted cavity. Therefore, in using a ligature for nævus of the face, it is best to pass the thread entirely subcutaneously, and to tie it at the common point, where it entered and emerged from the skin, so that no portion of the latter be destroyed. One ligature may be sufficient in slight cases; but in others several may be requisite to completely obliterate and destroy the diseased structure. The object to be kept in view in introducing the ligatures, is to effect the entire division of the diseased mass of vessels in several directions,—that, in fact, they be cut into several pieces; so that if, in the first instance, a ligature be introduced round a portion of the tumour, in the second operation one should be passed across; that each ligature should act on a fresh part; that ultimately the original mass may be divided into so many portions as will insure the consolidation of all the diseased tissue.

If the ligature be introduced on a needle at one point, and carried partly round the mass subcutaneously, and brought out at some distant point,—again introduced and carried in or through until it reach the original puncture, and the ligature then tightly tied at this opening,—no skin need be destroyed, nor mucous membrane cut or damaged. Each ligature should be tied as tightly as possible. It will soon cut its way free. The ends of the ligature should not be left too short; otherwise they may become buried in the wound in the subsequent swelling of the parts, and probably would cause suppuration to a greater extent than desirable, or even occasion some difficulty or delay, from being retained in the wound. The more tight the ligature is tied, the more speedy will be its release from the tissues which it grasps.

Vaccination should only be tried in very small nævi; the introduction of caustics, setons, the injection of perchloride of iron, each have their advocates; but none of these secure more effectual obliteration of the nævus than the ligature: they often fail to secure it as rapidly; they often set up more irritation than is requisite for the purpose in view; they sometimes cause the skin to slough; and they are more painful in adoption, much as, being slower in action, their application must really be often repeated before a satisfactory result is insured. galvanic cauterization has often been employed by the author; after a great number of experiments in the treatment of

such cases, he is satisfied that subcutaneous ligature will invariably be found quickest in action, most effectual in result, least objectional as regards scar, and, so far, less painful than any other kind of treatment. This is the case whether a nerve of the lip be large or small; unless the case be more suitable for the knife.

Cancer of the lip may be said to be confined to the lower lip, it is so rare in the upper: it is certainly more frequent in males than in females, the proportion being relatively large in the stronger sex.

The characteristics of cancer affecting the lip deserve particular attention. As a rule, we may say it is found to consist of the epithelial variety.

‘Epithelial cancer has its primary seat, with very rare exceptions, in, or just beneath, some portion of skin or mucous membrane. Its most frequent locality is the lower lip, at or near the junction of the skin and mucous membrane.’* Commencing sometimes as a small warty growth on the lip; sometimes as if the mucous membrane were excoriated, with the excoriation resting on an inflamed and thickened base; sometimes as a small, indolent tubercle: the condition, which does not at first excite suspicion, by degrees is seen to alter and the growth to increase. The wart grows in breadth and thickness; the excoriation becomes deeper, and rougher on its surface; the tubercle peels and ulcerates, scabs and peels again; until, sooner or later, the persistence of the mass, or the heat and pain of the part, attracts the more serious attention of the patient, and he then seeks for relief.

It is unnecessary here to enter upon the question of the pathology of cancer, or of its cachexia; the subject having already been discussed in the essay on CANCER, Vol. I.

We pass on at once to the consideration of diagnosis and treatment.

The diagnosis of cancer of the lip may at first sight appear simple enough; and yet a certain degree of caution must be observed, ere an opinion be definitely arrived at, as to the precise character of a hardened sore, or of a raised tubercle of the lower lip. It has occurred to the author, and no doubt to many others, to point out, that a previously supposed cancer of

* Paget's *Surgical Pathology*, vol. ii. p. 412.

lip was but a chancre : a chancre of the lip has been mistaken the more serious malignant sore. The surface under both conditions may be superficially excoriated ; the lip may be thickened at the part ; the sore may rest on a hardened base ; and there may be the additional suspicious circumstance of tendant enlarged glands in the submaxillary region. It must, however, be borne in mind, that cancer of the lip is somewhat slow in its progress ; that the cervical glands do not usually indicate the more grave implication of the constitution at an early period : whereas in chancre the glandular enlargement would be apparent in six or eight weeks, at the latest, from the first appearance of the sore ; and probably, if no specific treatment had been adopted, other evidences of secondary symptoms, such as eruption, sore-throat, &c., would be manifest within six weeks of the first infection. As a mistake in diagnosis between cancer and chancre of the lip has often been known to occur, a few remarks, in passing, on that subject will probably not be thrown away.

Cancer of the lip is a disease of advancing, if not of advanced life ; but man is often depraved, and experience teaches us that age need not be taken as a bar to the occurrence of a chancre of the lip.

A gentleman past the meridian of life, and on whose head time had more than snowed a few snowflakes, applied to the author with a condition of lip which had excited suspicions as to its cancerous character. The appearance of the sore itself was extremely deceptive ; scattered and isolated glands, slightly enlarged, and hard, were to be felt in the submaxillary regions ; and his position and circumstances in life rendered it most improbable that a chancre could have been contracted on such a part. This sore was found to be accompanied by a commencing faint lichenous eruption. The patient was put under a mild course of mercury, and rapidly recovered ; the hardness at the base of the sore entirely disappeared.

The characteristics of the two affections may thus be summed up : chancre of the lip may occur at any period of life, and is as often seen on the upper as the lower lip. Youth is the most probable period of the infection ; and we say, with regret, it is most frequently observed in women : it sooner or later forms a superficial sore, raised on a base of an almost cartilaginous consistence : the surface appears more like an excoriation than an ulcer ; or the surface may have cicatrised and the hard base remain. The glands of the submaxillary spaces enlarge at an early period, some six or eight weeks after the sore commences ; and though hard, small, and scattered

usually, they sometimes attain a large size. In one instance which came under the author's notice, the glands on the right side were very much enlarged, and on the inside of the right half of the lower lip was a large hard mass of cicatrix, the original sore of which had entirely healed some weeks previously. Secondary symptoms will sooner or later appear, unless early treatment be adopted.

Cancer of the lip is a disease of advanced life; affects the lower lip mostly; is most frequent in men; does not progress uniformly, and usually slowly; does not implicate the absorbents at so early a period as chancre; does never thoroughly heal up and cicatrise like chancre. The disease is usually of the epithelial variety of cancer, and varies a good deal in its method of commencement, in its progress, its growth, and its appearance; while the characteristics of chancre are uniform in most persons. Cancer always contaminates the absorbents, if allowed to run its course unchecked; ultimately it destroys, locally, all tissues in contact with its surface; the whole lip may be affected; large cancerous masses exist, at the same time, from the symphysis to the clavicle; and death ultimately follows.

The treatment of cancer of the lip usually resolves itself into removal by knife, or destruction by caustic. In the opinion of the author, it should entirely resolve itself into that of the knife; and the earlier such treatment is adopted, the better the prospect of prolonging life.

The surgeon having decided to remove a cancer of the lip with the knife, the patient should be seated in front, with his head steadily supported by an assistant; or he may lie on a sofa. If the disease occupy but a small portion of the lip, the mass may be removed entirely by a V-shaped cut through the healthy structure. The lip may be transfixed with a thin straight knife, and then cut upwards on either side of the diseased tissue; an assistant should hold the flaps as they are liberated from the diseased part. The flaps are to be brought together with pins and the twisted suture, or by sutures alone; care being taken that the margins of the prolabium, or mucous edge, be accurately adjusted. One, two, or three pins may be requisite; some surgeons prefer sutures without the aid of pins; and silver sutures answer the purpose very satisfactorily. If pins be used, as soon as they are fixed by suture, the sharp-pointed ends should be cut off, and the remaining ends so protected that they be not readily caught by the dress, &c. of persons in

attendance, otherwise the patient runs the risk of being accidentally very much hurt, for want of a little foresight. In all these operations the pins or sutures should be removed at the end of forty-eight hours at the latest, and the wound supported by adhesive plaster.


If a large surface of the lip be affected, and the disease be superficial, it may be readily removed by a semicircular sweep with the knife, or a cut with a pair of curved scissors; in either case the opposed edges of skin and mucous membrane should be brought together with sutures. In the incisions through the lip, the labial arteries bleed freely. With the aid of the pins and twisted suture the hæmorrhage is readily and effectually checked; with the semicircular incision a ligature or two may be requisite.

It will save some little inconvenience, and also pain, to have the sutures well oiled before they are twisted round the pins: by using this precaution, it will be easy to remove the sutures subsequently, as they are readily detached from the pins; when not oiled, they adhere to the pins and to the skin, and always occasion pain when removed.

The use of caustics, in cases of epithelial or other cancers of the lip, is but waste of time, and an unnecessary infliction of suffering; provided the disease be not too far advanced for removal by the scalpel. The knife should always be used when practicable; caustic, only when the disease is too far advanced for removal by the knife. But even then it is doubtful whether any application of caustic, by retarding growth or destroying material, compensates for the pain, often exquisite suffering, which is inflicted by its use. It is a question worthy of consideration, whether caustics of any kind act beneficially in any cancerous affections. If caustic be decided on, the most efficient is the chloride of zinc, applied in the form of a paste.

The advantages of operating early can only be appreciated when compared with the more rapid results of a case which is allowed to run its course unmolested. But under the most favorable circumstances, we can only expect the operation of removal to relieve for a time; the disease usually returns, and generally shows itself in the glands of the submaxillary region.

in time increase to a great size; the skin over them becomes and sloughs; an ulcerated and fungoid surface discharges profusely, and often bleeds largely: and life is thus lost. We cannot here more fully discuss the advan-



lower lip. They sometimes increase to an
On the margin of the lip they seldom project
membrane; the skin is usually free. They
raised from the surface; covered by very
usually semitransparent, but occasionally son
and of a venous hue, as if veins traversed
cavity.* They contain either viscid clear m
contents of the cysts termed ranulæ; or so
fluid of thick consistence. They are gener
their surface and shape may be irregular. '
large; though this may be the result of local
is usually required to remedy the evil before
inconvenience, or becomes unsightly.

Such cysts are usually innocent in their cha
cur in the part when effectually treated; nor a
unless of that variety alluded to as connected
These cysts are often not larger than small
to the size of a small walnut; painless; in
handled; and only inconvenient from size
They occasionally remain stationary for life;
small in size, and frequently discoloured; the
ones have the greatest tendency to increase, i
the secretion of a gland, which probably, ha
obstructed, has given rise to the formation of

The treatment to be adopted in such cases

soon subsides, and the patient is then able to eat and talk usual. No surgeon should be satisfied with simply puncturing such cysts; a puncture, or even a small incision, will usually close before the cyst is obliterated; and in a few days it becomes as large as it was originally. Even after the application of nitric acid, unless the opening be free, and care be taken to keep it so, until the cyst be entirely destroyed, the accumulation of fluid is very apt to recur. It is a good precaution to cauterise the edges of the opening with nitric acid for a few days consecutively, to insure its not closing until suppuration is fully established, so that the cyst be obliterated. Occasionally a cyst in the lip is complicated with some solid growth. Such growths are usually somewhat of a glandular nature,* and are favourable for removal. These tumours are not common. They resemble, to some extent, the mammary glandular tumours in their existence and character; or 'they may appear intermediate between the foregoing and those tumours which are found over, near, the parotid gland, and consist of mixed glandular and fibrous tissue.'†

Dr. Paget refers to a case removed by himself from the upper lip of a man; he describes a second case, which occurred under Mr. Lloyd's care, in a man who had a tumour in his upper lip for twelve years, when it was removed. The tumour was firm, slightly lobed, yellowish-white, smooth. In general it resembled the mixed tumours over the parotid; but its minute structure presented as perfect an imitation of lobulated or acinous gland-structure as a mammary gland.†

Cysts of the mouth may be congenital, or the result of obstructed ducts; ranulæ; or possibly bursæ between the muscles of the tongue.

Congenital cysts of the mouth are not frequent, though they are sufficiently often to be familiar enough to surgeons. Mr. James Hawkins long ago drew attention to their real character. Usually such cysts are single, and may increase to a large size; they may be multilocular and numerous, and occupy a considerable portion of the floor of the mouth. Whether single or multiple, they will usually be found between the lower jaw and the tongue; if large, pressing the floor of the mouth forwards upwards, and the submaxillary soft tissues outwards and forwards.

* A specimen, see Museum of St. George's Hospital; referred to also by Paget, vol. ii. p. 73.

† Paget, vol. ii. p. 262.

‡ Ib. p. 263.

An infant three weeks old was sent to the author by Mr. Frank then assistant-surgeon of the Second Life-Guards. The mother found child suckled with increasing difficulty during the last few days, in consequence of a swelling on both sides of the floor of the mouth, which was pushed upwards by the tongue, and had much widened the lower jaw. It had also encroached so as to press somewhat on the larynx and pharynx; respiration was impeded; deglutition was slow and difficult. Under the tongue, on both sides of the mouth there projected a lobulated irregular cystic mass; portions of which, especially on the left side, were so prominent, that they were pushed upwards by it, and the tumour bulged out between the floor of the lower jaw, to such an extent, that the infant could not close its mouth. Irregular lobulated swelling projected on each side of the throat, below the lower jaw.

The cysts projecting between the tongue and lower jaw were pellucid, the membrane covering them pale-coloured, and free of vascularity on surface.

A seton passed through a cyst on the left side soon allowed it to thin; purely transparent fluid oozed out through the punctures. On other threads were passed through some of the more prominent cysts, which were punctured. The partial reduction of the sublingual swelling enabled the infant to partake more freely of food, and for some days a slight improvement was observed in his condition. This was but transient; evidently deeper cysts became larger; gradual increase of dysphagia and dysphagia ensued; and the infant died, greatly emaciated, the sixth week after admission to the hospital.

On examination, it was found that this cystic formation involved the tissues between the mucous membrane of the floor of the mouth and the covering the submaxillary region; so that not only did the mass project upwards but downwards on both sides, in the submaxillary spaces. On a section of the tongue, the whole muscular structure was seen studded with cysts, millet-like in some parts, rather larger in others; some single, others multiple. The larger ones were generally multilocular, with bands running around their walls.*

Other forms or characters of cysts are occasionally observed in the mouth, sometimes between the gum and the cheek.

Mr. Paget describes a case of this nature, which occurred in a child in whom a soft elastic swelling pushed out the thin mucous membrane of the upper jaw, producing externally an appearance somewhat similar, at least to distension of the antrum. An incision into the cyst allowed the escape of nearly an ounce of turbid brownish fluid, containing crystals of cholesterine.

Cysts connected with defective development of teeth, 'epithelial cysts,' usually confined to the alveolar region of the maxillæ, are not uncommon. Their history and treatment are related by Mr. Salter, and will be found in the paper on the subject in this essay.

* The preparation is in St. George's Hospital Museum.



Obstructions of mucous follicles occasionally give rise to the formation of mucous cysts, already spoken of; the obstruction of the submaxillary or sublingual ducts may be productive of the cysts termed 'ranulae.'

Again, congenital sebaceous cysts are occasionally found among the tissues of the floor of the mouth.


A young woman was admitted into St. George's Hospital, under Mr. Hewett's care, with a considerable swelling on the right side of her mouth and neck. The swelling bulged into the mouth; had displaced the tongue to the opposite side; and protruded downwards in the submaxillary region. The swelling was painless, but inconvenient from its size; it had been many years forming, and it was doubtful whether it was congenital or not. A free incision through the skin projecting into the mouth, allowed the escape of a large quantity of a thickish yellow fluid, most offensive and putrid in smell, similar to the contents of a suppurating sebaceous cyst. By degrees the cavity contracted and healed.

This patient returned to the hospital about three years afterwards. The cyst had again filled, and projected a good deal below the lower jaw. It was now opened by the author, in the neck, through the skin, below the lower maxillary bone; and a large quantity of offensive sebaceous-looking fluid was evacuated; the opening was then passed through the cyst, and retained several days, until the cyst had become perfectly contracted. The opening soon closed, and as far as could be ascertained, the cyst appeared to have been obliterated. The patient has since again shown herself; but promised to do so on the slightest recurrence of swelling.

The diagnosis of cysts of the mouth is not often complicated: usually, fluctuation is very evident; as frequently, the surface of the cyst projects into and on one side of the mouth, so that it may be seen covered only by mucous membrane; the walls are thin and pellucid; the contents visibly transparent, or slightly tinged from some accidental circumstance or individual peculiarity.

In other cases these cysts present more obscure conditions: they may lie deep among the muscles of the floor; or may press upwards and interfere with the movements of the larynx or trachea; or a sanguineous tumour may be mistaken for a cyst, though sanguineous tumours are exceedingly rare in these situations.

Such serous and mucous cysts as we have referred to, may be treated without hesitation, and without much danger. If very large, it may be as well to draw off a portion of the fluid, in the first instance; and subsequently have recourse to setons, or injections of iodine. But as in any operation in the region of the mouth and neck, oedema may occur about the root of the tongue



knife ; but the attempt to remove a large and in
cyst is hazardous and unnecessary. Sir W. Ferri-
case which illustrates the difficulties a surgeon
in such a proceeding, and the danger the patient

We would strongly dissuade the attempt to
cyst in the region of the mouth ; not only are there
obstacles to the entire removal of the cyst from
operation must necessarily be a tedious one, at
risk of great loss of blood. By laying the cyst
all probability it will close up from the bottom
become so much diminished, that ultimately a scar
or openings, a seton, or injection of iodine, will
ration. A silver wire may be used as a seton, draw-
more towards the integuments than the
contents, by tapping, are proved to be clear
if opaque, dark-coloured, thick, or purulent, is
the safest, the most effectual, and the least tedious
adopt.

A large cyst opened by incision in the mouth
contain purulent fluid, may not contract very
sequence of matter bagging downwards in the lower
part of the sac. Under such circumstances, an
may be made under the jaw. But as a scar in the
in women, is always objectionable, such an operation
avoided unless absolutely requisite.

dingly, care being taken to ascertain first the character of fluid contained ; for should the contents be blood, the treatment should be different to that described.

Mucous cysts about the mouth are rare. When present, they are usually seen on the lips, near their free border. Instead of incision, when deeper seated they should be treated with ligatures ; when superficial and accessible, they may be excised out, or included in a ligature, like a *nævus*.

Sum up the evidence on the history of mucous or serous affections of the mouth, we find that, in early life, they either singly or in clusters ; are generally of a congenital nature ; and are found most commonly on the floor of the mouth, or, in after life, on the surface of the gums. Venous or hæmorrhagic cysts are generally observed in the lips and on their surface ; are usually congenital, and frequently stationary ; they differ from other cysts. Sebaceous cysts, contain sometimes a mixture of epithelial, thick, white, and often cheesy matter ; hair growing from the lining membrane (which is of the character of cuticle) ; or loose hairs ; bone or cartilage these have no peculiar affection for locality, are most numerous in their selection of spots for habitation, and may be found in any part of the mouth or its immediate neighbourhood. They are usually round, small, and, if immediately under the lining membrane, appear like a white marble embedded in the tissue. Their inclination is to grow slowly ; so slowly in some instances, that years may elapse before their presence becomes inconvenient. These may all be of the congenital nature. Mucous cysts, *ranulæ*, cysts of the lips connected with portions of solid or glandular growth, some serous cysts, some generally single, will be found to commence after birth and at different ages. They usually occur before very early life.

Growths of the lips and mouth, of a non-malignant, *i.e.* benign character, are not of many kinds, nor do they frequently occur. Labial glandular tumours have already been mentioned to. Occasionally a fatty tumour grows among the folds of the tongue, or about the floor of the mouth. The growth is usually simple, though when deep-seated it may require the surgeon, until he cuts down upon the growth. The tendency is generally to protrude towards the skin, rather than towards the mouth ; the constant movements of the

tongue being usually sufficient to direct the mass toward the least resistant surface.

A fatty tumour need only be removed when of an inconvenient size. A free incision over its surface, below the skin, will generally allow of its being readily dissected or drawn out.

The author removed one for a lady, forty years of age, who had been for some years of its existence, under the tongue and jaw, on the left side. Cutting freely down upon it, the lobes of fat were found to dip deeply into the floor of the mouth, round the mylo-hyoid muscle; but the whole was removed.

The chief danger attendant on such an operation is exposure to inflammation and suppuration to the intermuscular space about the root of the neck. The lady alluded to, had for several days considerable pain and difficulty in swallowing, but was ultimately recovered.

A solid mass is sometimes observed in the floor of the mouth of a calcareous consistence. Such a mass is often contained in one of the salivary ducts; it is but a salivary calculus. Occasionally, from neglect or indifference, one has been known to increase to a size which would be usually considered an intolerable annoyance and a source of constant disturbance to the patient's comfort. It can be readily removed by a slight incision over it, and the removal of the mucous membrane.

The condition, in infants, known as 'tongue-tie' has been spoken of in the essay on Diseases of the TONGUE.

TUMOURS OF THE JAWS, AND THE OPERATIONS FOR THEIR REMOVAL.

There are few points in Surgery which require more careful consideration, in order to arrive at a correct diagnosis, than the variety of tumours which affect the upper and lower maxillary bones. We propose to consider these tumours in the following order, viz.:

- | | |
|-------------------|---------------------------|
| 1. Cartilaginous. | 5. Osseous. |
| 2. Cystic. | 6. Vascular. |
| 3. Fibrous. | 7. Mucous, or Myxomatous. |
| 4. Myeloid. | 8. Cancerous. |

The operations for the removal of tumours of the jaws will form the subject of subsequent consideration.

Cartilaginous tumours of the bones of the face are rare, and the upper and lower maxillæ may be said to be almost free from their attacks.

Mr. Paget alludes to but one well-known case, which occurred in the upper jaw of a patient in Guy's Hospital. Mr. Beaumont, Professor of Surgery in the University of Toronto, has recorded an interesting case in the *Royal Med. and Chir. Transactions*,* which occurred in a boy seven years of age, affected the lower jaw, and was successfully removed.

In the Museum of the College of Surgeons is a wonderful specimen of one of these tumours, which implicates the greater portion of the lower jaw; it had been some eight years growing, and when the patient died, exhausted by its ravages, it measured two feet in circumference.†

The symptoms and prognosis of enchondroma are the same, when it affects either of the jaw-bones, as in other parts of the body. These tumours of the jaws may be removed with much confidence of relief. The operation cannot be too early decided on; delay may be so far dangerous, that a very rapid growth, by a short postponement, may really make the operation very severe and extremely hazardous.

In the removal of the smallest cartilaginous tumour, care should be taken that every particle of the surface of the bone from which it is growing be freely removed. In the removal of the larger tumours, it will depend on the involvement of the bone, how much of it has to be removed. The greater portion of one side, or even the whole, of the lower jaw, or the whole of the upper maxillary bone, if involved, must necessarily be taken away. The disease will probably be restricted to one side in the upper jaw; in the lower, it may more readily involve the greater part of the bone. The prognosis after operation is usually favourable; but it should not be overlooked that enchondroma sometimes recurs, and that in other cases the tumour grows rapidly, and large cysts are developed in its substance. Both these forms of enchondroma approach in character to cancerous tumours.

Cystic tumours of the jaws are not very uncommon.‡ We undertake their consideration in this place, in order to mark particularly the distinction which must always, pathologically, be made between the secondary cysts of bone resulting from

* Vol. xxxiii. p. 243.

† Museum Catalogue, No. 1034.

‡ See Stanley, *Diseases of the Bones*, p. 268; also, *DISEASES OF BONES*, Vol. II. p. 227.

degeneration of cartilaginous tumours, and the primary cystic tumours which attack the upper and lower maxillæ. The latter are seen to originate in the substance or from the surfaces of the jaw, under a variety of unexplained circumstances.

Cysts of the jaws are sometimes found to be connected with a diseased condition of a tooth-fang; or, more frequently, with a diseased action attendant on the development of the tooth-pulp—'dentigerous cysts.' (See p. 363.)

Specimens of simple primary cysts of the jaws, commencing in the substance of the bone, may be found in some of the museums of the metropolis. They expand the bone more or less, so as to take with them portions for their outer covering or walls, the remainder of which will be generally made up of a tough membranous substance. The contents are fluid, serous, or gelatinous, and of different shades of colour. 'This disease is usually of slow growth; and there have been instances in which the tumour of the jaw, formed by it, has acquired a large size.'*

In the Museum of St. George's Hospital may be seen a preparation (Fig. 351) which well illustrates the prominent features of a simple cystic tumour of the jaw. The tumour was removed from a woman forty-five years of age. It occupied on the right side the situation of the lower jaw, and extended from the second incisor to the condyle of the bone: it formed a large globular cyst, which occupied nearly the whole of the side of the face. It extended downwards over the upper portion of the neck; and inwards, displacing the tongue; it greatly interfered with speech and mastication. The principal part of the growth was upwards and outwards towards the malar bone, which was somewhat expanded and partly absorbed. In some parts the tumour appeared of bony hardness; in others very elastic. The integument covering it was not discoloured, and there was no enlargement of the neighbouring glands. The disease had been noticed about eighteen years before admission, when it formed a small hard incompressible lump, just over the angle of the jaw: it gave no pain; and for a long time its increase was extremely slow. About six months previous to her admission, it commenced to enlarge very rapidly. The cyst may be seen in the figure to extend from the symphysis to the right condyle.

The parietes are partly osseous and partly membranous. The entire substance of the jawbone on this side has disappeared, and is replaced by the foreign growth. When removed, the cyst appeared to be divided into several cells, which contained a transparent gelatinous fluid; though in the dried preparation there are but few indications of such divisions. The tumour was successfully removed; but the patient died subsequently from erysipelas.

In the Museum of the College of Surgeons there is a specimen of a cyst of the lower jaw, occupying the greater extent of the right side, and projecting

* Stanley, op. cit. p. 267.

wards rather than upwards. It is oval in shape, and multilocular; the cysts were filled with a glairy fluid. The diseased portion of the jaw was successfully removed.*

Whatever the origin of these cysts, they appear, on the one hand, independent of any tooth-irritation; and on the other, independent of any previous cartilaginous deposit. The diagnosis in such cases is not beset with much difficulty; and even if a doubt

FIG. 251.



of a cystic tumour of the lower jaw. (From the Museum of St. George's Hospital.)

As to the nature of the tumour, it is at all times safe and sound to explore it with a fine trocar.†

In the treatment of these tumours, it is a milder and a more

Museum Catalogue, No. 1033.

Dr. Giraldès has described a form of cyst in connection with the upper jaw, which he believes to be of very common occurrence, and to be formed by a change in mucous glands naturally existing in the lining membrane of thetrum of Highmore (see his *Recherches sur les Kystes muqueux du Sinus maxillaire*); but no such origin can be ascribed to those which distend the body of the lower jawbone in the manner above described.

justifiable course to make a free incision into the cyst, and subsequently to trust to reparative action, than at once to have recourse to removal of the diseased mass.

If the cyst be large, or even if it be ultimately requisite to remove it in its entirety, the previous incision will in all probability have much reduced it in size; when the parts, if not in a better, will certainly be in no worse condition for removal.

In both the cases related removal of the tumours was accomplished without difficulty. Two cases are mentioned by Mr. Paget, in which incisions were employed successfully; but in neither case did the bone appear diseased.

In a case under the author's care in St. George's Hospital, the cyst occupied a large portion of the left side of the upper jaw. Its anterior wall projected under the cheek, and bulged out the side of the face. A free incision was made into the cyst from within the cheek. The anterior wall was found to be partly membranous, and partly consisted of thin flakes of bone. A portion was readily removed, so that a free opening was left for the escape of the contents, which were of a glutinous consistence and brownish in colour. At the bottom of the cyst there was found projecting into its cavity, the extreme point of the fang of the canine tooth. On the removal of the tooth, the fang was found to be partially necrosed. The cyst filled and closed in a short time without any further treatment. The patient was about ten years of age, and when he quitted the hospital there was no trace of the disease left, nor the slightest disfigurement.

In all cystic diseases of the jaws, careful examination should be made of the teeth of the corresponding jaw; all diseased ones near the growth should be at once removed. One of the most frequent causes of serious mischief, and still oftener of exquisite suffering, is the too long abode of defective and decayed teeth in their sockets. We feel satisfied that the greater the experience in the treatment of diseases about the mouth, the more exacting will the surgeon become respecting the removal of all useless, defective, or decayed teeth or stumps.

Fibrous tumours of the upper and lower jaw are by no means uncommon; and in their general features, in their growth, and in their varieties, correspond to fibrous tumours of other parts. 'The favourite seats of fibrous tumours of bone and periosteum are about the jaws.*' Such tumours may originate in the substance of the bone, and in growing, expand it, as a crust, over the outer surface of the tumour. In the Museum of the College of Surgeons there is a preparation showing a fibrous tumour of the jaw, with a thin shell of bone over it.† They may grow

* Paget, vol. ii. p. 145.

† No. 1045.

from the periosteum, and embed the bone in their surrounding mass. They affect equally the upper and the lower maxilla. They may appear early in life.

Mr. Liston removed a large fibrous tumour of the upper jaw, in a woman twenty-one years of age, which had made its appearance four years previously.* It grew on the outer side of the jaw, and was removed six months after its first appearance, when about the size of the end of the thumb. In eighteen months, a return of the disease, the size of a hen's egg, was removed, with a portion of alveolus. The growth reappeared in two or three weeks, and attained a large size in two years, which rendered necessary the removal of the whole of the superior maxillary bone. The patient recovered.

Fibrous tumours of the jaws are usually of rather slow growth; but their growth may in exceptional cases be somewhat rapid. Their usual history is one of progressive and persistent increase; perhaps more rapid in proportion to their increase of size, but, in the main, rather slow than rapid. The size some will attain is best illustrated by the collections in our various museums; and especially the preparations collected by the late Mr. Liston, now in the possession of the College of Surgeons. In some respects, as to the size which they attain, they appear to vary from fibrous tumours of other parts, and especially those of the uterus; for fibrous tumours of the jaws are not usually seen to expend their energies in growth; they are not seen to stay their course, to degenerate, or calcify; or perhaps something may be said to depend on their position, for this is such that it can never permit of sufficient increase, or life, to attain that end, without first having grown to such a size as would occasion a fatal impediment to deglutition or respiration. They are usually painless in their growth, and innocent in character, as compared with cancer; but their locality, as they grow, renders them formidable as to disfigurement, as to discomfort, and even ultimately to life, if they be not removed before such growth be attained. It is wise, therefore, to pass the verdict of early removal, when we have to pronounce judgment on such cases; the earlier the operation is performed, the less formidable will it be. We cannot hold out a prospect of arrest; but we may certainly forewarn as to the inconvenience of delay, and we may positively predict danger from increase.

In their structure, we cannot point out any distinction between the fibrous tumours which attack the lower, and those found

* Museum of College of Surgeons, No. 1046.

connected with the upper jaw ; but, practically, there are some important considerations connected with the history of the latter, to which we would especially draw attention ; and a knowledge of which is most essential to any operator dealing with such cases. Fibrous tumours of the lower jaw are invariably more or less globular in form ; usually they have a uniformly smooth surface ; their tendency is to protrude externally, more than towards the mouth : if originating in the substance of the bone, perhaps they may be found equally surrounding it ; if originating in the periosteum, perhaps most prominent on the corresponding surface. Fibrous tumours of the upper jaw are often far different in their outward character, in their surrounding relations, and in the tendency they display, from position, to spread in different directions. A fibrous tumour supposed to be attached to the upper jaw, may really have its origin from the base of the skull, the sphenoid, or the ethmoid bone ; and from either point make its way through the various foramina and fissures, and into the various fossæ in and around the superior maxilla. Instead of a globular dense mass, as in the lower, the tumour of the upper jaw may be lobulated ; or, from its very position, and as a result of its insinuation into these fissures and fossæ, it may be moulded, as it were, into a mass of pedunculated bodies, somewhat analogous to the bulbous roots of a tuberous plant ; or commencing in the interior of the antrum, it may by growth or position be pressed through its anterior wall, and puff out the cheek ; or through its floor, and push down into the mouth ; or into the nares, and plug up the lachrymal duct ; or into the orbit, and displace the eyeball ; or into all or several of these spaces, and produce corresponding deformity. So that the diagnosis of the origin of a fibrous tumour of the upper jaw is often as complicated an affair as that of the lower is easy ; indeed in some cases, until the upper jaw be removed, it will be impossible to determine to which portion of the skull, or bones of the face, the growth may be attached.

The following case illustrates the force of these observations. A man twenty-five years of age was admitted into St. George's Hospital, under the care of Mr. Prescott Hewett, 'with a large tumour, of an irregular shape, occupying various regions of the left side of the face.' In the cheek, it formed a swelling of the size of a turkey's egg, and filled up the greater part of the superior maxillary region ; the outline of the bone was perceptible to the touch in a few places only ; the zygomatic arch was much more prominent and more curved than natural, having been pushed forward by the tumour, portions of which could be felt under the temporal muscle. The diseased structure was also found, in the shape of small flattened bodies, at the lower part of the orbit, lying im-

ately underneath the conjunctiva, and apparently quite movable; the bones of the inner and outer walls of this cavity, as well as those forming its circumference, were not affected or displaced; but it was impossible exactly to make the state of the bones at the lower wall, owing to the tumours which were there; the eyeball was not more prominent than natural. Portions of the tumour growth were detected in the left nasal fossa, from whence a small round growth projected slightly, at times, into the pharynx. The tumour overlapped the front part of the alveolar process, and projected beneath the lip.

About six years previous, the patient had observed, what was supposed to be a dyspnea of the nose, which was easily removed; but some little time after, the cheek began to swell, and gradually the tumour commenced to show itself in the various other positions described. Its growth was painless throughout. A year previous, caustic had been freely applied, with the idea of destroying it; two large cicatrices marked the places of its action. At various times, there had been extensive bleeding from the nose; these bleedings had somewhat retarded the patient, who was of a spare habit and pale; this paleness being attributed to a loss of blood, which occurred shortly before his admission. It was intended to remove the tumour by the usual incisions for the removal of the lower jaw. The bones having been divided with cutting forceps, the superior maxillary and malar were easily tilted out of their place; when it was found that the disease was not connected with the upper jaw, but was behind it. The lower portion, which was in view, was removed; some portions were lying in contact with the pterygoid process; some portions embedded under the temporal bone; other portions in the orbit. Before the operation could be completed, the patient became so faint, that it was found impossible to proceed with the operation, and he subsequently sank.*

The preparation of the parts shows the superior maxillary and malar bones, and portions of the tumours extracted from behind them; the bones are healthy, and altered in shape from pressure; the tumours present all the characteristics of a fibrous structure. The parts removed after death show a morbid growth, extending in the root of the left nostril, and especially on the inner edge of the alveolar process and under surface of the body of the sphenoid bone, to which small portions of tumour were found still attached. The sphenoidal sinuses were filled with diseased structure of a similar character, and were very much dilated; so much so, that at one point the bone had altogether disappeared, leaving a small hole, where the tumour was lying in contact with the dura mater. A small portion of the growth was also found at the upper and back part of the septum nasi, which was forced over to the right side and partially absorbed by absorption: here the mucous membrane was somewhat thickened; there was a small pendulous body, loosely connected to the velum palati, hanging by the side of the uvula. Small flattened growths of a similar nature, and of a bulbous shape, were found deeply embedded in the sphenoidal and temporal fossae, as well as at the back part of the orbit. None of these had any attachment to the bones; they were all connected to each other, and to the growths in the nostril by a slender pedicle, which passed in the direction of the spheno-palatine foramen; the growth in the orbit had reached this situation by creeping through the spheno-maxillary fissure. The bones of the orbit were quite healthy. The preparation is in the Museum of St. George's Hospital.

* *Med.-Chir. Trans.* vol. xxxiv. p. 43.

greater diameter, and situated entirely on the anterior surface, extending along nearly the whole portion removed. Its base was fibrous.* The other shows the ascending ramus, or process of the lower jaw, with a tumour of a similar character. A thirty years old had suffered from 'toothache' from nine months after a blow on the cheek. The first portion of the jaw, with the tumour, was removed about five months after it was first observed; its growth continued in the jaw and the cheek, and was attended by constant pain. The disease was apparently left at the first operation. A second operation appeared at the ramus, which was subsequently removed. The patient recovered.

Subject to this occasional tendency to recurrence, tumours may be classed as essentially innocent in proportion to many of those which affect the jaw. In the innocent we intend to convey the possibility of cure by operation, and, subsequent thereto, of the part and the system from recurrence of the disease.

And yet, as in other parts, so in these, we meet with fibroid tumours which present characteristics of true fibrous growths, but possess the peculiar property, that upon each removal there is a more rapid tendency towards recurrence, as well as rapidity in growth: from the true structure of each removal takes place, there is a gradual fibro-plastic or mere gelatinous tissue; until at length a rapid deterioration of constitution runs on, and the patient sinks, exhausted by a foul ulcerated mass. When cut into, the mass is a smooth, gelatinous, and somewhat elastic mass.

George's Hospital is a preparation showing, on section, a small focus of bone which in no part approaches the surface of the tumour.* Another specimen shows much bone radiating into the substance of a fibrous tumour from its base, and approaching in some parts very near the surface.

Fibrous tumours, usually of slow and painless growth, are often attended by suffering when they affect the jaws; and their growth may be sufficiently rapid to require the aid of the surgeon within a few months of their outset, to obviate the serious results of pressure and encroachment on the cavity of the mouth.

As a general rule,' says Mr. Paget, 'the vascularity of a fibrous tumour is in inverse proportion to its singleness and compactness of construction.'† Yet, in the region of the mouth, we constantly observe a tendency to recurring hæmorrhage; which may be attributed in some measure to the friction the jaws may be subject to, or the accidental bruising they may receive, in the daily process of mastication. This hæmorrhage is not usually very great at any one time; but it has this important practical bearing, that by its amount or by its frequency it deprives the patient of much blood; he becomes blanched and debilitated; and if relief be long deferred, he becomes little capable of undergoing an operation, which of itself often exacts the immediate loss of a large amount of blood. A case already related illustrates the principle here enunciated.

Myeloid tumours of the jaws.—What has already been said relative to the general features of fibrous tumours originating in the jaws will to some extent apply to myeloid tumours affecting the bones. They will perhaps be found to affect those parts most as frequently as the fibrous tumours. The Museum of George's Hospital contains several specimens of myeloid tumours, connected with the maxillary bones.

The general characteristics of these growths have been well summed up by Mr. Paget. 'Myeloid tumours,' he says, 'usually arise singly; they are most frequent in youth, and very rare after middle-age; they generally grow slowly, and without pain; and generally commence without any known cause, such as injury or hereditary disposition. They rarely, except in persons, become osseous; they have no proneness to ulcerate or

* Series ii. No. 155.

† Op. cit. vol. ii. p. 134.

three weeks, and was successfully removed.

The diagnosis of a myeloid tumour is always without microscopical examination of its structure. Its disposition has already been fully described by its origin and growth, in connection with the joint, mistaken, as in all other parts which it attacks, for fibro-cellular growths. Its characteristic features are marked in its intimate structure; for an account refer the reader to the essay on TUMOURS, Vol. I.

The nature of myeloid tumours of the jaw permits of entire removal in treatment; entire removal of the portion of bone to which it may be attached, is the only justifiable course to be pursued; and it is recommended or sanctioned. As it will survive when once started, it should without hesitation be removed whenever detected. The removal does not ensure safety. There is no doubt that myeloid disease is recurrent. The experience of late years has confirmed this fact.

Osseous tumours of the upper and lower maxilla are commonly met with, but occur sufficiently often to be especially interesting, in connection with the mouth. They will be found to occur in two characters: simply as outgrowths from the bone to which a base more or less narrow or broad:

Museum of the College of Surgeons is a preparation
ws the true character of the hard ivory-like tumour
s. It is seen growing from the angle of the lower
hich it has an attachment by a broadish base.*
imilar specimen may be seen in the Museum of
e's Hospital.† The removal of these out-growths,
hey implicate the upper or lower jaws, presents no
fficulties than does the removal of fibrous or other
excepting, that if deeply nodulated and irregular on
e, they may be entangled in the fibres of the ad-
sles, and require much careful, it may be tedious,
before they can be cleared from the surrounding soft

emains to be considered the other condition referred
h the bones of the face, or any portion of them, take
liar process of growth, thickening, and occasionally
rgement.

seum of the College of Surgeons contains specimens
of this diseased action in bone, and they demon-
t an amount of hideous deformity may occur in the
this horrible disease—horrible, inasmuch as it ap-
ectly beyond the control of medical treatment; and
ed in a part readily removable, entails on the wretched
lingering malady which, though slowly, yet surely, in
. continues its growth until its mechanical interference



patient's confidence, to suggest anything but of the tumour. Of course this recommendation the amount of bone implicated by the disease can be *entirely removed* by the knife and saw, be otherwise practicable, the removal of the likely to be perfectly satisfactory in its result, stated,' says Mr. Stanley, 'that absolute se reproduction of an exostosis, can be obtained by the removal of every part of its circumference. A portion of the exterior of the exostosis, with its capsule, be left, reproduction of the tumour is not an improbable occurrence.'*

Experience has fully confirmed the observations of Mr. Stanley, and the caution which they inculcate is, not the removal of a portion, when the whole mass of bone may present obstacles to its entire resection. The hypertrophied condition of bone may hereafter be subject to constitutional treatment, is a fair question. The disease is not common, and opportunities for testing the efficiency or benefit of drugs, externally or internally, are fairly tested. The disease has certainly some constitutional character than simple osseous condition is one which has been known only as the prerogative of hereditary transmission. In treatment, Mr. Stanley states, that 'medicine is of no use in the disease.'

geon, even a portion may be cut away with present if not permanent benefit. 'I know, in cases where only part of the hypertrophied bones have been removed, the wound has healed handsly over the remaining portion of them; and it has not in such instances appeared that the operation was followed by any increase of the disease.' *

Vascular tumours of bone are so rare, and of the jaw so few instances have been put on record, that we even venture to express some hesitation in taking for granted that those described may not have been tumours of a cancerous nature. The entire removal of the disease is the only treatment to be recommended. As described, the disease appears to commence in early life, and to be painless; but its external characters are such that no very accurate opinion can be formed of its nature, unless it present on the surface the deep-red colour produced by enlarged vessels.

Mucous tumour, or myxoma.—Virchow has drawn attention to this variety, and Mr. Moore fully described its characteristics in his article on TUMOURS (Vol. I. page 525). They are rare in the jaw, but have been met with. They may be removed like other tumours, and are not liable to return.

Cancerous tumours of the maxillæ, and cancerous ulcerations of the gums, remain to be considered.

Cancer of the bones of the mouth is by no means uncommon; it is generally of the medullary form, occasionally of the osteoid variety. Cancer of the gums is more rigidly confined to the epithelial character. Scirrhus, in its onslaughts, appears to regard these regions; for though found as a secondary condition in bone, it has not fallen to our lot to notice it in the bones of the face; the experience of others confirms this observation.

Cancerous growths are observed more frequently to affect the superior than the inferior maxilla, and to commence most frequently in, or around, the walls of the antrum. The history of life, and the anatomy, of medullary and osteoid cancers are, written elsewhere in this work, apply in every minute particular to the origin, the growth, and the structure of cancers

* Stanley, *On Diseases of Bones*, p. 5.

a cancerous tumour of the mouth. We observe it in children, in persons of middle life, in the elderly. It commences often without pain, but it rarely continues long without suffering; and in many instances the pain arising from the progress of these bones is one of unmitigated torture. The appearances of these tumours differ in many ways in each case.

When the disease is confined to the lower jaw, it may be more readily and earlier detected than when it is medullary, it displays a softish pulpy mass, more or less projecting, it may be, on the outer side, and on the chin or cheek; or on the inner side, and pushing into the throat; or it may uniformly surround the bone.

A case illustrative of the latter condition was (1863) in S. J. under our care. It was rapidly running its course. The patient, forty-six, observed a swelling on the outer side of the lower jaw previous to admission. The tumour rapidly increased, and involved the bone, from the symphysis to the first molars on each side, backwards and downwards to the hyoid bone. The skin over it was soft and partially adherent, especially at the lower and left side. The surface was soft, red, and pulpy; much as if the disease had nearly reached the surface. Some of the teeth on the left side were becoming loose. Under such conditions, it was thought best not to interfere. The patient died about three months after his admission.

Occasionally the masses which project into the mouth early into ulceration, discharge very offensive matter, and have a ready tendency to bleed. If teeth become loose and are removed, a fungoid growth shortly fills the sockets. The external parts, if the tumour projects,

frequently bear evidence of the constitutional contamination; and the cachexia of cancer is usually well marked.

When the upper jaw is the seat of cancerous growth, there is often, in its early stage, some difficulty in arriving at a correct diagnosis as to the nature of the tumour. It is often but little exposed to view; it may be entirely embedded in an osseous chamber; or compressed and moulded by osseous walls, which, in the commencement, retard or direct the course of its growth. It may project into the nostril, block it up, and, perforating the septum, pass through to the opposite side; it may commence in the anterior wall of the antrum, and project forwards under cover of the cheek; or from the outer wall, and proceed downwards over the alveolus, and upwards into the temporal fossa; or from the interior of the antrum, and make its way uniformly into the nares, into the mouth, into the orbit, outwards, and forwards, and backwards. As it grows, the bones adjacent to it become expanded to some extent; then implicated or absorbed. If it presses forwards, the cheek and upper lip become distended and puffy; the fungoid growth within increases, and protrudes sometimes, at the angle of the mouth; the eyelids become oedematous, and by degrees are closed; or perhaps, prior to this, the eyeball is pushed forwards or outwards; and the cornea, no longer covered by the lids, ulcerates or sloughs, and the contents of the globe escape. Nothing can be more hideous than the advanced condition of a medullary cancer of the upper jaw, presenting often a mass half as large as the head itself; ulcerated or sloughy on its surface; discharging horribly stinking pus, mixed with blood; and persecuting the sufferer with unabating and often agonising pain.

Rapidity of growth is a strong point in the chain of evidence which is to decide the character of such tumours. But we have already observed, that rapid growth is seen occasionally in tumours of a non-malignant character. In medullary tumours, rapid growth is the rule; early attainment of size a marked feature; and the rapidity of growth commensurate with increase of size. The constitutional deterioration also is continuous. The early contamination of the skin with the tumour should surely point out to the surgeon the extreme foolhardiness of attempting to remove a mass of disease, the extent of which it is impossible to define; and, consequently, the satisfactory result of which is rarely practicable, with due regard to the patient's benefit.

the mass; but upon each occasion alarming hæmorrhage disease was evidently medullary cancer, increasing rapidly, uncertainty as to its origin, it was recommended that no permitted. The child died within two months of our first

The face was greatly disfigured, by considerable protrusion outwards and forwards, by the expansion of the nostrils, and a fungoid mass from the left one.

The dura mater, between the sella turcica and the ethmoid on its attached surface, and readily separated from the bone portion of bone, as well as the cribriform plate and crista, darker coloured, and less compact in structure than natural filled with a whitish medullary tumour. It had projected extent, that it had separated the nasal bones and cartilage each other. Inferiorly, it had partially protruded through of the left nostril, and had separated the palate process maxillæ and palate-bones, sufficiently to admit the point their edges. The septum nasi was absorbed to a great extent walls of the orbits; and thus the tumour projected into orbit, occupying the whole of that of the left side, where eyeball outwards, on a level with the anterior orbital margin had also penetrated into each antrum. Posteriorly it had openings of the posterior nares; was in close contact with the basilar process, and the anterior surface of the upper pressed down the soft palate considerably; and had extended almost as low as the epiglottis.

The above case illustrates the progress of medullary cancer in childhood; the following shows how similar are the cases in advanced life; and both show how in each the surgeon was attempt to remove them. A woman aged fifty was admitted to Hospital in July 1861, under the author's care, for a tumour from the antrum, and involving its anterior wall and the months previous she had a decayed tooth removed from the on the left side of the face and head; immediately after

skin was not implicated in the growth. The bones surrounding the were so soft, that they were readily cut with a scalpel. The tumour protruded largely into the mouth; it involved the palate, the inferior turbinated,moid, the sphenoid, and the superior maxillary bones. Through a portion of the sphenoid and sphenoid bones there was an opening communicating with the interior of the skull. The growth, with the portion of upper jaw that remained fixed with it, is in the Museum of St. George's Hospital.*

A patient, aged thirty-eight, was admitted to St. George's Hospital under the care, Jan. 1870. In July 1869, she first noticed swelling of the right

FIG. 252.



Portrait of a patient with malignant tumour of the upper jaw, and deficiency of the lower jaw.

eyelid, with neuralgic pains over the right temple and brow. After a month the tumor began to protrude. In September, the right cheek began to bulge out, and the right side of the nose to enlarge. The sight of the right eye had been

The author is indebted to Mr. Charles Hunter for the post-mortem exami-

If a cancerous tumour of the lower jaw be in this condition, if it is situated near the symphysis, and the patient desires its removal, with the whole depth of the bone in which it is attached, may be attempted; for if its limits may be ascertained, and the tumour disarticulated. But if it involve the bone near the angle, we cannot say how deeply it may extend, or to what degree it may infiltrate the tissues at the root of the tongue, or how far beyond the reach of the scalpel.

After all, it will be found that much depends on the part of the surgeon who has to remove a cancerous growth of the jaws: the result depends on the individual features of each case, and is entirely out of the question to attempt a strict rule relative to treatment. At best, however, intervention is in the large number of cases most unsuccessful; few, we hope some benefit is conferred.

We cannot do greater justice to the subject than by translating the words of Monsieur Giraldès: ' Dans le traitement du cancer de la mâchoire maxillaire, il est des cas dans lesquels le chirurgien doit intervenir, et dans lesquels il est prudent de s'abstenir. Lorsque le cancer est à un volume, et surtout lorsque la constitution du malade n'est pas altérée, il peut intervenir, et dans ce cas, il peut arrêter le mal pendant quelque temps, et prolonger la vie des malades. Si la tumeur est volumineuse, et si la constitution du malade soit encore altérée; si la cachexie cancéreuse est prononcée, le chirurgien doit encore intervenir, et par deux raisons: premièrement, il peut arrêter pour quelque temps la marche d'une maladie qui tend à progresser: ensuite, parce qu'il ne peut avoir affaire à un

ns and the other soft tissues of the mouth, if affected are usually attacked by the epithelial variety. Comradually on the surface of the gums, it runs its course, that species of cancer elsewhere. By degrees the surface spreads; the borders of the ulcer present their thickened and elevated edges; not only the cheek, but of the mouth and tongue become secondarily affected. In this situation is generally very painful; much is occasioned by movements of the cheek or tongue; every effort is entailed by constant escape of saliva; great inconvenience and often difficulty is experienced in articulation.

Operation in such cases is justifiable. Epithelial cancer of the mouth appears to partake of a more malignant nature than it assumes elsewhere; or rather, we might say, its location renders it an obstacle to the requisite supply of blood, consequently the life of the patient is sooner terminated. The treatment to be adopted is palliative: locally the use of opium to obviate the distress occasioned by offensive discharges; constitutionally, anodynes to lull the pangs which dart through the parts affected.

OPERATIONS.

Operations for the removal of the lower or upper jaw, or parts of them, have been fully described in all treatises on Plastic Surgery, we feel that there is nothing new to require the following observations. Our notice of the diseases of the jaws would, however, be incomplete were we not to lay down for the reader the general rules he should observe, and the order in which he should proceed, when called upon to operate on either of these bones.

Operations in diseases of the lower jaw may involve the removal of a small portion of the bone; of a large portion of the bone; or division; of one half—from the symphysis to the other end; or rarely of the whole bone, with both articular processes. There is considerable difference in the method of proceeding under these various conditions; but chiefly as regards the nature and extent of the incisions required in each.


A small tumour may occupy only a portion of the anterior surface of the lower jaw, and the growth be found connected with the bone along its edge without affecting the bone to a greater depth;

extending perhaps further laterally than downwards. It may have involved the alveolar process to such an extent, that the corresponding teeth are loose or have been pushed out. A case in which a tumour (fibrous) affected the lower jaw to the extent of the four incisors, came under our care a few years ago. The tumour was firm and painless, and involved about half the depth of the jaw. The canine teeth were removed; a saw was then used to cut down on each side of the tumour, through half the depth of the bone; then, with a pair of cutting forceps, the upper half of the bone, between the lateral cuts, was taken away, with the tumour still adherent to it. Sir W. Fergusson relates a case in which he removed a much larger portion, in breadth, of the lower jaw, by a similar operation.* He justly observes, that the advantages of being able to leave the lower half of the bone are very considerable; for the lower jaw retains its shape, and false teeth can be readily adapted to it, when the parts are healed. When, under these circumstances, the tumour is small, no external incision is requisite. In the case alluded to, Sir W. Fergusson was obliged to expose the tumour, by making incisions from the angles of the mouth to the base of the lower jaw. If the operator can possibly avoid cutting through the lip, he should by all means do so: and it will be found that, by drawing the lower lip down, separating it from the bone, and with retractors pulling upon the angles of the mouth, a considerable surface of the anterior part of the bone may be exposed, without any external incision. Instead of two incisions through the lower lip, when a tumour has to be removed with the anterior portion of the jaw, one incision in the median line may suffice: it can be carried down below the chin without any obstacle or objection; and this allows the flaps to be drawn well over to either side, so as to expose a considerable portion of the bone. The downward extension of the incision also affords a ready escape for the future discharge from the wound.

When a tumour involves the whole depth of the bone, and this has to be removed with the morbid growth, the external incisions having been made, the flaps must be cleanly dissected from the tumour, to expose a sufficient surface of healthy bone on each side. In order to enable the surgeon to cut where the tissues are sound, the teeth should be removed where the bone

* Fergusson's *Practical Surgery*, p. 668.


is to be divided. The anterior surface being denuded, the posterior has to be freed from attachments of muscles, &c. Prior to passing the knife behind the bone, the tongue must be secured; a strong ligature should be passed through its apex, and held by an assistant. Of course this is only requisite when the muscles are detached at the symphysis. If this precaution be neglected, the tongue is apt to fall back on the division of its muscles, and the patient runs the risk of suffocation. The ligature may be removed a few hours after the operation; in the mean time it should be held by an assistant, or otherwise secured on the face. A large fibrous tumour of the lower jaw was removed in our presence, with more than half of the bone, without any attention being paid to the tongue; the patient appeared choking, when the tongue was seized with a pair of forceps, drawn forwards, and immediately secured with a needle and thread, and the patient again breathed with comfort. If the tumour be very large, and irregular on its posterior aspect, it is better first to cut through the jaw at its sides, and then subsequently detach it and the tumour from its posterior adhesions. The isolated portion of the jaw can then be turned forwards, and the mass dissected from the soft tissues at the floor of the mouth, commencing within from above downwards. The method of dividing the jaw is simple, if properly performed. The outer shell of the bone should be notched with a fine saw, and then the bone cut through with a pair of strong bone-cutting forceps. When a tumour implicates the side of the lower jaw, or runs near to or involves the ramus, the external incision, to expose the bone, will differ from that described. If it be requisite to disarticulate the bone, the external incision should commence just above and over the corresponding temporo-maxillary joint; be carried down to near the angle; then turned forwards to the side of the chin and (if the tumour extend far forwards) towards the border of the lip, and terminate within a little of its free margin. The integuments of the side of the face included in this line of incision are now to be dissected upwards. The incision has crossed the facial artery; and as the flap is reflected off, the artery must be divided; it had better at once be tied, to save loss of blood. The bone immediately in front of the diseased mass has then to be divided, as described. The portion to be disarticulated must then be seized and drawn forwards; some force may be requisite to effect this. As the bone is drawn outwards, the mucous membrane and muscles are



be neglected. If disarticulation be not required of the tumour not extending to the corner, the incision need not be carried quite so high on the face; but, in other respects, should be made as just described. The bone, however, will be supported behind, as well as in front of the tumour. The stages of the operation are comprised in separating the vessels and in bringing the edges of the wound together. Silver sutures are probably most useful here, and are introduced at frequent intervals, so that the portions of the side of the face be accurately approximated. The extreme vascularity of the tissue requires rapid union, and a speedy recovery from this frightful wound. The inner surface of the tongue is supported with a fold of wet lint. The dissipated vessels adapt themselves to their novel duty, and in a few hours are capable of supporting the tongue so that it can perform its usual offices, without fear of suffocation towards the pharynx. The chief inconvenience of the jaw which is left is, that it is not supported inwards. But if a thin cap of metal be adapted to the operation, so that the upper and lower teeth be brought into contact, it may be applied, as soon as the operation is completed, to great advantage. It steadies the parts, and enables the patient with the administration of food. This latter part may easily be sucked in between the lips, and the teeth on the sound side. The jaw should

s to the opposite angle of the jaw, and then upwards to the condyle. This is a formidable operation. If practicable, a considerable facility may be obtained by dividing the symphysis immediately after the exposure of the surface of the whole bone. The operation will be found to accelerate, and render much less painful, the separation of the muscles and soft tissues from the inner surface of the bone and tumour, and especially aid in the rapid disarticulation of each condyle.

An extensive incision necessary for the removal of a large tumour of the lower jaw unavoidably passes across and divides the superficial branches of the portio dura; the result of which is a loss of power over the cheek, and a drawing over of the mouth towards the opposite side. This disfigurement, at first, is generally considerable; but appears usually to improve by time, though it is entirely disappears. Mr. Syme succeeded in removing a tumour confined to the ramus of the jaw by a modified incision on the side of the face, commencing over the tumour, and extending down towards the angle, and then forwards, without entering into the mouth. The ramus, with the condyle, was fully removed.* During operations for removal of a tumour connected with the lower jaw, the patient should sit up rather than lie down. The blood thus escapes externally, more readily than into the throat. Chloroform may be given with safety, but requires to be administered cautiously, with a wet towel held near the nostrils. We should not do



extent of the external incisions are very in
tumour is small, and it is supposed to be
surface of the antrum, or projects from th
an incision through the middle of the up
sponding nostril, continued upwards from t
ala for an inch or more along the side of
the surgeon to dissect off a considerable fla
large portion of the superior maxillary bon

When it is supposed requisite to remo
superior maxilla with a large tumour, a se
described will be requisite. This second in
through the cheek, from the inner surfac
mouth, obliquely to the prominence of the
tumour be very considerable, and implicate
incision can be run outwards over the zy
may thus be secured, when the soft parts a
from the anterior surface of the bones.

If, when the requisite incision is made,
ciently exposed, it should appear that the
the inner wall of the antrum, or growing fr
the nasal bone should be cut through, an
one side, without entirely separating it from
ments. Thus the nostril will be laid fully
can perhaps be removed without cutting :
this measure be successful, the nasal bone
flap of skin brought over it, and secured w

If the disease is found to affect the lower

ter, on the side of the median line which is affected by the disease. Then the bone immediately below the margin of the orbit, and from the nostril to the outer edge of the malar bone, must be cut through with saw and bone-forceps, and the attachments of the soft palate separated with the scalpel. The portion of bone to be removed should then be seized with strong forceps, when, by means of a few additional cuts with the knife, the surgeon will be able to dislodge the lower portion of the maxillary bone, with the tumour attached.

If it be requisite to remove the whole of the superior maxilla, the external incision which extends in the direction of the zygoma must be continued backwards sufficiently to allow of the exposure of the zygomatic process of the malar bone. When the flap of the cheek has been freely dissected upwards to the margin of the orbit, the floor of the orbit must be cleared carefully of the attachment of the inferior oblique muscle, &c. The zygomatic process of the malar must be notched with the saw; the alveolar ridge similarly treated; and then the bone-cutting forceps employed. One blade of the instrument should be introduced into the nostril, the other at the inner and inferior portion of the orbit; great care being taken that the eyeball is not injured. The blades of the forceps should cut through the nostril and floor of the orbit to the spheno-maxillary fissure; the malar bone should be cut through its middle, right back to the anterior extremity of the same fissure; and the hard palate, on the same side of the septum nasi, should be cut back to the posterior border. Thus the tumour and bone will now only be held by a few attachments of soft tissue, which the knife easily separates. The bone can then be removed with slightly forcible manipulation, or by twisting it in the grip of a pair of strong forceps.

The hæmorrhage in such an operation is usually greater from the flap than from the rest of the exposed surface; but generally subsides in a short time, so that many ligatures are rarely requisite. The hollow left by the removal of the bone should be partially filled with wet lint, and the edges of the flap secured as already described. In such operations the patient should sit in the upright position, that he may readily get rid of the blood from his mouth. Chloroform should be administered by an experienced person; for its action can only be kept up by its being applied on a sponge or towel.

Tumours of the upper jaw vary so materially in size, in pro-

the artery, and to preserve the former in soft parts. He believes that he has obtained deformity by so doing; while, if he had not, deformity would have resulted.

He has also adopted an operation, which is a resection of the upper jaw. This operation is the origin and growth of the tumour is in the maxillary bone, but cannot be reached with the bone.* Instead of removing the superior part, requisite external incisions are made down to the bone, the bone sawn through in the same incisions, partially detached forcibly turned inwards, dividing it from its connections with the nasal bone. The tumour is then removed from behind, and the latter replaced in its original position. No deformity is the place where this fracture or bending may occur.

Langenbeck's observations, and the cases which he has so lengthily for us to introduce here; we must refer the reader to the original.†

DISEASES OF THE PHARYNX

Congenital defects of the pharynx are rare, and occur are associated with such other conditions as to render them incompatible with life; so that their importance attaches to their consideration.

Inflammation of the pharynx is generally associated with a similar condition of the fauces and soft palate.

The general conditions indicative of this form of inflammation will be considered under the head of œdema of the glottis, the essay on DISEASES OF THE LARYNX. Difficulty in swallowing; laboured respiration; swelling of the throat, often observed externally; swollen and rounded uvula and margins of the palate, hiding from view all behind them, are the prominent symptoms. The tongue soon becomes brown and dry, the mouth covered with sordes. The patient dies suddenly in a spasmodic effort to respire, or gradually sinks. Usually death takes place at an early period, and often sooner than has been anticipated. A patient, well to-day, is attacked in the evening with sore throat and difficulty in swallowing; these symptoms are soon followed by great obstruction to respiration, entire loss of sleep, rapid pulse, and hot skin; in the course of three days or four at latest, with all his faculties clear, and while able to direct by motion or writing those around him, the patient's head falls back, and he is dead.

If the parts be examined after death, there will be found œdema of the sub-mucous areolar tissue, around and in the substance of the constrictors, and travelling down around the œsophagus as low as the cardiac extremity. Often the areolar tissue is the seat of purulent infiltration. The treatment of such cases is most frequently hopeless. As long as the patient can swallow, as long as the mischief stops short of purulent infiltration, there is hope of recovery, and encouragement to persevere with the administration of stimulants and nutritious food, if they can be swallowed; or with enemata of wine, opium, and beef-tea. Laryngotomy should be performed when danger of suffocation becomes evident, or if the patient is suddenly attacked by spasm of the larynx threatening immediate death. Even if the benefit from laryngotomy be not permanent, it affords much immediate relief to the distress of the symptoms, and certainly prolongs life in most cases.

W. S., aged forty-five, was admitted on the 29th September 1845, into St. George's Hospital, with difficulty of breathing and swallowing, and all the symptoms of diffuse cellular inflammation, extending down the pharynx and œsophagus. On the 2nd October, at five A.M., he was attacked with a paroxysm of dyspnoea, and became quite black in the face and insensible. Laryngotomy was at once performed; and after artificial respiration had been kept up some time, consciousness returned; but it was found that the patient was utterly unable to swallow anything, and he died the following day. The effusion of lymph extended from the cellular tissue round the glottis downwards around

Abscess of the pharynx is an occasional occurrence, a formidable evil; and is often as a cause with some disease of the cervical vertebræ. It is not taken to arrive at a correct diagnosis of such a case readily be mistaken for a polypoid or other growth on the walls or cavity of the pharynx.

The following case well illustrates the importance of an abscess, the size of a pigeon's egg, was situated between the cervical vertebræ and the back of the pharynx, but the front did not cause any material projection of the posterior wall. In connection with the abscess, a second small cyst was found as to form a nipple-like swelling in the pharynx; it completely closed the orifice of the glottis. The abscess was between this process and the body of the abscess admitted the tip of a finger; and the whole swelling was freely movable on its extremities and sides. This abscess occurred in a patient who had suffered from dyspnoea for three weeks. The symptoms became very urgent three days before death; was continued at intervals; but the slightest exposure to cold, again brought on a recurrence of threatened suffocation, attended by an inspiratory sound. There was no attendant difficulty of deglutition. The disease was found after death.†

Abscess may arise from an injury to the pharyngeal wall by the passage or the impaction of a foreign body. In the Museum of the Hospital is a preparation which shows an irregular ulcerated abscess, on the posterior wall of the pharynx, which was found in a young woman who died from its effects, and who, previous to her death, swallowed some pins, one of which was lodged in the pharynx, and produced the mischief which led to death.

times observed to produce considerable destruction of the soft tissues, and its progress is very difficult to control. The symptoms generally are, offensive breath; offensive purulent secretion from the nostrils, and excoriated nasal apertures; constant cough; and the external characteristics of a strumous diathesis. The ulceration is readily seen when the mouth is opened; is not deep, but irregular, and the edges of the ulcer usually well defined; with crusts of dried secretion, often highly offensive, sticking in patches to its surface. There is not much pain complained of; perhaps some, less or more, is experienced when food is taken.

This characteristic form of ulceration is extremely obstinate in duration, and often extends to the tissues of the soft and hard palate, as well as to the nares. In the case of a little girl in St. George's Hospital, the mischief spread most awfully from the pharynx to the nose and mouth, and from the mucous surface to the deeper tissues; until the nose fell in, and the greater portion of the palate, soft and hard, was destroyed, leaving the child a dreadfully disfigured object.

This strumous form of ulceration is certainly found to exist in children more frequently than in adults. The treatment is to be directed chiefly to constitutional measures, though local applications may also prove beneficial. It should consist of the various tonics, which are beneficial in the management of scrofulous cases generally, aided by good living, great attention to cleanliness and pure air, and, if within the means of the patient, a sea-side residence; the Isle of Thanet is especially to be recommended.

Syphilitic ulceration of the pharynx is very commonly observed, especially in the lower classes of the metropolis. It is constantly seen to attack the posterior wall of the pharynx; but is usually accompanied by ulceration of the velum or tonsils, or some other secondary evidence of the prevailing poison.

Its character may be suspected by the age of the patient, the period of its duration, and other symptoms which may be present; i.e. rupial sores in the skin; nodes of the skull or other bones; pains in the limbs or joints; general reduction of health and flesh; and last, not least, the confession (not always to be obtained) of a former primary sore, with perhaps a mark recent habitation.

Such an ulceration will usually yield rapidly and satisfactorily to treatment. It granulates, contracts, and cicatrises under the influence of iodide of iron, or iodide of potassium alone; and if

622. The pouch projects downwards behind the upper part of the œsophagus; it is apparently composed of a portion of the mucous membrane and sub-mucous tissue protruding through the muscular coats of the pharynx, and does not show any muscular fibres entering into its formation. The upper portion of the pharynx is enlarged, and the pharyngeal muscles are hypertrophied. There is a stricture of the œsophagus.

FIG. 253.



Back of the pharynx. (From a preparation in the Museum of St. George's Hospital. Ser. ix. No. 14.)

The preparation was removed from the body of a man aged sixty-three, who for several years had suffered from difficulty in swallowing. After the conclusion of a meal he was in the habit of returning small portions of food. For two or three years previous to his death he was subject to attacks of inflammation of his larynx, from which he recovered under treatment. His death was the result of inflammation of the lungs. The dysphagia was always supposed to be owing to stricture of the œsophagus.

Tumours attached to the pharynx are not very common. When they occur, they are generally apt to be somewhat pendulous, and are often attached to a portion of the walls by a narrow neck. They have been found to consist of fat, pus or fibro-cellular tissue. But the character of the growth is perhaps of less importance as regards the immediate treatment, than the shape, the size, the position, and the attachment of the mass.

The tumour was found to be a large, pendulous, fatty mass, attached to the pharynx, and extending downwards to the œsophagus for a distance of two inches, being attached by fibrous tissue, covered by mucous membrane, the epiglottis, and also to the upper part and side of the pharynx by its weight so dragged upon the epiglottis, that the laryngeal aperture was not practicable. The mass, except the pedicle, was hanging loose in the pharynx and œsophagus.

In the Museum of the College of Surgeons are two cases of the pharynx, successfully removed during life. One was a fatty mass, apparently fatty or gelatinous, like a nasal polypus, attached by a narrow pedicle behind the tonsil. A second† is a fatty mass, and was attached by a narrow pedicle to the wall of the pharynx.

Other cases have been observed in which a fatty or fibrous structure, having encroached upon the pharynx, without becoming pendulous, has been seen to grow and insinuate themselves under the pharyngeal membrane; and as they increased in size, so did the aperture of the glottis, and interrupted the passage of food to the stomach.

The treatment of tumours located in the walls of the pharynx is often attended by much difficulty; their removal is an operation of much hazard, and removal should be had recourse to, and should be undertaken at as early a stage as possible, and not attempt to impress upon the reader.

The only hope of benefit must be from the removal of the pendulous form of tumour. The removal of a tumour attached throughout to the walls of the pharynx is a very difficult operation.

m, and the pedicle cut through with a knife. If the mass appears very vascular, or large vessels are detected through the pedicle, a double ligature may be first passed through the base, and when securely tied the mass may be safely cut off beyond it. A ligature through the mass is far preferable to one round it; for when the mass is removed by the knife, the latter is apt to slip off, and some hæmorrhage might ensue.

The chief, and probably the only risk the surgeon has to run in an attempt to remove a polypoid mass from the pharynx is that of suffocation, produced either by the mass itself filling the fauces and pressing down the epiglottis, or by spasm of the larynx, the result of irritation from the tumour coming in contact with the surface of the laryngeal mucous membrane. The character of the tumour recorded in the *Transactions of the Medical Society* by Mr. Holt be considered—its attachment, its size, and the extreme feasibility of its removal, in a great part or in its entirety, *supposing* there was no suffocation; if it be remembered that the tumour could not be drawn into the mouth without the aperture of the larynx being stopped up, but that otherwise the mass might have been drawn forwards, and the greater portion removed, would it not be justifiable, we ask, under all such circumstances, that, prior to an attempt being made to remove the tumour the precaution should be taken to perform laryngotomy, and to introduce the tube into the larynx as a safety-valve for the patient to breathe through while the operation progresses? We should have no hesitation to perform this operation under circumstances such as were attendant upon the case related, and prior to a proposed removal of a tumour; nor have we any hesitation in strongly recommending the adoption of this precaution under similar conditions. The operation of laryngotomy is simple; it is so safe in its results, provided the patient is in a healthy condition, that no danger of irritation could be apprehended from the introduction of the tube. Dr. Marshall Hall operated himself, and witnessed several other instances in which the operation had been performed on patients under the care of the late Dr. Marshall Hall for the relief of suffocation; and in no case was there the least approach to danger from the operation itself, or from the presence of the tube in the larynx. Under circumstances similar to those related by Mr. Holt, laryngotomy would insure the safety of the patient's

usually proves fatal within a few days of l

Dilatation of the œsophagus, and *pouche* tube, are morbid conditions of occasional the former have been put on record by Ro Barker.†

Contraction of the œsophagus is a very usually an insurmountable difficulty to the majority of cases rapidly reduces and dest

Contraction, or 'stricture' of the œs result of several distinct conditions. I membrane; cicatrices after injury; pre neighbouring tumours; thickening and walls; or, lastly, and most frequently, ca the tube, are the chief causes of stricture

In the Museum of the College of Surgeons is a shows stricture of the œsophagus from a fold of below the cricoid cartilage, the œsophagus gradual meter, and then dilates. The mucous membrane a transverse sharp-edged and projecting folds, which p the circumference of the tube. The surrounding tis if the result of cicatrix. There is no history to th paration (No. 1080) shows a very similar conditi child, probably the result of some acrid substance l preparation removed from the body of a boy ten year state of the œsophagus, subsequent to the action o during life. With respect to the changes subsequen poison over the mucous surface of the œsophagus

ple stricture of the oesophagus, from thickening and contraction of its is occasionally met with, and apparently without satisfactory explanation. ston exhibited at the Pathological Society a specimen possessing some ting features. The stricture existed at the upper part of the tube, was in inch in length, and only capable of allowing the passage of a goose- The contraction had existed for a number of years; there was considerable hypertrophy of the constrictors of the pharynx, especially of the superior The upper horns of the thyroid cartilage were so approximated as to ut a space of $\frac{5}{8}$ ths of an inch, being a diminution of about an inch in the h of the natural interval between them. The patient had been under re of Cruikshank and John Hunter, and lived to between seventy and

is unnecessary to multiply instances of these forms of con- tion; there are, however, certain circumstances in con- on with their progress, which will form the subject of deration when their diagnosis and treatment are dis- d.

mplete obstruction to the passage of food from the pressure tumour on the oesophagus is very rare. Occasionally some e of difficulty in swallowing is experienced, from the nce of a tumour of a cancerous nature; or from the id, when very large, or when it wraps round the trachea; om an aneurism or any other kind of swelling in the

idence of supposed pressure from a tumour, must, how- be carefully weighed, before a decision be arrived at cting its influence in this respect. Dr. Wilks † has very

or early life. Rokitsky states that 'cancer generally occurs in an isolated form, i.e. without the presence of the disease in other organs.'* Such is the case, but the rule is made too general. Every reader may satisfy himself that we differ with the author on reasonable grounds, by reference to the *Transactions of the Pathological Society*.

The usual symptoms of stricture are, first, a difficulty in the passage of solid food to the stomach; this gradually increases; sooner or later the patient is obliged to restrict himself entirely to food of a fluid nature; he finds that this he can take only in small quantities, until at last he sips but a few drops; and even this small quantity may be rejected before it has been attempted to swallow it.

A patient with stricture of the œsophagus, in the advanced stage, he visibly deteriorates in physical power and strength day to day; the circulation becomes feeble, there is a sensation of cold on temperate days, and he is more susceptible to all changes of the atmosphere.

If there be ulceration in the upper part of the pharynx be also implicated, there is generally a foul smell in the patient's breath, easily detected when entering the room. The patient is often exhausted by retortion, and the secretion spat up is occasionally very offensive. He complains of weight in the earlier period of the disease, and when the disease is advanced he can no longer take food; but gradually he becomes

ounding friends may be little prepared for such an announcement.

Occasional complications occur in the progress of stricture of the œsophagus to which our attention is requisite; they render the disease more distressing to the patient, but generally hasten the termination of his sufferings. The epithelial variety of ulceration is apt to spread superficially without destroying the tissues deeply; and though a large surface may be implicated, the patient so affected will survive longer than one affected by the medullary form of cancer. With the latter form of disease the surrounding tissues become implicated at an early period, and thus the passage may become much constricted soon after the stricture is suspected. Ulceration of the tube above or about the diseased mass will generally take place, and cavities in the cellular tissue around will be constantly found communicating with the ulcerated points of the œsophagus. Occasionally the destruction makes its way outwards, until several openings are formed externally about the neck, and allow the escape, not only of matter, but also of the fluids taken as nourishment. Or the ulcerative action may run in a different direction, burrow about the muscles of the neck, and effect a fistulous communication with the trachea. The latter condition is made evident by fluids, taken through the mouth, being coughed up through the glottis.

In other instances, almost instantaneous death will occur from sudden and violent hæmorrhage, dependent on the ulceration of a large artery.*

The symptoms of stricture of the œsophagus are so marked, that it might at first sight appear almost impossible to be deceived regarding its occurrence. But an hysterical spasmodic condition of the pharynx or œsophagus is occasionally met with, and might be mistaken for organic mischief, without the precaution and judgment brought to bear on the case. It is therefore prudent to weigh carefully the symptoms of all cases of supposed stricture of the œsophagus, in order to be able to discriminate between actual stricture, and that of

* A case is recorded in the *Transactions of the Pathological Society* (vol. ix. 1894) of death by hæmorrhage from ulceration extending from the œsophagus to the right subclavian artery. And a second instance is found at page 202 of the same volume, in which death was the result of perforation of the superior mesenteric artery, by extension of ulceration from the cancerous stricture. The vertebral artery has been known to be perforated under similar conditions.

apparently unobserved. There is little, if there is there the early indication of loss of constant desire for food if it could only be swallowed.

The diagnosis of stricture of the œsophagus is easy and sure. It comes on after mid-life and is accompanied by a constant craving for food, but an inability to swallow it. He longs for it, and would at any time swallow a large meal, if it could possibly pass to the stomach. 'Nothing of the pangs of hunger,' observed a patient of our profession, when dying of cancerous pharyngitis. 'Day and night I suffer horrible thirst, but I cannot get food from the stomach; but I can swallow.'"

When a pouch of the pharynx or œsophagus is suspected, if, after partaking of food, there is a fullness in the neck, or if, after a time, a part of the food is returned. As the pouch is capable of holding a large quantity of food there is not so much difficulty in disposing of the food as in stricture; for a part may pass into the stomach and the rest into the pouch. A patient suffering from this condition of the tube does not emaciate, and may live long after affected by stricture; several instances on record of persons living to a good old age.

The treatment of stricture of the œsophagus is simple, and in a very few words.

If hysterical, general constitutional treatment will cure it, without much difficulty, in a comparatively short time.

If permanent, from cicatrix, or other disease, the treatment is more complicated.

emata of strong beef-tea and laudanum ; and at the same time, the allowance of whatever fluid he relishes, either to wash it the mouth, or to swallow. The enemata may be repeated with great benefit every four or six hours, according to the slings or wishes of the patient. Such treatment will prolong it some days ; and it may be a matter of very considerable importance to the surviving members of a family, that a life could be prolonged over a certain period, not far distant, when a medical attendant is first consulted.

We cannot recommend the use of bougies in the treatment of stricture of the œsophagus. In its early stage, curiosity may perhaps require to be satisfied by an exploration, so that the exact seat of the stricture be ascertained ; and to such a proceeding there is no great objection. But, beyond this, very little, if anything, is gained by such treatment. In attempts made to pass a bougie through the stricture, the end of the instrument has been known to find its way into the mediastinum ; into the cavity of the pleura ; and into the cavity of the pericardium. In each instance the instrument was used by skilful hands ; but, in all the cases, rapid death supervened upon the severe symptoms which followed the withdrawal of the bougie. It is wise in such cases to refrain from interference without the certain prospect of good ; for in these cases, as in all, needless surgery is bad.

GEORGE POLLOCK.

A FEW years ago the name of Diphtheria was little known in England. It is highly probable that some of our older writers refer to this disease as it appeared then, as now, in an epidemic form. In the period England had been free from its acquaintance with its symptoms and consequences were wholly derived from observations made in France, which had been known and studied for a number of years. It must be confessed that a lingering doubt still exists in the minds of many on this side the Channel, whether it differs from our own croup, or might not be produced merely by an unusual prevalence of croup, or by an epidemic of scarlatina. Careful observation has shown that it is distinct from both these diseases, and that it is in many respects so close that no doubt can be given to one not conversant with the disease. It holds a place intermediate between them. It has a febrile character and epidemic influences of its own, and a specific exudation of the other; it combines the symptoms which belong specially to the fever, with those which belong wholly to the inflammation: and it is produced so either by the general exhaustion which precedes or less all epidemic diseases, or by the suffocation of the exudation of fibrine, exactly like the croup.

of each other, and so completely merged into one another, that it has been difficult to separate them, and to distinguish between the effects produced by each. It has not unfrequently happened that in the same family, within a very short period, one member has suffered from diphtheria, and another has had an attack of scarlatina, and each has been so marked as to leave no doubt on the subject: but if the cases were not so clear, or if, as has often happened, the question of diversity were never raised at all, the two sets of symptoms would be set down to the same cause, and false inferences drawn as to the true character of the disorder.

This circumstance has no doubt given greater vagueness to the descriptions of the older writers than would otherwise have been the case, but the conclusion is unavoidable that cases occurred in their practice also, and that doubts have often arisen as to whether it was or was not a form of scarlet fever. They were not conversant, as we are, with the exudation-process, and did not discriminate between patches of lymph resting upon the surface of the mucous membrane, and a slough involving its superficial layer. They were, indeed, surprised that when the layer was detached, the membrane was found healthy beneath it; but pathology was not sufficiently advanced to draw intelligent conclusions from such observations, and we can gather little from the past that can be of use in the consideration of the special character and treatment of this disease.

History.—The first observations which, in this country at least, led to a true understanding of the nature of that false membrane which so eminently characterises diphtheria, were made with reference to sporadic croup by Dr. Home,* about the middle of last century. He first recognised the existence of a mass of fibrine, moulded to the form of the subjacent structure, and adherent to the mucous membrane, but lying quite upon its surface, ‘and generally lying loose upon it.’ Prior to this date such exudations were supposed to be part of the membrane itself, detached in the form of a slough. His descriptions, as well as those of Dr. Cheyne,† forty years later, prove that in the sporadic croup of their day the exudation was confined to the trachea, and did not spread to the fauces.

Inquiry into the Nature, Causes, and Cure of the Croup, by Francis M.D.

Essays on Diseases of Children, by John Cheyne, M.D.

G. IV.

K K

membrane, at that point where its presence
of the most important functions of life,
caused death by suffocation.

Bretonneau, in his Memoirs* read before
in 1821, was the first to give a scientific
disease; and its symptoms have under
that his report continues to be a very
usual forms which it presents, although
character of the disease has been now almost
Till very recently this view was maintained
French pathologists; and Trousseau has
to yield the point, that the empoisonment
hibited in the general symptoms, was not
of the local affection, and of the absorption
connected with the exudation. This question
one of mere abstract reasoning, which could
cise test of its truthfulness, yet reached the
field of opposing theorists; for it offered
for endeavouring promptly and effectually
tion by local applications, while it led to
such measures as were calculated to counteract
which it was assumed could be prevented
removal of the local affection. The practice
has continued more or less to form the
the modes of treatment, and with too much

which we arrive. The characters of the sign may seem very definite when a well-marked example is presented to us; if it be at all obscure, we are forced to look for other signs to aid our judgment. In the investigation of diphtheria we are met on the threshold with a very prominent fact, which gives it a place at once in our nosology as a blood-poison; we must at present call it, while confessing our ignorance of its blood-poisoning means. It comes to us under the form of an epidemic; it attacks one individual after another; it is confined to a given locality; singles out a certain proportion of the community for its victims; and then disappears. We know not how it came, and how it went; and the answer must be necessarily somewhat vague. Sad experience has proved that persons who have been accidentally inoculated, so to speak, with the virus, have caught the infection and suffered from the disease. And as with other inoculable diseases which are also conveyed from the sick to the healthy through the air, here without actual contact, so with this there seems no reason to believe that in certain investigated instances the infection has been so propagated; and we can hardly without dissent to the proposition that it may be so in all, though it is not always, perhaps not often, possible to trace the source of the infection, and the means of its transmission. It matters not, however, whether the theory of the contagionists or the non-contagionists be proved the true one; the fact remains, that diphtheria has all the known characters of

genuine diphtheritic inflammation, and made wide enough to embrace them, being a pathognomonic sign be wanting, yet still a part of the epidemic; and quite as essential to the standing of the disease as the more manifest signs. The name is more truly applicable.

Definition.—Diphtheria may, then, be defined as a blood-poison propagated by transmissible contagion in the majority of instances, either by the discharges of the sick, or in consequence of changes as yet unknown to us. This poison appears a very few days from the date of exposure, and is followed by a general febrile action throughout the system, and a local inflammation limited to the membrane of the throat. Quite exceptionally, deep-seated inflammation is the consequence of it; but most commonly, it is followed by inflammation of the glands of the neck, which are very tender: these accidents, however, are not essential to the disease in the same sense as the specific inflammation of the membrane.

Albuminuria.—In addition to this, albuminuria is very common in the severer forms of the disease. In scarlatina, the presence of albumen is a sign of inflammation of the kidney; and we conclude that in diphtheria the vitiated or poisoned blood stimulates the kidney so as to produce this state of congestion. We have no clue to unravel further the mystery. In the present case, the urine presents no trace of albumen during the disease.

ll, when it does not accompany any other form of sore
r to the same extent any other febrile disorder.

is.—Another phenomenon accompanies or rather
on attacks of diphtheria with sufficient frequency to
certain relationship between them; but as yet we
firm that paralytic affections may be apprehended
al convalescence from diphtheria—after, indeed, the
got perfectly well, and nothing seems wanting to re-
the due performance of the nutritive functions. This
e gravest of the sequelæ, if it be true that patients
alysis of the heart* when no lesion whatever can be
r death. The paralysis most frequently affects only
s of deglutition and of speech, but does also some-
ude nearly all the muscular tissues throughout the
sight becomes impaired through loss of the adjusting
he eye,† the pulse fails, apparently from diminished
ergy in the heart, the legs and arms become partially
and sensations of tingling and numbness are com-
or actual anæsthesia exists.

itic fever.—The primary action of the poison when
to the system is an alteration in the character of the
is we infer, in the first place, from the sense of mal-
febrile action, in some instances very slight indeed,
e very severe, which can be traced before any local
ion occurs. This fever alone may kill, and when it
is always by its assuming an asthenic or typhoid
the patient either becomes daily weaker, and ulti-
s of exhaustion, or the case is marked by low mutter-
m, with a tendency to slough about the inflamed
embrane, excessive prostration, and rapid sinking; in
the pulse is feeble and very frequent. It is to be
ind that this is the invariable character of the disease,
flammatory the fever may in the first instance seem
the second place, we infer the existence of blood-
rom the occasional appearance of purpura and sanious
in fatal cases, and the constant sequel of intense
en in comparatively mild cases, during convalescence.
eculiarities are very striking, because the depression
rom the attack is out of all proportion to its severity,

ia, its Symptoms and Treatment, by Wm. Jenner, M.D., pp. 42–59.
ia, by E. H. Greenhow, M.D., p. 229.

of active inflammation. And such no d
specific, and not an ordinary inflammatio
in its character and consequences from t
tion of mucous membrane, and bears some
is more commonly restricted to serous
from it most, perhaps, in its tendency to
all perfectly familiar with the rapid spre
blush, from a single point, over the wh
peritonæum, with its sticky, glazy, and fi
see the same exudation forming a thick s
the heart, or lining the entire cavity of th
we have the same character marking t
parts involved do not extend as far as the l
In many cases the patches of lymph are
inflammation is confined to the back of th
number of instances the trachea, and
involved; not unfrequently the membran
attacked more or less extensively; more
travels down the œsophagus; and in a few
lachrymal duct to the surface of the eye. T
seem to be much more in harmony with
of serous inflammations, and there is but
respect between diphtheria and ordinary s
either be limited to the fauces, or may be
a catarrh, ending in general bronchitis.
exudation varies much in the same mann
of inflamed surface is small, the effusio

e fungus had given an explanation of the whole pathology of diphtheria. Competent observers have, however, shown, not that the exudation may exist without any trace of the parasite in its substance, but also that the very same form of vegetable life may be seen occasionally in some of the secretions of the mouth when no symptom of diphtheria, and no exudation of pus, had been present.

The ordinary consistence of the effusion is such that it is often a false membrane; but it is sometimes scarcely more coherent than a paste, and contains more granular matter and cells than the mucous membrane in its structure. The mucous membrane on which it rests is almost always redder than usual; and when the fibrinous exudation is removed, it looks raw and irritable, as if denuded of its epithelium; it is generally also dotted over with bloody points where the adhesion has been closer, or the exudation has torn the mucous follicles, and torn the tissue in its removal.

Diphtheritic exudation is not, however, confined to the mucous membrane of the throat, although probably always to be found there in a true case of diphtheria. It is to be seen very distinctly on the skin when the cuticle has been removed by a blister; and it is alleged that the contagion of the disease has been communicated in this way to persons who have had blisters applied to the skin while living in rooms where others have been suffering from diphtheria. Any ulcerated or abraded surface, in persons labouring under the disease, may similarly become coated with a false membrane; and not unfrequently the mucous membrane of the pudenda in female children is the seat of a similar exudation. Suppuration of an unhealthy kind sometimes attacks the mucous membrane beneath the exudation, which melts away, leaving a gangrenous odour, leaving a slough or a foul ulcerated surface behind; and then the whole course of the disease may be changed, and the patient may fall into that low typhoid condition which is believed to be caused by the absorption of the pus.

The mere fact of exudation having taken place, apart from its being an indication of the severity of the attack, claims our earnest attention, from its occasional bearing on the issue of the case as a local complication. It is not the extent of the surface affected which excites apprehension, but the importance of the narrow chink through which air is drawn into the lungs, and the readiness with which a very small amount of pus, or a very slight deposit on the membrane covering

results from apnoea is identical in both diseases, the exudation, so unusual in inflammation of the lungs, is also apparently identical, but, so far as the diseases are different. The diseases appear to be two distinct communicable diseases, the one having not the one a simple inflammation, the other a general disturbance in the one case only ; in the other the local inflammation and the difficulty of the other the local disorder is only one of many of a previously existing condition which the fever is caused.

Treatment.—The fatal termination of a disease may follow either as a remote or immediate consequence of itself, or in consequence of the obstruction to exudation ; and the treatment must vary according to these circumstances seems to be the most important. Like the other acute specific diseases, diphtheria follows on general principles, inasmuch as no specific remedies have yet been discovered which have the power of destroying blood-poisons on which they severally depend. Such remedies exist. The condition of the patient during incubation of a fever, while an infinitesimal quantity of poison has been absorbed, must be very different from that in which the poison is again generated in the system, and is passing off from the patient's body. Remedies are brought within its range : and a remedy which is not a blood-poison, and which is not a

terial can only be got rid of by elimination; fresh blood must be formed to take the place of that which has undergone a process of deterioration. On these broad principles must our treatment be based with reference to the fever itself; the local case demands separate attention and different remedies.

In some rare cases the patient is at once prostrated by the rity of the fever; he has a brown tongue, a quick and feeble e, probably purpurous spots on his body, or a sanious discharge from the nose or fauces, and occasionally muttering delirium. Such cases rarely show any signs of rallying; and the use of stimulants affords the only chance of saving the patient or enabling him to recover from the shock of the attack.

If not thus prostrate from the very first, yet generally from an early period the tendency of the disease is to assume a low sthenic type; and depressant remedies must be carefully avoided. In some cases the local inflammation is severe, and the fever is proportionately acute, and for a few days mild salines may be indicated; but even such treatment must not be persisted in too long, and the depression which follows must be either forestalled by the administration of stimulants, or the difficulty of swallowing must not be permitted to interfere with the quantity of nourishment taken; patients who escape the first severity of the disease are still exposed to the danger of lingering convalescence, or a gradual exhaustion of the vital powers; and hence the importance of a sustaining mode of treatment. Chlorate of potash, hydrochloric acid, and muriated tincture of iron, are the constitutional remedies on which, according to the testimony of most writers, the greatest reliance is to be placed, combining them with diffusible stimuli or tonics, as the particular case seems to require. Each has been vaunted in as almost specific, and each has notably failed where it has been misapplied. Chlorate of potash acts so completely as a stimulant in some cases of cancrum oris, that it is not unreasonable to suppose it may act beneficially where there is any analogous irritation of the mucous membrane; in the purely non-ulcerated form of diphtheria, it is less likely to do good. Hydrochloric acid, either alone or in combination with the chlorate, is well known as a gargle and drink in scarlet fever. It seems to act locally as a local stimulus to the throat, and partly as a general alterative, especially in those forms of the disease in which the appearance of purpura or sanious discharge proves that important changes have taken place in the condition of the blood.

Iron in all its forms is so distinctly the remedy which tends to restore the blood to its healthy condition, that its use cannot be dispensed with in a disease of which blood-changes are the commencement, and anæmia the usual termination. The tincture of the perchloride has appeared to act most beneficially in some cases of erysipelas, and its local astringent action seems to render it especially suited for the treatment of this disease. In some cases it has certainly answered even beyond expectation.

3. Foul and unhealthy suppuration not only aggravates the general disorder, but may lead to the supervention of blood-poisoning of another kind. Pyæmia, as it is called, must be regarded as a most dangerous complication of diphtheria. Ulceration of the mucous membrane shows that the vitality of the patient is low, and calls for increased activity in the administration of stimulants and tonic remedies, while every attempt is made to convert the foul suppuration into a healthy sore, to prevent the possibility of the absorption of unhealthy pus into the system. If secondary abscesses begin to form, the liberal employment of wine and brandy, with large doses of opium, gives the only chance of recovery. The extent to which both opiates and stimulants are borne in such cases is quite surprising.

4. The local disorder claims our attention more especially when it spreads towards the larynx and trachea. The idea that the extent of the exudation was the cause of the severity of the symptoms has been entirely abandoned, and with it that meddling activity which deemed it necessary to apply caustics or astringents several times a day to the throat. An abraded surface, whether of cuticle or of mucous membrane, is speedily covered with the diphtheritic exudation in the severer forms of the disease, even when no contact of parts is possible; and it would seem perfect madness to apply an escharotic which tends to denude the adjacent membrane of its epithelium, and prepare it for the fibrinous exudation which is certain to take its place. Such treatment, however, has been adopted, under the mistaken idea that local means could check the constitutional malady. The circumstances which seem to demand local applications are the rapid spread of the exudation, and the existence of ulceration. The application must be astringent, not escharotic; a stimulant to the diseased surface, not a destroyer of its vitality. By occasional sponging with a solution of the perchloride, or a very dilute mineral acid, or a weak solution of lunar caustic,

ly hope to prevent the exudation reaching the larynx, or operation assuming an unhealthy character; but the power of means is allowed on all hands to be very limited. When the larynx is also involved in the exudation, dyspnoea, insufficient aëration of blood, add very materially to the dangers of the patient, and the probability of a fatal termination and the question naturally presents itself, whether any, and how much, benefit may be anticipated from the operation of tracheotomy. It is by no means easy to give in few words a satisfactory answer to this inquiry, or to lay down rules which may be sufficient to guide the practitioner in deciding on his course. To turn to statistics, we find that the fatal termination is not confined to any great extent, although in all probability some have been saved by the operation, which must otherwise have been lost. But it is manifestly impossible to frame a list of cases in which it has not been performed which shall be an exact counterpart to those operated upon; and without a basis of comparison, the knowledge of the exact number of deaths and recoveries after operation is valueless. It is asserted that of late years the mortality in France after tracheotomy is not nearly so great as formerly; but this may be due not so much on the results being more favourable, as on a more hopeful series of cases being selected. In such circumstances we must be guided more by general principles than by mere experience, and the rules for our guidance must be admitted to be founded partly on conjecture. It may be assumed, then, first with tolerable confidence, that when the general symptoms indicate that the attack is comparatively mild, while the danger of suffocation is imminent, tracheotomy does give a prospect of life in cases otherwise all but hopeless, and that it certainly does give prompt and certain relief to the suffering immediately caused by dyspnoea, than which nothing is harder to bear. But no surgeon ought to undertake the operation, in such circumstances, without fully explaining that relief is afforded only, from impending suffocation is its object. Usually, when the dyspnoea is less intense, it becomes a question whether the relief to the breathing may not help forward the process of cure which nature is working out; or it may be said, whether the existing amount of dyspnoea does materially hinder the recovery, and render it more uncertain. French surgeons are disposed to answer in the affirmative; but the same opinion is shared by many among ourselves; but the

result, the issue of the operation having been per-
apt to tell unfavourably on the issue. In addition
must remember that the operation may even fail
in consequence of the trachea and bronchi being
false membrane. In such cases it can only
intense dyspnoea and impending suffocation, and
longing of the patient, or the friends, to have
to procure relief. Could we know with any degree
how far the exudation extended into the tracheal
tubes, we should have most valuable information for
decision. Auscultation should with this view be
fully practised; but it must be confessed that this
is a matter of extreme difficulty, and one in which
an experienced stethoscopist may fail to elicit any
trustworthy information.

5. During the prevalence of the epidemic may
occur, which assume more or less its specific character
marked by the presence of small shreds of lymphatic
An ordinary sore throat at such times does not
course, but without manifest exposure to infection
assimilated to diphtheria. These generally require
treatment, and will in all probability end in recovery
the aid of medicine. We have only to remember
to depression which accompanies all the well-marked
of the disease, and therefore in all cases we must
any lowering remedies. No good can result from the
application of escharotics, while some risk is run


CROUP.

General characters.—With the disease known by the name of croup, or tracheitis, as it has been sometimes called, the cynanche trachealis of Cullen, English medical men have been long familiar. Although occurring in particular localities at certain times with greater frequency than elsewhere, it presents none of the true features of an epidemic; it is much more truly endemic. It appears chiefly during the cold weather; it may visit the same locality year after year, wholly disappearing during the interval; but it does not spread to adjacent houses or villages, its victims being comparatively few, although several persons are usually attacked at or about the same time. In all these respects it has much more affinity with pneumonia than with such a disorder as diphtheria. Like the latter, however, it is marked by one very striking feature, viz. the existence, in a large number of instances, of false membrane as an exudation from a mucous surface.

With our present knowledge it is impossible to assign any reason for this peculiarity. There is no specific poison apparently introduced; the mucous membrane lining the larynx and trachea is simply inflamed; and in many cases the exudation is purulent or creamy-looking; but in the majority it is serinous, contrary to the usual laws governing the inflammations of mucous membrane. Why this is so, we know not; for an acute attack of bronchitis or lobular pneumonia, both not uncommon in children of the same age, is not attended by the same sort of exudation from the mucous surface of the small tubes. It is usually only when the inflammation attacks the trachea, and is localised there, that false membrane is formed; but cases are occasionally met with in which the same material may be found lining the bronchial tubes down to their minutest ramifications. It is scarcely to be hoped that we shall ever attain to a knowledge of the circumstances which determine why one exposure to cold is followed by inflammation of the parenchyma of the lungs, another by that of the lining of the all tubes, while in a third the membrane of the trachea is the seat of the inflammation. It is this tendency to localisation which constitutes the specific character of each

The attack ordinarily sets in with acute fever and a hard dry cough, which has a very peculiar sound, dependent on the changes which from the larynx and trachea. The sound of the cough, that, when a child produces this brassy to the attendant is tempted to conclude, without that the disease is croup. There is very general in swallowing, any feeling of soreness of throat to the windpipe, and not reaching the fauces instances, however, the inflammation is more general membrane is seen on the tonsils and the back of the throat as ordinarily happens in diphtheria. The breathing is hurried, partly as a consequence of the fever because air enters the lungs with difficulty. As the disease proceeds, this difficulty increases, and the dyspnoea, which become every hour more distressing, paroxysms often follow a fit of coughing, but come on without any apparent cause. In the later stages shreds of false membrane are often seen, and cough sometimes follows its removal; but in the severe disease the relief is only temporary—the exudate as soon as it is rejected, the breathing becomes oppressed, the skin acquires a dusky hue, and is covered with cold perspiration. After several ineffectual struggles for breath, the brain seems to give way by the circulation of imperfectly-ventilated blood, and the patient sinks into unconsciousness.

larynx, we almost at once assume that in an adult we do with laryngitis, in a child with croup, when febrile signs are present, and there is nothing visible about the larynx to account for the dyspnœa. Indeed we need hardly go far in distinguishing these two diseases, because if false croup be present, its appearance in the expectoration will at once remove any doubt. From bronchitis and pneumonia are distinguished by the cough, by the existence of dyspnœa, the fact of this dyspnœa being traceable to the larynx. True or false croup is much more sudden in its onset, more violent in character; the one disease is essentially febrile and inflammatory, the other spasmodic, free from fever, and dependent on nervous irritability. Between diphtheria and croup the relations are very close; and although it be quite true that the natural site of the exudative inflammation is in the former disease, in the latter the trachea, yet there are instances in which it presents the name given to the attack must depend on the consideration, whether the case occurred sporadically or was one of such a number as could only be caused by an epidemic influence. The cases which cannot be determined by the above rule are comparatively few; and during an epidemic, the cases of croup may very generally be distinguished, even when the epidemic element impresses on them, to a certain extent, its own characters and features. The chief symptoms on which must be relied on for this discrimination are also those on which the treatment of the two forms of disorder is in great measure based; they are those which characterise the one as a spasmodic and the other as an inflammation. How much there is in the distinction those only can tell who, regardless of theories, observe the actual condition of patients under either form of disease, and can feel assured, as they certainly may be, of the effects of depletion and starvation in the one case, of the absolute necessity for supporting and sustaining in the other. The condition of the skin and the pulse help us more in this matter than any other set of symptoms; fever is associated with more burning and pungent heat of skin, with greater frequency and less force of pulse from the earliest period of the attack, and is not associated with perspiration. But though these point the way to reference to treatment, it is by constant and repeated observation alone that we learn in each case how far the type of disease is of a sthenic character, and are able to determine how far the patient will bear for a little while mere watching.



may be. The state of the skin and pulse, and breathing, lead us most naturally to salines as the best means of accomplishing this end. In healthy and robust, a few leeches to the throat, in emptying the vessels and checking the extent to which these remedies are employed, more on the constitutional strength of the patient. In diphtheria prevailing at the time, their use is still more sparing, because of the known tendency which accompanies it, and seems to modify the time being, any coincident cases of sporadic croup, or any such complication, the mere fact of sequestration, with its exhausting and debilitating action, must put us on our guard against allowing it to be too much lowered by depletion, and demand a plan of treatment at a comparatively early period in the susceptible and rapidly-changing structure. The intention of this part of treatment is, if possible, to arrest the febrile action before such an amount of effusion has poured out as may run the risk of producing permanent disease. We do not hope or expect to arrest it altogether.

Mercury has so long enjoyed a reputation as a powerful solvent of fibrinous exudations, that it is almost always used in croup; but one decided objection presents itself, it is too slow to be very available. If the patient is strong enough to be brought fully under its influence, the chance in favour of recovery is already considerable, and

DISEASES OF THE LARYNX

IN Laryngeal Diseases the ordinary symptoms, such as cough, difficulty in breathing and in swallowing of the voice, &c., often fail to indicate the precise seat of the malady.

Moreover, irritation of or pressure upon the other nerves by tumours, aneurisms, &c., as well as various morbid conditions of the nervous system generally, may produce symptoms which more or less closely simulate those produced by actual disease of the larynx itself. Hence it happens that some aid to diagnosis must be obtained before treatment adopted is likely to be as successful in its results as is necessarily uncertain in direction. Such aid is afforded by the Laryngoscope. This instrument was suggested by Riccati, and repeatedly suggested—many years ago, but it has only recently come into general use. Its great value as an aid in the diagnosis and treatment of diseases of the larynx, thus established, was not yet generally recognised and acknowledged when the First Edition of this Work was published.

Historical details relative to the invention of the laryngoscope would be altogether out of place in these pages.* It may, however, be stated, that Dr. B. G. Babington appears to claim to be considered the first successful laryngologist.

* The history of Laryngoscopy is more or less completely discussed in the following works:—*The Use of the Laryngoscope*, by Morell Macdonald, 2nd edit. London, 1866. *The Laryngoscope in Diseases of the Larynx*, by G. D. Gibb, M.D., 3rd edit., 1868. Art. by Mr. Windsor in the *British Medical Review*, Jan. 1863. Art. 'Laryngoscope' in the *Dict. encyclop. médicales*. Paris, 1868. *Klinik der Krankheiten des Kehlkopfes*, by Türk, Wien, 1866. *Die Laryngoskopie*, by Dr. Bruns. *Tul. Lehrbuch der Laryngoskopie*, by Dr. Tobold. Berlin, 1863. In many of these works may also be found much valuable information relative to the comparative practical uses of the laryngoscope, as well as to the diagnosis and treatment of Diseases of the Larynx. Constant reference has been made to them in the following pages.

† *London Medical Gazette*, vol. iii. p. 555. London, 1829.

er, it is impossible to omit an expression of grateful acknowledgment of the labours of Professor Czermak, formerly of * To him, probably more than to any one else, we are indebted for the earliest complete demonstration of the general utility and value of laryngoscopical examinations, and the methods of making them, and still more, perhaps, for the rested and painstaking manner in which, both on the continent and in this country, he has published his results, and fully taught his method.

LARYNGOSCOPY.

Laryngoscope essentially consists of the laryngeal speculum together with some contrivance, by means of which a light can be thrown upon it, when introduced into proper position in the pharynx.

Laryngeal speculum is a small plane mirror, fixed to one end of a slender but strong stem of sufficient length. The extremity of the stem is fitted into a light handle, either directly, or in such manner that it can be removed or added at will. The mirror may be of polished steel, speculum or silver; or it may be made of silvered glass (looking-glass mounted in a thin metal frame. The metallic mirrors readily warmed, retain their heat well, and are so thin as to occupy but little space. On the other hand the silvered glasses are much less liable to get tarnished or scratched; no real disadvantage arises from their comparative thickness, there is no difficulty in keeping them of such temperature as to prevent the deposition of moisture upon them. They are, therefore, as a rule to be preferred.

In shape the mirror may be circular, ovate, elliptical, or quadrilateral with rounded angles. The three first named forms are commended by Türck,† the last by Czermak.‡ For all laryngeal examinations the circular and quadrilateral forms are best, and they answer equally well. In cases, however, in

Zeitschr. der Ges. der Aertze, No. 17. *Der Kehlkopfspiegel und seine Ver-
wendung für Physiologie und Medizin*. Leipzig, 1860. *Du Laryngoscope*.
1860. A translation of this work was published by the Sydenham
Society in 1861. Also many subsequent contributions to Foreign and British

Méthode pratique de Laryngoscopie, par le docteur Türck. Paris, 1861;
cit. p. 38.

Laryngoscope, par le docteur Czermak. Paris, 1860.

form with the diameter of the mirror drawn at junction an angle of from 120° to 125° . Morec should be so inclined as to look somewhat downwards when the stem is held horizontally. It is advantageous to be of such material that it may be bent in one direction or the other, so that different degrees of inclination may be obtained for the mirror according to the requirements of the examination.

For the illumination of the speculum when either direct or reflected light may be employed in either case, if the light should not be very powerful the illumination of the rays is desirable.


Direct sunlight answers perfectly, but is seldom used. Some other source of light *must* frequently, and is commonly employed. Diffused daylight is rarely, if ever, employed. The light of a lamp concentrated through a glass globe filled with water,—both lamp and globe being behind the observer, the light coming over his head,—was formerly recommended by Türck.* A somewhat improved arrangement is advocated by Dr. Wainwright.† But the inconveniences of any such arrangement are obvious. Moura-Bourouillon describes, in his 'Laryngoscopy,'‡ an instrument which he names 'laryngoscope.' This consists of a large biconvex lens which brings to a short focus the rays of light from a lamp placed to the stand of which it is so connected as to be moved and moved in various directions. This and

ment is commonly used in Paris, and is recommended by Krishaber.* The modification consists in the addition of a concave reflector behind the light, and a shade which surrounds the lamp to such an extent as to protect the eyes of the observer from any direct rays. The condensing lens, the reflector, and the shade, are all attached to a metal ring, which is capable of being adapted and fixed by screws to almost any lamp.

Various other contrivances have been from time to time adopted for the direct illumination of the laryngeal speculum. But all are open to the objections that in their application the direction of the light is too little under the ready control of the surgeon; and too much depends upon the maintenance of the patient in proper position. He cannot be examined when recumbent. Moreover, the head of the surgeon is apt to obstruct the light if its source is behind him; and the lamp may be awkwardly in the way if placed in front of him. These and other objections are more or less completely avoided by employing reflected light for the illumination of the speculum. Hermann first suggested the use of a modification of Ruete's large ophthalmoscopic reflector for this purpose. And some such method as that which he originally suggested still appears the most convenient and most generally applicable. At any rate it is the method commonly adopted in this country and in Germany.

The source of light is placed on one side of and somewhat behind the head of the patient, who should, if practicable, be seated near the corner of a table. The rays are then reflected from and concentrated by a circular concave mirror duly arranged immediately in front of the face of the surgeon. The mirror should be three or four inches in diameter, and should have a focal distance of about twelve inches. It may be perforated or not in the centre. If perforated it may be placed in front of one eye, and the view may be taken through the perforation. If not perforated it may be arranged in front of the forehead and nose, and between the eyes; or lower down in front of the nose and mouth. In either of these arrangements both eyes may be used; in the former looking below, in the latter above the reflector. The perforated reflector should theoretically give the more perfect view. But practically the

* See Krishaber in the *Dictionnaire encyclopédique des Sciences médicales*, t. 'Laryngoscope.' Paris, 1868.



a strong spectacle frame, an elastic frontal spring passing over the vertex. The first adapted for the perforated reflector; the second perforate reflector in front of the forehead; the reflector is worn lower down. Czermak's fixing the stem of the reflector at right angles to be held between the teeth may be convenient. In any case the reflector should be connected either by a ball and socket joint, or in such a way as permits it to be turned easily in any required position.

As a source of light, any lamp which burns a steady, full bodied flame, may be employed. A Bunsen or reading lamp, or an Argand gas burner, will serve well for all ordinary purposes. But it is desirable that the lamp be so arranged that it can be raised or lowered and moved from side to side. Mackenzie's 'laryngoscopic lamp,'* which readily admits of vertical and horizontal movement, is admirably adapted for the consulting room.

The light of the lamp may be advantageously concentrated by means of one or more lenses placed in front of it and a concave reflector or white reflecting surface behind it. Of all the so called 'light-concentrating lamps' I believe Mackenzie's is the best for the purpose. It consists of a metallic cylinder having a shoulder at one end which bears a plano-convex lens two or three inches in diameter. The cylinder replaces the glass chimney of the Argand lamp.

d plano-convex lens, and a concave reflector attached to a metal ring, which by means of a spring clip or screw can be easily fixed to the brass collar of the lamp glass. Tobold's apparatus contains two lenses. It is very efficient in working; it is heavy and cumbersome, and can only be applied to a lamp especially adapted for it.

For purposes of demonstration, and sometimes in operating, it is advantageous to have command of a more brilliant light than any ordinary lamp can afford. Such a light may be obtained by means of the electric, magnesium, or oxyhydrogen burners. Of these the last named, or some modification of it, is the most easily available and most manageable. The apparatus employed at the Hospital for Diseases of the Throat answers admirably. The so-called 'Medical Lantern' of Dr. Thudichum,* which he especially recommends for use in rhinoscopy, is usually adapted for laryngoscopical and other examinations. In this apparatus the hydrogen flame is conveniently replaced by that of a spirit lamp.

In cases in which the diagnosis is doubtful, or in the treatment of which difficult operative measures are needful, it is sometimes well to wait until sunlight is available. In making examination by sunlight, a plane rather than a concave reflector should be used. The latter throws too brilliant a light; moreover the rays concentrated by it may possibly burn the patient if proper care is not taken.

In all cases in which sunlight, or any other powerful light is employed, the eyes of the surgeon should be protected by a shade, or by the hollow conical eye protectors,† which are described in use in the chapter on RHINOSCOPY (p. 258, fig. 227). When a comparatively feeble light is employed it is advantageous to darken the room in which the examination is made.

The general arrangements for making an ordinary laryngoscopic examination by means of the reflector and artificial light are represented in Fig. 254.

In the lamp, the mouth of the patient, and the eyes of the surgeon should be as nearly as possible in the same plane. If the laryngeal speculum is held in the right hand of the surgeon, the lamp should be placed on the right side of the patient, and

* Made by Orchard of Kensington. See Thudichum *On Polypus of the Nose* (Lancet, 1869), p. 5. Churchill, 1869.

† Made for me by Messrs. Weiss.

vice versa. If, as may sometimes be convenient, the larynx is placed above and behind the head of the patient, the same plane must be preserved; but the speculum may be held with equal advantage in either hand without risk of obstructing the light.

The patient should if possible be comfortably seated opposite the surgeon in a nearly erect position. His body should be somewhat forwards, and his head should be inclined slightly backwards; but should not be thrown so far back as is often done. Sometimes it is advantageous to employ a 'head rest,' which may either be fixed to the back of the chair, or may have a separate stand. When in proper position, the patient is directed

FIG. 254.



to open his mouth as widely as possible, and to breathe quietly and deeply. His tongue may be protruded, and held by himself or the surgeon, the fingers and thumb being covered by a soft cloth or handkerchief to prevent slipping; or it may be kept in the mouth, and gently pressed forwards against the lower incisors, an attempt being made at the same time to depress and flatten, or render concave its posterior part. Sometimes one plan, and sometimes the other is found to be the more effectual. Meanwhile the surgeon arranges his reflector so as to throw the light directly on the open mouth of the patient. When he has succeeded in illuminating the middle of the soft palate, the uvula, and posterior pharyngeal wall by a bright disc of light, he may proceed to introduce the laryngeal speculum.

The handle of the speculum should be held somewhat loosely 'like a pen' between the thumb and the fore and middle fingers.

one hand or the other. The speculum should be slightly warmed previous to its introduction, either over the lamp or by momentary immersion in hot water, in order to prevent deposition of moisture upon its surface.* Care must be taken, however, that it is not made too hot. Its temperature may be easily tested by the hand or cheek of the surgeon. The speculum should be introduced with its reflecting surface directed downwards and forwards. It should be carried backwards through the cavity of the mouth, deliberately, confidently, and steadily, and then applied with gentle but firm pressure against the uvula and neighbouring portions of the soft palate. Uncertain, hesitating, and hasty movements of the instrument are liable to give rise to objectionable titillation. In its passage through the mouth it should not come in contact with the tongue, nor indeed with any other parts than those against which it is to be placed. As a rule it should not be carried so far back as to touch the posterior pharyngeal wall. Sometimes, however, especially in certain exceptional cases, this may be done with considerable advantage. The hand of the surgeon must be kept somewhat down and well towards its own side, so as not to obstruct the light. The third and fourth fingers may rest against the chin or cheek of the patient. The stem of the instrument should lie in or near to the corresponding angle of the mouth. In a large proportion of cases no difficulty is encountered, and the patient suffers comparatively little inconvenience.

When the speculum is thus placed in position and well illuminated, it exhibits to the surgeon images of those parts of the larynx upon which the rays reflected from its surface are made to fall. It must be borne in mind, however, that as a necessary consequence of the position of the mirror in relation to the eye of the observer on the one hand, and to the larynx of the patient on the other, the parts appear in the image as though reversed antero-posteriorly. And they are thus conventionally represented in all ordinary laryngoscopic diagrams.

This so called 'reversal' or 'inversion' of the image is illustrated by Fig. 255, in which B diagrammatically represents

* Various expedients have been suggested with a similar view. Thus, Dr H. G. Wright contrived a speculum, the temperature of which was maintained by a wire placed behind the mirror and heated by galvanic action. Dr Buzzard recommends that the surface of the mirror should be covered with a of glycerine. But no plan answers better than the simple one above-mentioned; nor is any other so easy of application.

the parts as seen when simply looked down upon, and A the same parts as they appear reflected in the mirror. No practical difficulty or inconvenience results from this apparent inversion.

FIG. 255.




Diagram illustrating the antero-posterior reversal of the Laryngoscopic image. The small excrecence situated on the right vocal cord (n) might be falsely supposed to be on the left from the appearance presented in the mirror (A).

By varying the inclination of the mirror, and slightly changing its position from time to time, the following parts, or rather certain portions of them, may be successively brought into view; the base of the tongue and glosso-epiglottidean ligaments; the epiglottis and aryteno-epiglottidean ligaments; the cartilages of Santorini, and the arytenoid cartilages; the true and false vocal cords; the ventricles and anterior wall of the larynx; more or less of the anterior wall of the trachea; and if the glottis is very widely open and the light very good, even the bifurcation of the trachea, and a small portion of the right bronchus. These several parts, however, cannot all be seen with equal facility. Numerous rules have been laid down as to the

position and degree of inclination of the mirror best adapted for the special examination of each particular part.* But it is needless to repeat them. A correct knowledge of the relative positions and natural appearances of the parts, and a consideration of the simplest laws of catoptrics, together with a certain amount of practical experience, will enable the laryngoscopist to bring into view and recognise one part after another so far as may be practicable. Repeated attempts are often necessary before even partial success can be attained. And sometimes even the utmost perseverance only results in failure. But this seldom rarely happens. The possible sources of difficulty are numerous, and the actual difficulties may be great; but they may generally be obviated or overcome, at any rate to some extent, by patience and judicious management. Those who suffer from chronic laryngeal affections are as a rule the best subjects for laryngoscopic examination—not the worst as might be supposed. Their desire for benefit affords a stronger stimulus to patience and self-control than can actuate the healthy; and, moreover, they are usually already accustomed to expose their fauces for inspection, and to submit to local applications, which indeed the normal sensitiveness of the parts may have been diminished.

The most common sources of difficulty are as follows: general nervousness and excitability of the patient; fancied or real inability on his part to open his mouth sufficiently widely, and awkwardness in the management of his tongue; relative largeness, or some swollen condition of the tongue; malformation or deformity of the soft palate and neighbouring parts from ulcerative disease, and subsequent contraction of cicatrices; elongation and thickening of the uvula; enlargement of the tonsils; redundancy of the epiglottis; and, lastly, extreme sensitiveness of the uvula and soft palate, either natural or resulting from existing inflammation or ulceration.

Difficulties arising from the first three sources may be overcome almost invariably by patience and judicious management on the part of the surgeon, and a little practice on the part of the patient. Hastiness in manner, and in method of proceeding, should be avoided; and every effort should be made to encourage, reassure, and inspire confidence. By his own example

Krishaber, Art. 'Laryngoscope' in *Dict. encyclop.* Op. cit. p. 497 et seq.
 * Op. cit. and *Méthode pratique de Laryngoscopie.* Paris, 1861.

the tongue may generally be kept out of the way. If these methods fail, and the tongue still rises to obstruct the view of the soft palate, &c., the patient is directed to make a few deep inspirations, and to breathe through the mouth only; the tongue is then depressed, and the larynx raised, and a favourable position is obtained. Sometimes a broad spatula may be advantageously used. But the introduction of any unnecessary instrument should be avoided.

Occasionally by sliding the finger along the speculum being at the same time introduced, an effort at vomiting is produced, during which the tongue is depressed and the larynx raised, and a momentary view may be obtained.

In cases in which the soft palate is unusually small and thin, the speculum is apt to slip behind them, and to become partially hidden. In such circumstances it is better to hold the instrument as far as possible in its proper position without touching against the soft palate.

An elongated and thickened uvula may obstruct the view hanging below, and turning round the inferior margin of the speculum. The employment of a larger speculum, and manœuvring of the instrument serves to obviate this difficulty for the time being, and it may be necessary to use the finger to push the uvula up and out of the way.

Pendency or projection backwards of the epiglottis is a frequent source of considerable difficulty. Gibb states as the result of his observations that some such abnormal condition is to be found in eleven per cent. of otherwise healthy individuals.* It depends upon undue elongation of the glosso-epiglottidean ligaments or folds of mucous membrane, together with relaxation or weakness of the muscular fibres they contain. The degree of obliquity observed varies greatly in different cases, and to some extent in the same case under different conditions. In some instances the epiglottis hangs almost horizontally over the entrance to the larynx; but more frequently its deviation from the normal erect position is comparatively slight, or even only occasional. Laryngoscopic inspection is of necessity proportionately impeded. The greatest difficulty is often experienced in obtaining a satisfactory view, and sometimes even partial success appears to be impossible.

In all cases of pendent epiglottis it is necessary to carry the speculum further backwards and downwards, and to make its reflecting surface look more forwards than is desirable in the normal condition of parts. Sometimes its lower border must be placed against the posterior wall of the pharynx. When the speculum is in position, the patient should be directed to utter a series of short, high, *staccato* notes, or to imitate a shrill *falsetto* laugh. During the emission of such sounds the epiglottis is suddenly raised, and jerked forwards. Opportunity may be thus afforded for a succession of brief glimpses of the interior of the larynx, which may suffice at any rate to determine the diagnosis. Another method which sometimes proves partially successful is as follows. The head of the patient is thrown back as far as possible, and his tongue is projected and held by himself. The speculum is introduced and inclined slightly backwards and downwards, and the surgeon having lowered his head looks into it as it were from below upwards. At the same time with his free hand he presses the pomum Adami of the patient backwards and somewhat upwards.† There can be no doubt that the larynx may often be moved into comparatively favourable position for inspection by external manipulation; but the method is liable to occasion much discomfort, and sometimes considerable pain.

In some instances in spite of all attempts the epiglottis still

* Op. cit. p. 44.

† See Krishaber. Op. cit. p. 493.

persistently hides the larynx from view. And in such cases it may be desirable to raise it mechanically by instrumental aid. Forceps, tenaculums, and hooked sounds of various kinds have been devised for this purpose by Bruns,* Voltolini, Fournié, Lewin, Mackenzie and others. But all such instruments are difficult for the surgeon to use, and still more difficult for the patient to tolerate. It is only in very exceptional cases that they can be employed with advantage. The epiglottis is extremely sensitive; but it bears firm pressure and a steady but gentle pull better than the uncertain titillation of a hesitating touch. If any attempt at seizure be made, the instrument should be carefully warmed previous to introduction.

Difficulties arising from unusual sensitiveness of the uvula and soft palate may generally be overcome by patience and dexterity on the part of the surgeon, and a little practice on the part of the patient. During the earlier attempts the speculum should not be maintained too long in contact with the parts. It should be at once withdrawn if decided retching is produced; and short of such effect, if the irritability is great, it is better to apply the speculum repeatedly for a moment or two, and remove it as often without caring to obtain any view until the parts have become as it were accustomed to the presence of the instrument. When once efforts at vomiting have been excited, there is little probability of success until a future occasion. Preparatory measures of various kinds have been from time to time suggested; some of these are probably useful; others have been proved by experience to be altogether useless. Many owe their accredited efficacy to the confidence with which their exhibition has inspired the patient. The internal administration of bromide of ammonium or bromide of potassium in full doses has been strongly recommended and extensively tried, but with variable results. Gargles containing bromide of ammonium are said by Gibbs to produce 'moderate anæsthesia' of the fauces in twenty-four hours.† Türck reports favorably of the repeated application to the fauces of a mixture (recommended by Bernatzik) which con-

* Bruns appears to have been the first to have used an instrument of this kind; and he did so in the case of his own brother who suffered from polypoid growth in the larynx. The epiglottis was elevated by means of a pair of flat-bladed forceps, and the growth was successfully removed. See *Die erste Ausrottung eines Polypen in der Kehlkopfshohle*. Dr. Bruns. Tübingen, 1862. Also *Die Laryngoskopie*. Dr. Bruns. Tübingen, 1865, p. 257.

† *The Laryngoscope in Throat Diseases*. Op. cit. p. 46.

sists of three grains of hydrochlorate of morphia, one drachm of alcohol, and half an ounce of chloroform.* But all such applications as this would appear more likely to inflame than to soothe the sensitive parts. Alum gargles, or solutions of alum or some other astringent in the form of spray are often useful. Occasionally a few whiffs of chloroform may be advantageously inhaled. But by far the best and easiest plan, so far as I know, is to direct the patient to suck a little ice immediately before submitting his throat for examination. The result is generally satisfactory.

Besides the various sources of difficulty thus discussed, there are others which depend upon inexperience and want of dexterity on the part of the surgeon. These can only be obviated by careful practice. The surgeon should not only familiarize himself with the management of the laryngoscope, but also with the relations and appearances presented by the healthy living parts when viewed by its aid. In colour especially, the mucous membrane of the larynx seen during life in the laryngeal mirror looks very different to what it does when examined after death on the *aut mortem* table. It is very possible for the tyro in laryngoscopy to mistake the natural tint of some parts for a certain degree of inflammatory redness.

The general appearances and relations of parts as seen in the laryngeal speculum need no special description. They may be learnt by the careful study of a larynx removed from the body, or by *auto-laryngoscopy*, or by the examination of the larynx of some living subject. But with regard to the colour presented by the different parts in a state of health, the following observations may be made.

The lingual surface of the epiglottis appears of a yellowish or pinkish drab colour. Its upper border is decidedly yellow. Its aryngal surface, especially the 'cushion,' varies from a pinkish yellow to a deep pink; sometimes it appears bright red—so bright, indeed, as to suggest the idea of the existence of an inflammatory condition. The aryteno-epiglottidean folds are pale pink. Stoërk accurately describes them as being about the same colour as the gums. The mucous membrane covering the arytenoid cartilages is still pink, but of a somewhat deeper tint. The false vocal cords are perhaps slightly deeper pink still. The true vocal cords are distinguished by their glistening pearly whiteness; but some-

* *Allgem. Wiener Med. Zeit.*, p. 98. 1863.

ing the laryngeal mirror in a position so as to
of the surgeon free; and of depressing the
time the laryngeal mirror is introduced. It
useful; others are rendered unnecessary by the
assistant. For descriptions of all such contrivances
may be made to the special treatises already cited.

GENERAL REMARKS ON THE TREATMENT OF LARYNX.

The laryngoscope affords most valuable aid
diagnosis, but also in the treatment of the
the larynx. Guided by the view obtained
enabled to make such local applications as may
with accuracy and precision, and to perform
operations—such as scarifying the mucous mem-
abscesses, removing growths, &c., with certainty.
The use of the laryngoscope has altogether
doubts formerly entertained as to the practi-
cating brushes, probangs, and other instruments
larynx.* Such doubts have in past time un-
vented the adoption of methods of treatment
cases, might have been eminently successful.
it would appear that the danger is in the over-
and that the tendency is rather to carry out too
vigorously, or to rely upon it too exclusively.

Remedial applications may be made to part

Solid substances may be applied either in the mass or in powder. If applied in the mass, some form of caustic holder is requisite. No instrument answers so well, and at the same time is so safe, as a piece of moderately thick aluminium or silver wire mounted in a slender handle, and hollowed into a shallow cup, or roughened at the extremity. The roughened extremity may be dipped into various substances, as nitrate of silver, chloride of zinc, &c., while in a state of fusion; and a sufficient quantity may be taken up in a bead-like form, or as a thin coating. The wire may be easily bent at any angle requisite, and there is no danger of any considerable portion of the substance breaking off and falling upon parts it was not intended to reach. Such danger might arise during the use of some of the various complicated caustic holders that have been devised for similar purposes. By aid of the laryngoscope a small caustic bead, prepared in the manner described, may be directly applied to any particular point; or if the wire is heated to a sufficient extent, the general surface may be wiped over.

Powders can only be applied to the mucous membrane generally. They may be inhaled through a tube, as recommended by Fournié, or, far better, blown in by means of Achuss's insufflator* or some modification of it. The insufflator, duly charged, is introduced into the back of the fauces, with the extremity turned down towards the larynx. The patient is then directed to make a slow steady inspiration, and while he is doing so the india-rubber ball of the insufflator is suddenly compressed. Under such circumstances it is obvious that a considerable portion of the powder must be diffused over the mucous membrane of the larynx. The powders that have been used with most advantage in this manner are tannic acid, iron, acetate of lead, sulphate of zinc, sulphate of iron, calomel, and in some exceptional instances, nitrate of silver. It is especially needful to add that in every case the substance used should be very finely pulverised. Sometimes it may be advantageously diluted with two or three or more parts of sugar milk.†

Liquids, or substances in solution may either be applied by

This instrument is figured in the *Essay on DISEASES OF THE NOSE*, p. 278. Studley, *American Medical Times*, March 2, 1861.

useful of such solutions may be enumerated the silver (ʒij. or ʒij. to ʒj. of distilled water), potassium (ʒij. or more of the tincture to ʒj.), pure carbonate of copper (a saturated or weaker solution), or honey may be advantageously mixed with in considerable proportion, in order to give them render them more adhesive. Iodine (gr. xx.) potassium (gr. v.), and olive oil (ʒj.), form an anodyne recommended by Dr. Marcet.† The 'glycerinum' (B. P.) may also be used in some cases. By means of all such substances may be applied with a precision to any particular part. In some instances, acetic acid, chromic acid, and other powerful escharotics thus applied to ulcers, warty growths, &c., may be of advantage. At the moment the brush is to be applied to the larynx, the patient should be directed to take a deep inspiration in order to dilate the aperture as well as possible. In cases in which the epiglottis is pendent, it is difficult to get the brush beyond it into the larynx. In such circumstances the laryngeal syringe may be used, inasmuch as its extremity may be guided to raise the pendent valve to a sufficient extent.


The inhalation of 'atomised fluids,' or sprays, for the treatment of diseases of the air passages, though previously, was first shown to be generally practicable by Girton in the year 1858.‡ Since then the method

has been fully established, and various improvements from time to time effected in the apparatus employed.* 'ray-producers' at present in ordinary use, consist of two (Bergson's tubes) fixed at right angles to one another; of one, placed horizontally, is connected at its distal extremity with some apparatus by means of which a strong current of air or steam can be continuously projected through the other, placed vertically, dips into a bottle or other vessel containing the medicated fluid. The proximal extremity of each tube is terminated by a very fine orifice; and

FIG. 257.



FIG. 256.



Arm of Siegle's steam spray-producer.

Siegle's steam spray-producer on stand, with Beigel's face-screen attached.

ice of the former tube is immediately over that of the

When a current of air or steam is made to rush forcibly from the orifice of the one tube over that of the other, it causes the medicated fluid to rise, and breaks it up as it issues, and discharges it in an exquisitely fine spray. In Dr. Andrew Clark's own instrument (as well as in the various modifications of the apparatus used by different makers), air is employed, and the current

Matthieu's modification of Matthieu's, which is worked by means of a pressure-pump: and acts on a principle quite different from that described above. The fluid is forced through a fine orifice into a small chamber, and is then forced against the interior of a vulcanite 'drum.' It thus becomes broken up into a fine spray. Op. cit. p. 90.

like the handball spray-producers: and of importance, the atomised fluid is mixed with a current of warm steam instead of cold air. Of different solutions, the strength being in different circumstances, may be advantageously inhaled by spray. Among them may be especially mentioned—the medicament being in each case in alcohol specified to an ounce of water:—Alum (gr. i. to gr. ii.), acid (gr. j. to gr. xx.), perchloride of iron (the tincture ℥ x. to ℥ xxv.), as ‘astringent’, or chloride of ammonium (gr. x. to gr. xxx.) borax, or iodide of potassium (gr. iij. to gr. i.), and watery extract of opium (gr. $\frac{1}{4}$ to gr. $\frac{1}{2}$), morphia (gr. $\frac{1}{8}$ to gr. $\frac{1}{3}$), or fluid extract of hyoscyamus (℥ iij. to ℥ x.), as ‘sedative’. Very weak watery solutions of iodine, and of sodated soda), tar water, lime water, weak alkaline water, the saline waters of many mineral springs, solutions too numerous to mention have all been used in different cases with variable advantage. The handball spray has recently been very strongly recommended for the treatment of various affections of the throat by Dewar and others. But experience seems to show that caution is requisite in the use of this method, though doubtless beneficial in some cases, in many

only less efficacious, but far more mischievous than was anticipated. Solutions of nitrate of silver and other powerful metallic salts are unsuitable for application by means of the spray-producer.

Many substances may be inhaled with very great advantage in the form of vapour or gas. In various acute or subacute affections nothing is more soothing and beneficial than the almost constant inhalation of warm steam, either plain or charged with some narcotic or anodyne element. The 'hemlock inhalation' of the British Pharmacopœia is very useful; it is rendered less disagreeable and perhaps more efficacious by the addition of a pinch or two of fresh dried hops; or the hops may be infused alone in the hot water; or a portion of opium may be added; in some cases vinegar is useful and pleasant. In chronic cases, according to their nature, the vapour of turpentine, or some aromatic terebinthinate, or iodine, or calomel may be inhaled with very great benefit. For the inhalation of steam, plain or medicated, various so-called 'inhalers' have been devised. Among those ordinarily used may be mentioned Nelson's, Maw's 'double-valved inhaler,' Mudge's, and Beigel's made by Robbins. The two first named are furnished with pieces of sponge which are intended to be imbued with the medicament to be used; but practically it is far better to remove the sponges and to mix the medicament with the hot water. The two last inhalers involve somewhat more respiratory exertion on the part of the patient. But they have the advantage of acting on the 'hookah' principle; the air to be inspired passes through the fluid, and thus becomes thoroughly warmed and charged with vapour. But by far the most perfect inhaler yet devised is that made by Messrs. Maw under the direction of Dr. Mackenzie, and by him termed 'the Eclectic Inhaler.' The advantages of this apparatus are so obvious that it will probably to a great extent supersede those hitherto in use. Bird's inhaling pipe is worthy of mention, as likely to prove very useful for the exhibition of certain vapours, &c.*

Scarification of the mucous membrane of the larynx may be practised with considerable advantage in many cases. The instruments requisite and the method of operating have been already described in the Second Volume of this Work (p. 470). In all cases, if practicable, the instrument (which must be very

* Made by Maw. See *Med. Times and Gaz.* vol. i., 1869, p. 672.

The direct galvanisation of the vocal cords is recommended in many nervo-muscular affections of the most efficient apparatus for this purpose is the galvaniser of Mackenzie. One pole of the battery is with a properly curved metal wire terminated in a sponge which is placed in contact with the inner surface of the larynx. The other pole is connected to a handle bearing a sponge which is applied to the skin. When the two parts of the instrument are brought into contact the circuit is completed by pressing upon a small handle of the laryngeal portion of the instrument. The galvaniser may be applied several times at a time, being kept in for a few seconds each time.* (There have been devised with similar intent, but the Mackenzie appears to have found most favour with practitioners. It is recommended by George Johnson, Smyth, Marcet, Gibb, and others.

CATARRHAL LARYNGITIS.

Acute Catarrhal Laryngitis is a formidable, and dangerous affection. It varies greatly in intensity, and in the degree of danger to which it gives rise. It may attack almost any period of life, but is most frequent in adults of plethoric habit.

Exposure to cold and damp, especially after exertion of the voice, and when the general power is weak,

The patient complains of dryness, soreness, and a sense of constriction about the upper part of the throat, together with tenderness on pressure over the larynx, and pain and difficulty in swallowing. The feeling of discomfort is constantly referred to the *pomum Adami*, or its immediate neighbourhood. Respiration soon becomes impeded; the patient experiences a sense of oppression, and from time to time feels it needful to make an effort to breathe deeply. His voice becomes thick and husky; and occasionally he tries as it were 'to clear his throat' by a dry, harsh, or half suppressed and painful cough. Accompanying these symptoms there is more or less general febrile disturbance.

As the malady advances, the symptoms increase in severity; and the patient becomes restless and anxious. The face is flushed, the skin hot and dry, the pulse hard and quick. Respiration becomes more and more seriously impeded. Inspiration is difficult, protracted, and painful, and is accompanied by a wheezing, whistling, or harsh throttling sound. The chest is not fully expanded; and there is diminution of the respiratory murmur. Expiration remains comparatively easy; for the warm, moist air from the lungs irritates the inflamed and sensitive mucous membrane much less than the colder, drier air, inspired from without. The dyspnoea is constant in so far as it depends upon constriction of the glottis from swelling of the mucous membrane; but it becomes greatly aggravated at intervals in consequence of spasmodic action of the muscles. The voice, at first husky and hoarse, soon becomes low in tone, or 'cracked' and uncertain, and finally is altogether lost, the patient speaking only in a whisper. In some cases there is but little cough. In others the cough is frequent, and harsh, husky, or stridulous in character. Cough may be provoked either during inspiration by the contact of cold dry air, or during expiration by the presence of little masses of secretion. The sputa in the earliest stages are very scanty, and consist of little more than saliva; later they become somewhat more abundant, but are still semitranslucent, though more or less tenacious and viscid, containing laryngeal mucus as well as saliva; still later they are blue and greyish, and sometimes are slightly streaked with red. After any of these have been coughed up a painful sense of dryness about the part is often experienced. The small, agglutinated masses which are occasionally coughed up, notably come, according to *Krishaber*, from the ventricles of

THE LARYNX.

swallowing, almost invariably the symptoms, becomes more and more fully progresses. '*Difficulty of breathing*' cause is visible in the fauces and the chest, '*breathing*' for which no adequate cause in the thorax,' are, in the words of St. Asaph, the earliest of the symptoms that but rarely excite alarm.*

As the disease advances to the next stage, and no relief is afforded, the distress of the patient becomes intense, and he gasps for breath. His anxiety and restlessness increase. He cannot lie down; or if, exhausted, he starts up involuntarily, gasping for breath, his face becomes pale and livid, or even cyanotic; sweat pours from his forehead; he is clammy; his pulse becomes weak and irregular, drowsiness, and perhaps delirium supervene, he dies suffocated—either almost suddenly, or by a process of comparative duration.

The changes which give rise to the symptoms that attend this disease during life by aid of the laryngoscope, and which can only be ascertained with certainty on post mortem examination.


On inspection the mucous membrane of the larynx is seen to be of an unnatural scarlet red colour, and is thickened or swollen. Occasionally superficial ulceration is recognised here and there; but neither deep ulceration is observed, or only extremely rare. The swellings, as a rule, are especially manifest in the lower part, in which the mucous membrane is thick, and is surrounded by abundant loose submucous areolar tissue. The epiglottidean folds, the mucous membrane of the aryepiglottic folds, the vocal cords, and the laryngeal aspect of the epiglottis, in these appearances in the most marked degree. When the disease is in its most advanced stage, and the speaking, they are affected in extent in the order named. The mucous membrane of the trachea, on the other hand, being comparatively thin, and surrounded with dense subjacent tissues, is less swollen, and is only slightly reddened. Sometimes it is marked by redness. Sometimes it seems only to have lost its

The swelling soon becomes erect, rigid, and prominent. It is not only unfitted to fulfil its natural valvular office. Some-
times may be felt by the finger as a smooth, rounded, 'cherry-
tumescence. Occasionally it gives rise to sensations
such as those produced by the presence of a foreign body.
The patient 'feels as though he had something in his throat.'
The condition of the epiglottis serves in great measure to
increase the difficulty in swallowing, and the sense of suffocation
experienced when the attempt is made. The dysphagia
is further accounted for by the pain caused by the neces-
sary movement of the inflamed and unduly sensitive parts.
The swollen condition of the mucous membrane and con-
striction of the glottis manifestly give rise to the
difficulty of breathing; and the temporary paroxysms
of dyspnoea are caused by muscular spasm incidentally
and mentally excited.

Post mortem examination shows that the swelling depends upon
a congested, infiltrated, and oedematous condition of the mucous
membrane and submucous tissue. The redness of the parts
during life is often partially or wholly lost—inasmuch
as the mucous membrane shares in the general pallor of death.
No purulent, or viscous exudation is usually found
on the surface of, or to some extent the membrane. In some cases the
trachea and bronchi are also affected; and the lungs are
inflamed, or even pneumonic.

The diagnosis of acute catarrhal laryngitis is not difficult. It





fauces. In laryngitis, on the other hand, the an in the upper part of the throat (if even an gether insufficient to account for the extre breathing and swallowing, and the alteration the voice. It must be borne in mind, however or at any rate cedema of the larynx, may coe mation of the tonsils and mucous membrane c pharynx.

In simple pharyngitis there is little or no swallowing is painful and difficult, and pain larynx is pressed backwards.

The *course* of acute catarrhal laryngitis is u duration depends upon the intensity of the i constitution of the patient, and the nature an treatment adopted. If unchecked it often pr fourth or fifth day, and sometimes much mor celebrated General Washington died of thi twenty-four hours. Sir Thomas Watson allud in which a fatal result ensued in twelve hours.

Mr. Gray mentions the case of a man who was 'admit Hospital complaining merely of sore throat. He walked seemed in such good health that he would not have been circumstance of his having come some distance from the c hours after his admission, the house surgeon was summons him, as he was said to be dying of suffocation. He went in the patient quite dead. Post mortem examination reveal

In some instances, however, the malady runs a less rapid course, and the patient survives until the eighth or ninth day. In others, and these happily not a few, the inflammation subsides either spontaneously, or as the result of treatment. In such case the effects may disappear almost altogether with comparative rapidity, or the malady may assume a chronic form.

Treatment.—The treatment requisite depends upon the severity of the malady, and the stage it may have reached. Measures which are urgently demanded in some cases are in others unnecessary, and even dangerous. And the mode of treatment that might be most efficacious at the onset of the malady, if adopted too late, may only hasten the fatal termination.

In the earlier stages, and in cases in which the symptoms are comparatively mild, perfect quiet in a warm moist atmosphere, soothing inhalations either of simple or medicated steam or spray, and the administration of salines, with perhaps antimonials and tincture of aconite or some sedative, according to circumstances, constitute the treatment requisite. But in all cases careful watching is necessary; for dangerous symptoms may supervene at any time, and even almost suddenly.

In the more severe forms of the malady, similar but more energetic measures must be rigorously enforced. Soothing inhalations may still be advantageously employed, but their use must be more constant. Antimony and tincture of aconite in small but frequently repeated doses should be administered, especially in cases in which there is much general inflammatory fever.* Free mercurialisation has been strongly advocated, but its efficacy appears doubtful; and much precious time may be lost before the system can be affected. General bloodletting, so highly lauded in past times in the treatment of this disease, may be resorted to in the earlier stages. But satisfactory evidence in favour of this proceeding is altogether wanting. Sir Thomas Watson says:—

‘When there is high inflammatory fever present, and the skin is hot, and the pulse firm and full, and the cheeks red, and the lips florid, you may bleed your patient with decision and advantage. But if his powers are beginning to sink

* The following formula may be recommended:—Vin. antim. tart. ℥ ij. to ℥ iij., tinct. aconiti ℥ j. to ℥ ij., liq. ammon. acetat. ʒss.; to be taken in water or camphor julep every quarter of an hour or every half-hour, until some obvious effect on the circulation, &c. is produced, and then to be continued less frequently.

rather than less blood in the system. Local means of leeches to the neck or upper part, cupping to the nape of the neck, may sometimes be used. So also may blisters or milder counter-irritation. Should not be applied immediately over the larynx.

When the mucous membrane is so swollen that respiration and swallowing to a serious extent are impeded, by means of a sharp pointed guarded bistoury, a small incision, designed instrument, is often followed by speedy relief.

A somewhat remarkable case illustrating the effects of cold came under my observation in December 1866. A gentleman of Essex, fancying he heard poachers in the neighbourhood of the night, went out carelessly dressed, and concealed himself for several hours in a ditch. The next morning he was seized with intense laryngitis; and he died suffocated within eight days. Afterwards the brother of this gentleman, depressed in spirits and almost exhausted, was exposed for some time to cold and wet. He was then attacked by symptoms similar to those from which his brother suffered. The next morning I saw him in consultation with Mr. Jordison of South Ockenden. The soft palate, uvula, and aryteno-epiglottidean folds were in a similar condition. The swelling was extreme; breathing was carried on with great difficulty, and was almost impossible. I at once freely scarified the affected parts with a bistoury, and made several small incisions, and caused the patient to inhale steam of water. A considerable quantity of serous or seropurulent blood was discharged. In the course of half an hour the patient was much relieved, and swallowing was accomplished without difficulty. The swelling of the parts rapidly subsided, and the next day again, the patient could breathe and swallow with comparative ease, almost naturally—but he still suffered from considerable

Sir D. Gibb (following Dr. Horace Green) strongly advocates the application of a solution of nitrate of silver (three or four scruples to an ounce of water) in cases of acute laryngitis. The application may be made by means of a full bellied camel's hair brush on a curved aluminium wire stem, or by 'the laryngeal fluid pulveriser.' 'The effect of this proceeding is some considerable amount of burning heat, associated with comparatively little spasm, and sometimes dyspnoea, the last two persisting for may be a few seconds only. . . . The relief experienced, and the amelioration in the general symptoms is observed in periods ranging from half an hour to four hours, and the dyspnoea subsides very speedily.*

If, as too often happens, the malady advances in spite of all such measures as those above suggested, or if the difficulty of breathing is too urgent to admit of their adoption, tracheotomy must be at once performed. In no class of cases probably is the value of tracheotomy as a life-saving operation more obvious and more pronounced. No definite rule can be laid down as to the period at which it may be desirable to operate. The surgeon must decide in each case when the proper moment has arrived, and must not hesitate to urge in the strongest terms the importance of every minute. There is far greater danger in delaying the operation too long, than in performing it unnecessarily early. The operation itself, if properly performed, is attended by little or no risk.† But it affords immediate relief from the direst suffering. It obviates altogether the recurrence of those frightful paroxysms of dyspnoea, in any one of which the patient may die suffocated. It affords perfect repose to the suffering parts, and saves them from the irritation of the hard-drawn breath. It thus allows opportunity for the subsidence of inflammatory action, and the absorption of effused material. It may not cure, but if performed in time, it certainly ensures a period of safety and freedom from distress, during which the cure may be effected by natural processes.‡

* *On Diseases of the Throat and Windpipe*, by G. D. Gibb, M.D., 2nd edition, p. 196.

† See an essay by the present author: 'On Some of the Difficulties and Errors of Tracheotomy,' in the *Practitioner*, April 1869.

‡ Out of 49 cases of tracheotomy for acute laryngitis of this kind which have come under my observation, or of which I have collected particulars, life was saved in 30, and death ensued in 19. In many of the fatal cases the operation was obviously performed too late; hence the result.

nevertheless bear in mind that until the dead it can never be too late to attempt even if it should seem that the last breath of the operation should still be completed, and made to restore life by artificial respiration crowned such efforts in cases in which the not been too far exhausted by slow asphyxia

Chronic Catarrhal Laryngitis rarely occurs of one or more attacks of acute or subacute such, however, it is a comparatively common respect it differs remarkably from 'Chronic Granular Disease of the Larynx,' which, preceded by any acute inflammatory symptom contrary is chronic in character throughout its

Chronic catarrhal laryngitis is much more common among adult males than among females is due in great measure, no doubt, to the exposure to vicissitudes of weather to which men are more exposed, as well as to the greater demands made upon the vocal organs from the nature of their occupation. It results in cases in which the acute symptoms are very severe; or in which they have subsided, and the patients resume, without immediate notice, their usual habits before the actual structural changes are sufficiently repaired. Hence the necessity of watchfulness, and judicious treatment during the

ing sound emitted during inspiration in acute laryn-
forced respiration, however, is sometimes accompanied
tral sounds, which may be readily heard by means of a
pe placed over the larynx or trachea. The cough is
'tearing' and painful, nor harsh, ringing, and 'brassy'
acute stages: though often hoarse, and somewhat
it is not distressing, and suffocative paroxysms very
ever occur. Deglutition is easy and painless.

It would appear that there is nothing absolutely dis-
in the symptoms of chronic catarrhal laryngitis; for
lar symptoms are associated with other totally different
is to be hereafter discussed. Nevertheless, it is very
t that a correct diagnosis should be made, inasmuch as
ment requisite, as well as the probable issue, may differ
erially from what would be indicated in at any rate
the other affections alluded to.

laryngoscopical inspection the mucous membrane of the
seen to be thickened and unnaturally red, and covered
l there by patches of greyish opaque mucus. The
ng and redness may be either more or less uniform, or
and much more pronounced in some parts than in
In order of frequency the posterior aspect of the
s, the aryteno-epiglottidean folds, the interarytenoid
anterior aspect of the epiglottis, the false vocal cords,
true vocal cords, are the parts usually affected. In
the acute inflammatory attack is most frequently

or some other effect of the inflammatory process or otherwise interfere with the necessary action. Whatever may be the explanation in any paroscopy has amply proved (contrary to what is supposed) that the voice may be more or less of a different character, or impaired in power, although the vocal cords retain their integrity and healthy appearance is also true. The vocal cords may be visible to some extent in various different ways, as hereafter described, without any very serious detriment to the speaking, if the voice is not affected.

The *course and duration* of chronic catarrh of the larynx varies greatly in different cases. Sometimes it subsides in the course of two or three weeks or more. If neglected, it remains obstinately permanent, and a more or less permanent thickening or hypertrophy of the mucous membrane results. As a general rule, it is amenable to treatment. In some cases of long duration, large growths become developed. In others, tracheal and pulmonary complications arise. In all cases, the prognosis is favorable to fresh attacks of acute or subacute inflammation.

Treatment.—The treatment most likely to succeed in simple catarrhal laryngitis, which has become chronic, depends somewhat upon the stage to which the disease has reached, and the severity of the symptoms. In all cases, however, that any other than local measures are indicated, indeed beneficial, except in so far as they

solutions of alum, tannic acid, or perchloride of iron, or of zinc four or five times a day—or even of common oxide of ammonium, or chlorate of potash—is all that is y. Sometimes, however—and indeed I believe in all xording to some authorities—it is better to resort at the application of a strong solution of nitrate of silver 3j. of water) by means of the brush, or sponge pro-

e more advanced stages, when the affection has become ly chronic in character, and all reasonable hope of eous subsidence, or of improvement under milder treat- as passed away, the efficiency of the application of the of silver solution is most marked. The whole surface interior of the larynx may be wiped round with the or by aid of the laryngoscope the application may be to those parts only which are especially affected. If age is used, as originally recommended by Dr. Horace precision of application is of course impossible; but the lic contraction excited may serve to squeeze from the a considerable quantity of the solution which may diffused over the whole surface. But, as before stated, sh is far preferable to the sponge. The application is ade in accordance with the directions already given.

spasmodic distress, sometimes though not always pro- very speedily subsides. The good effects are almost

tageously applied over the larynx kept up by means of the strong

In cases of very long standing which the tracheal and bronchia affected by chronic catarrhal balsamic and terebinthinate vapour Turpentine, tar, the balsams or their alcoholic solutions, gum benzoin or similar substances, may be mixed with hot water in an appropriate quantity and inhaled from time to time. These substances of this kind should be placed in a wide mouthed bottle and a current of steam should be made.* Another method consists in placing the patient in the room, in which the patient inhales with the remedial vapours, either balsamic substances over a spirit lamp or over hot coals. This method is especially recommended by Pidoux, who states that by interrupted inhalations, chronic laryngitis have been cured to the good effects sometimes produced by the oil of burning paper ('l'huile de papier brûlé') or the inhalation of the smoke of cigars imbued with some arsenical or camphorated oil.

Tar water may be inhaled in weak solutions of chloride of zinc. These saline solutions appear to be of the class now under discussion, and in which the membrane is less extensive, and the inflammation is less intense.

Internal remedies sometimes are indicated in catarrhal affect

* The following formulæ are given:—
1. Balsam of Peru ʒij.; mix. 2. Pyro-acetic spirit ʒij. of Peru ʒij.; mix. 3. Æther ʒss., spirit of balsam of Tolu ʒij.; mix. 4. Æther ʒij. warmth of the hand is sufficient for volatile oils.
'Memoranda,' by J. A. Symonds, M.D., *British Medical Journal*, 1869.

† *Traité de Thérapeutique*, par Trouessart, 1869.

‡ Ibid. Vol. i. p. 172.

membranes generally, as, for example, decoction of senega with ammonia, the various balsams, chloride of ammonium with tonics, &c.


Men who are subject to catarrhal and other chronic affections of the larynx should wear their beards; and women should be advised to take due precaution for the protection of their necks, especially when they are exposed to cold, or vicissitudes of temperature.

CHRONIC GLANDULAR LARYNGITIS. FOLLICULAR DISEASE OF THE LARYNX. DYSPHONIA CLERICORUM.

Chronic Glandular Laryngitis is a comparatively common affection.* It is most frequently, but by no means solely met with among those who are subject from time to time to continuous exertion of the voice, as clergymen, barristers, singers, and others. It essentially consists in more or less general enlargement and hypertrophy of the glandules and follicles of the laryngeal mucous membrane, the result of an inflammatory condition which usually commences very insidiously, and always progresses very slowly. It is often, but not invariably associated with a similar affection of the faucial, nasopharyngeal, and pharyngeal mucous membranes.

The most frequent cause probably is, as already indicated, continuous exertion of the voice. It is obvious that in prolonged speaking, reading, or singing, the demands made upon the vocal organs are not only greater than during ordinary conversation, but also in some degree different. In ordinary conversation the parts are subject to no great strain; and more or less frequent intervals of rest are afforded, during which the mucous membrane can recover its normal condition. But in the continuous and strong exertion of the voice constantly made by public speakers and singers, the mucous membrane is especially liable to become irritated by the forcible contact, and rapid passage over it, of cold dry air drawn in at each inspiration through the mouth, and not warmed and moistened by passing through

* This affection appears to have been first accurately and fully described by Horace Green of New York, to whom also is due the credit of having espoused and insisted upon the proper method of treating this and other lesions of the larynx by the topical application of solutions of nitrate of silver. See *A Treatise on Diseases of the Air Passages*, by Horace Green, M.D. New York, 1846.



the laryngeal mucous membrane. Now this subject to a very great extent of motion, and considerable strain, during forced vocalisation its glands are especially liable to be stimulated; and the morbid condition undereventuate.

It is important to bear in mind, however, glandular laryngitis may, and indeed often does, occur in persons who have not been subject to any such exertions as those above referred to. In such cases cold and fog (especially if there is any obstruction to the nasal passages) and the constant inspiration of air charged with dust, fumes or particles, appear to be the most frequent causes. It is stated that he has seen this malady 'in a variety of persons, in photographers, and in persons much exposed to acrid chemicals in confined chambers, and it is something quite remarkable.'* It is possible that a constitutional tendency may favour the development of the malady. Indeed, Trousseau, Chomel, De laennec, and others attribute its origin to the 'herpetic diathesis.'

The *symptoms* are :—alteration of the voice, hoarseness, in sustaining it—these are by far the most prominent symptoms; more or less discomfort in the throat, never amounting to pain but occasionally to burning, and sometimes a sense of heat about the throat, and a constant desire to clear the throat by 'hemming.' There is little or no regular cough; and the ex-

gether, and the patient speaks in a husky whisper. As rule the alteration in the voice is most pronounced in morning, before the vocal organs have been at all and again in the evening, or after any considerable of the voice, when the parts have become fatigued. As the patient will feel that he is able to carry on conversation with ease ; but directly he begins to read or continuously in a loud voice he finds it impossible to proceed only does so with great difficulty, sense of effort, and great fatigue.

section, the mucous membrane of the fauces and back of the throat is usually seen to present a more or less raw and granulated appearance ; and the enlarged glandules, may be) surrounded by a little halo of undue redness, recognised studding the surface. *Laryngoscopical examination* shows that the mucous membrane of the larynx is affected ; and in some cases, as already indicated, the inflammation is confined to the larynx. The parts most frequently obviously affected are, as might be expected, those in which the glandules are most abundant. These parts are the mucous membrane covering the arytenoid cartilages, the inter-folds, the base of the epiglottis, and the ventricles and saccus of the larynx. In cases of long standing, points of inflammation may sometimes be observed, especially about the base of the epiglottis. In others the glandules appear to be not enlarged, but distended with opaque yellowish material



known to be the most efficacious, consists in the brush of solutions of nitrate of silver or perch at intervals varied in frequency according to . The nitrate of silver, I believe, is, as a rule, the best. Sulphate of zinc, sulphate of copper, and tincture have also been recommended, and indeed it may be that it is often well to vary the application. That which is perfectly in one case, or at one period, proves to be in others, or at other periods and under changing conditions in the course of the same case. Any points of irritation may be seen should be touched by means of the solid nitrate of silver. Between such applications should be inhaled at frequent intervals during the day. Solutions for the purpose are those of common salt, of ammonium, iodide of potassium, and in some cases weak solutions of perchloride of iron. Certain solutions, especially such as contain the sulphides of sodium, have also been strongly recommended, as well as solutions of sulphurous acid.

A method of treatment, which appears to be effective in this form of laryngeal disease, consists in the use of medicated lozenges. The precise and careful use of Fournié † prove (contrary perhaps to what might be expected) that certain portions of liquids swallowed, especially if swallowed slowly, or allowed almost insensibly to find their way to the pharynx, may become applied to the upper part of the larynx and even diffuse themselves to some extent over the surface. The lozenges that appear to prove most


ly to reassure the mind of the patient, for he is but too oppressed in spirits, and apt to think his malady far more than it really is.

PHTHISICAL LARYNGITIS.

Larynx is found to be more or less seriously affected in a great proportion of cases of tubercular phthisis.* In some of the earliest recognisable indications of the commencement of the malady appear in the larynx. In others, when the disease is fully established, the laryngeal symptoms are the only or not the only source of the distress from which the patient

In others again, the condition of the larynx gives rise to symptoms which may to a considerable extent mask or simulate other physical signs and symptoms of incipient, and even of advanced pulmonary disease; and thus the diagnosis may be rendered very difficult.† Hence the importance of making laryngoscopical examinations in all cases of suspected phthisis, and especially in such as present indications of advanced mischief.‡ There can be no doubt, I think, that most of the so called ‘cures of consumption’ said to have been effected by local applications and inhalations (in so far as the results given may be considered reliable) have occurred in

* states that he found ulceration of the larynx in 63 out of 193 cases of phthisis which he examined. *Researches on Phthisis*. By P. C. A. Louis, M.D. *Soc. Trans.* Lond. 1844. According to the records of postmortem



...dotted granular appearance, which is usually first most conspicuously manifest on the posterior aspect of the glottis. This appearance is probably associated with the deposit of tubercular material taking place in the mucous membrane at many distinct points. Somewhat later, the mucous membrane covering the arytenoid cartilages, and the arytenoid and interarytenoidean folds are affected, and the mucous membrane presents granulation-like elevations. At the same time the thickening of the mucous membrane may be general. This thickening is an important and almost characteristic feature of the disease. It is to some extent due to increased deposit of tubercular material as well as to inflammatory infiltration of the true vocal cords, as a rule, become implicated in the disease at an early period, and usually on one side first. The epiglottis also partake of the general thickening, and often becomes inflamed; the cords; and these latter lose their clearness and elasticity. As the malady advances ulceration occurs in one or both of the cords, usually first on the posterior aspect and lower part of the glottis. At or about this period the voice becomes hoarse, and breathing is carried on with a sense of effort and difficulty and pain; the cough (before this time often little more than a more or less frequent dry cough) becomes constant, and sometimes painful; the sputum is increased in quantity, and somewhat changed in character, becoming yellowish, and occasionally slightly bloody; and sometimes, according to the position of the ulcer, it may contain pus or blood.

[illegible]



described. Postmortem examination not infrequently shows a more or less pronounced affection of the larynx ; and during life there have been scarcely any laryngeal symptoms beyond slight hoarseness, or some other alteration of voice.

The *course and duration* of phthisical laryngitis is determined not only with the special character and type of the disease, but also still more notably with the degree and severity attained by the pulmonary disease, and the morbid condition at the period at which the larynx begins to suffer. Thus, in the first place, as already stated, the disease may begin in the larynx, and there may be no other evidence whatever of the presence of tubercle in any other part of the system. In the second place, the larynx may appear to become affected simultaneously with the lungs, and the disease may go on almost *pari passu*. In the third place, the affection of the larynx may be consecutive to advanced pulmonary disease, and the organisation of the lungs, and may even supervene almost at the very termination of the case.

In cases belonging to the first class indicated, the disease may be comparatively rapid, and the patient may die soon after the onset of the laryngeal disease ; or the affection may come on gradually. More frequently the progress of the malady is slow ; and sometimes even actual recovery of temporary character is effected. But sooner or later the malady appears afresh, or

Treatment.—It would be altogether out of place to discuss in these pages the constitutional treatment requisite in tubercular affections generally, and in pulmonary phthisis especially. It is only to say, that similar constitutional treatment is indicated in phthisical laryngitis. It must ever be borne in mind that this malady is but a local expression, so to speak, of a general morbid tendency, to modify which all possible endeavours must be made. At the same time, the part affected is important, and the associated danger and suffering may be great, that no measures must be neglected by means of which the symptoms may be alleviated and the danger averted, if only for a time. Experience amply shows that in such respect local treatment is often very beneficial.

Even in the earliest stages it does not appear that any topical applications are likely to be efficacious in arresting the malady; rest to the parts affected, and avoidance of all sources of irritation, may afford opportunity for general improvement under constitutional treatment. But when once ulceration has commenced the need for local treatment arises, and indeed may become most urgent. Soothing inhalations of the steam from water in which hemlock, hops, stramonium or opium have been infused, often afford much comfort. So also do spray inhalations of anodyne solutions. Similar solutions applied by brush are sometimes very useful. They may not cure, but they give the temporary relief the sufferer so often and so justly looks for. An excellent formula quoted from Krischer * is as follows: Extract of opium and extract of belladonna equal parts, dissolved in forty parts of cherry-laurel water. The beneficial effects of this application, although transient, are immediate and almost invariable.

Small ulcers may often be advantageously touched with the solid nitrate of silver: a protective covering from the irritating influence of the passing breath is thus formed for them. But the application of the solution of nitrate of silver to the mucous membrane generally, appears in many cases to do harm rather than good. Marcet recommends repeated scarification of the thickened parts of the mucous membrane.†

In cases in which the dyspnoea is great, and clearly dependent upon the condition of the larynx—in cases especially in which there is great difficulty in swallowing, and consequent

* Op. cit. p. 674.

† Op. cit. p. 95.

complete relief afforded by the operation in some cases, the hopeless character of which there could be no doubt, should never hesitate to repeat it in similar cases. That though life may not be saved, it may be prolonged, and suffering may assuredly be diminished during the operation. Further, it may be added, that cases do occur at a late time in which the whole disease is in the last stage, the accompanying symptoms and wasting closely resembling general phthisis, and the physical signs are so feeble as to afford no reliable indications. Three cases have been under observation, in each of which the patient had been considered to be dying of 'hopeless consumption.' In each case, more or less complete recovery speedily followed the tracheotomy.

SYPHILITIC LARYNGITIS.

Syphilitic affections of the larynx are comparatively rare. They vary in character and importance with the stage of the disease at which they arise, and also with the general health and condition of the patient.

During the *Secondary Stages* of Syphilis the mucous membrane of the larynx may be affected in a manner corresponding to the cutaneous eruption present; or it may be affected either independently, or in direct continuity with

Without such aid the precise condition of the larynx in this case can only be inferred from the symptoms; a considerable degree of uncertainty necessarily attends the diagnosis.

Inflammation of the laryngeal mucous membrane often occurs in connection with syphilitic roseola of the skin. It may either originate from the fauces to the epiglottis and upper part of the trachea and thence spread over the whole mucous surface; or it may arise in isolated, and more or less well defined patches on the laryngeal mucous membrane in the parts affected presents on superficial inspection a dusky red or even purplish hue, and may be, a slightly elevated or swollen appearance. Distinctly elevated and better defined patches (*mucous patches*) are sometimes seen in association with papular, squamous and especially with tubercular affections of the skin.

Symptoms accompanying these comparatively simple conditions are not, as a rule, severe; nor are the results likely to be serious. The voice is generally altered to some extent: it becomes weak, wanting in tone, husky, or somewhat hoarse.


But there is neither dyspnoea nor troublesome cough; there is no local pain, nor any difficulty in swallowing that can be referred to the affection of the larynx.

The peculiar character of the voice so constantly noticed in secondary syphilis, and by some considered almost pathognomonic, would appear to depend in great measure upon one or more of the conditions of the larynx thus described, rather than upon the concomitant affection of the palate, fauces, and laryngeal mucous membrane, the existence of which is readily obvious, and to which alone the explanation is usually attributed. Such, at any rate, is the conclusion inferred

by the results of my own observations, as well as by those of others. It must be borne in mind that an affection of the laryngeal mucous membrane of the larynx which is only superficial, and which the eye of the observer appears slight, may nevertheless, directly or indirectly, hamper to a considerable extent the movements necessary for the production of the natural voice.

On this subject, see especially the brochure of M. Dance, *Sur les éruptions cutanées dans la période secondaire de la Syphilis*. Paris, 1864.

Hardt and Roth state that in eight out of fifty-four cases of secondary syphilis under observation in Würzburg Hospital, the hoarseness was produced by mucous patches or condylomata of the larynx, which could be distinctly seen on inspection with the laryngoscope. *Arch. für. path. Anat.* Bd. xxi. Heft 1.



during the secondary stages, except in cases where the general health and strength are much broken down. There has been continuous or frequent exposure to the ordinary causes of the disease.

Occasionally small isolated spots of syphilis are similar to those often observed in the mucous membrane of the laryngeal mucous membrane.

During the *Tertiary Stages* of syphilis the larynx is affected in one way or other, and in many cases to a serious extent.

Papulo-tubercular Elevations of the Mucous Membrane are means rare. They vary in size, and also in shape. In some instances they simulate warty growths of moist surfaces, in others they more or less closely resemble the tubercles met with on other mucous surfaces.* Sometimes they subside under appropriate treatment, but in many cases they increase in size and necessitate removal by operation. Sometimes they form the starting points of ulcerations, as described in the next paragraph. They have been observed on the true vocal cords, as well as on various parts of the laryngeal mucous membrane. The chief symptoms are hoarseness, or some other marked alteration in the voice, and sometimes occasional fits of dyspnoea.

Tertiary Ulcerations of the larynx are comparatively rare. They may commence either superficially in

formation of abscess. They may occur singly, and in any part; usually, however, several spots are attacked, either simultaneously or one after another. The epiglottis is generally affected, and in a large proportion of cases is the part most extensively destroyed. Sometimes its edges are gradually eaten away; sometimes its substance early becomes perforated, and the perforation extends until the edges are reached. In such cases semidetached portions, flapping loosely, occasionally give rise to much embarrassment. The false and true vocal cords are often affected—in some cases by comparatively superficial, but most by deeply penetrating ulceration. The mucous membrane covering the arytenoid cartilages, and the posterior part of the larynx generally, is comparatively less frequently and less early attacked by syphilitic than by phthisical ulceration; but it is liable to become involved sooner or later. The cartilages (and especially the arytenoids) may become carious or necrotic, as the result either of deep extension of ulceration commencing in the mucous membrane, or of perichondritis arising independently of any such superficial affection. Portions of dead cartilage may become detached, and acting as foreign bodies, may give rise to the most serious symptoms; or they may be expectorated, or removed by operation; or they may remain at any rate partially fixed, and in such case may become the foci of abscesses, and the excitants from time to time of the most dangerous paroxysms of spasmodic dyspnoea.

The *symptoms* vary according to the extent of the ulceration and the part especially affected. There is invariably marked alteration of the voice, sometimes hoarseness, sometimes complete aphonia, and sometimes one or other of the various intermediate conditions which are readily recognisable but difficult to describe or name. There may or may not be dyspnoea; and the dyspnoea, if present, may be slight, or of the most distressing character. In some instances, and especially in those in which the cartilages are affected, there may be occasional attacks of severe dyspnoea, with intervening periods of comparative or even complete ease; and sometimes suffocative spasm of the larynx may occur.

Two or three years ago, a patient (in the Venereal Ward in Guy's Hospital) who had suffered from occasional and slight but evanescent attacks of dyspnoea, was suddenly seized with such urgent and distressing difficulty of breathing, that he rushed from the ward to seek help. He reached the bottom of the stairs, and there fell dead from suffocation. On postmortem examination, a detached portion of necrosed cartilage was found in the glottis.

more liable than the latter to 'go down the windpipe' to the slight degree of difficulty, and the absence of dysphagia, sometimes observed in certain cases of this kind, in which the epiglottis has been entirely destroyed, are very remarkable.

The *course and duration* of these tertiary affections of the larynx, as may be readily understood, vary greatly in cases, under appropriate treatment cicatrization occurs; but in such it constantly happens that very serious complications result either from the loss of substance or from the subsequent contraction of the cicatrices. The voice is, as a rule, permanently lost. Breathing and swallowing may be rendered more difficult. Attacks of acute laryngitis, attended by more or less of dysphagia, are liable to be excited by comparatively slight causes, and may lead to a fatal result if timely relief is not afforded, or if the patient has not been previously secured by the performance of tracheotomy.

The *general diagnosis* of syphilitic affections of the larynx is not often difficult. The history of the case, and the syphilitic cachexia, together with the coexistence of more unmistakeable signs or symptoms (such as the presence of cicatrices about the palate and fauces, cutaneous eruptions, nodes on the tibiae, &c.) generally serve to indicate the nature of the malady.

But beyond the general indications afforded by the history of the case, and by the presence or absence of other

eriment of 'specific' treatment before expressing any opinion. Antisyphilitic remedies and local treatment certainly effect a cure in the one case; and in the other, though necessarily useless, they cannot do any great harm.

Deep and extensive ulceration of the more advanced syphilitic disease of the larynx not only gives rise to dysphagia, but on laryngoscopical inspection may present appearances, which more or less closely resemble those of cancerous disease on the one hand, and epithelioma on the other. It may be easy, but it is always important to determine the nature of the malady, as well as the extent of the disease. For, as need scarcely be stated, the treatment requisite and likely to prove more or less successful in syphilitic origin, might be absolutely injurious in those cases with phthisis, and altogether useless, or perhaps even useless, in epitheliomatous disease. The prognosis obviously depends in great measure upon the determination of the origin of the local affection. The distinctive features presented by syphilitic, phthisical, and epitheliomatous ulceration of the larynx may be stated as follows:

Specific ulceration usually attacks the epiglottis first. It progresses rapidly, and is emphatically destructive in its progress. It involves the submucous tissues at a comparatively early

sent an irregular, somewhat worm-eaten and The accompanying expectoration is general thinner, and more mucopurulent in character ulceration, and much more abundant than i

Epitheliomatous ulceration of the larynx portion of cases commences on the phary mucous membrane covering the arytenoid o and comparatively rarely in the interior of associated with very considerable and irr due to the development and increase of Its surface presents a dirty greyish appear are elevated. It progresses very slowly. The pectoration, at any rate during the earlier sta and often more or less sanious in character. stages it becomes abundant, more or less p times tinged with blood.

The other local and constitutional signs ar in syphilitic and phthisical cases may aid altogether wanting in cases in which the lar simply epitheliomatous in origin.

Lupous, and other Strumous ulcerations of cannot, strictly speaking, be classed as ph occur,* and are with difficulty distinguish syphilitic origin. In some cases of the kin under my observation, it has seemed that probably been due to congenital *scabie

any exposure to cold and wet, should be sedulously avoided. Mercury in one form or other is the only remedy which any reliance can be placed; and the appearance of laryngeal symptoms may generally be accepted as an indication that it is desirable to get the system under the influence of mercury as speedily as may be practicable and safe. The use of calomel vapour bath may be especially recommended in these cases, inasmuch as beyond its general efficiency it affords a special facility for the occasional inhalation, and thereby the application of the mercurial vapour. The value of such inhalation, however, is more pronounced in cases in which inflammation has commenced.

Secondary ulcerations of the laryngeal mucous membrane (whether in continuity with others about the fauces or pharynx, or appearing independently) may often be advantageously treated by local applications. But such applications must be regarded simply as adjuvants to the general treatment indicated, and not as in any measure rendering such general treatment necessary. The frequent inhalation of the spray of weak solutions of perchloride of mercury with chloride of ammonium, sulphate of copper, very often appears to be beneficial. In some cases the insufflation of calomel, or an occasional inhalation of the vapour of calomel, may be recommended. In other cases the application of solid nitrate of silver or sulphate of zinc, from time to time, to the ulcerated surface does good. Where there is much irritability, however, the inhalation of warm



advantageously applied by the brush. Cases in which the internal administration and plication of mercurials in conjunction with iodine are indicated. The more powerful medicines may sometimes be beneficially alternated with the exhibition of chlorate of potash in the form of solution, spray, or powder, and combination with tonics. Swollen and oedematous mucous membrane may be freely scarified to advantage; and any spot at which an abscess has formed may be punctured. If any portion of necrosed cartilage should be seen, an attempt should be made to remove it by means of appropriate forceps.

In every case in which breathing becomes difficult, whether in the secondary or tertiary stage, tracheotomy should be performed without delay. Experience proves that in syphilitic disease of the larynx, operation may be regarded as fairly successful in averting impending danger, but also in affording a means of recovery for the more or less complete restoration of the organ.* The operation is especially called for in cases of paroxysms of spasmodic dyspnoea have occurred. Laryngoscopic examination shows that the ulcer is deep, and some portion or other of the cartilage is exposed. In such cases recovery is almost always possible.

as that tracheotomy becomes necessary after partial or complete cicatrization has taken place, in consequence either of the resulting contraction of the parts, or of an attack of laryngitis incidentally occurring. The damaged parts are liable to a low subacute or chronic form of inflammation, which at any moment may become exacerbated and give rise to urgent symptoms. In some cases the passage gradually becomes more free in consequence of the absorption of effused material; but such, if it should have been necessary to perform tracheotomy, the canula may be removed at an earlier or later period. In other cases the contraction increases, and the tracheotomy must be worn during the remainder of life. In some cases it may seem desirable to divide the cicatrices by incision, and to attempt the dilatation of the air passage. As far as I have been able to ascertain, no very satisfactory result has hitherto been obtained in any case in which the effusion has been in the larynx itself. Much good however has been effected in several cases in which the contractions have been situated above the larynx. Some encouragement to such attempts may perhaps be derived from the successful results which have been obtained in instances in which the contractions have resulted from the effects of injuries, although, it is obvious, the conditions in such cases must necessarily be somewhat different.*

These local symptoms are preceded and attended by constitutional symptoms (such as feverishness, rigor with erysipelas generally. On inspection, the tongue of the fauces is seen to be unnaturally red, and sometimes more or less dry and shining. By laryngoscopy, the mucous membrane of the larynx is found similarly affected. As the malady advances the mucous tissue becomes infiltrated; the swelling increases, and the character is pronounced; the glottis becomes more encroached upon, and respiration is checked and more impeded.

Unless the malady is checked in its progress by treatment afforded from the urgent symptoms, the patient passes into a 'typhoid' state, and dies either from exhaustion or from exsiccation. Such is the common result.

On post mortem examination the mucous membrane of the larynx appears inflamed, swollen, and oedematous. In some instances it is of a dirty greenish colour in places. The mucous tissue is infiltrated with seropurulent exudate, and there is sloughy. Small collections of pus in the areolar tissue are occasionally seen round the trachea. The tracheal mucous membrane is also often found inflamed. The mucous membrane of the fauces is similarly affected. The base of the tongue and the tonsils sometimes show a more or less extensive sloughy appearance.

Treatment.—The general treatment must be of a stimulating and supporting kind; and no effort must be made to suppress the local symptoms.

administration of an emetic followed by a purge. But the propriety of so doing must be decided by the general condition of the patient, and the progress the malady may have

Locally, the application of a strong solution of nitrate or perchloride of iron may be recommended in the early stages. In the more advanced stages, especially when there is much cedema, free scarification of the mucous membrane, and frequent inhalations of warm soothing vapours and steam afford the most marked relief. If the dyspnoea becomes extreme, tracheotomy must be performed. It is true that from the nature of the malady there is often but little hope of a successful issue; at the same time there can be no doubt but that even in the most advanced stage a chance of prolonging life is afforded that could otherwise be obtained; and that by it at any rate the last moments of the patient may be rendered comparatively free from suffering.

It may be worthy of remark that apart altogether from any other disease or affection as that thus described, it not infrequently occurs in connection with that in cases of erysipelas of the scalp and face, there is often soreness of the throat, slight difficulty of breathing, and hoarseness of the voice, such as weakness or hoarseness.

DIFFUSE CELLULAR LARYNGITIS. ACUTE CEDEMATOUS LARYNGITIS.

The inflammation of the areolar tissue of the larynx is a comparatively rare, but very formidable malady. It corresponds to diffuse cellular inflammation not infrequently met with in other parts of the body; but it is especially dangerous on account of the impediment to respiration and swallowing to which from its situation it necessarily gives rise. It is closely allied in nature to erysipelatous inflammation, from which, however, it differs in the fact that in it the submucous membrane is primarily attacked, and the mucous surface is left unaffected, or only becomes secondarily affected. In erysipelas, on the other hand, the mucous surface is first affected, and the submucous tissue is only secondarily or concomitantly involved.

Diffuse Cellular Laryngitis may either occur in association

The result ensued in 10 out of 15 cases of tracheotomy performed on cases of erysipelatous laryngitis which have come under my observation, or I have collected particulars; in 5 cases the patients made good

with previously existing diffuse inflammation of the neck, in which case it seems to be a general affection; or it may come from the mucous tissue of the pharynx and pass through the neck generally. It may cause death from suffocation or from considerable extension to other parts.

In this malady the appearance of the disease is preceded, and its onward progress is attended by more or less severe constitutional depression, and great general depression. At the commencement the patient complains of having felt unwell for some time, of headache, lassitude, lowness of spirits, &c. Soreness of the throat, which becomes severe, is noticed early; and as the malady progresses breathing becomes more difficult, weight and oppression at the chest, and may be slight hacking cough, attended with expectoration of a little whitish glairy mucus. The dyspnoea increases; the fauces and tonsils become inflamed, and a dusky red colour; and sometimes a white exudation upon them. Somewhat later the swelling of the jaw becomes enlarged; the neck becomes so much enlarged that it can only be opened with difficulty. When external pressure is made on one side, a constant and profuse discharge of pus issues from the neck becomes swollen, and the system is so depressed that respiration is more and more difficult, and is rendered almost or quite impossible. In a typhoid condition, and dies exhausted. Frequently happens, he is killed by the disease. In some cases, however, a moderate dyspnoea proves fatal at an early period in the progress of the malady; and the œdema increases very rapidly, and the patient dies. Either of such results may ensue, and without any perceptible general sweating.

So far as I have been able to ascertain, the post-mortem examination has hitherto been in the case of this kind; and therefore the details of the life cannot be described. They may

ed and swollen, and dusky or paleish in colour. The of the mucous membrane differs altogether from the scarlet redness of acute catarrhal inflammation. The parts of the larynx are more or less completely concealed
sw.

post mortem examination extensive inflammatory œdema submucous tissue of the larynx and neighbouring parts rially found. In some cases one side is much more l than the other. When cut into, the areolar tissues ; an infiltrated, and sometimes more or less sloughy ap-æ. The infiltration is serous, seropurulent, or purulent acter, according to the period at which death has occurred, e circumstances under which it has taken place. The of the neck are found in many cases to be infiltrated to er or less extent with serous, seropurulent, or fibrinous ion. In cases in which there has been much swelling, they present almost a brawny consistency. Sometimes the tion extends upwards behind the pharynx, sometimes ards alongside the trachea and œsophagus, even into the tina. Occasionally the infiltration in the neck is puru- character, and in some instances the areolar tissue is r and putrid.

tment.—The same general treatment is required as that ended in erysipelatous laryngitis. But in this malady pplications of nitrate of silver, &c. are far less likely to

view of diminishing the tension and consequent pressure upon the trachea and œsophagus.

If such means fail to afford relief, and the dyspnoea is urgent, tracheotomy must be performed, although the difficulties attending the operation may be great, and the chances of a successful result may appear small.

An excellent illustration of the value of tracheotomy in cases of this kind occurred some few years ago in the practice of Mr. Pollock at St. George's Hospital.

The patient (a butcher, aged forty-three) was admitted with severe and extensive diffuse cellular inflammation of the neck. Urgent laryngeal symptoms soon supervened. Tracheotomy was performed, and the patient made a good recovery in a comparatively short space of time.*

OTHER FORMS OF LARYNGITIS. EXANTHEMATOUS LARYNGITIS; TRAUMATIC LARYNGITIS, &c.

Consecutive inflammation of the larynx occasionally occurs in the course of various general maladies, other than those the laryngeal complications of which have been already described in detail. It is especially liable to occur in measles and typhoid fever. It is more rarely met with in smallpox and typhus, and still more rarely in scarlet fever.

The *Laryngitis of Measles* is catarrhal in character, and, in most instances, appears soon after the nasal catarrh. Respiration may be more or less embarrassed; and sometimes there is remarkable harshness of the voice, and frequent harsh cough. In young children the breathing is occasionally stridulous, and the cough ringing. The consequences are very seldom serious; but it is well to bear in mind that they may possibly become so. In some rare instances, dangerous, and even fatal œdema of the larynx has ensued; and in some, the œdema has persisted after the general symptoms have subsided. In such cases the laryngeal symptoms have appeared early, and from the first have been very acute. As a general rule, the mucous membrane of the larynx is simply affected in a manner corresponding to the cutaneous eruption; and the symptoms subside as the eruption disappears. It is not often that any local treatment is necessary or desirable, beyond the application of warm poultices

* See *Lancet*, September 1863, p. 276.

and fomentations to the neck, and the inhalation of warm soothing vapours. If, however, the symptoms are severe and persistent, the same treatment may be required as that recommended in cases of acute catarrhal laryngitis (see p. 539).

The *Laryngitis of Typhoid Fever* is ulcerative and destructive in character.* It may either commence during the earlier or middle stages of the malady, concomitantly with the ulceration of the intestinal glands and mucous membrane, and speedily give rise to serious symptoms, or even to a fatal result; or its existence may first become manifest during the period of convalescence and after all febrile symptoms have subsided. The ulcerations which occur during the earlier stages are generally situated either on the aryteno-epiglottidean folds, in the neighbourhood of the arytenoid cartilages, or, according to Rokitansky, about the lateral walls of the larynx. They may give rise to more or less severe functional disturbance—often to cough, and difficulty of swallowing, and, somewhat less frequently, to difficulty of breathing. The difficulty of breathing usually corresponds to the amount of inflammatory œdema that may have occurred. It is sometimes difficult to determine how far the cough may depend upon laryngeal, and how far upon bronchial affection. It is probable that in a large proportion of cases, if the patient survive the general malady, these ulcerations heal favourably, like the corresponding intestinal ulcerations. But in cases in which the symptoms become manifest at a comparatively late period in the course of the malady, or in which ulceration goes on insidiously, and perhaps almost unnoticed until seeming convalescence has taken place, the perichondrium and cartilages (and especially the cricoid) are very liable to be implicated to a greater or less extent. Under such circumstances the consequences are often very serious. Necrosis

* This affection is the *laryngo-typhus* of the Germans. It appears to have been first definitely and fully described by Louis, in whose treatise, entitled *Recherches sur la Fièvre typhoïde* (Paris, 1841), much valuable information on the subject is contained. Trousseau especially refers to the laryngeal complications of typhoid fever, and gives some excellent illustrative cases in his *Clinique médicale*, Vol. i. p. 299; Paris, 1868. Chomel, Rokitansky, Wilks (*Med. Times Gaz.*, 1862, p. 276), and others also fully discuss the pathological history, and examples of typhoid disease of the larynx. The appearance of the ulcers is characteristic that Louis says that, if observed in a patient who had died of some acute disease, they would be sufficient to indicate almost certainly that the affection had been typhoid fever. *Op. cit.* p. 321.

of the cartilages may result in association with typhoid fever only from gradual penetration of the ulcerative process also in some cases as an effect of some deep-seated inflammatory condition commencing in or about the cartilages themselves.

A remarkable instance came under my observation in Guy's Hospital. I was acting as Ward Clerk to the late Dr. Addison. A boy, ten years of age, had apparently recovered from typhoid fever. One morning he complained of pain in the throat. A day or two afterwards he expectorated some purulent matter. The next day his neck was observed to be swollen and emphysematous. The emphysema rapidly spread over the whole body, even to the scrotum and testes, which became enormously and peculiarly distended. Two days later he died. On post mortem examination a small ulcerated opening leading to a necrosed portion of the cricoid cartilage, was found below the glottis on the right side. Through this opening evidently the air had escaped during the typhoid fever, and had thence spread over the body. This is the only other case of any kind has ever come under my observation in which emphysema was so extensive and so general. The cavities as well as the superficial parts of the body were affected; and indeed it appeared that death had resulted from the impediment to respiration caused by the thoracic emphysema.

In a certain proportion of cases in which the symptoms are not severe, recovery may take place without any local treatment beyond the application of warm fomentations to the neck and the inhalation of warm soothing vapours. The general treatment of typhoid fever must, of course, be carried out. In some cases, however, in which the symptoms are urgent, tracheotomy should be performed. It is undeniable that the successful results hitherto obtained, do not appear to be very encouraging. At the same time the success which has resulted in some cases justifies resort to this operation, which indeed alone offers a chance of safety.

In the *Laryngitis of Smallpox*, the mucous membrane of the larynx is affected in a manner which varies with the stage of the malady, and corresponds more or less closely with the obvious external appearances. In the earlier stages, pustules may arise analogous to those observed on the cutaneous surface. Neumann, Türk, and Krishaber, all state that they have seen, by aid of the laryngoscope, true variolous pustules.

* I find that out of thirty-five recorded cases in which tracheotomy was performed on account of 'typhoid' disease of the larynx, in twenty-four cases recovery ensued. In eleven life was saved; but in several of them the voice was permanently impaired, and the canula could never be dispensed with.

surrounded by circumscribed inflammatory areolæ, upon the true vocal cords, as well as on other parts of the laryngeal mucous membrane. The symptoms are, hoarseness, cough, and more or less uncomfortableness or pain about the larynx. About the eleventh or twelfth day, at the period at which swelling of the face and extremities is especially liable to occur, œdema of the larynx may come on; and respiration may be more or less seriously, or even fatally impeded. During the still more advanced stages of the malady, at or about the period at which abscesses are occasionally formed in other parts, perichondritis and subsequent necrosis of cartilage, with all the attendant symptoms and dangers, may ensue.

In *Scarlet Fever* the larynx may either be affected during the earlier stages by extension of inflammation or inflammatory œdema from the pharynx; or, in common with other parts, it may become œdematous as the result of consecutive kidney disease. The true scarlet fever eruption very rarely indeed, if ever, invades the larynx. The local treatment necessary in each case may be gathered from what has been already stated in regard to similar affections arising in connection with other maladies.

It may here be remarked that in *Bright's Disease* the larynx may participate in the general œdema and this may prove dangerous or fatal from the obstruction to respiration to which it gives rise.

Gouty Laryngitis may possibly be considered to possess small claim to be described as a distinct form of disease. But it is worthy of remark that in gouty subjects chronic laryngeal affections are by no means uncommon. Sometimes they are very troublesome, and even distressing to the patient; and they are always difficult to treat successfully unless the existence of the general condition upon which they depend is fully recognised, and modified by appropriate constitutional remedies. In many cases they occur in association with gouty bronchitis and tracheitis, but even in some of these the laryngeal symptoms may attract most attention, and give rise to most trouble and danger. In certain rare instances acute laryngitis of very formidable character has been observed to occur in connection with an attack of gout.

remarkable instances, their presence could be detected only on post mortem examination. Since the laryngoscope came into use, however, a large number of cases have been brought to light, and many more at present unpublished have been discovered, in which growths in the larynx have been recognised during life, but have been removed by operation. There is probably no other class of diseases in which the value of the laryngoscope has been so significant. It would appear not unlikely that this circumstance, with the interest naturally attaching to cases of disease of the larynx, has attracted to this means of leading to the now prevalent idea that such growths are more common than is actually the fact. They are more frequent than was formerly supposed; but they do not constitute only a very small proportion of the diseases of the larynx that come under every day observation. It is true that such growths are to be 'met with in two-thirds of the cases of maladies of the larynx except the chronic.* My own experience would lead me to believe that they are much less frequent even than this.

The new growths in the larynx met with vary in character, size, and precise situation, and in the responding extent, in the symptoms to which they give rise, in the degree of danger they immediately in-
cur, in the degree of danger they immediately incur at any period of life; in some instances they are congenital; in others they do not appear until late in life. They are almost equally common in the two sexes.

aining causes of their appearance in the majority of cases. In some instances, however, they may be of syphilitic origin; in the so called new growths may be simply the thickened folds of surface, or the 'vegetation-like' granulations of chronic laryngitis.

To give an approximative idea as to the relative frequency of the different kinds of new growth met with in the larynx may be derived from the following table, which has been carefully compiled from the published records of cases, from the results of my own observations, and from the descriptive catalogues of anatomical museums.


Nature of growth.	Number of cases.	Number in which the growths were recognised during life.	Number in which they were only found after death.
Papillomatous or warty	110	65	45
Fibrous	23	13	10
Cellular. Sarcomatous, &c. . .	52	47	5
Glandular. Adenomatous . . .	6	2	4
Carcinomatous	14	9	5
Mucilaginous	4	—	4
Papillomatous. Cancerous . .	19	8	11
Clearly indicated	16	14	2
Total	244	158	86

Papillomatous or Warty Growths, as will be seen by the above table, constitute a large proportion of the new growths met with in the larynx. In structure and appearance they bear a general resemblance to growths of this nature found in other parts; they may appear either as little warty elevations, consisting of comparatively few, and sometimes almost isolated papillæ, or as closely packed and numerous villous or filipendulous outgrowths, or as densely massed 'cauliflower-like' aggregations of elongated and enlarged papillæ, rounded or flattened at their extremities. They most frequently, perhaps, spring from the mucous membrane of the anterior wall of the larynx, somewhere near the insertion of the true vocal cords. In many instances, they take origin from the boundaries of the ventricles, and in some from the true vocal cords; but they may arise from almost any part of the laryngeal mucous membrane. As a rule, they

FIG. 258.



Papilloma of larynx springing from border of left ventricle. Larynx opened in front.



compared with those met with in many of
was sufficiently large, however, to lead to
some cases these growths increase very rapid
extent as to fill altogether, and sometimes c
cavity of the larynx.

A very remarkable instance is recorded and figured b
case of the same kind, of four or five years' standing, in
growths after having laid open the larynx, it appeared
mass as was removed could ever have found room in a
laryngeal mucous membrane was literally covered by gr
size and density. The patient nevertheless made a r
now a very fair voice.

In most cases the rapid increase of the
general inflammatory condition which may
tion with their presence, and the consequ
respiration, render tracheotomy necessary,
operation, lead to a fatal result at a compara
But in some instances patients suffering from
of this kind have been known to live for year
it is probable that the larynx has been kep
for breathing purposes by the detachment an
portions of the growth from time to time
Türck records a case in which the whole g
have been thus got rid of. §

I have at the present time a little girl under my ca
whose trachea I opened three years ago on account of
and with the view of subsequently removing the growth
ever, objected to any further operation. The child's vo

ding to some observers, it would appear that these
atous growths occasionally undergo a process of pulpy or
degeneration.* In such case it is easy to understand
y may become broken down and expectorated.

us and *Fibrocellular Growths* are met with in the larynx
ss frequently than the papillary growths above described,
erous instances are on record.

growths are smooth on the surface, hemispherical,
, or pyriform in shape, and on laryngoscopical inspection
pear somewhat paler in colour

the mucous membrane of the sur-
g parts. In substance they are
less firm and compact. In
e, they resemble growths of
nature developed in other parts;
le, they are solitary; they may
e with broad base, or peduncu-
If pedunculated, the pedicle is
y less firm and more vascular
e growth itself. In a large pro-
of cases these growths are com-
ly small, and do not each exceed
of a pea; but they may attain
able dimensions. In an in-
ferred to by Rokitansky, the
was as large as a pigeon's egg.†

pedunculated fibrous growth of me- Pedunculated fibrous polypus
ze is represented in Fig. 259, of the larynx arising near
taken from a specimen in the the anterior extremity of the
of Guy's Hospital.† right ventricle of the larynx.



FIG. 259.

arger pedunculated growths generally take origin from
ther of the true vocal cords, or from some immediately
g part of the ventricular wall—and in the latter case

L. Causit, *Étude sur les Polypes du Larynx chez les Enfants*. Paris,

hr. d. k.k. Gesellsch. d. Aerzte z. Wien, 1851, quoted by Krishaber.
730.

ration No. 1703²⁰. Tracheotomy had been performed in this case;
tient had for a long time been comparatively comfortable, but voice-
day the tube got displaced. Spasm of the larynx, conjoined with
nical obstruction caused by the growth, proved immediately fatal.

V.

P P

most frequently near the anterior extremity of the ventricle. They may, however, arise in other parts of the larynx, but they very seldom do so; in scarcely any instance has a growth of this kind been found connected with the mucous membrane covering the arytenoid cartilages.

Adenomatous or Glandular Growths are occasionally but rarely met with in the larynx. As a rule, they take origin from the mucous membrane covering the arytenoid cartilages, the aryteno-epiglottidean folds, or the base of the epiglottis,—or in general terms from those parts of the mucous membrane which are naturally most abundantly supplied with glandular structures. In this respect they differ remarkably from all other laryngeal growths of nonmalignant character. They consist essentially of enlarged and hypertrophied glands and follicles surrounded by more or less thickened submucous connective tissue and mucous membrane. Occasionally they appear to contain adenoid structures of new formation. They may be sessile or pedunculated. When sessile (localised glandular hypertrophy of the mucous membrane), they often present more or less lobulated appearance; and in some instances they attain very considerable dimensions.* When pedunculated they are usually more or less pyriform or globular, and the peduncles are proportionately longer than those of the fibro-polypi. When seen during life, by aid of the laryngoscope, these growths are generally of a deepish red colour, but they often become pale as they advance in age. They are remarkable for the rapid changes in volume they are liable to undergo under various circumstances. Thus a slight attack of catarrhal laryngitis is almost sure to be attended by speedy and considerable increase in the size of the growth, which may, however, again become smaller as the inflammatory condition passes off.

Cystic Tumours of various kinds are met with from time to time in the larynx; but instances are comparatively rare. In some cases they may possibly result from the degeneration or breaking down of myxomatous growths, as suggested by Coe.

* See case quoted by Ehrmann. *Histoire des Polypes du Larynx*. Strasbourg, 1850.

vier;* but more frequently, probably, they owe their existence to causes corresponding to those which lead to the formation of similar cysts in other parts. In some cases the contents are serous, or serosanguinolent; in others mucous, or synovial in character. They may occur in almost any part of the larynx, and in some cases have attained considerable size.†

An interesting, and I believe almost unique example of mucous cyst containing the posterior aspect of the epiglottis, came under the care of Dr. J. B. D. myself in Guy's Hospital some few years ago. The patient, a boy eleven years of age, had suffered two years previously from a severe sore throat, and ever since had experienced more or less difficulty in speaking. This difficulty gradually increased. His voice soon became hoarse, and by-and-bye he began to suffer from frequent and severe attacks of asphyxia, which often came on during sleep. On laryngoscopic examination, the epiglottis was not to be seen presenting its ordinary form; but instead of it appeared a large rounded swelling, projecting downwards and backwards, completely covering in and concealing the glottis. The tumour was of a pinkish colour, shining, and somewhat translucent in appearance. It could be moved by the finger, and was elastic, but very tense to the touch. To ascertain that it contained fluid, I made a free incision into it, by means of a sharp-pointed bistoury. A sudden gush of thick, glairy, mucous matter took place; and after the momentary dyspnoea which occurred immediately, the patient was relieved of all the symptoms from which he had suffered so severely. He made a rapid recovery, and left the hospital the next day. I examined him laryngoscopically four years afterwards (as well as from time to time in the interim), and could discover no indications of any recurrence of the disease, nor indeed any very clearly perceptible mark of my incision.‡ A somewhat similar kind is recorded by Delorme, in which two cysts, containing albuminous contents, were found on the posterior aspect of the epiglottis of a man forty years of age, who died asphyxiated.§

It is worthy of remark, perhaps, that *true hydatid cysts* are rarely to be found in the larynx in some few very exceptional cases.||

For instances of *Cartilaginous and Osseous Growths* in the

larynx, see the case reported by Krishaber.

A remarkable case described by Gibb (op. cit. p. 154), in which the tumour apparently sprang from the right ventricle, and filled nearly the whole

Transactions of Royal Med.-Chir. Soc. Vol. xlvii. 1864.

Bulletin de la Société de Médecine de Paris, janvier 1808.

Gazette médicale, fév. 1835; and Andral, *Anat. path.* Tome ii. p. 490, and Ryland, *On the Diseases and Injuries of the Larynx*, Lond. 1837,

larynx are on record; but the accounts given, so far as I have been able to make out, are not altogether satisfactory.*

Besides the growths of the various kinds thus described others of mixed form and character are occasionally met in the larynx. Thus the fibrous, fibrocellular, and glandular elements may vary in relative proportion in such manner to such extent as may render it impossible to assign the particular growth under examination to any definite class. Again some growths which superficially appear papillomatous may have firm fibrous bases. And, further, the condyloma syphilitis, the thickened elevations of phthisis, and the protuberances resulting from localised chronic inflammation in some cases can hardly be distinguished from what more properly perhaps, ought to be regarded as 'new growths' in the strict acceptance of the term.

A very interesting specimen was almost accidentally

FIG. 260.



Extroversion of the mucous membrane of the left ventricle of larynx, simulating laryngeal polypus.

with some short time since in the post mortem room at Guy's Hospital, in which there was extroversion of the mucous membrane of the ventricle of the larynx. Fig. 260 conveys a somewhat feeble idea of the appearance presented. The everted mucous membrane could easily be replaced in its proper position and when this was done there was nothing abnormal in the appearance of the larynx. The patient had no laryngeal symptoms.†

Cancerous Growths in the larynx for the most part *epitheliomatous* in character; but instances of *malignant lary cancer affecting the larynx* are wanting.

Epithelioma of the larynx generally commences on the laryngeal aspect of the mucous membrane covering the ary-

* See Ryland. Op. cit. p. 231. *Edin. Med. Journ.* Vol. xxxv. Krish. Op. cit. p. 769. Türk. Op. cit. Fall lxxxi. p. 321.

† The case is fully described by Dr. Moxon, *Trans. Path. Soc.* 1868, p. 18. The preparation is in the Museum of Guy's Hospital. Preparation 1683.

id cartilages, and subsequently invades the interior of the

But in some instances it commences inside the larynx ; such cases, as a rule, it first appears on the mucous membrane of the posterior and lower part, over or near the internal of the cricoid cartilage, in the form of small irregular growths, which gradually increase in size and soon ulcerate. Instances of epithelioma commencing in this situation, in the manner described, have come under my own observation. In each the precise position and character of the growth was recognised at a comparatively early period ; and in each case the result of the case established the correctness of the diagnosis. The distinctive features of advanced epithelioma of the larynx, as seen by aid of the laryngoscope, have been already mentioned (see p. 562).

Symptoms to which ‘new growths’ or ‘polypi’ in the larynx give rise vary in character and severity with the dimensions and precise relations of the particular growths under consideration ; and also with the intensity of the inflammatory action in which in many cases may be excited by their presence, and the rate may be sooner or later associated with it.

Difficulty of breathing, alteration or extinction of the voice, and, it may be (especially in the case of cancerous growths) difficulty of swallowing ; these are the symptoms—as they are also the symptoms, varying only in intensity and frequency—in all serious affections of the larynx, as well as in other maladies in which the larynx is only indirectly affected.

It is very rarely indeed, if ever, that any one of these symptoms manifests such peculiarities as may be considered sufficient to indicate the presence of a ‘new growth.’

In many cases when the patient first comes under observation the difficulty of breathing is so urgent, or becomes urgent so rapidly, that tracheotomy must be performed before any exact diagnosis can be made. The difficulty of breathing may depend upon the mechanical obstruction caused by the growth, or upon spasm of the larynx excited by an incidental contact of the growth with some unaccustomed, or already irritated part of the highly sensitive boundaries of the glottis. A growth may exist for a long time almost harmlessly, moving in accordance with the respiratory movements of the larynx, and at once may get into such position as to excite the severe attack of spasmodic dyspnoea. This is especially

likely to occur in the case of pedunculated growths. It may occur in other cases, if from any cause an attack of ordinary laryngitis should supervene. Sessile growths by their gradual increase may gradually constrict the aperture of the glottis, and thus proportionately impede respiration, without giving rise, at least during their earlier stages, to any such suddenly serious and dangerous results as those thus indicated.

In the case of growths situated below the glottis it may happen that expiration is almost as much impeded as inspiration.

In some cases by means of the stethoscope placed over the larynx a sound produced by the movements of the growth during respiration (the 'bruit de drapeau' of the French) may be detected.

The voice is affected to a greater or less extent in almost all cases; and if the growths are small and situated on the vocal cords, the affection of the voice may be the only symptom which the patient complains. The voice may be simply changed in tone and quality and become harsh; or it may be the 'cracked pot' or 'Punch,' sound; or it may be reduced to hoarse whisper; or it may be altogether extinguished. In some exceptional cases the voice, at first seriously affected, has been noticed to improve to a certain extent with the increase of the growth.

In a considerable proportion of cases there is little or no troublesome cough. But if an attack of catarrhal inflammation should occur, the cough becomes especially distressing; and fits of spasmodic cough may be excited from time to time in a case in which a pendulous tumour is liable to get into obstructive position. Occasionally the patient becomes conscious of the presence of some cause of obstruction, and tries 'to cough it up.'

Slight mucous or mucopurulent expectoration may attend the cough; sometimes portions of the growths may be mingled with the expectorated material;* and, as already stated, in some exceptional instances considerable portions, and even the whole of the 'new growths' have been got rid of in this manner.

Among the various affections which may give rise to symp-

* It may be interesting from an historical point of view, to quote the words of Ehrmann, writing in the year 1850:—'Il n'existe, il faut le dire, qu'un signe certain de l'existence de cette maladie; c'est l'expuition de quelques-unes des cellules du polype.'—Op. cit. p. 31.

similar to those thus referred to as caused by the presence of new growths in the larynx, may be enumerated—chronic laryngitis with cedema; spasm of the larynx, from whatever cause; laryngismus stridulus; nervous aphonia; pressure upon the pneumogastrics or therecurrent branches by tumours, aneurisms, &c.; retropharyngeal abscesses; pharyngeal polypi hanging down or projecting into the larynx, &c. It is, however, in the present day altogether needless to discuss in detail the distinctive differences that may or may not be presented by the symptoms in these several affections. In all cases in which time and opportunity are at command the diagnosis may be readily determined by aid of the laryngoscope. In cases, however, in which the symptoms are very urgent, whatever may be their cause, relief must be afforded and safety secured by tracheotomy or otherwise before any full enquiry can be entered upon: laryngoscopic examination may then be made under favourable circumstances at any convenient subsequent period. It occasionally happens in the case of growths taking origin low down, that they almost disappear under the vocal cords or into the ventricles during inspiration, and only come thoroughly into view during expiration.

It is worthy of note that in some cases, particularly in children who are difficult subjects for laryngoscopical examination, valuable, if not conclusive indications may often be obtained by digital exploration. The effort at vomiting commonly excited during the introduction of the finger into the pharynx causes the larynx to rise momentarily, and thus to come within easy reach.

Treatment.—Measures should be adopted for the removal or destruction of all nonmalignant new growths in the larynx as soon as practicable after their presence and probable character have been clearly ascertained. For it may be asserted without fear of contradiction that no patient can be considered safe who has in his larynx a new growth of any considerable size—that is to say, unless his safety has been secured by tracheotomy. And small growths, as already stated, are liable to enlarge more or less quickly, and often unexpectedly; and even while still small they may favour the supervention of dangerous inflammation.

In some cases it is absolutely necessary, and in others it may be desirable, to perform tracheotomy before proceeding to any other operative measures. Very many cases, however, have

DISEASES OF THE LARYNX.

which, without tracheotomy, growths have been removed from the larynx, not only with the best ultimate results, but also without having involved the patient even in any serious risk.

The advisability of performing tracheotomy as a preliminary measure must be determined by the urgency of the symptoms, the probable difficulties and dangers of the proceedings which are to be carried out.

A great variety of different methods have been devised and adapted for the removal of growths from the larynx. They may be divided into those in which the operation is performed by aid of the laryngoscope through the mouth and natural passages, and those in which access is first obtained to the interior of the larynx by means of incisions in the median line through the skin, cartilages, &c.

The following table, compiled from published records of cases and the results of my own observations, may serve to convey some idea as to the relative frequency with which the different methods of operating specified have been deemed expedient, as well as the degree of general success that has attended the efforts hitherto made.

TABLE OF CASES OF OPERATIONS FOR THE REMOVAL OF NEW GROWTHS FROM THE LARYNX.*

Methods adopted.	Total number of cases.	Completely successful results.	Partially successful results.	Deaths.
A. OPERATIONS THROUGH THE MOUTH AND NATURAL PASSAGES :				
Direct application of caustics or powerful astringents	16	12	4	
Removal or crushing by forceps	37	33	3	1
Removal or crushing by wire snare	32	28	3	1
Removal or destruction by the galvano-caustic wire or pencil	5	3	2	
Removal by knife or scissors	20	14	5	1
Simple puncture	4	4		
B. OPERATIONS AFTER EXTERNAL INCISION INTO THE LARYNX, ITS CAVITY BEING MORE OR LESS COMPLETELY LAID OPEN				
	24	15	5	4
Total	138	109	22	7

The above table must be taken for what it may be worth. I

In the cases specified as 'completely successful' the voice was restored. In those specified as 'partially successful' the voice remained more or less impaired. In the

certainly appears to indicate the methods of operating that have hitherto been found most successful. But it cannot for one moment be regarded as affording any trustworthy guide to the method that ought to be adopted in any particular instance that may come under observation.

In deciding upon the course to be pursued, it is necessary in every case to take into consideration not only the size, precise location, and character of the growth, but also the age, development, and condition of the patient, together with his general and special powers of endurance. If success is not obtained by the method first adopted, another may be tried. Several methods were put in practice in each of many of the cases included in the above table. But in every instance the case is referred to the method which finally proved successful.

A. Operations through the Natural Passages.

1. *The Application of Caustics and powerful Astringents to growths in the larynx*, may be accomplished by the several methods already described (see p. 529); in the case of small isolated papillary growths, such applications several times repeated, are not unlikely to prove efficacious.

2. Small fibrous, fibroid, and fibrocellular growths, and small large papillary growths may, in many cases, be successfully seized by means of properly adapted *Forceps*, and either torn off and removed in mass, or so crushed as to lead to their destruction. In the case of multiple growths, the repeated use of the forceps is often necessary. Portion after portion may be removed either at the same sitting, or during a series of sittings at intervals, according to circumstances. This method is especially advocated by Mackenzie;* and the laryngeal forceps, which bear his name (Fig. 261), are, I believe, the most generally applicable, and therefore the best of all the instruments yet devised for such purposes. Forceps made upon the principle illustrated in Fig. 262 are also very efficient. Some of these forceps are made to open laterally, and others antero-posteriorly, like those represented in the figure. In some cases binary laryngeal forceps, such as those figured and described

cases, with two exceptions, the growths were epitheliomatous or cancerous in character.

Op. cit. p. 114.

in the chapter on Foreign Bodies in the Larynx (see Vol. I p. 524) have been used with perfect success.

3. For the removal of laryngeal growths by *the Wire Snare* is necessary to employ a slender and appropriately curved instrument carrying a wire loop, which can be quickly and easily drawn home when once it has been made to encircle

FIG. 261.



Mackenzie's Laryngeal Forceps.

growth. The instrument successfully used by Dr. Walker Peterborough in the first case in which a laryngeal growth removed from the larynx during life in this country was a modification of Gooch's double canula.* But the double tube is cumbersome and altogether unnecessary. Gibb's 'Laryngeal Écraseur'† is a far more perfect instrument—indeed, not

FIG. 262.



Laryngeal Forceps, opening antero-posteriorly.

could be better adapted for its purpose, unless it be some modification as that suggested by Dr. George Johnson.‡ The instruments bear a general resemblance in construction and mode of working to Wilde's Aural Polypus Snare, and Hilton's Polypus Snare (see p. 301, Fig. 230); but they are, of course, different in shape and more slender. In using this instrument

* See *Lancet*, November 9, 1861, p. 444.

† Made by Weiss; see Gibb. Op. cit. p. 138.

‡ See *Transactions Royal Med.-Chir. Soc.* Vol. li. p. 173 et seq.

wire loop is made to encircle the growth, and then pulled tightly home. The growth is then forcibly pulled or twisted from its attachment, and, as a rule, is brought away in the loop. This method is strongly advocated by Sir D. Gibb,* and Dr. George Johnson.† It is especially applicable in cases in which the growths project in such manner and direction as to render them easily caught, and in which they are soft or pliable in texture, and not very large.

4. *The Galvanocaustic Method* was first practised by Middel-dorpf, and has since been adopted by Voltolini, Türck, Bruns, and others. It has been still more recently recommended by Dr. Reichel of Breslau.‡ This method consists in encircling and cutting off the growth by means of a platinum wire, which is so arranged in an appropriate carrier that when in position it can be intensely heated by means of a galvanic battery with which it is connected. The difficulties of this method are sufficiently obvious, and they, together with certain other practical objections to it, are clearly and forcibly stated by Bruns,§ after some experience of its use. Its advantages in certain cases are equally obvious; and it would appear to be especially applicable in cases in which single fibrous growths exist, the peduncles of which are too strong to permit them to be safely torn away by means of the snare or forceps. A touch of the heated wire may be sufficient to dispose of small papillomatous growths.

5. *The Knife*, properly curved and guarded, or the *Laryngeal Scissors*, may be used for the separation of firmly attached growths which may have been seized but cannot be pulled off by the snare or forceps. Such means were adopted in several of the cases successfully treated by Bruns.|| Several different forms of laryngeal guillotine have been devised. Of these Matthieu's is probably the best. It has been used with success in several cases.

6. *Simple puncture* by the guarded bistoury or some specially contrived instrument (such as Mackenzie's laryngeal lancet) has proved completely efficacious in several instances in which the growths were cystic in character; and in such cases this method

* Op. cit. p. 138.

† Op. cit. p. 173 et seq.

‡ *Klinische Wochenschrift*, No. li. 1869. Berlin.

§ *Die Laryngoskopie und die laryngoskopische Chirurgie*, p. 244 et seq. Tübingen. 1865.

|| See Table, Op. cit. p. 254.

may be recommended. If in any case the cyst should fill again, it may be desirable on a second occasion to remove a portion of the cyst wall, or at least to make a more extensive incision than the previous one.

It is scarcely needful to add that all these several methods of removing laryngeal growths require during their execution the guidance afforded by the laryngoscope.

*B. Operations for the Removal of Laryngeal Growths after Exposure of the Cavity by External Incision and Section of the Cartilages, &c.**

In cases in which the growths are numerous, or very large, or single but firmly attached, and in those cases again in which the patients are young, and ill able to bear the introduction of instruments through their narrow natural passages, there can be no doubt, I think, that the easiest, most certain, and at the same time the safest method of operating consists in laying open the cavity of the larynx by external incision, and then removing the growths by scissors, *écraseur*, or galvanocautic wire. The results of experience † encourage the further and more confident adoption of this procedure in all such cases as those referred to. It affords opportunity for the more immediate, complete, and effectual removal of the growths than can be obtained in any other manner. It is by no means difficult of execution, nor does it involve any immediate risk. Such bleeding as may occur can be readily checked, for all the parts are fully exposed; and freedom of respiration may be secured by the preliminary performance of tracheotomy—if, indeed, this operation have not previously been rendered necessary.

A good illustration of the success attending this method is afforded by the case of a girl, thirteen years of age, under my care some years since in Guy's

* It is worthy of note, that the first recorded operation of this kind was performed in the year 1844, by Ehrmann, who, without the aid of the laryngoscope (not then in use), diagnosed the existence of growths in the larynx from the symptoms and the appearance of portions in the sputa. He was justly so confident of the correctness of his diagnosis, that he boldly opened the larynx by median section, found the growths, removed them, and cured his patient. *Laryngotomie dans un cas de Polype du Larynx*; Strasbourg, 1844. Also, *Op. cit.*

† See especially the excellent treatise of Dr. Charles Planchon, *Faith cliniques de Laryngotomie*; Paris, 1869. In this treatise a large number of illustrative cases are given.

Hospital. Tracheotomy had been performed four years previously on account of the urgent dyspnoea from which she then suffered. On laryngoscopic examination, masses of warty growths were distinctly seen. These were removed by the scissors after exposure of the laryngeal cavity by section through the thyroid cartilage, cricothyroid membrane, and other structures. Nitrate of silver was freely applied to the bleeding surfaces. The growths had occupied the whole cavity of the larynx. The divided parts were brought together. The patient made a speedy recovery, and the tracheotomy canula which she had worn for four years, was removed ten days after the operation. I saw her the other day, three or four years after the operation. She was in good voice, and no appearance could be discovered of any recurrence of the growths.* In two other cases in which I have performed similar operations, the results have been so far equally satisfactory.

A remarkable case is recorded by Balassa of Pesth. The patient, a young woman, twenty-one years of age, had suffered from more or less difficulty of breathing, and had been aphonic for more than two years. On laryngoscopic examination growths in the larynx were seen. Balassa laid open the larynx by section through the thyroid cartilage, and removed five portions of new growth. The patient made a good recovery, and her voice returned. Less than a year afterwards, she again lost her voice, and experienced increased difficulty in breathing. Balassa a second time opened her larynx through the thyroid cartilage in the situation of the old cicatrix. Three or four portions of new growth were removed. The patient again made a good recovery; and as the result was altogether relieved of her dyspnoea, and regained her voice.†

In operations of this kind the thyroid cartilage and cricothyroid membrane, may be divided in the middle line probably with impunity. But it is desirable to avoid, if possible, cutting through or otherwise damaging the cricoid cartilage.

MM. Pratt and Follin have each succeeded in removing tumours from the larynx after having divided the thyrohyoid membrane, and superficial structures by transverse incision.‡ The objections to this method, arising principally from the necessary section of the hyoidean muscles, as well as from the imperfect extent to which the cavity of the larynx is exposed, are sufficiently obvious.

In the case of epitheliomatous and other cancerous growths, it would appear to be worse than useless to attempt removal, without first fully opening the larynx and exposing the whole extent of the morbid structures. Whether any such attempt is justifiable the circumstances of the case must decide. As a

* See *Guy's Hospital Reports*, 3rd series, vol. xii. p. 541.

† *Wiener medizinische Wochenschrift*, November 1868, quoted by Planchon. 2. pp. 62 and 79.

‡ *Bull. Gazette des Hôpitaux*, 1857. Follin, *Archives générales*, fév. 1867. ed by Planchon. Op. cit.

general rule tracheotomy becomes necessary sooner or later; and it is probable that when once the character of the malady is recognised the earlier the operation is performed the better.

NERVO-MUSCULAR AFFECTIONS OF THE LARYNX. NERVOUS OR FUNCTIONAL APHONIA. HYSTERICAL APHONIA. PHONIC PARALYSIS, &c.

Nervous, or Functional Aphonia.—Numerous cases come under observation from time to time in which the voice is lost, and which also, more or less difficulty in breathing may occasionally occur, although no visible defect or disease can be discovered in the larynx itself on the most careful and complete laryngoscopic inspection. It may be seen that the several parts do not act properly, and that is all. Why they do not so act cannot satisfactorily be made out. Sometimes when the patient tries to utter a sound the glottis may be seen to remain widely open and no sound is produced. Sometimes one of the vocal cords appears to come into proper position, the other remains motionless, and apparently paralysed. Sometimes the vocal cords seem to approximate naturally at the commencement of the attempt to speak, and then separate without vibrating, so that the air from the lungs is forced against them. Other odd phenomena may be observed by aid of the laryngoscope.

Alluding to cases of this kind, Sir Benjamin Brodie states (and his statements are supported by general experience and observation):—

‘This affection takes place suddenly, continues often for many months, or for one or two years, and then disappears as suddenly as it began. A patient thus affected may, when under the influence of strong mental emotion, be herself speaking in her natural voice, when for some time before she had spoken only in a whisper. Her recovery may be permanent, or she may relapse into her former condition. And again, this affection is not unfrequently met with in the male sex, especially in those of the clerical profession, probably because they often lead very sedentary lives, and also because they are called upon to speak in public in a tone raised above the ordinary standard.’

Cases of the kind under discussion are nowadays commonly classed as examples of some so called ‘hysterical affection of the larynx,’ although in many cases no other indications of hysteria are present. There can be no doubt but that in a large proportion of instances the sufferers are either delicate

* *Lectures on Certain Local Nervous Affections*, p. 51.

omen, or highly sensitive 'nervous' subjects of the
 At the same time it must be admitted that the true
 y and proper explanation of the affection are at present
 r wanting. Türk * and Gerhardt * on the Continent,
 kenziet† in this country have of late especially attempted
 precision and definiteness to our ideas respecting the
 phenomena. They have also contributed somewhat
 knowledge of the determining causes of the affection.
 treatment that should be adopted when it is fully
 ed.

imes the aphonia comes on as the result of exposure to
 r extra exertion of the voice. In such case, as suggested
 urdt, it may be to some extent analogous to the facial
 due to *coup de vent*. Sometimes it appears to be simply
 t of fatigue and incapability for renewed effort. Some-
 remains after an attack of 'sore throat' has altogether
 ff; and in such case it may be the effect of indisposi-
 it were, on the part of the muscles to resume those
 functions, from the performance of which they have
 nporarily restrained by reflex sensibility. Sometimes,
 he determining cause may be mental emotion, from the
 effect of which the nervomuscular apparatus of the
 may have been unable to recover itself. But much
 equently the explanation remains altogether undis-

ffection now under consideration must be carefully dis-
 ed from those forms of laryngeal paralysis with aphonia
 eness which are due to the pressure of tumours, aneu-
 sc. upon the motor nerves of the larynx.

ment.—In some cases, and almost always in such as
 to depend simply upon local indisposition to exertion
 g from the effects of exposure to cold, fatigue, &c., re-
 akes place naturally after a certain period of rest. But
 y cases, and most constantly, perhaps, in such as are
 scure in their origin, the aphonia may remain persistent
 weeks, months, or even years; and then, either as the

cit. p. 430, et. seq.

how's *Archiven*, vol. xxi.

Toarseness and Loss of Voice in Relation to Nervo-Muscular Affections of
 ur; London, 1868. See also Lagarde, *De l'Aphonie nerveuse*, Paris, 1865;
 haber. Op. cit.

IN some cases the introduction of the *ice* together with all the attendant formalities, required. Türk appears to have been the first 'method of cure' of aphonia.* Trousseau alludes to several instances.† Three have come under my observation. One of these may be worth mentioning. It affords an excellent illustration of the results in cases of this kind the voice may sometimes be regained.

A young lady about nineteen years of age was brought to me who appeared most anxious about her condition, believing a serious disease existed. I was informed that the patient had been in bed for more than six months, and that all sorts of treatment had been tried but with no good effect, &c. After a great deal of time and many abortive attempts, I at last succeeded in putting her into position. I then, in a matter of course tone, requested her to say 'Ah, ah.' Pushing away my hand, she immediately and in a fair natural voice, 'How can I, with that thing in my throat?' she added, 'Oh, dear! I've spoken!' It is almost needless to say the cure was complete. But supposing that it had happened that the patient had been in process of application at the time the cure this would have been!

In some cases the inhalation of strong stimulant vapours, as those of ammonia or chlorine solution; in others the application of solutions of sulphate of copper by means of the sponge, is followed by restoration of the voice. This method found especial favour with Trousseau, and is illustrated by the good results obtained in

Another method of treatment especially relied on by Macle consists in the direct galvanisation of the vocal cords. It can be most easily and perfectly accomplished by means of Mackenzie's laryngeal galvaniser or electrode. There can be no doubt of the efficacy of this method. But—without any idea or intention of impugning for one moment the genuineness of the numerous examples from time to time published, in which cases aphonia of long standing have been stated to have been cured single, or by a few successive 'applications of galvanism to the vocal cords'—it may nevertheless be asserted that the application of galvanism possesses no *specific* virtues in such cases as are under consideration. It simply affords a convenient and effective method of giving the necessary 'fillip' to the laryngeal apparatus. Equally good results have been obtained by passing the interrupted current of an electromagnetic battery through the back of the tongue.

The method of treatment most recently advocated consists in manipulating (that is pinching and pressing backwards) the larynx; and many cases are recorded illustrating its efficacy.* In conclusion, it may, I believe, be stated without fear of contradiction that the one thing especially necessary is, if possible, to arouse or startle the larynx into proper activity, and it matters comparatively little by what means this effect is accomplished. It must be borne in mind that in these affections there is no perceptible organic disintegration or wasting of the tissues. The larynx appears to be sufficiently exercised for purposes of nutrition during their ordinary respiratory movements, which, as a rule, are carried on normally and without interruption.

Idiopathic Atrophy of the Muscles of the Larynx, and Muscular Atrophy as the result of disease are very rare affections. The examples that have at present come under investigation are so few that no satisfactory conclusions can be deduced.† It may well, however, to bear in mind that such cases may from time to time present themselves.

In every case of aphonia, hoarseness, or other alteration of the voice, in which on laryngoscopic inspection one vocal cord is found to be more or less completely paralysed, and in

* See H. K. Oliver, M.D., in the *American Journal of Medical Sciences*, vol. 1870.

† See Türck. Op. cit. p. 203. Mackenzie, *On Hoarseness, &c.* Op. cit. p. 71.

which, at the same time, the parts generally are natural in appearance, search should be made for some cause of pressure upon, or interference with, the motor nerves of the affected side of the larynx. If the vocal cord rests motionless during attempted vocalisation, attention should be especially directed to the recurrent nerve. If, however, the vocal cord is adducted and abducted more or less normally, but is not duly rendered tense, it may be that the superior laryngeal nerve, or rather its cricothyroid branch, is interfered with. Paralysis of the laryngeal vocal cord not infrequently indicates the existence of an aneurism of the arch of the aorta, or some intrathoracic tumour.

SPASM OF THE LARYNX. SPASM OF THE GLOTTIS.

Spasm of the larynx, attended by the most alarming and sometimes fatal dyspnoea, may arise from various different causes. Many such causes have been already discussed in the preceding pages. Others may depend upon the effects of injuries, or introduction of foreign bodies into the airpassages (see Vol. pp. 458—478, et seq.). Others again may depend upon irritation produced by the pressure exerted upon the laryngeal nerves by aneurisms or tumours, whether they be 'new growths' or simple enlargements of some natural organs, as the thyroid body, lymphatic glands, &c, and whether situated in the neck or within the thorax. Intrathoracic tumours pressing upon the trachea or bronchi by some reflex nervous influence, hysteria, tetanus, and certain cerebral affections in some cases altogether inexplicable, may also give rise to spasm of the larynx.

However difficult it may sometimes be to ascertain the cause, the symptoms of this formidable affection are always unmistakable.

Treatment.—If the spasm does not very quickly and very completely subside under simple treatment—and even in absence of all satisfactory explanation of the cause of the attack—tracheotomy should be performed without hesitation. There is more danger in waiting too long than in operating too soon: whatever the result, the conscientious surgeon may always be assured that it is better to have made an unnecessary incision into the windpipe than to have allowed his patient to die of want of one.

ARTHUR E. DURHAM

DISEASES OF THE INTESTINES.

ABDOMINAL ABSCESSSES.

ABSCESS originating in the substance of the anterior abdominal wall, or in the sub-peritoneal cellular tissue, in region of the loins, or pelvis, and totally unconnected with any bone, or with ulceration of the bowels, is not a very common occurrence; but as we occasionally meet with such a case, it is requisite to allude to its peculiarities and treatment. An abscess confined to the substance of the muscular parietes is most frequently the result of a bruise, and may be treated as such in any other position. But one commencing in the cellular tissue of the loin, or iliac region, is generally obscure at the commencement; frequently produces much constitutional disturbance in its progress; and is very often fatal in its results. Abscesses occurring in these situations are sometimes slow in formation; but in other instances form with great rapidity, and acquire a large size within two or three weeks. According to the rapidity of their progress will be the acuteness of the symptoms which accompany them. They are as frequently seen in children as in adults. Without careful examination, they may be readily mistaken for psoas abscess dependent on diseased vertebrae, or for some kind of tumour growing in the cavity of the abdomen. In children, from the resistance of the abdominal and psoas muscles, and the drawing up of the thigh on the belly, disease of the hip-joint might be suspected, if the conditions be not particularly examined into.

A very sickly child was admitted into the Hospital for Sick Children, under the care of Sir W. Jenner, with a large prominent roundish swelling on the left side, between the region of the kidney and Poupart's ligament. There was very little indication of pus, or fluid, in the swelling; its shape and consistence evinced the impression that it might be a malignant growth. As some doubt existed as to its character, a fine trocar was passed into it by the writer, and a thick pus immediately escaped through the canula. A small opening was made through the parietes into the abscess, and allowed the free escape of

most prominent point, and cautiously extended to some pus escaped, exploration was not carried farther, and a poultice to be applied. On the second day, a sudden flow of pus from the wound; relief to all the urgent symptoms followed; and recovery shortly.

Acute and large abscesses occur in the sub-peritoneal tissue about the region of the ovaries after the birth. For an account of them, and their treatment, refer the reader to Vol. I. p. 129.

Abscess of the abdominal region is often connected with an artificial aperture in the alimentary canal, either in the intestine. Such an abscess may arise from perforation of the bowel, the result of simple ulceration; or the introduction of a foreign body in the intestine; often from ulceration of the gall-bladder by a gall-stone; or from stricture of the intestine; and most frequently from some cancerous growth in the alimentary tube.

When such an abscess occurs, its contents consist of pus and faeces mixed together; but as the opening in the bowel has been so small, that the contents have escaped; sufficient, however, to cause secondary mischief. When the abscess is opened, the escape of mere pus alone does not justify an opinion that the disease is cured; neither is, nor has been, implicated. It will usually, some days after clear pus has been discharged from the opening, that faeculent matter is observed to be coming away with the discharge.

cal attendant may be daily watching for an opportunity to exit to pus at some favourable point, the swelling may only subside, and disappear by evacuating the contents, though some fresh opening, into the bowel, the pus being urged per anum.

When of an acute character, the symptoms of a 'fæcal abscess' are very marked. Sudden pain is referred to the part affected, often of a very severe character; constipation often precedes the attack, or accompanies it. The pulse and tongue change early in the mischief; the former becomes rapid; the tongue loaded and creamy, and soon dry and brown. There is great anxiety and restlessness; much thirst, and a hot skin. There is very frequently, early in the attack, distinct tenderness of the part implicated; and, in addition to extreme tenderness and an intolerance of pressure, general indications of diffused peritoneal inflammation. The most frequent site of such abscesses is the region of the ilio-cæcal valve—right iliac fossa; but they may occur in any part of the peritonæum.

Sometimes the formation of pus is so rapid, and the mischief in the peritonæum is so general, that the patient dies within a few days of the first onset of pain. Mr. —, late House-Surgeon of St. George's Hospital, walking in Hyde Park, was suddenly seized with excruciating pain in the right iliac fossa. Though but a very short distance from his lodgings, he was obliged to be carried home. When seen by the author in the evening, he was in intense pain in the right iliac region, with a certain amount of fullness and great tenderness on pressure. The skin was hot; the pulse rapid, and there was great anxiety of countenance. The symptoms were in no way relieved by the treatment prescribed in consultation with the late Dr. Bright. He proved fatal in a few days. On examination after death, a large abscess was found to occupy the right iliac fossa; the boundaries were formed by the caecum of the intestines and the parietal peritonæum. The contents were pus. The appendix cæci was found ulcerated through at its extremity. No other substance could be found, to account for the ulceration by its pressure in any other way.

It is desirable to note that when abscess is the result of an perforating the wall of the cæcum, much will depend on the exact seat of that ulcer; viz., whether it be in the part covered by peritonæum, or on the posterior surface where the caecal wall is in contact with cellular tissue. In the former instance, such an abscess is frequently rapidly fatal on account of the immediate effects of acute peritonitis: or adhesions of the opposed peritoneal surfaces may have occurred prior to the formation, and the abscess become limited by such adhesions,

cular smooth-edged small opening was found in the wall of the posterior of the cæcum, communicating with a large abscess in the right ilia with numerous sinuses running in various directions. The ulceration to be due to an attack of fever.

Occasionally an abscess in this neighbourhood is more tardy in its progress ; and although matter continues to collect, the patient, having passed over the first acute stage of the attack, may soon improve in health and appetite ; the tenderness of the abdomen, or distension of the bowels, if either have existed—commences to subside ; and a lump remains in some portion of the abdomen, marked by external fulness, and often by a well-defined surrounding redness. The skin over this part becomes more red and shining than of the immediate neighbouring surface ; then darkens ; and becomes thinner at one or more spots ; and either spontaneously or is punctured, when pus readily escapes.

It is a safe and very judicious proceeding to open an abscess, as soon as it may be suspected to have formed, and be within reach of the knife. If the incision be delayed, matter makes its approach towards the surface, it will happen, during this process towards self-liberation, that pus works its way in several directions ; it has been kept back by the resisting action of the abdominal muscular fasciæ, that it has burrowed wide and deep : so when an external opening has been long postponed, by nature,

and late in the progress of the case, long sinuses will often be found, which lead from the opening in the abdominal wall to the surrounding parts, and thus render the ready escape of matter difficult. Under such circumstances, after a time, other abscesses of matter point, either in the groin, in the thigh, or at the ischiatic rami. Many a patient will, however, sink after the opening of a fæcal abscess: not only has he to contend against the effects of profuse discharge, but he has the extra evil of a too ready and early escape of intestinal contents from the opening in the bowel.

Notwithstanding the severity of such cases in their general result, and the frequently fatal results they entail, we occasionally find a patient gradually recover, as it were, from a state of death, and after he has endured the opening of a great number of consecutive abscesses; the sinuses slowly contract; the discharge diminishes by degrees, and ultimately ceases; the orifices of the abscesses cicatrise one by one. The patient, however, recovers with a crippled limb, as a general rule.

The suppurative action usually implicates the psoas and iliacus so much, and often the adductor muscles, that the movements of the thigh are subsequently restricted, nor can the limb be extended to its natural limit.

The general treatment of this class of cases is not to be laid down as simple, nor conveyed in a few words. It depends entirely on the course of the case.

In the early stage, i.e. of pain, constipation, and fever, constitutional measures and local treatment must be combined. Internally, and hot fomentations, perhaps leeches, externally, are usually most beneficial at first. Opium may be given in full doses, and often repeated. It may be requisite to combine small doses of calomel with it. Purgatives, as a rule, should be strictly avoided. Under the influence of opium alone, combined with calomel, the bowels will act as early as their condition will permit; and quite soon enough to answer nature's purpose, *if they will act at all*, without the aid of purgatives.

When the patient survives the shock of the attack of inflammation, so soon as matter is suspected, the part should be explored, and, if possible, the pus let free. Under the influence of chloroform alone, can a satisfactory examination be made, in a case such as we are now considering; and we would, under conditions which indicate, though they do not absolutely require, the presence of pus, urge the use of chloroform, that a

as well as in the local dressings. A constant amount of nutritious food and stimulants are wanted to meet the depressing and lowering effects of profuse secretion from the abscess. It is often surprising to observe the quantity of water required, and will take with impunity, under these circumstances. The bowels must be carefully watched; they are often relaxed, and this diarrhoea, if unchecked, tends to exhaust the patient. If collections of matter should point, they ought at once to be let out; or if superficial, from one abscess to another, they should be slit.

The usual conditions found after death, in the case of an abscess, are general adhesions of the intestines to the surrounding parts in the immediate neighbourhood; some portions of the peritoneum form the boundaries of the abscess-wall. The space between the intestine and the abscess is generally very clear; often no pus is observed; but when more than one exists, the abscess is commonly the result of ulceration *from* the abscess.

Most commonly the mischief which produces an abscess independent of cancer, arises near the ileo-cæcal junction. The inflammation of the appendix is one of the most frequent causes.

In the *Transactions of the Pathological Society* (vol. vii. p. 85), a case is recorded in which perforation of the cæcal appendix was occasioned by the pressure of an intestinal concretion. Another case is recorded (p. 85), in which a fistulous orifice in the abdominal parietes formed a circumscribed cavity, which communicated with the interior of the duodenum, and indirectly with the gall-bladder. This was the result of a gall-stone ulcerating into the duodenum, and acceding to the colon.

the Pathological Society, vol. i. p. 265; another, of abscess in the abdominal walls in connection with cancer of the stomach, vol. xi. p. 122; and a third, of fistulous openings through the abdominal parietes communicating with the transverse colon and stomach, the result of cancerous disease, vol. viii. p. 221.

The treatment of fecal fistulæ dependent on cancer does not call for much remark. The patients often linger much longer than might be expected, provided they can take plenty of nourishment; but they often succumb much more rapidly than anticipated by those in attendance: it is therefore best to warn the relatives of the sufferer of the possibility of sudden collapse in this disease.

INTESTINAL OBSTRUCTIONS.

The causes of intestinal obstruction are so various; its occurrence is fraught with so much danger; its symptoms are so severe and distressing; its diagnosis is so obscure; and its treatment so uncertain, and so often unsuccessful, that it is with the greatest diffidence we undertake the consideration of such an important subject as the pathology and the symptoms of intestinal obstructions, and the measures to be employed for their relief.

In entering upon an inquiry beset with so many difficulties, it has appeared to us a very important point, at the outset, to endeavour to classify the causes of obstruction, so that, in some measure, we may be able to determine, according to the symptoms, whether the obstruction depends on an acute strangulation entailing great and imminent danger to life; or on a slowly altered condition of the intestine itself, or of the adjacent vessels. Under the first condition the treatment must be decided; and to prove effective, must be early applied. In the second, there is generally more time to consider the prospects of life, and the advantages of interference. In the first, without prompt and entire relief, death generally follows rapidly in the wake of the onset of the symptoms. In the second, while we are calculating our measures, the symptoms may even subside; and their sudden cessation be followed by perfect recovery.

We therefore propose to divide the causes of obstruction into, those we may term sudden; such as are productive of acute rapidly fatal results, if not presently relieved: and 2nd, the more slowly acting; productive of symptoms, chronic when

thorough examination may be instituted, and exploration made if deemed advisable.

Great attention and care in treatment is necessary, as the pus has an outlet. Strict cleanliness is requisite, to prevent excoriation from the discharge, and also to avoid bed-sores. As the discharge is often extremely offensive, the chamber becomes insufferable, if not constantly and carefully ventilated. A free use of disinfectants is requisite in the room as well as in the local dressings. A constant and liberal supply of nutritious food and stimulants are wanted to meet the wasting and lowering effects of profuse secretion from the abscess. It is often surprising to observe the quantity of wine patients require, and will take with impunity, under these conditions. The bowels must be carefully watched; they are often apt to become relaxed, and this diarrhoea, if unchecked, tends to reduce the patient. If collections of matter should point in other parts they ought at once to be let out; or if superficial sinuses lead from one abscess to another, they should be slit up.


The usual conditions found after death, in cases of faecal abscess, are general adhesions of the intestines and viscera to the immediate neighbourhood; some portions of which form the boundaries of the abscess-wall. The communication with the intestine is generally very clear; often several may be observed; but when more than one exists, the others are commonly the result of ulceration *from the abscess into the gut*.

Most commonly the mischief which produces faecal abscess independent of cancer, arises near the ileo-cæcal valve: ulceration of the appendix is one of the most frequent causes.

In the *Transactions of the Pathological Society* (vol. vii. p. 210) is recorded a case in which perforation of the cæcal appendix was occasioned by the pressure in it of an intestinal concretion. Another case is detailed (vol. p. 85), in which a fistulous orifice in the abdominal parietes opened into a circumscribed cavity, which communicated with the interior of the colon, the duodenum, and indirectly with the gall-bladder. This was probably the result of a gall-stone ulcerating into the duodenum, and accidental rupture of the colon.

Fæcal abscesses and fæcal fistulæ occur in other parts of the abdominal wall; they are most frequently connected with the large intestine, and are generally the result of some cancerous obstruction, attended by ulceration of the bowel above, which, extending to the integuments, allows a partial escape.

A case of fæcal abscess and fistulæ communicating with the cæcum by cancerous disease of the large intestine, is recorded in the *Trans*



by ulceration, from the gall-bladder.

3. Twisting, or 'dislocation,' of the bowels observed in the large, but often in the small.
4. Loops, formed by bands of false membranes, at both extremities; by diverticula, adhering to some portion of the viscera or abdominal wall, the fimbriated processes, or other parts, by contracting adhesions, so as to form a loop, for intestine to become entangled in the result of peritoneal inflammation.
5. Mesenteric pouches; foramen of Winslow, or peritoneal sheaths,* the result of old adhesions.
6. Invagination—often caused by worms, in the small intestine. It is hardly necessary to mention, that all hernia are excluded in this consideration.

The more slowly acting processes produce in which the symptoms are at first not urgent, relief sometimes occurs without assistance when all hope of life may have vanished, are

1. Constipation, habitual or accidental.
 2. Inflamed, thickened intestine, the result of chronic inflammation.
 3. Chronic peritonitis (tubercular) and abscess.
 4. Tumours pressing on the bowel—hydatids.
 5. Simple stricture of intestine, the result of other causes.
 6. Cancer of the bowel, producing contraction.
- Although we have divided these usual causes

In many cases the symptoms may subside from the acute the chronic, or from a quiet state become very severe. It not be taken as a fact in every case, that acute symptoms do, or do not, depend on any one of the latter causes of obstruction; or a more chronic state of symptoms on any of the conditions noted in the first division; but as a general rule, may venture with some degree of assurance to assert, that several symptoms which will mark the cases coming under the first division are urgent, acute, and rapidly result in collapse and death, if not relieved by nature or art; while the greater number comprised in the second division are attended by symptoms which come on by degrees, may last many days, often weeks, or even months in some cases; are generally by so much less urgent than the former in character, and permit of delay in treatment as regards surgical interference; allow time for consultation; are often capable of medical relief; and, if surgical treatment be considered requisite, offer many more features favorable towards the prolongation of life, than do the acute cases of obstruction. It must be borne in mind, that constipation, or entire stoppage of defæcation, is not an absolutely requisite symptom to indicate fatal obstruction of the intestine; in suspected organic obstruction, should we presume to say that the disease has yielded to our treatment, even though diarrhoeal action of the bowels comes on; it will probably last for a short time, or the quantity of fluid passed will be small in proportion to the frequency of the motions. A relaxed state of the bowels in stricture is usually indicative of ulceration, and may occur, in old contracted stricture of the intestine, within a very short period of death.

1. Congenital malformations of the intestinal canal, giving rise to immediate obstruction, are almost entirely confined to the rectum and its external aperture. These deformities, and their treatment, will be found described in the essay on **THE SURGERY OF CHILDHOOD**, and therefore require no further notice here.

Another, but very rare condition of stricture, has been occasionally found in the duodenum of infants; the two following cases illustrate these structural alterations.

A child, when born, presented no unusual symptoms for the first twenty-four hours; vomiting then came on, and continued, with short intermissions, until the third day, which took place some thirty-eight hours after birth. The bowels were never relieved during life. The only disease found was a stricture of the

under the care of Dr. West, apparently in much pain; convulsions and throwing about her legs and arms. The mother had vomited frequently, rejecting everything she took; there was constipation. The child died in a few days.

The stomach presented a constriction, beyond which looked at first like the stomach, but ended sharply at a fold of mucous membrane; beyond this followed intestine. A probe inserted into the bile-duct, from without, passed the surface of this septum, and appeared just below the constriction, which was detected to be duodenum; and what appeared to be the membrane at the further end of the pouch, proved to be the ileum, with a slight obliquity, across the calibre of the bowel. The constriction was almost exactly in the middle, by a small hole in the membrane. The valve possessed all the ordinary appearance of a mucous membrane.†

These strictures, so similar in position and relations, were probably the result of an abnormal action of the valve of the duodenum; that in the second case was obstructive as the infant grew, on the principle of the membrane or soft tissues, the result of disease, and acquired a tendency to contract towards their normal position.

Such cases are interesting, but are quite beyond the scope of treatment.

2. Obstructions, the result of foreign bodies in the intestines, have already been considered in the *OF THE ABDOMEN*, Vol. II. pp. 465 et seq.; and we add them to complete the causes of obstruction in the small intestine.

3. Twisting, 'dislocation,' or displacement of the intestine, producing obstruction to the passage of its contents, is a common accident, but demands very careful

an obscure, but its treatment is a matter of very great consideration.

incarceration may be the consequence of a rotatory movement, and of this there are three varieties :

1. A portion of intestine may have become twisted upon its axis ; and we then find that even semi-rotation causes such approximation of its parietes, that they touch, and close up passage. This can probably only occur in the colon ; and adding to cases on record, only in the colon ascendens. Accumulation of gas, and unequal filling of different portions of intestine, appear, as far as we are able to judge from the cases which have been noticed, to be the cause. Such an occurrence is scarcely conceivable in the small intestine, on account of the uniformity of its calibre, the absence of angular turns, and its loose position, as every rotation of one portion about its axis would be counterbalanced by the rotation of the next segment.

2. The mesentery may be the axis, and the intestine will be rolled up upon the former ; *i.e.* the entire mesentery, or portion of it, is twisted into a cone ; and in proportion to the number of its rotations, more or less of the intestine will be engaged after it. In this case, we must take into consideration traction and the pressure which the dependent mesenteric forms with the base whence its point rises. This variety scarcely occurs anywhere but in the small intestine and its mesentery.

3. One portion of the intestine, either single or double—a loop—may afford the axis round which another portion with its mesentery is thrown, so as to be throughout in contact with the circumference of the axis, and thus to compress it like a ferule. This variety is evidently a higher degree of the first, in which a portion of the intestine is merely compressed from before backwards, and, as it were, flattened down. A coil of small intestine, sigmoid flexure, or the cæcum, may form the axis.

The last two varieties occur, like the first, chiefly at an advanced period of life. In early life, a predisposition to the obstruction may be caused by a congenital malformation of the mesentery, by large herniæ, or by small herniæ when there is adhesion of the intestine.*

It is a matter of some importance to be reminded of, that the

* Rokitansky, *Pathological Anatomy*, vol. ii. p. 52.

seat of *most* incarcerations *and of all twists*, will be found to the posterior unyielding wall of the abdominal cavity; for only in that direction that pressure exerted on the intestine effect its incarceration; the occurrence of a similar re anteriorly is inconceivable, on account of the smoothness and yielding nature of the parts.

The symptoms of twisting of the intestine, especially of the sigmoid flexure (which is the most common variety), are usually urgent from the first; great pain is suddenly experienced at a small circumscribed spot of the abdomen; obstinate constipation usually setting in from that date. If the sigmoid flexure is incarcerated, there soon follows very considerable distension of the abdomen, often distinctly observed to be chiefly confined to the left side. The distension of the abdomen is generally greater when the large gut is affected, than when the small intestine is implicated. In the latter conditions, enemata may be injected into the rectum very readily, and may return tinged by fecal matter. In twists of the sigmoid flexure, we have injected fluids into this division of the bowels; if retained for a time, they do not return; the fluid has been thrown into the dilated sigmoid portion, and cannot again be evacuated. Vomiting is generally present, and often very copious and constant.

The following cases illustrate the history and pathological conditions of these accidents.

A man, aged twenty-eight, having suffered for five days from irrepressible constipation, died five hours after admission to the Hospital-ship *Dreadnaught*. The sigmoid flexure occupied nearly the entire portion of the front of the abdominal cavity, and was in a gangrenous state. The bowel was immensely distended; obliteration of the muscular bands, and rupture of the serous coat having taken place. This portion of the gut had been originally well developed, and had now become strangulated, from falling over on itself towards the right side, causing a twisting and strangulation. The patient on previous occasions, been subject to constipation.*

A man, aged sixty-five, was admitted into St. George's Hospital with a circumscribed swelling, which occupied the greater part of the abdominal cavity, and apparently extended into the pelvis. Fluctuation could be distinguished over the whole of the tumour, the surface of which appeared smooth. No solid matter could be detected in any part of it. Percussion gave a resonant sound in the upper part of the abdomen only, and there but to a small extent. There was a tendency to diarrhoea. A day or two after admission, peritonitis came on, with great pain in the abdomen and constant sickness. He died on the fifth day. The greater part of the peritoneal cavity was occupied by the

* *Pathological Transactions*, vol. i. p. 103.

coloured tumour, which had displaced the various viscera, and encroached on the chest. This mass was formed of the sigmoid flexure enormously distended. The dilated bowel was connected to the left iliac fossa by a pedicle and by an exuberant meso-colon, which was twisted upon itself, and had occasioned partial obstruction of the gut: the communications respectively into the colon above, and the rectum below, being small and tortuous. The sigmoid cavity was distended with fluid fæces, and contained one or two masses. The mucous membrane was of a dark livid colour, but not ulcerated. Extensive evidences of peritonitis were present.

A man, aged fifty-five, was admitted into Charing Cross Hospital, nine days after the onset of almost complete obstruction of the bowels. A displacement of the bowel prevented the termination of the ileum, cæcum, and ascending colon. The cæcum was pressed upon and strangulated by the termination of the ileum and twisted mesentery. Just at the part where the pressure was exerted, a mass lay in a calcareous state, as large and hard as a marble, lay immediately above the strangulated bowel, and had considerable influence in exaggerating the obstruction, as every effort to force the contents of the gut onwards tended to press the gland into the opening. Close above the constriction the internal surface of the bowel was extensively ulcerated, and perforation was about to take place. In consequence an attempt was made during life to relieve the obstruction by operation in the left loin, but without success, as the obstruction was above the point where the incision was made in the bowel.*

A boy was seized with a violent pain in the stomach, and with sickness. The symptoms continued, varying in intensity, with complete constipation. He died on the ninth day. The intestines were matted together. The duodenum, jejunum, and ileum were greatly distended; the lower part of the small intestine was highly inflamed, and two loops of it were quite black from congestion. The mesentery of these loops had been twisted on itself, and had occasioned strangulation of the gut attached; and while in this state, the folds had become attached to an intestinal diverticulum proceeding from the small gut to the linea alba, about one inch below the umbilicus, and thus gave rise to an additional point of mechanical obstruction to the circulation and contents of the bowel.†

These cases are apt illustrations of the usual forms of 'twist' of the small bowel, such as occur, 1, in the sigmoid flexure; 2, about the cæcum; 3, of the small intestine. As a rule, such twists, when they occur in the small bowel, are usually nearer its lower than its upper extremity.

Rokitansky has very justly observed that the predisposition towards incarceration from twisting of the gut is dependent on congenital, or acquired long, loose, flabby mesentery or meso-colon. But the mere existence of such a state, which allows of free movements of the bowel to take place, is hardly sufficient to account for sudden strangulation of the parts taking place. It appears to us that something is requisite to establish a twist; and that this something is to be found in an accumu-

* *Pathological Transactions*, vol. ii. p. 222.

† *Ibid.* vol. vii. p. 205.

fluid feces from above can enter the sigmoid it rarely happens that much is able to escape.

If, in the dead subject, the large intestine is artificially with water, and the mesentery of it is abundant, this portion of bowel is first seen to descend, and then gradually rise up towards the diaphragm. In this movement there is a slight tendency observed of the intestine on its mesentery; this when the mesentery is open, and no restriction applied to the bowel, it is however, been able, by simply distending it with water, to produce actual twist; but perhaps our subject's was restricted; and no doubt some other cause interfered with it besides distension and free movement. This depends on the motions of the body, motion of the bowels, presence of the other viscera and abdominal organs. The following particulars of a case bear strongly on the point of our subject.

A man, aged forty-five, was attacked on the 10th of November. This ceased on the 12th. On the 13th he was complained of deep pain in the region of the bladder; paroxysmal and severe until his death. Along the wall of the abdomen an unusually hard and broad ridge could be felt, extending from the stomach to the bladder. Death took place on the 15th.

The peritonæum contained a considerable quantity of fluid. In the epigastric region was seen the transverse colon, with the mesocolon adherent to the abdominal parietes, on the left of the umbilicus. The rest of the abdominal cavity was entirely occupied by the small intestine; it lay obliquely, and had forced the diaphragm high up.

and this without any marked disposition to again become twisted. *He justly remarks, that usually the tightness of the twist is in proportion to the extent of the twist of the bowel.** A tube passed up the rectum after death in this case with some little trouble introduced into the sigmoid flexure, beyond the point of obstruction, and could empty the bowel. When this was effected, and the patient rolled over on the side, the twisted bowel righted itself.

The facts just quoted point to this rule in treatment, that a twist of the sigmoid flexure is just possible without opening the abdomen, provided the long tube can be introduced into the distended gut, its contents drawn off, and the twist corrected by the altered position of the bowel. But no operation for the ultimate relief of the patient will be successful, without the intestine be unloaded first, and the twist then reduced. It may be stated as an axiom in these cases, that once formed, the twist prevents the escape of the contents, and the contents of the twisted portion maintain the distortion; and to remedy the obstruction, the contents must be removed.

It will be observed that in one case already mentioned,† the covering of the intestine was found ruptured, after five days of constipation. This rupture had no doubt occurred previous to death. In an experiment made to distend the large intestine with water, the peritoneal coat in several places ruptured, before the muscular appeared inclined to give way to the pressure. This rupture of the peritonæum appears to commence over the transverse colon, and other portions are subsequently affected if the distension is continued.

This tendency of the peritonæum to rupture in sudden distension of the bowels, in cases of obstruction, points towards the importance of early interference, if any operation be considered beneficial. When peritonitis has set in, in consequence of rupture of the serous membrane, or any other cause, operative interference would probably be too late to hold out any prospect of success.

Obstruction of the intestines occurs from a portion of the intestine being strangulated in a loop, or bound down by a cord or false membrane; or it may be constricted by a diverticulum, or adherent by its apex to some opposed surface of the viscera; or the Fallopian tube, attached by its fimbriated process to the point of peritonæum; or by a thickened and elongated portion of omentum fastened down at the ends by adhesions; or other accidental circumstances. All these conditions

Pathological Transactions, vol. x. p. 153. † See the first case on p. 603.



be separated as regards their pathological treatment may be considered under one of the obstructions which are due to the products of inflammation, but most of them are very uncommon in old age, and are of a different nature, whereas strangulation by a mesenteric polypus is in advanced life.

Females are rather more liable to interstitial inflammation of the generative organs; for the appendages of the generative system are points for adhesive inflammation, and entail so much more danger. In most of the conditions now under consideration, the same is alone implicated.

The history and pathology of internal obstructions by false membranes and adhesions produced by inflammation are illustrated by the following cases:

A preparation in the Museum of St. George's Hospital of a short band of lymph, adherent by one extremity to a portion of small intestine; from this the band is seen to be free, and then is immediately attached to the mesentery supporting the latter piece of bowel. When the lymph contracted, it was so pressed on the portion of intestine by the band, that entire stoppage of the bowels was produced.

A young lady, aged nineteen, died after a few days of vomiting. The omentum was firmly adherent to the abdominal wall. This was the result of old adhesions. The stomach was greatly distended. At the lower part of the abdomen, dipping into the pelvis, a large portion of sphacelated omentum was about five inches in length, and consisted of a

ly tinged with blood. At the lower part of the belly several convolution small intestine were highly congested and dark-coloured; this to the extent of a foot in length; and one portion of it was almost black in colour. This portion was found strangulated by a band of false membrane, which was fixed by one extremity to the point of an appendix, by the other to the mesentery. The appendix was attached to the lower portion of the ileum, and about an inch in length, and communicated with the bowel. The appendix and the band with its attachments, formed a complete ring about an inch in diameter.

A man, aged twenty, died after fourteen days' complete constipation. Near termination of the ileum, a portion of it, nearly two feet long, much distended and much darker than the rest, was firmly constricted by a narrow band (thicker than whipcord), which passed from the vermiform process to the mesentery close to its mesenteric attachment.*

In a case in which a portion of small intestine was strangulated through a perforation in the great omentum, perforation of the bowel had taken place, and had prevented the escape of feces.†

A man, aged sixty-eight, was admitted into St. George's Hospital on March 7, 1841. He had been lifting some heavy iron about eight days previously, when he felt a sudden strain, and immediate severe pain in the loins and belly. He passed a restless night; the next morning there was slight action of the bowels. He complained of pain chiefly in the belly and dragging at the umbilicus; there was also constipation from the second day of pain. Four days after the attack he first vomited; but now everything taken was rejected. The skin was cold, and eye sunken; there was blueness of surface, and the pulse was small and weak; the tongue dry and brown; abdomen enormously distended and tympanic. No stricture could be detected through the rectum. An enema-tube could not pass readily beyond a short distance. Mr. Johnson decided to open the ascending colon, with a hope to relieve the distension; but the bowel, when laid down upon, was found collapsed. The question of making an exploratory incision in the abdomen from the front was considered, but decided against. The patient died on the eleventh day. The small intestine was very vascular. In the lower part of the ileum, close to the ileo-caecal valve, a band crossed the small intestine, forming a ring around the gut and the commencement of the mesentery. The origin of the ring was observed to spring from the sigmoid flexure, which was drawn over to the right iliac region so as nearly to touch the caecum. The tissue forming the ring was loaded with fat, closely resembling the structure of an epiploic appendix. The other appendices were very long and broad, and some were perforated at their base, and presented an incipient condition of such a ring as had embraced the intestine. There could be no doubt that this ring was either formed in an appendix, or was formed by the adhesion of two neighbouring appendices. The gut, where it was bound by the ring, was deeply marked by the stricture. From the stomach to the situation of the obstruction the bowel was greatly distended, but more particularly at the lower part, where for about a foot in length it was stretched so tight that it could not be drawn under a stream of water thrown in to wash it out; and was nearly black from congestion. The constricting band was not adherent to the intestine, and on removal from the body the gut could be easily moved backwards and

* *Path. Soc. Trans.* vol. ii. p. 62.

† *Ibid.* vol. i. p. 259.

IN A MESO-COLIC OR MESSO-COLIC POUCH IS NOT

A specimen of meso-colic hernia, which had not been noticed until the death of the patient (1849); having been removed from the body of a man, half of the transverse colon was deflected in a longitudinal middle line, to the brim of the pelvis, and pushed forward, which projected on each side of the displaced colon. The intestines were to be seen; but were found concealed in the pouch to be a large pouch formed by a fold of the meso-colon; but the convolutions could be detected through it at the jejunum entered the pouch at the upper and posterior end, passed out below and on the right side, about two inches from the junction of this portion of the intestine. There was no obstruction of the bowel; nor did the displacement appear to have caused any inconvenience or disorder.

A second similar case was attended by strangulation and death. A man, aged twenty-seven, was seized with vomiting. He vomited everything taken, and the symptoms were most severe. He died about forty-one hours after the attack. The descending colon was found lying on the right side, and the small intestines were contained in a large meso-colon, and situated on the left side of the corresponding ileum passed out of the pouch about two inches above the point the bowel was contracted, thickened, and gangrenous.

6. Invagination, or intussusception of the intestine, the result of irritation caused by worms; of tumours of the mucous membrane; and of other accidental causes, is to be detected or explained.

Invagination may frequently be observed in the post-mortem examinations of children. It also often takes place in the adult.

It is certain, that when obstruction occurs, it is the result of extensively invaginated portion of gut; and that when in mortem examinations several intussusceptions are met with, they are but slight, and have not been productive of urgent symptoms during life.

The Museum of St. George's Hospital is a preparation, which shows an intussusception caused by the presence of a tumour growing from the wall of the bowel and projecting into its cavity. A process of sloughing has nearly separated the prolapsed portion from the rest of the intestine, and so almost set free the tumour. The patient died of peritonitis.

FIG. 263.



Intussusception, caused by the traction of a fibrous tumour of the intestine.

Another preparation in the same Museum displays a portion of small intestine intussuscepted, apparently from the irritation set up by an *ascaris lumbricoides*, which may be seen coiled round the prolapsed portion. But many other parts of the small bowel were also invaginated; and when the patient died, the state of the peritoneum was not suspected.

be quite exceptional, even if it ever occurs common and often fatal.

A perpendicular section through an invagination on each side, and lying parallel with each other intestinal wall; a transverse section would show the intestine, one within the other. In each perpendicular or transverse, the opposed intestines will correspond in character; peritonæum, and mucous membrane be in contact. The outer surface of the external layer is peritonæum; the lining of the inner layer is mucous membrane; but between these there are two layers of mucous membrane, and two likewise of peritonæum, one between the inner and middle layers, and in the space between the middle and outer layers. The portion of mesentery dragged in by the inner layer of bowel as it invaginates. The limit of the invagination above is at the antimesenteric surface of the bowel, and consists of a fold formed by the folding in of the intestine at that point. The lower limit is inside the tube, and can only be seen when the gut is laid open. The portion of intestine between the invagination is a single layer; the invagination which slips into the former, consists of two layers. The portion of mesentery will be found between the two layers, its size and shape varying with the length of the intestine. The lowest extremity of this invagination is almost drawn to a point, and is near the external

mesentery, being always in a state of tension, drags upon the side of the prolapsed bowel; it will therefore be found, usually in an extensive prolapse, that the prolapsed end of the gut, its inner extremity, is turned in the direction of the fixed mesentery; from which circumstance the piece of gut presenting the prolapse will be found slightly curved on one side, and thus the whole mass assumes a crescentic form.

Skitsansky, in referring to this condition, makes the following important remarks. 'Firstly, that the invaginated portion does not lie parallel with its sheath, but always offers greater curvature than the latter, the inverted tube being pressed on its concavity into tense transverse folds. Secondly, that the orifice of the invaginated portion of bowel does not lie in the axis, or in the centre of the sheath, but towards the side; and that, following the traction exerted upon the mesenteric fold that belongs to the inverted intestine, it is directed towards the mesenteric wall of the sheath; that the opening is not circular, but represents a fissure. This presents a diagnostic sign for the examination of intussusceptions of the rectum, which are within the reach of manual exploration.*

Invagination may go on to involve a large portion of the small, and the greater portion of the large intestine. When an intussusception increases in size, the *involution* is that of the small bowel below the commencement of invagination; if this commences at the lower part of the ileum, more of the upper small intestine does not disappear, but the prolapse grows in length at the expense of the inferior portion of the intestine, cæcum, ascending colon, &c.; each of which in turn would become involved in, as the prolapse increased.

We do not attempt to explain the causes of intussusception.

On all probability, an empty state of intestine, and some exciting cause under such conditions, are essential to its production.

If invagination increases beyond a slight extent, a change will soon occur in the condition of the implicated piece of bowel; which, we may predict, places it beyond hope of reduction, accidental or applied. From the pressure exerted on it and its mesentery, congestion, followed by inflammation, soon

* *Path. Anat.* vol. ii. p. 55.

portion within the ring. This is the first step in the effort by nature to remedy the evil, now rendered irreducible that presently the prolapsed portion may be cast off by itself; and that the escape of the contents of the bowel may be provided against, by the union of the upper and lower parts of the otherwise severed intestine.

Peritonitis of a severe character usually occurs when a large portion of intestine is involved, and soon carries off the patient, but should he survive this attack, he may yet recover. The involved portion may slough, and be passed by stool. The patient may rise from his bed convalescent in a few weeks.

The further history and treatment of intussusception are illustrated by the following cases.

A boy, nearly five years of age, complained of soreness about the navel, the result of ascarides. The next day he was seized with sudden pain in the lower part of the abdomen, and strained several times at stool, without effect. He soon afterwards vomited. He continued to go to stool at half-hour intervals. In the afternoon he experienced constant desire to go to stool, and passed a clot of blood. On some of these occasions he passed dark, bloody, and slimy matter. There was great restlessness, and the abdomen was more swollen above than below the umbilicus. There was tenderness over it on pressure. The symptoms continued unrelieved until the fifth day. A considerable intussusception had taken place, the ileum through the ileo-cæcal valve; the appendix was also involved. About 3½ inches of bowel protruded into the cæcum. The involved portion had a deep red colour. It was so firmly grasped by the ileo-cæcal valve that any effort to reduce it appeared likely to tear the bowel.*

te invagination of intestine; sudden pain; constant and urgent desire to stool, with little faecal matter passed, but blood clots or bloody mucus escaping in small quantities; more or less vomiting; great distress, both general and local; and death following in a few days. But death may be somewhat sudden.

A sailor, during a railway journey, was observed to place himself on the floor of the carriage, but had no attention paid him by his fellow-passengers. At the journey's end he was found dead. The abdomen presented marks of severe peritonitis. A fold of intestine near the termination of the ileum, six inches length, was intussuscepted and gangrenous. Probably in this instance the mischief was of some few days' duration.*

In other instances the symptoms appear to extend over a much longer period, and ultimately prove fatal; or if the strangulated bowel slough off, may terminate favourably.

A man, aged twenty-five, suffered from attacks of colicky pain in the abdomen, and gradually lost flesh. One day he became suddenly worse, and on the following day died; nearly four months from the supposed date of the commencement of his attack. About six inches of ileum, caecum, and the first part of the ascending colon, had passed into the large bowel beyond. There was no appearance of gangrene in any part. The colon contained much mucus and coagulated blood. The mucous membrane was turgid and purple. The convoluted layers of the intestine were firmly adherent to each other by dense bands of false membrane.†

Recovery after the passage of the strangulated portion of the bowel through the rectum is the only fortunate result that we can anticipate in a case of chronic intussusception, and fortunately is a result which occasionally takes place; nor does the loss of the bowel appear to be attended by any subsequent ill consequences, *i.e.* contraction or stricture of the gut, so far as we have been able to investigate the subject.

A boy, aged five, was taken ill four months previously to the time at which a mass of bowel was passed. At first he suffered from simple fever, but afterwards experienced much distress. He complained chiefly of pain in the region of the bladder; but there was no tenderness. The urine was sometimes retained for a long time. The bowels were very much confined, sometimes for twenty hours. There was occasional vomiting. He was never known to pass blood. *Eight inches of the ileum, the caecum with its appendix, and about four inches of the colon, were passed by stool.* The tissues were almost black, and sensitive. Diarrhoea preceded the discharge of the separated bowel. At the end of six weeks from this event the little patient seemed quite well, and the bowels acted regularly.

* *Path. Trans.* vol. i. p. 77.

† *Ibid.* vol. vii. p. 193.

and following day enemata were returned without any twelfth day some feculent matter was brought away ; and after a dose of castor-oil. On the nineteenth day he had last consisted of **bright florid blood**, mingled with many intestine. The two previous motions were said to have same character, and to have contained many 'skins.' At weeks the patient returned to his usual occupation.*

Peritonitis may be the result of perforation during the separation of the sloughing process death is generally rapid, and attended by much suffering.

The average number of deaths from intussusception at all ages, must be very small. In five years there were 10 cases in the Hospital for Sick Children, in-patients 46,298—total 48,218. Only one death of the intestine occurred among this number.

It will appear that, although invagination with the causes of acute strangulation, many of which these symptoms are not very acute, and may even terminate in recovery, though at a sacrifice of the bowel.

Invagination may be suspected when there is pain, sudden in its attack, and confined in its seat, and urgent desire to stool, without any relief, of blood and mucus ; and sometimes a swelling in the shape, may be felt in the ileo-cæcal region ; and vomiting ; but all evidence of swelling is absent. No case is known to have terminated fatally.

tion of calomel when peritonitis sets in, and leeches to the in pain. The proposal to open the abdomen should not be retained. The cases recorded will justify this opinion. And general measures give no relief, we have only the one hope that the involuted portion may be thrown off by slough, thus the obstruction be removed.

Distension of the lower bowel with fluid or air has been recommended, and even stated to have been successfully applied in cases of urgency; but however promising this kind of treatment may appear, it must be recollected that, even when the abdomen has been opened, it has been difficult, and in some cases impossible, to reduce the involution without tearing the bowel. A case is, however, recorded by Dr. Baldwin of Tallahassee, Florida, of recovery in supposed invagination, effected by injecting a very large quantity of fluid into the rectum.*

We have now to consider the more chronic conditions productive of obstruction.

Habitual constipation may become accidentally prolonged to such an extent as to cause anxiety; or the accumulation of fecal matter may be mistaken for a tumour, so circumscribed occasionally is the collection in the large intestine. The time which may elapse while the bowels are entirely closed, and the amount of feces which may be collected therein, would surpass belief, were it not that the experience of those now living confirms the statements made by writers on this subject.

A child, aged seven, on recovering from an attack of fever, was affected with torpor of the bowels. This gradually increased to such an extent, that six years treatment failed to have any effect. When admitted into the Hospital, under the care of Mr. Gay, *nothing whatever had passed from the bowels for three months.* The health had not suffered; the appetite was good. The abdomen was greatly enlarged, and was forty-nine inches in girth; and there was considerable inconvenience to respiration. There was considerable prominence on the right side, as if the colon and sigmoid flexure were chiefly loaded.

A catheter was passed into the rectum, and after dilating it, an enema tube was passed high up. The contents of the bowel were washed out by a stream of warm water kept constantly playing upon them, for half an hour at a time. A large quantity of hard black feces, like cinders, was brought away. These enemas repeated several times reduced the girth of the abdomen to twenty-four inches.†

The surgeon is usually consulted at the latter period of

* *American Journal of Medical Sciences*, Oct. 1852, p. 568.

† *Path. Trans.* vol. v. p. 174.

obstinate constipation, which has at last arrived at complete obstruction of the intestine; and as all power of propulsion is then lost, the contents have to be removed mechanically. This can only be done by the aid of repeated warm-water enemata; or by means of some kind of scoop or short lever, with which the lowest masses of hardened fæces should be carefully dislodged.

Purgatives in such cases are not of much use, until the load is somewhat reduced in the lower bowel.

The following very instructive case is mentioned, in a letter to Dr. Burne, by the late Dr. R. Williams, Physician to St. Thomas's Hospital. A lady, aged thirty-five, was attacked with severe gastrodynia, and most obstinate constipation. Paroxysms of the pain recurred at short intervals, gradually became more frequent, and at length quotidian. Each meal was followed by pain as severe and continued, that she limited herself to dry biscuit and brandy-and-water; and this was commonly rejected. Nothing gave relief but opium, of which, for months, she took not less than sixty grains daily. Her bowels, independent of the astringent effects of opium, were always greatly constipated; and no quantity of salts, oil, or senna in the least moved them; elaterium, croton-oil, or other powerful purgatives, immediately inverted the action of the stomach, and were rejected. Calomel was the only medicine which would act upon the bowels, and this not until her mouth was affected.

From this cause she frequently had no evacuation for *six weeks* together, and in one year, when much weakened by repeated salivation, so that it was necessary to defer the administration of the calomel as long as possible, she had only one evacuation every *three months*, or *four in the year*.

At the times her bowels acted she suffered immensely, her pains being more severe, if possible, than those of labour. The fæcal matter when passed was enormous in quantity, healthy in colour, and was formed into large round lumps or scybala, each certainly not less than a large foetal head, and so numerous as often to fill a common-sized pail.*

We may, perhaps, have travelled somewhat from the path of our especial calling, in introducing the latter cases, illustrative of chronic obstruction of the intestine. But the accumulation of large quantities of fæces is often productive of a strange and anomalous train of symptoms; often such as indicate structural obstructions; or simulate morbid growths in the abdominal cavity. Experience, discrimination, and careful examination are often requisite to insure a correct diagnosis as to the true nature of these collections: and without extreme caution, the surgeon may be baffled in his attempts to unravel the condition attendant on a loaded intestine, accompanied by obstinate constipation. With a gut greatly distended by fæcal matter, and

* Burne, *On Habitual Constipation*, p. 28.

In the symptoms of organic obstruction, it will frequently be seen that the bowels act freely, and yet without relief, or reduction of the size of the abdomen. It is but the overflow of the sluice; the head-stream is dammed up; the canal is loaded to overflowing.

Obstruction, the result of injury, we need but allude to. The conditions and treatment have been already described; see Vol. II. p. 626.

Constipation, occasionally amounting to obstruction of many days' standing, or even of a fatal character, is apt to occur in cases of peritonitis, attendant on tubercular deposit in the subserous cellular tissue, or the result of other causes; a condition often accompanied by ulceration of adjacent and adjacent surfaces of the intestine.* A direct communication is taken place between two or more opposed surfaces of bowel: an abscess is found within the abdomen, communicating through ulcerated openings with several portions of the intestinal canal; there is great irritability of the stomach, and rejection of all food; an unyielding abdominal wall, an irregularity of its surface; general but not exquisite tenderness; and constipation of, perhaps, some days' standing. Such symptoms might, at first sight, convey the idea that obstruction was due to some of the causes previously enumerated. But in such cases as we have now under consideration, there are usually general indications of tubercular cachexia long before symptoms of obstruction set in. The condition of the bowels is, however, as a rule, extremely uncertain in these cases: occasionally confined, but more frequently relaxed. With adhesion of peritoneal surfaces, and ulceration through the coats of the intestines, obstruction is rare; and is rather the exception, unless the calibre of the bowel is contracted by the pressure of abscess, or diminished by false membrane, or by a media, stretched across the tube. Looseness of bowels is a usual condition in the cases just considered.

4. Stricture of the intestine is the most common cause of obstruction.

Strictures of the intestine occur under various conditions, independent of cancerous deposit in the walls of the gut. The chief causes of simple stricture are due to the action of caustic

* An interesting case is recorded by Dr. Bristowe, *Path. Trans.* vol. viii. 200.

or some escharotic accidentally or intentionally swallows of some other form of ulcer which, in healing, has α the diameter of the gut.

Stricture of the duodenum is uncommon. We have a few instances of it. In the Museum of the College of is a specimen of a stricture of the duodenum; the str half an inch in length. It appears simple in its chara has no history attached to it.*

The following case well illustrates the effects of an escharotic substance introduced into the stomach. A woman, aged forty, accidentally swallowed a wine-glassful of 'Burnett's solution' (chloride of zinc), and was seized with violent pain and vomiting; but under treatment recovered immediate effects of the poison. A month afterwards, pain and vomiting returned; and in about two months after she was admitted into St. M. hospital, under the care of Dr. Markham. All food taken was rejected; died about four weeks after her admission, and some three months after the fluid. In the pyloric portion of the stomach, about an inch and a half the valve, the stomach was so contracted as only to permit the passage of a bougie. The constriction was clearly the result of a cicatrix.†

A young lady had accidentally some 'Burnett's fluid' given her in a large dose of medicine. Acute pain and vomiting were the immediate result; many months life was despaired of; by degrees the stomach became tolerant of food, and was able to retain small quantities of fluid nourishment at frequent intervals. It is now some five years since the accident; the greatest care is requisite on the part of the patient as regards her diet; she can only be taken in a fluid form, in small quantities; and is general at intervals of four hours. Solid meat, or indiscretion in quantity, once produces vomiting, and the contents are rejected. The health in other respect good. No doubt some contraction has taken place from

cicatrising. At the commencement of the ileum a cicatrix had formed, and reduced the calibre of the intestine to such a degree that the point of the little finger could barely be inserted. Below, the bowel was contracted; above, it was considerably dilated. One slough, opening through the intestine, allowed escape of feculent matter into the peritoneal cavity. The intestinal wall at the strictured part was an inch thick. Throughout the large intestine were numerous cicatrising ulcers. The history of the patient did not throw any light on the case. For some years she had been liable to sudden attacks of pain and constipation.*

A man, aged fifty-four, had been poorly for twelve months. At the commencement of that period he had suffered from an attack of fever, since which he complained of pinching pain in the bowels; the abdomen became swollen; and he lost flesh. He could take very little food; was constantly sick, but the bowels acted once or twice daily. A constriction of the ileum was found at the junction of the upper with the middle third, dependent on great thickening of the walls of the bowel. The stricture was an inch long, and barely admitted the tip of the finger. Immediately above this, the bowel formed a pouch, the parietes of which were inflamed almost to perforation. In this pouch were found thirty-three plum-stones, and sixteen cherry-stones, all perfectly black; and half a dozen recently-swallowed orange-pips. There was an oblique inguinal hernial sac, into which was evident, from the position of the parts, that the strictured portion of the intestine had been in the habit of passing.†

These two cases are interesting when we consider the causes of the strictures described. In the former, though there was extensive ulceration of the bowel above and below the stricture, the symptoms, and the nature of the latter, incline us to believe that contraction of the bowel from cicatrix was the primary disease. In the latter case it is most probable that inflammation, produced by the bowel frequently slipping into the old hernial sac, was the original cause of the stricture, quite independent of the attack of fever, or any ulceration of bowel following that attack.

In allusion to this consideration, viz. whether the ulcerations of the intestine in fever are subject to contraction, and are thus productive of stricture, Rokitansky observes that the cicatrices have occasionally been observed thirty years after the fever had occurred; and he adds, 'it is singular and characteristic of this ulcer and its cicatrix, that they never in any way give rise to a diminution of the calibre of the intestine.'‡

Of the healing of ulceration, the result of tubercular inflammation, he further observes: 'In consequence of the contraction of the ulcer, a cicatrix forms on the surface of the intestine,

* *Path. Trans.* vol. iv. p. 153.

† *Ibid.* vol. x. p. 154.

‡ *Path. Anat.* vol. ii. p. 73.

substance, the approach of the edges is ren the deeper layers of the tissue, which takes mucous membrane, is frequently condensed i which form corded projections into the intest lace with one another, and not unfrequently calibre of the intestine, in the shape of valvul thus giving rise to a stricture in the colon form. This mode of regeneration is more closely resembles that following the destru phageal mucous membrane by mineral acids.

A case of stricture of the sigmoid flexure, near its jun is reported by Dr. Peacock.† The contraction was appare of an old ulcer. There was much thickening, puckering internal tunics. There was no appearance of any canc diately above the stricture the cavity of the intestine v ulcerated aperture of sufficient size to admit the passage on the anterior and inner side; a portion of fecal matter cavity of the pelvis.

5. Obstruction may also be the result of st on cancerous deposit in or about the walls of t frequently, it is occasioned by tumours or cyst bowel: such as fibrous tumours of the uter hydatid growths, or other masses, which ori croach upon, the abdominal cavity. The ultim of these causes of obstruction are equally fu ment must of course vary according to the life may be prolonged many months under

Cancer of the intestine occurs most frequently in the large intestine. When the small intestine is affected, usually the disease is first attacked it secondarily, having originated in some cancerous viscus. The nature of the deposit varies much in different cases: true scirrhus, medullary deposit, or villous growth, will each be met with; but the latter less frequently than the former, and true scirrhus less often than the medullary deposit of cancer; but, as Rokitansky justly observes, they 'may be combined with one another, from their first origin, or consequently.'

Of the colon,' he continues, 'is almost exclusively the seat of cancerous degeneration; but there is a gradation in the probability of its different sections to the affection. The rectum is most frequently attacked; in second order, the sigmoid flexure; the remaining portion of the colon but rarely. . . . Carcinoma occurs as a primary affection of the intestine in three ways. Firstly, in the mucous membrane, as carcinomatous thickening of the erectile tissue, into which the former has previously converted—fungus; secondly, more frequently in the submucous cellular tissue, as round nodulated accumulations; thirdly, most commonly as an annular deposit of the cancerous tissue in the submucous cellular layer.* And he adds, that 'cancerous stricture of the intestine is the most common variety of stricture that results from alterations in the internal coats, and at the same time the one that advances to the highest degree.'

The following case illustrates the history of such a stricture:

A man, aged fifty-eight, was admitted, under the care of Mr. Birkett, into the Hospital, with constipation, which had existed for a week. Twelve days previously the patient had been attacked with diarrhoea, sickness, and pain over the whole of the abdomen. The diarrhoea ceased, but the sickness and pain recurred at intervals. He lost flesh. The bowels were sometimes costive, at others relaxed. He became gradually worse in every respect, and then sought admission into the hospital. At this time he had constant nausea; frequent vomiting every two or three hours, and always refusing food. The abdomen was tense, but not tympanitic. There was swelling in the right inguinal canal, but no hernia could be detected. Two days after his admission, as the symptoms were not relieved, Mr. Birkett cut open the old hernial sac; but no intestine was found in it. The symptoms became gradually more distressing, and the patient died about four days after admission. The right flexure of the colon was constricted, as if a string

* *Path. Anat.* vol. ii. p. 97.

by the deposit of medullary cancer external to the :
The patient died of acute peritonitis.†

Advanced age generally appropriates to intestinal obstructions which take their deposits. From the pylorus to the ileo-cæcum to the rectum, there is no portion attacks. Occasionally they are found to : somewhat oftener the jejunum and ileum lower portions of the large intestine. If : found in youth, the rectum is almost inva mischief.

The pressure of a fibrous tumour has b obstruction. An ovarian cyst may occa bowels.

A woman was admitted into the Middlesex Hosp Moore, suffering from constipation, and at the same large ovarian cyst. The constipation did not yield t of the cyst was withdrawn, when entire relief was a obstruction.

Spasmodic strictures, as they are term on disordered or costive bowels; and need

Before entering upon the consideration obstructions of the intestines, it appear attempt some kind of estimate of the peri different kinds of obstruction may occur; rally the respective symptoms demanded

rather general, conclusions: that obstructions occur in the following order of frequency, from the various causes mentioned, in youth, middle age, and old age: viz. in youth, from internal strangulation by bands of lymph, omentum, or adherent diverticula; from adhesions of various coils of bowel to each other; from intussusception; from foreign bodies taken by the mouth; from cancer, rarely, and hitherto found in the rectum only:—in middle life, from twists of large or small intestine; from concretions, intestinal concretions, and foreign bodies; from intussusception; from simple stricture; from mesenteric hernia; from internal strangulation by bands, &c.; from peritonitis, resulting in abscess; from simple constipation; from cancer:—in advanced life, from cancer; from thickened intestine; from the result of old reducible hernia; from intussusception; from simple stricture; and lastly, from internal strangulation.

The following table indicates the relative frequency of different causes of obstruction irrespective of age. The results are taken from Mr. Hinton's valuable communication on intestinal obstructions.*

In 135 cases, the following were the causes of obstruction, in order of their frequency:

and uterus	1	In first col.	26
ure of ileum	1	Doubtful	8
of small intestine	2	Peritoneal adhesions, tubercles, &c.	9
al hernia:		Stricture of sigmoid flexure	10
anal, high up	1	Ditto colon	11
phragmatic	2	Ditto rectum	11
so-colic	2	Intussusception	24
curator	3	By bands, adherent diverticula, uterine appendages, &c.	36
accumulations	3		
of sigmoid flexure	4		
ions, calculi, foreign bodies	7		
	—	Total	135
	26		

From whatever cause obstruction has its origin, the symptoms are commonly so similar, that a correct diagnosis as to the seat of the block is not to be expected. We may, however, by careful inquiry and examination, approach near it. Whether occasioned by intestine entangled in a loop, or by red omentum; by a cicatrix, or simple stricture; by a foreign body; or by a twist contracting the gut—the symptoms do not vary much in character, though they will in degree

* *Association Medical Journal*, 1853, p. 431.

and in the period of their commencement and sequence. Constipation and pain, vomiting, and tension of the abdomen, each and all may be present; and should be present, to constitute the conditions of absolute obstruction.

But we are of opinion that intussusception should be excluded entirely from the above general observations. The usually early and marked symptoms of invagination are sudden, often acute pain, as severe as it is sudden; constant desire to pass motions, and perpetual attempts made to this purpose without any relief. Most commonly there is discharge of blood mixed with mucus, or pure blood in a fluid or clotted form. From no other cause of obstruction have we seen such perpetual desire to stool in the early part of the attack; nor, with such attempts, the passage of blood and mucus. Sometimes a small quantity of feculent matter is mixed with the mucus but often only enough to colour the discharge. Intussusception, again, is the form of obstruction which occurs earliest in life. We have known it occur in an infant at the breast between five and six months. Mr. Hinton mentions a fatal case at the age of two and a half months. In men affected with intussusception, similar symptoms are observed. The disease is more rare in old age; the oldest of which we have a record was in the fifty-sixth year.

The symptoms attendant on obstruction from other causes vary according to the completeness of the strangulation, or block. In all internal strangulations, by bands, &c., the symptoms are usually very acute; the pain is sudden, sharp, even agonising occasionally; vomiting sets in early, and is usually incessant; the distension may not be as great as in the more slowly operating causes of obstruction, since little food can be taken; but still the small intestine becomes much loaded; there is generally great tenderness on pressure, for usually peritonitis is not long absent; there is early evidence of grave constitutional damage, for the portion of intestine, tightly blocked down or encircled at the strictured part, soon thickens, inflames, or may blacken and mortify.

In twists of the large bowel; simple stricture, or cancerous contractions; foreign bodies; loaded intestine; abscess of the abdomen; and other chronic causes of impediment, the symptoms may vary to some extent, and in most of these instances may present themselves gradually.

Constipation, more or less difficulty in defæcation, small thin or flattened motions, constitute the primary and leading symptoms of commencing stricture. The belly becomes distended by degrees, as the obstruction increases; pain arises upon the increase of contents; often transient periods of hæmorrhœa occur. Vomiting is frequently absent till quite a late stage. Of course something depends on the position of the obstruction. If it occurs in the upper portion of the intestinal canal, vomiting occurs early in the case, and distension of the abdomen is less marked. If in the lower bowel, sickness is late, but great accumulation may occur in the large gut, the peritoneal coat be in consequence ruptured before death, and no timely relief be afforded.

The examination of the general surface of the abdomen does not often convey much, if any, idea of the exact seat of the obstruction, unless it be caused by a tumour in that cavity. Occasionally, with a twist of the sigmoid flexure, the chief swelling has been observed on the left of the umbilicus; but little dependence can be placed on such evidence for a satisfactory solution of the exact cause or seat of the obstruction. The examination of the rectum often conveys more satisfactory information. The obstruction may be often here detected by the finger, or bougie; water thrown up may be immediately returned, or a small quantity may be retained, sufficient to indicate that a stricture, not to be detected by the finger, exists higher up.

A long tube passed up the rectum may frequently detect the actual obstruction; but it is often apt to deceive us as to the seat of the stricture. It is, however, always desirable, in all cases of obstruction, to attempt its introduction as far as possible without any force, and to inject warm water frequently and plentifully, if any fæcal discharge accompanies the return of the injected fluid.

The emaciation which usually accompanies cancerous deposits in other parts, the existence of fæcal fistulæ in the abdominal wall, the bladder, or the vagina, are symptoms of frequent occurrence in cancerous stricture. Hepatic pains of various years, or months, the symptoms of former mischief about the gall-bladder, point towards the supposition that a biliary calculus may have ulcerated into the duodenum, and produced the obstruction by blocking up the canal. Con-

often so early in the scene, even in a healthy, that the medical attendant has little or the patient to acquiesce in, measures which becomes, therefore, a matter of grave importance for our guidance under such perplexing

The subject is not a new one, and yet little taken of it in systematic works on Surgery. The writings of Callisen, Littre, Amussat, Phillips, Hawkins, Curling, and others, have of late attracted attention to its consideration; whilst the results from our hospitals, and the results of modern practice have to a great extent sanctioned surgical formation of artificial anus, when other means failed.

In the management of this class of cases, should be scrupulously avoided. This is a precaution to most rigidly at the commencement of the disease, if there be one golden rule to guide us in all cases of obstruction, it is this: that opium should be given by the mouth, and aperients only by the rectum. There are a few exceptions; but experience has taught us that ever this rule has been departed from, subsequent regret has arisen. In the early and moderate cases, small and repeated doses of calomel may be given, or opium if peritonitis be suspected, or if the bowels are not moved. Leeches and hot fomentations may be applied to the abdomen, especially if great pain and acute tenderness be present.

ata injected in the usual manner were retained, and never
ed. Subsequent to death, it was found that a long tube
be inserted beyond the twist into the dilated bowel; but
sequence of a flap-like entrance at the commencement of
atter, the fluid injected into it could not be returned.
memata used in such cases should be warm water, their
anical action being of far greater importance than their
e properties.

the management of every case of supposed intestinal
action, every variety of solid food should be rigidly pro-
ed. Life should be entirely sustained by fluid nourish-

men situated low down in the sigmoid flexure, or upper
an of the rectum, the obstruction will be more readily
ed, as the accumulation in the larger bowel increases.
ach conditions it is prudent, if practicable, to overcome
mediate effects of obstruction, by passing a small gum
ter through the stricture, and injecting warm water into the
; or by introducing the finger gently and slowly into the
above, to dilate slightly the contracted part. In the latter
apt much care is necessary. The passage of the finger
id not be hurriedly or hastily effected. The stricture in
e parts is commonly the result of cancer; the parts are
quently thickened, often ulcerated, and brittle under
pulation. The duration of the disease has allowed it,
ably, to implicate the whole structure of the surrounding
and has often involved the peritonæum in the diseased
a. As the bowel, from the load above, is pressed down far
e than its natural relations, the peritoneal reflexion in
sequence comes much nearer the external aperture; a
len dilatation of the diseased bowel, a rapid thrust of the
er through the obstruction, can only be effected at the ex-
e of some laceration of the diseased tissues.

man was admitted into St. George's Hospital, under the care of Dr. Nairne,
constipation of some days' standing. Treatment having failed to procure
tion of the bowels, the author was requested to make an examination of
state of the rectum. This was found pushed down so that the orifice was
level with the lower edge of the nates. A stricture was readily felt, within
at distance of the anus, pressed down, however, by an immense collection
cal matter. A small gum catheter was introduced without difficulty, and
ed the passage of some fluid fæces and flatus, with great velocity. The
finger was subsequently passed very slowly through a rather resistant con-
tion of the bowel, the result of large deposit around it. The man shortly

We have now to discuss the condition justifiable to operate in obstruction of the bowels, when other efforts have failed to procure relief.

The operations for the relief of obstruction are of two kinds. In one (Littre's), the abdomen is opened, and the peritonæum is cut. In the other (Amussat's), the peritonæum is opened in the loin, external to the peritoneum.

The duration of constipation is not *the* indication for such interference. Death takes place in constipation, whether perfect strangulation of the bowel has occurred, or a band of false membrane, or a hernia, or a slipped through some contracted aperture of the mesentery. In such a case, constipation is a most important symptom. It is not *the* symptom which indicates the absolute necessity of active interference, but the delay. In common or cancerous stricture, constipation is not often complete, and the accumulation of feces may be great, and months may elapse without a motion, and death may result from such an occurrence. Constipation is alone *the* symptom which warns the surgeon to operate. In constipation following peritonitis, the bowels may remain locked up for a month, or even longer, and finally yield to treatment or time, and death may result. Constipation in the latter instances cannot

never so small a quantity, the means to encourage them must be persevered in. No immediate operative measures should be sanctioned.

It is but rational to suppose, that in invagination of the intestine, operative interference holds out no prospect of relief, immediate or remote. The pathological conditions already discussed must satisfy the reader that such a conclusion is founded on substantial reasons. The common occurrence of invagination in early infancy; the frequent, often severe, peritoneal mischief which accompanies many cases; and the prospect of slough and separation of the invaginated portion, are the chief facts which strengthen the above conclusions. The following results are interesting: In *twenty-four cases* of intussusception, occurring at various periods of life, from two months and a half to fifty years of age, there died without any relief *thirteen cases*; some as early as three days, some as late as forty days, from the commencement of the symptoms. In the *other eleven cases*, the invaginated pieces of intestine sloughed and were passed by stool. Of these *eleven cases*, *two* died soon after the passage of the slough. *Nine cases* entirely recovered. The patients who recovered were ill for periods varying from five to thirty days respectively; their ages varied from six to fifty years.*

The injection of air or water into the rectum has been proposed and adopted in supposed cases of intussusception; nor can we say that such treatment has not been attended with advantage; for in well-authenticated cases with symptoms of invagination, this treatment appears to have done occasional good. The proof, however, is wanting, that actual invagination existed in such cases. This form of treatment cannot be supposed to possess any advantageous influence in the reduction of invagination of any extent or of some duration. The too indiscriminate employment of air, or of water, for the purpose of distending the bowel, with a view to reduce the prolapsus—to push it up, as it were—may act most injuriously; the peritonæum covering the intestine may be ruptured in consequence. But applied with discrimination, the use of warm water enemata alone is a measure which frequently will afford relief, tempo-

* These particulars are taken from cases published by Mr. Hinton of Blaina, in an able article on 'Intestinal Obstructions,' in the *Association Med. Journal*, 1853, p. 451.

rary, if not permanent, in many conditions of constipation, and should be always had recourse to; not with an expectation that it will unravel an invaginated piece of gut, but that, if the diagnosis be incorrect and constipation depend on other and remediable causes, benefit may accrue, and perhaps recovery be insured thereby.

In the treatment of all other forms of obstruction dependent on structural derangement or organic disease, medicines are of no avail. If the obstructing cause be not removed, or if the condition of the intestine be not relieved by some effective measures, the patient will surely die, and in a comparatively short time.

‘Internal strangulation,’ observes Rokitsansky, ‘when diagnosed, most imperatively requires an operative proceeding, for the purpose of disentangling and arranging the intestines, and for the division of the strangulating structures.’* Mr. Phillips after careful study of the subject, arrived at the conclusion ‘that the interference by surgical operation is justifiable when three or four days have passed without any relief from ordinary means (provided the constipation be complete, and vomiting of fæcal matter continue), because it affords a greater chance for the preservation of life than ordinary means.’†

Mr. Curling is of opinion ‘that the success which has attended the operation of lumbar colotomy, in persons weakened by organic disease and want of nourishment, shows that the operation is not so formidable and dangerous as is commonly supposed.’‡

The alternative of cutting into the abdomen for the relief of obstruction must always be regarded as a choice of evils. This is the intrinsic worth of the operation. Nothing better can be said of it. On the one hand, a patient is suffering from the effects of internal displacement of intestine, which will surely destroy life in a very few days; or from a disease which will, by causing obstruction, certainly prove fatal in a few weeks, unless the displacement be set right, or the effects of the disease be obviated by an artificial outlet above the stricture.

It becomes manifest, when the anatomy of obstructions is fully considered, that one of two operations must be adopted, the relief is to be attained and life preserved; viz. cutting into the

* *Path. Anat.* vol. ii. p. 54.

† *Med.-Chir. Trans.* vol. xxxi. p. 35.

‡ *British Med. Journal*, 1863, p. 111.

peritoneal sac, and endeavouring to remove the cause of strangulation; or cutting into the colon in either lumbar region without entering the cavity of the peritonæum.

It is next a matter of importance to consider under what conditions either operation should be undertaken; in what manner each should be performed; and what should be the subsequent treatment of patients subject to such operations.

The operation of opening the abdomen through the peritonæum is requisite for the relief of internal strangulations of the small intestine, or obstructions caused by foreign bodies or calculi, or supposed strictures of the small intestine. That of opening the colon in the lumbar region can only be useful when obstruction occurs in the large bowel.

In internal strangulation dependent on any of the causes already referred to, an opening into the peritoneal sac should enable the surgeon to ascertain the cause of obstruction; or in many cases to divide the constricting band, or dislodge the intestine from the gripe of a fissure in the omentum.

The operation of opening the peritonæum will prove less successful under the following conditions: 1st, in cases of stricture of any portion of the small intestine; 2nd, in cases of obstruction from foreign bodies or calculi lodged in the intestine; 3rd, in cases in which peritonitis has set in from too long neglected strangulation.

The last condition alluded to indicates the importance of early interference. The formation of an artificial anus of the small intestine offers the only prospect of prolonging life. In the removal of a foreign body, provided the operation be recovered from, it may reasonably be anticipated that the artificial anus will close. In a case in which a portion of the ileum was strangulated by small intestine coiling round it, the peritoneal cavity was opened, the strictured part was relieved, and the patient recovered.*

The conditions which justify this operation are, a sudden attack of pain, followed by desire to be relieved, but with no action of the bowels, and persistent vomiting; and if little or no tenderness on pressure should exist, constipation must be complete, and must have exceeded a period of twenty-four hours, in company with all the above symptoms. It must be borne in mind that the amount and character of the pain are very im-

* *Prov. Med. and Surg. Journal*, June 11, 1851.

opening the peritonæum.

In strictures of the rectum, sigmoid and colon, the operation for relief is confined to that from whence the intestine may be opened, by peritonæum. The advisability of this must be fully weighed before it is necessary, for there is usually less urgency of symptoms require it than in strictures higher up; of a week or two may be permitted to the patient be carefully watched, before operation is requisite.

The conditions which demand it are, unyielding constipation, great distension and commencing irritability of the stomach, or

The conditions which forbid us to attempt it therefore should withhold surgical interference, feeble and quick pulse, cold general tenderness of the abdomen, suppurative conditions enumerated above.

The operation of opening the peritonæum in cases of strangulated or obstructed intestine, and the formation of an artificial anus in the walls of the abdomen, was first suggested and adopted by M. Littre, in 1776, though the abdomen had been opened by him in 1776, and an artificial anus formed in the

que l'on ne refermerait jamais, et qui ferait la fonction

This operation, originally proposed for the relief of imperforate anus, is that to which we can alone have recourse to in any form of internal strangulation, or obstruction dependent on mischief confined to the small intestine.

It is not desirable to adopt this operation in obstructions confined to the large intestine. The operation which insures the opening of the ascending or descending colon, commonly called the operation of Amussat, is alone to be adopted in these cases.

The operation for opening the peritoneal sac, to explore its contents, requires but little to be said respecting the method of proceeding. The surgeon having decided upon the operation, should have the patient placed on a conveniently high bed or table. The bladder should be emptied by a catheter. The incision should commence an inch below the navel, and be extended down about four inches, or sufficiently low to allow of the obstruction being thoroughly investigated.

The first incision should extend down to the linea alba, which should be next carefully cut through at one point, that the peritonæum may come into view. A broad director should now be passed under the conjoined aponeurosis, and the latter then divided, so that the peritonæum be fully exposed. A small opening must be carefully made through the latter, sufficient to allow the introduction of the first and second fingers. These being passed into the sac of the peritonæum, to serve the purpose of a director, and upon them a blunt-pointed bistoury can safely be introduced, in order to extend the incision in the peritonæum, either upwards or towards the sides, as necessity or convenience direct.

When the abdomen is laid open, search must be made, slowly and carefully, for the constricted piece of intestine. The dilated bowel may be followed down, if there is difficulty in doing so, until arriving at the point of obstruction, until the cause of mischief is reached. It is impossible here to enumerate all the difficulties which may now be encountered; it must be sufficient to point out that bands of lymph or adhesions should be carefully separated, or divided; or the strangulated intestine withdrawn from any loop or fissure in which it may have become

* *Hist. de l'Acad. des Sciences*, 1710, p. 36.

should at once be closed, the edges held by silver-wire sutures. The subsequent operation should consist of perfect rest and applications to the abdomen; and opium mouth in one or two grain doses every four hours if the patient does not sleep. Sleep should administer a dose due: for sleep alone, it indicates the repetition of opium. Its continuation until the patient is quite restored and the motions are natural and regular administration of opium to this period may hinder a natural and healthy action of the bowels, and are most instrumental towards recovery.

When an obstruction is known to be situated in the sigmoid flexure, the descending colon, or caecum, must be opened; when suspected in the transverse colon, the ascending colon, or caecum, must be opened. The exact seat of stricture may occasionally be ascertained by the introduction of a finger or sigmoid flexure, it is often made manifest by the swelling, or hardness of the part. The stricture is often external, when seated in other portions of the large gut, it often becomes mere speculation at which the obstruction is seated.

The chronic nature of the symptoms, the constipation of faeces, point to the large gut as the seat of the obstruction. Whether the obstruction lie in the transverse

M. Amussat is alone due the credit of having revived colotomy—an operation not only fallen into disuse, but considered as dangerous and impracticable by some very able writers.

The virtue of this operation is, that it enables the surgeon to find the intestine and open it, where it is uncovered by peritoneum, and where it lies in front of the quadratus lumborum, and merely separated from that muscle by cellular tissue. The situation of the external incision, on either side, in the lumbar region, between the last rib above, the crest of the ilium below, a vertical line running from the end of the last rib, the crest of the ilium in front, and the edge of the longissimus behind. The colon in this space is fixed to the abdominal wall by the reflexions of the peritonæum, and lies in contact with the quadratus lumborum. The kidney is situated rather above the seat of the incision. There is no colon here; and if the colon be much distended, the space between the folds of peritonæum will be very narrow. It is at this part that the intestine should be

opened. Amussat has judiciously recommended a transverse one. The advantages are thus fairly stated: 1st, that it makes the operation easier and more certain, and avoids the danger of wounding the lumbar vessels and nerves; 2nd, that it facilitates the opening of the intestine without wounding the peritonæum; and 3rd, it enables us to establish the artificial anus anteriorly.*

The incision should commence in front of the longissimus muscle and be carried forwards to the extent of about six inches. After the integuments having been divided, the muscles are to be cut through until the intestine is exposed. This is a difficult or troublesome proceeding. The intestine, especially if loaded, presents its muscular surface in the bottom of the wound, uncovered by peritonæum. The bowel should be

hooked up by a curved needle; two or more points then be secured by threads to the margins of the wound, the gut opened. Usually a great gush of fluid fæculent matter takes place; ample provision should therefore be made to receive it as it flows out, to prevent the bed becoming greatly

* Amussat's *First Memoir*, p. 241.

Should the patient survive the operation the wound is of next importance. In the inconvenience and even some distress may prolapse of the gut through the artificial and the latter be a free one.

But as the wound in the integuments has to contract as time advances, and the margins become adherent to the former, the openings gradually diminish in proportion to the contraction, and usually cease ultimately when the orifice is fully contracted. Indeed, should the patient's life be found as a rule, that the surgeon's attention is towards maintaining the freedom of the wound sufficiently to insure the escape of the contents. In the first instance the escape of faecal matter from the wound may be controlled by the use of a plug attached to a small shield, on which a piece of sheeting may be placed, and fixed in its position. As the wound continues to contract, it will be necessary to substitute for the ivory plug a piece of sponge, which may be introduced and worn in the opening, and changed morning and evening. Its use has usually control the escape of air and faecal matter.

Sometimes the patient will experience some accumulation of faeces between the strictured

ating pain is often experienced, when the bowels act, the passage of fæcal matter over the ulcerated surface ; has been found that by establishing an artificial opening at the seat of disease, and thus diverting the passage of the stool and leaving the stricture at rest, great comfort has been afforded upon the sufferer.

Darling was the first to advocate this operation under such circumstances, and he has frequently adopted it with great relief to the patient's sufferings while life lasted. He directs previous to the operation that the bowel should be filled with fluid in order to distend it to some extent, as in a distended condition, the bowel is more readily opened and secured than when empty and contracted.

Colmes* has published the particulars of a case in which he opened the descending colon for the relief of a patient in whom there was a fistulous communication between the sigmoid flexure of the colon and the bladder. The patients were immediately relieved, and the patient remained free from the passage of fæcal matter into the bladder for about fifteen months, when fæces appeared in the urine. He shortly after this, died ; and on post-mortem examination it was found that besides the communication between the sigmoid colon and the bladder, there had formed a more recent communication between the bladder and the cæcum.


Darling has either himself performed or recommended the operation of opening the descending colon in seventeen cases of diseased rectum, or obstructed bowels. In ten of these cases the patients survived over periods from two to eighteen months. In a case under the care of the author, in which complete obstruction had existed nearly three weeks, the patient survived the operation over two years and a half. The artificial opening always shown a tendency to close, but has been kept open by the use of a catheter. For some time past a portion of the fæces has passed the natural

The Operation of Tapping ; Paracentesis Abdominis.

When the abdomen becomes inconveniently distended with fluid contained either in the peritoneal sac or in an ovarian cyst, it is desirable to relieve the patient by tapping the cavity in which the fluid is lodged.

Previous to the operation, the patient may be placed upright on the edge of a chair, or recumbent on the side of a bed. A flannel towel, warmed, should be passed round and above the umbilicus, and a second one below ; the ends of these towels should be held behind the patient, to be tightened as the fluid

* *Trans. Med.-Chir. Soc.* vols. xlix. and l.



painful than if the trocar is thrust th
The trocar should be pushed nearly up to
and the stilette then withdrawn, care
canula does not slip out after it. The
freely if serous in character, and the cavit

If the patient become faint during t
finger may be applied to the orifice of t
escape, while some stimulant is adminis
ness passes off. The common trocar, wi
simplest implement to use in this opera
as any ; but of late a tube of india-rubb
the canula, so that the fluid may be ca
foot of the chair or bed which supports t
of the instrument is a matter of fancy ;
simple trocar is as efficacious and as c
modern complicated one.

It sometimes happens, that after a ce
has escaped, the flow suddenly ceases, wh
remaining in the cavity ; this is occasion
tine or some other substance coming in
extremity of the canula ; the obstruction
either moving the position of the canu
blunt-pointed but perforated tube, with
trocars should always be provided.

The large accumulation of fluid in the

mus, or below it, the operation had better be postponed or
L. Mr. Gay exhibited, at the Pathological Society, a
specimen of a portion of the jejunum, which had been punctured
in the operation of tapping, in a case of ascites ; but the patient
died of the accident, and died three months afterwards, appar-
ently from the disease which produced the ascites.* The wound
made by the trocar should be dressed with lint and plaster,
and a bandage applied over it, round the belly. In a few days
the wound usually healed.

GEORGE POLLOCK.

* *Pathological Trans.* vol. ii. p. 203.

AN escape of any viscus from the naturally placed is termed a hernia in this essay are exclusively restricted abdominal viscera. In common language rupture. Our object will therefore be to of hernia developed in the different regions their pathology and treatment.

The whole subject is divided into two parts.

The first embraces general considerations, statistics, pathology, and treatment of the

The second is devoted to an examination of regional varieties of hernia, their anatomy, etiology, diagnosis, prognosis, and the adapted to each kind.

Sex.—Both sexes are afflicted with hernia. Carefully considering the statements of writers on the numerical proportion in which the two sexes are afflicted with hernia, we must admit that we have not been enabled to arrive at any satisfactory conclusion. We however, broadly state the fact, that hernia is more frequently in males than females. Out of 96,886 applicants for trusses at the C

Frequency of hernia at different ages.—The only trustworthy relating to the various ages at which hernial protrusions first developed, have been collected by Mr. Kingdon. In reports of the City of London Truss Society for the years 1860 and 1861, he has tabulated 9,296 cases of inguinal and femoral hernia. All these patients passed under his personal examination, and the respective age of each individual when hernia developed itself was ascertained as accurately as possible by reckoning back to the age at which it was first noticed. Observers have been before this occupied in attempts to ascertain the ages when protrusions most frequently occur; they have uniformly noted the ages of the persons at the time of their appearance before them. Accurate data were, under these conditions, never obtained; that is, if a knowledge of the age at which hernia is developed is to be regarded as a desideratum. Hence M. Malgaigne makes it appear that there are fewer cases of hernia before thirty-five years of age than after. This is a fundamental error, unless national peculiarities

TABLE A.

Showing the ages of the patients when the hernia was first noticed.

	1860.				1861.				Kingdon's Tables—Report of Truss Society, 1861-62. Age at development.
	Inguinal.		Femoral.		Inguinal.		Femoral.		
	M.	F.	M.	F.	M.	F.	M.	F.	
5-10	478	27			563	37			1100
10-15	181	17			192	26			416
15-20	91	23	3	1	112	16	2		248
20-25	115	19	4	2	133	19	3	6	301
25-30	284	27	8	14	291	38	8	15	680
30-35	348	30	18	31	375	33	13	37	885
35-40	381	45	21	58	424	50	20	54	1053
40-45	334	48	19	55	399	37	14	70	976
45-50	320	38	18	59	373	41	16	68	933
50-55	271	31	30	47	268	24	11	49	731
55-60	258	11	18	32	272	10	15	30	646
60-65	167	12	12	26	224	11	11	27	490
65-70	147	6	9	16	151	4	8	15	356
70-75	97	6	10	11	110	3	7	12	256
75-80	59	5	8	2	44	1	1	3	123
80-85	32	1	1	1	54	3	3	7	102
Total	3558	346	174	355	3985	353	132	393	9296

Kingdon's statistics for 1860 and 1861, there were in the first five years 1,409 males, and only 107 females; while in the five years from 25 to 30 of age, the proportion had changed to 846 males against 207 females.

trustworthy data on this subject. The ger have been advanced, in some of which t high as one in eight of the male inhabit dom, and even one in five of the wh district, do not rest on any precise in exaggerated.

Frequency of hernia in relation to age.—] (Table A) show that out of 9,296 appli were under five years of age. This wou to show a much greater prevalence of l during any other period of life. But this when we take into consideration the nt any given population. Thus, if we tak as giving an accurate idea of the numeri at different ages in London, we must tween the number of cases of hernia the total number of cases of hernia, wit number of the population living at that a tion. Proceeding in this manner, we sh is no exact proportion maintained, yet t

* The following statements show the number c different ages, when the age of each patient was the time each one came under his observation; s were carefully ascertained by Mr. Kingdon, to sho

By advances towards the latter end of life. Thus, taking census of 1851, the total population of London of both at all ages was 2,362,236; out of these, 293,562 were 5 years of age: a ratio of 1 to 8·004. Out of the 9,296 from Mr. Kingdon's total, 1,516 were under 5 years of a ratio of 1 to 6·13. Taking the next decade, we find total number of the population between 5 and 10 to be 48, or in a ratio of 1 to 9·69 to the total population; the per of cases of hernia, however, is only 248, or in the ratio to 37·08 to the total number of cases of hernia. Again, it shown above that the absolute number of cases of hernia developed under 35 years of age was greater than that of those developed above that age, the proportion in Mr. Kingdon's table is 61 to 39. But in the census table the proportion of persons alive under 35 to those above that age is still higher, 69 to 31. Therefore hernia is more common relatively to number of the population above the age of 35 than under age. It appears from calculation on these data, that 41 per cent. of cases of hernia are developed under the age of 35, 59 per cent. over it.

The relative proportions of the different kinds of hernia at various ages and in both sexes, together with explanations of their occurrence, based upon anatomical and physiological facts, will be found in other parts of this essay.

The influence of occupations on the formation of hernia.—All persons, in every station of life, are liable to hernia. By far the majority of cases occur among the labouring class, which, standing as it does in all large towns, affords a favourable opportunity to ascertain whether the pursuit of any particular trade or occupation involves a greater liability to hernia than other.

Mr. Kingdon has inquired into this subject with his usual care, and the results of the investigation are printed in the Report of the City of London Truss Society.* He has produced a table in order to compare the twenty-five largest classes of occupation, arranged in the order of their magnitude from the census of 1851, with the numbers of each class who applied to the Society during three successive years. This indicates that the patients who seek relief on account of hernia bear a direct proportion to the numerical magnitude of the classes

* Report, 1861, p. 11.

to which they respectively belong, and not to the severity of the toil.'

The influence of hereditary conformation predisposing to the development of hernia.—A disposition to hernia may be inherited that is, the children of ruptured parents are frequently afflicted in like manner. Mr. Kingdon has taken great pains to ascertain the proportion of cases in which this hereditary influence exists. In the reports of the City of London Truss Society for 1860–61, he has published the result of his inquiries. Both sexes show an equal tendency to be thus influenced. The proportion, by calculation, seems to be about 84 per cent. This hereditary predisposition, paternal, maternal, or on both sides is manifested in the most marked degree with infants under one year; the cases being about 12 per cent. of the whole number in the first twelve months of life.

This fact points to two very important causes which give rise to hernial protrusions at this early age: first, to the arrested efforts of nature in closing the ventral orifice of the vaginal process of the peritonæum, and the obliteration of the sheath; and, secondly, to an abnormal elongation of the mesentery. Those two structural conditions belong to a class of anomalies very likely to be determined by hereditary influences.

Congenital defects of the parietes of the abdomen have been already mentioned as predisposing causes of hernia, and this subject will be further dwelt upon in discussing its various forms. Wounds, inflammatory lesions of the abdominal wall and other predisposing causes may be mentioned. It is a question, whether a portion of small intestine can reach the fundus of the scrotum, unless its mesenteric ligament be of preternatural length. For it is difficult to understand how the upper portion of the small intestine or the cæcum can descend so low as they are sometimes seen, without at the same time their mesenteric folds being elongated. But the fact to be accurately ascertained is, whether the mesentery is abnormally long antecedent to the descent of the hernia; whether, in truth, a morbid elongation of the mesentery be a primary cause of hernia or not. Doubtless this structure becomes lengthened and stretched as the result of repeated or continued descents of the intestine; but we are not cognisant of any facts to prove congenital condition of the kind above alluded to. At first sight, this circumstance regarding the elongation of the mesen-

tery may seem to have very slight practical value. Upon reflection, however, we shall arrive at an opposite conclusion. It has a very important bearing on the question of the radical cure of hernia, as it is termed. For if it be proved that the primary cause of any viscus protruding from the abdominal cavity can be traced to a morbid condition of its peritoneal retaining ligaments, the mere obliteration of the hernial sac can avail but little in producing the intended result of affording permanent immunity from the disease. That which often happens when a truss is used to support one kind of hernia would occur in this case also, namely, the development of another, perhaps even on the opposite side of the body; and thus any operation, even if successful in obliterating the first sac, would be of little value.

Persons in whom a hernial sac exists certainly state, that they are more troubled with the descent of a hernia if they get out of health. This circumstance leads to the inference, that when the tissues generally are weakened and relaxed by indisposition, those which should maintain the viscus in its proper situation, participate in the general morbid condition, and then permit the hernia to escape more readily.

It is equally certain, too, that persons constitutionally of a weak frame of body, whose contractile and fibrous tissues are deficient in tone and power, become more commonly subject to the development of a hernial sac as age advances, than those of an opposite conformation; always, of course, excepting those persons who have a congenitally open vaginal process of the peritoneum.

And further we may state, as the result of actual observation, that in both males and females of middle age the subject of hernia, the heavy abdominal viscera, the solid glands for example, are usually disposed in a much lower situation than their normal one; that, in fact, the abdominal viscera generally are not so firmly held in their proper places by their peritoneal ligaments as when no disposition to hernia is shown.

Between the muscular walls of the abdomen and the contents of that cavity a constant antagonism exists. The gaseous and fluid distension of the intestinal tube exerts an ever variable pressure against the abdominal walls, whilst they, in their turn, sit upon the inflated viscera; thus between these opposing forces a sort of equilibrium is maintained. The balance between them is, however, sometimes destroyed, and the parietes are no longer able to restrain the viscera within their normal limits.

Thus hernia is often associated with a great increase in the bulk of the viscera, from the rapid development of fat in the omentum and mesentery. A persistent laxity of the parietal peritoneum likewise, subsequent to distension of that membrane, occasioned by the gravid uterus, is a morbid condition which renders it liable to pass readily out of the abdomen, under the influence of any pressure from within, through the weak points of the walls, and it must also be reckoned among the predisposing causes of hernia.

The immediate cause of a hernia is certainly, in some cases, to be traced to a sudden and forcible diminution in the capacity of the abdominal cavity, the result of compression or contraction of its walls; in other words, the combined actions of the abdominal muscles, coincident with strained attitudes and postures of the body during the effort of violent muscular exertion generally, give rise to the sudden development of a hernia. We cannot, perhaps, accept as truth the statement of all the ruptured men who attribute their affliction to muscular exertion; but all surgeons who have had much hospital experience will be able to call to mind cases depending upon this cause.

A large proportion of the cases of hernia are undoubtedly of gradual development. In many of these, however, violent muscular contraction doubtless plays an active part. How often we observe hernial tumours in patients afflicted with dysuria. Is not hernia very liable to occur in persons labouring under bronchitis of one form or another? And although it must be admitted that more or less laxity and loss of restraining power in the tissues of these persons exist with regard both to the fibrous walls of the abdomen and the peritoneal ligaments of the viscera, yet this effective agent, muscular contraction, exerts its powers with less restraint and limit, and thus presses the viscera away from their normal situation.

The hernia consists of a part only of any of the abdominal viscera. Such of them as are permitted, by their peritoneal attachments, to change their relative situation within the abdominal cavity with the greatest freedom, most frequently escape or protrude through its walls. Thus portions of the small intestines and omentum form the contents of the hernial sac in the majority of cases, although a part of every abdominal viscus has been occasionally found therein.

Semeiology.—The signs which denote a rupture are more or

of the internal abdominal ring, and often along the track of the region of the abdominal walls termed the inguinal canal, if carefully examined. From a similar appearance tendency to hernia may be detected in the weakly and adult female at the crural aperture. In both cases the tumour becomes much more distinct if the patient maintains the erect posture and forcibly contracts the abdominal muscles. Then, a small swelling or 'lump' is felt, which is not permanent disappears under slight pressure or on assuming the recumbent posture. It reappears when the pressure is removed by standing erect, and it becomes more prominent when the abdominal muscles are put into strong action by coughing or voluntary effort.

In infants and children the tumour produced by a hernia is of considerable size when noticed for the first time; but it diminishes if the swelling be pressed, or the recumbent posture imposed. In youthful adults also a hernia is often noticed suddenly, generally in the inguinal region or scrotum. Happening under the influence of violent muscular exertion or forcible compression of the abdominal walls, is usually attended with more or less pain.*

The structure of the viscus which forms the hernia also modifies the signs and indications: thus, if the protrusion be solid, and the omentum escapes (epiplocele), the hernial swelling will be hard, resisting, and lobulated; if the protrusion be hollow,

they are liable to simulate some special kind of hernia, their differential diagnosis is given in other parts of this essay.

Prognosis.—By ordinary care and precaution, a person the subject of hernia is not in great danger of those accidents affecting it which imperil life. On the contrary, however, by neglecting to employ a bandage to prevent the escape of the hernia, or by making use of an instrument which fails to effect the purpose intended, namely, the complete retention of the hernia within the abdomen, the life of the individual is perpetually in jeopardy from the liability to inflammation or strangulation of the protruded viscus.

In proportion, then, as the palliative measures directed by the surgeon are carefully employed by the patient, the prognosis in any case of reducible hernia may be regarded as favourable, or the converse.

If the hernia be a small reducible enterocele, a truss may be used, which will prevent its descent for months, or even years. If the case be one of irreducible epiplocele, complicated with the occasional descent of small intestine, the prognosis must be regarded as relatively unfavourable, in consequence of the great difficulty in preventing the entrance of the bowel at the orifice of the sac, where it may become strangulated without any premonitory symptoms.

Nomenclature of herniæ.—The varieties of abdominal herniæ are named from the period of life at which the hernial sac is formed; the region in which the protrusion takes place or exists; as well as from the viscus which composes the tumour.

When named from the viscus forming the protrusion, a hernia is styled intestinal when any portion of the alimentary canal escapes. This is also termed an enterocele.

If a portion of omentum protrude, the expression omental hernia, or epiplocele, signifies its nature.

A combination of these two is called an entero-epiplocele.

The terms gastrocele, cystocele, have been applied to protrusions of the stomach or bladder from the abdomen.

The classification of hernia according to the regions in which it occurs is given in the second part of this essay.

The essential parts of a hernial tumour.—When an anatomist dissects a tumour caused by the protrusion or escape of any viscus from the cavity or region in which it is naturally con-

ed, his observation should be especially directed to three principal objects. These are :

The tissues outside the sac, or the ordinary structures of the region in which the hernia is developed ;

The sac ; containing

The hernia itself, or the protruded viscera.

Under these three sections we shall describe the essential elements of a hernial tumour : first, in the condition in which commonly exist, when the only source of trouble depends on the mere tendency to the protrusion of the hernia ; and, secondly, under those morbid conditions which affect certain parts of the sac ; the contents of the sac, the real hernia ; and the tissues outside the sac. Upon the judicious surgical treatment of these morbid conditions the salvation of life depends.

The inciting causes of hernia.—Protrusions of the abdominal viscera are the result of—

1. Wounds or lacerations of the abdominal walls ;
2. The weakening or destruction of the same parts by inflammatory processes ;
3. The existence at birth, and persistence afterwards, of a canal which is a prolongation of the peritonæum ; and,
4. The gradual expulsion of the parietal peritoneal membrane at weak parts of the abdominal walls, forming a pouch or receptacle for portions of the viscera.

1. In this essay I am not required to describe those of the first class, which depend on wounds, such cases having been treated of under the head of INJURIES OF THE ABDOMEN.

Hernia, however, occasionally occurs in the inguinal region, the result of the application of direct violence in its vicinity, without any wound of the integuments.

2. *The weakening or destruction of the abdominal parietes* in the effects of inflammation and its results give rise to the secondary occurrence of hernia. Thus, after the healing of abscesses in those parts, it is not very uncommon to see a hernial tumour developed at the site of the primary disease. It is important, therefore, as a prophylactic measure, to support the abdominal region by means of a suitable bandage during the healing of such abscesses, and for some time after cicatrisation.

3. *Congenital patency of the vaginal process of the peritonæum.*—

We must now consider that congenital condition of the peritonæum which allows a portion of the abdominal viscera to escape from its natural cavity and occupy an abnormal position in its immediate vicinity.

Towards the close of the last century, and at the commencement of this one, anatomists were very much interested in observing the changes in the situation of the testicles during the period of foetal life. Accurate observation established the fact, that the development of the testicles commenced in the lumbar regions of the foetus in utero, and that when these organs had reached a certain stage of perfection, they migrated from their primary locality, and, pursuing a course towards the pelvis, they at last reached the scrotum, the final point of their destination. In this progress, termed the 'descent of the testicles,' and whilst they are within the abdomen, they are placed behind the peritonæum, and partially invested by it. A prolongation from this serous membrane accompanies them into the scrotum, and receives the name of the vaginal process of the peritonæum.

Thus wrote Wrisberg in the year 1800: '*Testis semel in scrotum delapsus pluribus cingitur velamentis. Tria eam membranarum genera, quæ id præstant, Tunica vaginalis communis pro teste, epididymide et funiculo spermatico simul; Tunica vaginalis propria vasorum spermaticorum; et Tunica vaginalis similiter testi et epididymidi propria; cuilibet in arte anatomica tironi nunc cognitum est.*' *

A direct and uninterrupted communication exists between the cavity of the peritonæum and the interior of this sheath; so perfect, indeed, that the more movable abdominal viscera can pass, without impediment, from the one into the other. The line of demarcation between these cavities corresponds to, or is in relation with, that arrangement of the internal abdominal fascia termed the internal inguinal ring: and, adherent to the posterior surface of the vaginal process, near its inferior termination, is placed the testicle.

In early foetal life, and, in many instances, for a month or even a longer period after birth, this tubular process of the

* H. A. Wrisbergii, *Commentationum Med. &c. Soc. Reg. Scient. Göttingensium oblatarum et editarum*, vol. i. p. 179, § 5.

anæum extends into the scrotum. It lies in front of the spermatic cord and testicle; extends from the internal inguinal ring to the lowest end of that gland; and forms a membranous, cylindrical canal until it reaches the testis, when it expands into an elliptical cul-de-sac. Before birth, or soon after, this vaginal process of the peritonæum is divided into two portions—superior and inferior. The sheath contracts near the head of the epididymis, its surfaces adhere firmly at that spot, and thus two cavities are formed. The inferior one is termed the tunica vaginalis propria testis: for it is in immediate relation with a small portion of the surface of the testicle. The superior canal, termed the tunica vaginalis propria funiculi, lies in front of the spermatic cord.

When these parts are in a normal condition, the inferior cavity or vaginal covering peculiar to the testis remains throughout life as a closed serous sac, and contains a little serum. Under similar conditions the superior canal or vaginal covering peculiar to the spermatic cord is entirely obliterated. Its superior abdominal or ventral orifice is permanently closed; and although a trace of the existence of this canal is sometimes observable in adult life, it is merely a delicate fibrous cord, the remnant of the processus vaginalis peritonei. Occasionally, although very rarely, a fine membranous canal remains throughout a person's life sufficiently large to admit an ordinary sized probe. In the account of the post-mortem examination of Sir Astley Cooper, he stated that 'a minute serous canal, not more than a line in breadth when opened, was traced, extending from a depression at the right internal abdominal ring along the spermatic cord, to the cavity of the tunica vaginalis.' *

Sir Astley Cooper himself writes: 'I dissected a boy, six years of age, in whom the opening of the tunica vaginalis (vaginal process of the peritonæum) was still so large that I could pass a male catheter through it down to the testis.' And further on, in the same chapter, he relates a case of 'sudden descent of a hernia into a congenital vaginal process of the peritonæum, of the nature above described, in the person of a young man whilst in the act of lifting a sugar-cask.' † Several instances of a similar kind have come under the observation of the author of this essay.

* *Guy's Hosp. Reports*, 1841, vol. vi. p. 232.

† *The Anat. and Surg. Treatment of Abdom. Hernia*, 2d edit. chap. xvii.

Of the changes which take place in the vaginal process of the peritonæum.—Several distinguished anatomists have traced the changes which occur in this serous sheath to render its obliteration perfect. Without producing a literal translation of the writings of Seiler, I believe it will be interesting as well as useful to the reader to quote the substance of his observations.*

When the testis has reached the bottom of the scrotum, the inguinal canal, along which the cord passes, is still very short, scarcely one and a half to two lines long. In the fourth and fifth month the internal inguinal ring lies almost directly behind the external. The inguinal canal is developed in the later months, and soon after birth it is always longer. The spermatic cord now consists of the blood-vessels, vas deferens, nerves, and the vaginal canal, at the termination of which lies the testis and the larger part of the epididymis.

After the testicle has reached the fundus of the scrotum, the obliteration of the vaginal canal proceeds from the inguinal rings downwards to the superior border of the testis, so that only the proper vaginal membrane of the testis remains, as a rudiment of the vaginal canal—the tunica vaginalis propria testis. The time at which the closure of its ventral orifice takes place and the obliteration of the canal is completed cannot be well defined. It is usually closed, at least on one side, and generally at the upper part, from the internal inguinal ring to the centre or middle of the spermatic cord, even before the whole vaginal canal has contracted, especially that portion which is embraced by the internal inguinal ring. The first stage of the obliteration of this canal commences with this process.

In the second stage, the walls of the vaginal sheath unite together entirely, as far as the superior end of the testis; or it first closes in the neighbourhood of the testis, so that the central part still remains open.

The third stage is accomplished when the canal is partially entirely closed. This portion of serous membrane is converted into a flat band, which afterwards becomes connective tissue, but which is rather closer and finer than the rest of the same tissue of the spermatic cord.

* Anton Scarpa's *Neue Abhandlungen ü. d. Schenkel- u. Mittelfleischbrüche*, bearbeitet mit einer Anleitung zu der Zergliederung d. Leistengegend &c. Kupfer- tafeln von B. W. Seiler, vol. ii. p. 374, &c. Leipzig, 1822.

fourth stage, this stripe of connective tissue always thinner, and at last entirely disappears; or there remains only a slight trace of it below, above, or in the

of the vaginal process of the peritonæum at birth.—In the of new-born infants some portion of the vaginal canal exists. In 21 Seiler found 4 in which it was open on the left; 5 in which it was open on the right side; 4 on the left of these 13, 5 in which the abdominal aperture was open on either one side or the other. In 5 of the 21 the canal was closed above and below, but in the centre of the inferior part was closed, but in the upper part, at the internal abdominal ring, a portion remained wide open. Likewise, in other examinations, he found the inferior part closed towards the middle of the spermatic cord, or even at the internal abdominal ring; more rarely the centre of the canal, and still more rarely the upper part only. He found, in 53 new-born infants, 23 in which the canal was closed on both sides; 11 in which it was open on the right; 6 on the left.

Seiler found the following relations: in 13 new-born infants the canal was open on both sides in 8; in 6 in its centre between the abdominal orifice and the testis; in 2 in its lower part, in which also the abdominal orifice of the right testis remained open. In the remaining 5 it was open at the abdominal fold and the testis, but only on the right side.

He states that, as a rule, the complete closure of the vaginal canal takes place from the twentieth to the thirtieth day after birth.

He himself dissected numerous fetuses of full age, and was surprised to find in what a large majority of them the canal continued patent either on one side or both sides.

With a knowledge of these anatomical facts, we can readily understand how it happens that a canal or receptacle exists at the place to which a portion of the intestines may enter. When this takes place, the hernia of infancy exists. It is a circumstance of very common occurrence to see an infant soon after birth with an enlargement of the scrotum, which varies in size, and sometimes disappears altogether. It attains its largest dimensions when the infant exerts its abdominal muscles; it

slowly decreases and is lost sight of during repose, the bent posture, or in sleep. This tumour of the scrotum is to a hernia of small intestine which escapes from the vaginal cavity into the prolongation from the great peritoneal membrane, known by the name of the vaginal process of the peritonæum. This serous sheath, which, from its structure allows of considerable distension, is thus converted into a hernial sac. The attention of pathologists was first drawn to this by Haller in 1749. His observations were confirmed, and the nature of the disease was still further elucidated, by Hunter and Percival Pott. Haller employed the term *hernia congenita* to express that variety in which the intestine and testis touch each other, or are contained in the same sac; by this name the disease has been distinguished since the time of his publication. It is, however, a most inappropriate inasmuch as the hernia does not exist either during intra-uterine life or at birth. A congenital imperfection, it is true, allows of the descent of a hernia soon after birth; and therefore M. Morgagni calls it the '*hernia of infancy*.' But even this is not sufficiently definite. I prefer the designation, *hernia of the vaginal process of the peritonæum*.

4. *Of the slow and gradual development of the hernial sac.* The peritoneal membrane is capable of very great but gradual distension. This capacity is illustrated by cases of serous dropsy and of ovarian growths. In the same manner as the membrane yields to the general pressure of accumulating fluid so it dilates under the influence of local pressure into a distended culum, or sac, until it is sufficiently capacious to contain a large part of the alimentary canal. This condition we see in those cases of enormous double scrotal hernia, which sometimes even below the knees.

Every hernial sac is composed of a body, or central part, above which is the neck, and below the fundus. The orifice, abdominal or ventral aperture (*le collet* of the French) is the point of immediate communication with the cavity of the peritonæum. To the tissues forming the margins or borders of this opening particular attention must be directed. They constitute, by their rigidity and unyielding texture, the principal impediment to the reduction of the hernia, in most cases it is, therefore, this part of the sac which requires to be cut in order to replace the protruded viscus within the abdomen.

When the reduction cannot be effected without opening the hernial sac.

Development of the hernial sac.—The evolution of the hernial sac has been very completely traced and described by Drs. Jules Cloquet* and Demeaux.† I shall here briefly as possible the chief facts related by the latest anatomologist.

The peritonæum, depressed by the pressure of the abdominal viscera, traverses the walls of the abdomen, it presents first, the shape of a digital depression; then, of a ring, and next, that of a finger of a glove. These forms are, somewhat dependent upon the region in which the hernia is developed. During this period the entrance to the sac is larger than the fundus, and in these conditions a protrusion of the herniated viscus cannot take place. But when the fundus of the sac has reached those tissues which offer resistance, it dilates, becomes rounded, and assumes a spherical shape; in this condition the entrance is more contracted than the fundus, and under these conditions strangulation takes place.

There are three periods or states in which the orifice and neck of the sac should be examined: 1st, of their formation; 2nd, of their dilatation; and 3rd, of their contraction.

1st period of formation.—It is important to describe the formation of the mouth of the sac, since, from certain conclusions which are useful in practice may be deduced. From the formation of the hernial sac it is demonstrable that the descent of the peritonæum is a condition of more important stretching of the membrane.

When the hernial sac is complete, the portion of peritonæum which forms it may be represented by a plain membrane three or four inches in diameter and ten in circumference; whilst one half of this surface is placed at the fundus of the sac at its dilated part, the periphery, folded and puckered like the mouth of a closed purse, would be placed on a level with the rest of the narrowest part; that is to say, at the orifice of the sac.

**Recherches anatomiques sur les Hernies de l'Abdomen*, 4to, Paris, 1817; and
 †*sur les Causes et l'Anatomie des Hernies abdominales*, 4to, Paris, 1819.
 ‡*Recherches sur l'Évolution du Sac herniaire*, &c. 8vo, Paris, 1842.

phenomenon takes place. The solution of the intestine by the muscular ring, upon the study of this stage of the process, is an advanced stage, the orifice of the sac is the focus of organisation which sufficiently explains all the phenomena which follow. This is the focus.

The second period, that of organisation, is the series of phenomena which take place at the orifice of a perfectly-formed hernial sac. At this stage puckering only exists in such a condition as to form a ring, but, at a more advanced period, the folds form adhesions together, in consequence of contact of their serous surfaces, and the sac exists independently of other organs. It is a new organ, annexed to the peritoneum, and need of being supported by the ring in this moment the orifice has an evolutionary character. It becomes the seat of very remarkable changes which can be observed at different periods, and explained.

The serous surface is not alone the seat of changes relating to the organisation of the orifice. The connective tissue changes of no less importance. The adipose tissue in this part diminishes.

M. Demeaux considers that the organisation of this annularuration begins in the sub-peritoneal connective tissue, and that at a certain time the peritonæum itself undergoes a change; becomes vascular, and the two structures united together adhere closely. This induration of the internal surface of the orifice of old hernial sacs is due to an annular thickening of the peritonæum, nearly limited to the boundaries of the mouth of the sac, which is thus reduced to much less dimensions than any other part of it. Preparations demonstrating these conditions, with the exception of the vascularity, are preserved in the Museums of the Royal College of Surgeons and of the different hospitals of London.

A layer of fibres, which interlace in every direction and resemble the tissue of the dartos, is said to exist around the orifice of the sac. If these be contractile fibres, they must exert considerable influence to prevent the reduction of hernia, and may be considered as *one* impediment to that result. Are they not, probably, an excess of development of the contractile fibres of the connective tissue? M. Desprez considers them to be an exaggeration of the normal state. This dartos-like layer, having contractile properties, may also play its part in producing angulation of some varieties of hernia.

The third period is one of contraction. Most writers on hernia have remarked the constant tendency which the orifice of the sac has to contract, and even to become obliterated, as soon as the herniated organs cease to act upon it.

This process is demonstrated in the cases of adhesion of the bouchure of the sac in infants; closure of its orifice in adults; and obliteration of the sac by persistence of omentum in its position. This termination, the most desirable of all, does not instantly occur. A gradual contraction may take place without the obliteration being complete; it is this disposition especially which gives rise to such formidable accidents.

At this period, if the orifice be examined anatomically, it is easy to prove that it has been the seat of a new transformation.

As the proportion as the tissues around the orifice become thicker and contract, their vascularity diminishes; and this layer, originally resembling the dartos, shows the firmness and resistance of fibrous tissue. In the second period, the orifice, susceptible perhaps of contracting spasmodically, may also admit of slight dilatation. In the third, all dilatation becomes impossible; either the margins of the orifice resist the pressure

of the viscera, or the viscus which passes through the orifice becomes strangulated.

The tissues composing this annular contraction of the mouth of the sac often become nearly as hard as cartilage; and this change especially occurs in old cases, when the hernia has been allowed to descend for a long time; for then the boundaries of the orifice are placed in the most favourable position for the process of contraction and induration to take place. But when the hernia is not reduced, the constant pressure of the viscera dilates the mouth of the sac as well as the fibrous rings.

Of the two kinds of hernial sac.—The ‘hernial sac’ is always a prolongation of the parietal peritonæum from the abdominal cavity into the neighbouring structures. Its development and formation, however, depend upon very opposite causes. Two kinds require to be described; the intrinsic and essential distinction between them depending upon their mode of development.

1. That kind which, being congenital, is merely a serous canal or sheath, the vaginal process of the peritonæum, upon which a hernia takes place and escapes into it, when it becomes converted into a hernial sac. The physiological designation for it would be THE CONGENITAL HERNIAL SAC.

2. That kind which is the result of a slow and gradual process of relaxation, and is produced by a stretching, yielding elongation of the parietal peritonæum, before and under the pressure of the viscus itself, which constitutes the hernia. For the sake of brevity, this kind may be termed THE ACQUIRED HERNIAL SAC.

The first kind, or congenital serous canal, which may be converted into the congenital hernial sac, sometimes exists throughout life simply as a diverticulum of the peritonæum, without a hernia entering it; but the second kind, the acquired hernial sac, cannot exist unless a viscus, almost entirely invested by peritonæum, pushes the parietal layer of the peritoneal membrane before it.

In other words, the first kind of sac is peculiar to a person born with any portion of the vaginal process of the peritonæum open, and who then possesses a receptacle for the re-irruption of a hernia; and, if the ventral orifice and cavity of this sheath should not become obliterated soon after birth,

pture may occur at any subsequent period of life; but the second kind being an acquired formation, some length of time necessary for its production, and it cannot exist at all without the continued and effective influence of the hernia itself.

Pott, writing of the two varieties of oblique inguinal hernia, the congenital and ordinary, expresses his belief 'that common ruptures, or those in a common sac, are generally gradually formed, that is, they are first inguinal, and by degrees become total; but the congenital are seldom if ever remembered by the patient to have been in the groin only.'*

The paramount importance of a due appreciation of the physiological differences between the one kind of hernial and the other will be demonstrated in many points of view, especially with regard to the surgical treatment of hernia.

The treatment of reducible hernia.—The surgical treatment of all kinds of reducible abdominal hernia is divisible into two categories:

1. The employment of palliative or prophylactic measures.
2. The adoption of some proceeding designed to effect a permanent cure.

1. *Palliative measures.*—To this class belong all contrivances which prevent the descent of the hernia. These are the recumbent posture, bandages and trusses of various descriptions. We only judge from the class of cases observed in the hospitals, and selected from among the labouring population, we should form a very inadequate estimate of the advantage which accrues to sufferers with hernia by the employment of well-adjusted trusses. We know of adult persons in comfortable circumstances who, having no need for toil, have taken precautionary measures to prevent the recurrence of the rupture, and have consequently been free from everything of the kind for several years after. Some have even been enabled to discontinue the use of the truss entirely, after wearing it some months. It is, however, a condition of the utmost practical importance that the case be treated immediately that the disposition to the formation of rupture is detected. When the hernia enters the vaginal process of the peritonæum, and that the sheath is converted into the hernial sac, we may effect a cure

* *The Surgical Works of P. Pott*, edit. 1808, vol. ii. p. 120.

by attempting to call forth the processes adopted by nature to close its orifice and obliterate that canal; or, in other words, to procure adhesions of the serous surfaces of this peritoneal diverticulum.

On the other hand, in the slowly-forming hernial sac the object in making use of a truss should be, in the first instance to prevent the development of the peritoneal sac. This may be accomplished by applying support to those weak parts of the abdominal walls through which the parietal peritonæum is forced. When, however, the sac is once developed—and this is commonly the period at which a surgeon is consulted—measures should be immediately adopted to assist in arresting its dilatation by preventing the descent of the rupture.

The expediency of judiciously pursuing the mechanical treatment of every variety of hernia cannot be too strongly urged upon the laity by the profession. In both sexes it should be carefully conducted the moment that the slightest protrusion shows itself. Whether the hernia occur in infancy, youth, middle age, or at later periods of life, if properly watched and judiciously supported, it usually gives but little trouble; many cases it is even cured. But, on the contrary, if it is neglected, increase in bulk, and, sooner or later, diseased state of the rupture, often leading to the death of the individual, will almost infallibly occur.

The only exception to be made to this rule applies to the rare cases of hernia into the vaginal process of the peritonæum in which the abnormal situation of the testis interposes a practical difficulty to wearing a truss, the necessary pressure of which occasions intolerable pain by compressing that organ simultaneously. But even such cases should not be abandoned as hopeless of cure, without a reasonable attempt being made to afford relief.

The practice of leaving cases of rupture in the hands of mere tradesmen cannot be too strongly censured. Among the poor we constantly observe the lamentable effects of this proceeding. Ill-shaped trusses are applied; the springs, being too feeble, allow the hernia to descend behind the pad, where it becomes compressed; or they are too strong, and their pressure induces absorption of the abdominal parietes, on which the pad presses. Frequently a truss suitable for supporting a femoral hernia is applied to one of the inguinal kind, and *vice versâ*. That which is worth doing at all is always worth doing well. This

junction receives strong confirmatory testimony in the treatment of ruptures by mechanical means.

Characteristics of trusses.—The requisite and essential qualities of a truss are lightness, firmness, elasticity, so that it shall retain the required form or shape, suitable adaptation to the configuration of the wearer, and sufficient strength of spring to prevent the escape of the rupture from the abdomen.

The instrument consists of a pad or cushion attached to a metallic spring, with straps, so arranged that its movement during the varied postures of the body may be restrained.

Through the kindness of Mr. J. A. Kingdon, Surgeon to the City of London Truss Society, we are enabled to give the results of his extensive experience in the employment of trusses.

He considers that the circular spring truss is the most suitable form, in the majority of cases. Bandages which are not elastic do not afford sufficient support to the hernia in every posture of the body. They are necessarily unsafe on this account, as they become lax in the stooping posture of the wearer—the position of all others in which the hernia most easily descends, because of the relaxation of the pillars of the external ring. The curve of the spring and the relative position of the pad with it should be appropriate to the configuration of the wearer. A single piece of metal should form the spring and foundation of the pad. As far as practicable, the spring of the truss should pass around the bony rim of the pelvis, fitting closely to the figure, and should lie out of the region of the latæi muscles. For, unless it be so placed, their alternation in progression produces a corresponding movement of the pad. If these muscles be largely developed, extending upwards to the very edge of the pelvis, the curve of the spring would be wide at the shoulder, so that its bearing or resting point may be on the base of the sacrum.

For a single pad truss the free end of the metal spring should be beaten out flat and thin, and so ground as to cling round the opposite hip,—an arrangement which materially aids in steadying the truss.

The form of the spring may be designated as after the French model or the German. The former resembles the coil of a watch-spring, and is very elastic and clinging; the latter almost exactly fits the outline of the body in its state of repose.

It is almost inelastic, and very hard. The French is always pressing inwards, even when the wearer is at rest. The German scarcely presses at all when the abdomen is soft, but resists with power when any expulsive force makes the abdomen swell. In practice, the best shape for the spring is one which forms a medium between these two extremes.

The pad or cushion should be of moderate dimensions. For the adult, it should not exceed two and a half inches in length and two inches at the widest part. Its superior edge should follow the upper line of the spring, which falls a little from the shoulder or bend, where it lies in contact with the hip. The inner surface should be directed slightly upwards, but this inclination must depend upon the prominence, or otherwise, of the abdomen, as well as, in some measure, on the anatomic relations of the pelvis to the spine. The proper shape for the cushion or pad, and the materials of which it should be constructed, may be varied to accommodate particular cases, or to accord with the views of different inventors. Generally, the wearer discovers, after a little experience, which kind of pad is most free from annoyance. That pad, however, is the best which maintains perfect and unintermitting retention of the hernia. Every pad should have attached to it two studs, one near its junction with the spring, and another at its lower point. To the upper one the transverse strap passing from the free end of the spring is attached. The lower stud is used with the thigh-strap, which should be always worn. It is loosely fastened on to the spring of the truss near its shoulder, and should fall along the hollow beneath the buttock. In the erect posture of the wearer this strap should be moderately tight. It prevents the pad from shifting from its proper position, and should never be discarded.

These appear to be the general principles by which a surgeon is to be guided in the selection of a truss. Particular modifications suitable for special kinds of hernia will be noticed in other places.

Of the so-called radical cure of a hernia.—It would be idle to encroach upon the limits of this essay by describing in detail every method employed to effect a cure of hernia, from the most remote periods. A mere notice of them will suffice. They were—castration; cauterisation of the orifice and neck of the sac; ligature applied around the neck of the sac; incision of the sac and subsequent healing of the wound by granulation; excision

and scarification of the sac; detaching the sac from its adhesions and returning it into the abdomen; immediate and forcible compression of the sac by a bandage or truss; cold applications; stimulating injections, especially tincture of iodine. These proceedings have been long abandoned, in consequence of the bad results which frequently attended their employment. We must, however, add the treatment recently adopted, namely, the invagination of the hernial sac. Every modification of the operation is usually designated by the name of the inventor: there is the method of Gerdy, of Wutzer, of Rothmund, of others.*

The principle of this operation is expressed by the words, 'invagination of the hernial sac.' That is to say, the fundus of the sac is pushed up into its mouth, and retained there until adhesions have formed between it and the entire circumference of the external orifice. By this means it is hoped that obliteration of the external orifice of the sac and of its cavity may become permanently established. It is an indisputable fact, that cases of reducible inguinal hernia have been permanently cured by the performance of an operation on the plan of Wutzer. We must admit, however, the data by which to establish the proportion of successes to failures; but that a considerable number of the cases operated upon did fail, is certain.

Kingdon has very kindly given me the notes of sixteen cases in which an operation for the radical cure of the rupture was performed, either by London, provincial, or colonial surgeons. The patients applied to him for relief at the City of London Truss Society. In most of these persons, at the time they were used to be under the observation of the operator, the rupture was 'cured;' for reports of some of the cases appear in the medical periodicals of the day. In some of these patients the rupture was larger than before the operation, and greater difficulty was experienced in retaining it within the abdomen. In others, a protrusion existed on both sides, the second having developed after the operation.

Through the intervention of Dr. Hermann Weber, Physician to the German Hospital, I have received a valuable communication from Dr. Otto Weber of Bonn, who was formerly clinical assistant to Professor Wutzer. The number of persons upon whom the Professor operated amounts to about fourteen. Be-

*For the most complete account, especially of the German cases, is to be found in *Lehre von den blutigen Operationen*, Lieferung 50.

tween 1852 and 1856 the operation was performed but on the 'Klinik,' during which time Dr. Otto Weber was assistant to Wutzer. The patient was a teacher, forty years old. He quitted the hospital 'cured,' but the hernia returned in spite of his having worn a truss.

Wutzer is still of opinion—1. That when the operation properly performed after his method, it is not attended with danger. 2. That he has succeeded in fixing the plug of skin with invagination of the hernial sac by inducing adhesion between its internal surface and the interior of its neck. That in consequence of this, if the patient continues to wear a truss (for life), a return of the hernia may be avoided.

However, Dr. O. Weber writes, that he has never seen of the so-called 'cured cases' radically cured, but that—1. That the plug of skin is, by degrees, entirely drawn out again. Secondly, that the true herniary apertures, the external and internal rings, are not closed by the operation; and, thirdly, that an imperfect cure may be effected by means of a permanent closure, by adhesion of the internal walls of the neck of the hernial sac, and thickening of the surrounding connective tissue.

The cases which Dr. O. Weber had an opportunity of examining some time after the operation, showed a slight protrusion of the intestine into the inguinal canal; and there have been a perfect return of the hernia if trusses had not been worn. It appears that, at first, the invaginated skin became adherent to the inguinal canal, but without the participation of the hernial sac; that the latter probably becomes merely compressed by the invaginated skin passing by its side. In most favourable cases the inner walls of the hernial sac adhere together, but so loosely that they soon become separated by pressure.

The results of the examination of the bodies of persons who died upon whom the operation had been performed, are important. Streubel has published some. Dr. O. Weber examined the body of a man who died of pneumonia in 1856, upon whom Wutzer operated in 1840. The hernia had returned after the operation. There was not a trace of the invaginated skin in the inguinal canal, and the peritonæum did not show any signs of previous inflammation. The hernia and its contents did not differ from the ordinary cases of the same kind.

Dr. O. Weber thinks that Wutzer's method might be per-

diffied in such a manner as to become effective. But, first, considers it is necessary to show upon the dead body that fundus of the hernial sac can be actually invaginated into orifice, and not merely pushed up a certain distance before integuments.

But, instead of describing in detail the operation of Wutzer and its modifications,* let us inquire, what is the object the surgeon has in contemplation when proposing to perform an operation for the radical cure of a hernia? In order to be actual and permanent, it must, of course, produce the invagination of the hernial sac; the closure of its ventral orifice; the strengthening of those weak parts in the walls of the abdomen through which the rupture protrudes; and to these conditions must be also added, an improved tone of the peritoneal ligaments of the viscera, by which the power they exert in retaining the viscera in their normal situation is restored. Unless the operation perfectly and completely accomplishes these ends, failure must most certainly ensue sooner or later.

Another intention which the successful issue of the operation is expected to fulfil, is to enable the sufferer to dispense with the use of a truss, and thus to be free from an intolerable burthen. To recommend a dangerous operation, therefore, which may prove only so far successful as to impede the descent of a hernia, but yet leaves the patient under the necessity of continuing to wear a truss to prevent a recurrence of the rupture, is surely scarcely justifiable. It is contended by the advocates for the operation that the dangers to which the patient is subjected have been too prominently set forth. This may be true. Few persons have indeed died from peritonitis, from other causes, in proportion to the numbers upon whom the operation has been performed. But they all were subjected to great risk, and because they happily escaped the fatal complication, that is no ground for the inference that others would do so also. The facts only show that the operation may be done without exciting peritonitis in every case.

The surgeon, in the attempt to carry out his object, proposes to produce an effect in imitation of the processes of nature during the early periods of life; for there are facts to de-

* Operations intended to effect the radical cure will be described with the treatment of the special kinds of hernia.

monstrate that the cavity of the vaginal process of the peritonæum, which is, in so many cases, an apt receptacle for any protruded viscus, may become obliterated even after a hernia has passed into it in early infancy. But where is the evidence to prove that an acquired hernial sac becomes thus obliterated by similar natural efforts? The only instances, perhaps, are those in which the orifice of the sac is plugged by adherent omentum; and such cases are very rare.

We therefore believe that we shall not err in enunciating the principle that the cases of inguinal hernia selected for the performance of all operations for the radical cure should be those in which the protruded viscus has descended into a patent vaginal process of the peritonæum, and that all other kinds should be rejected as unsuitable; and that the more efficiently the proposed methods accomplish the ends effected by the processes of nature, the more worthy of confidence they become. Thus, allowing nature to guide our procedure, we must make it a rule to select those cases in which her efforts have failed; and by acting as her handmaid, we may reasonably hope to arrive at a successful result.

We may conclude these observations with the following quotation from the work of Mr. Lawrence: 'The subject of an incarcerated rupture submits to an operation in order to save his life. But he whose hernia is reducible, endangers his life to get rid of an inconvenience; and the operation affords no greater prospect of entire recovery than he had without it. For after he has undergone an operation, at the hazard of life, the complaint may return; and the only protection against relapse is to wear a truss' (*Treatise on Ruptures*, chap. vi.).

The morbid conditions of the hernia, and impediments to its reduction.—A hernia is said to be reducible when the protruded viscera can be returned into the abdominal cavity; it is irreducible when they cannot.

The impediments to the reduction of a hernia may be classed in three groups: those produced by the tissues outside the sac; those occasioned by the sac itself; and those which exist within the sac.

There are also primary or immediate causes which render hernia irreducible, as well as secondary. Among the first may include all those impediments to reduction which depend upon organic conditions, and relate directly to the hernial sac,

the muscular contraction which influences the size of the ominal rings and the tension of the tissues around the ce of the sac ; the condition of the orifice of the sac itself ; the development within the sac of adhesions or omental

he secondary are those which are partly exclusive of the ial tumour, and occur, as it were, accidentally, and are ely due to individual peculiarities ; as, for example, the lopment of a thick layer of fat around the hernial tumour ; great mobility ; its diminutiveness ; tension of the sac from umulation of the fluid or gaseous contents of the bowel ; erent omentum lying in front of the intestine ; and some r conditions.

he morbid conditions of the hernia which give rise to more ess serious difficulties may be classed under one of the owing states : 1. Permanent irreducibility. 2. Distension nflation by fluid, gaseous, or solid contents, termed obstruc- . 3. Inflammation. 4. Strangulation.

A hernia becomes permanently irreducible, after many s' existence, on account of its size. In most cases this con- on is only attributable to neglect of the use of a truss. The est double rupture I have seen was in a bricklayer, fifty-five rs old, whose bodily health, strength, and conformation were ther respects very good. The left hernial tumour was the er, although it had only existed about three years, whilst right had been there twelve. The lowest border of the our very nearly reached to a level with the patellæ. Its umference in the largest part measured thirty inches. dhesions also, when formed between the contents of the sac, etween the sac and its contents, prevent the reduction of protrusion.

. Distension ; choking ; obstruction ; rendering the hernia ducible. This particular state or condition of a hernia is ibuted to the accumulation of the solid, fluid, or gaseous tents of the alimentary canal within that portion of the el which constitutes the protrusion. From this cause the age of the stercoraceous contents of the canal are arrested ; alimentary canal or tube becomes choked up with its own etions and the egesta of the stomach, giving rise to local bles and constitutional disturbance.

The doctrine of obstruction is of great antiquity, and for a long period that morbid condition of a hernia we now term 'strangulation' was believed to originate in this cause alone. 'During nineteen centuries,' writes M. Broca, 'the existence of an accumulation of faecal matter in the hernia had been admitted, or rather supposed: but when it was looked for, the accumulation was not to be found. Thus theory fell to the ground on the first observation.'

The local symptoms of an obstructed hernia are—pain, a flatulent state of the tumour, increase of size in comparison with its usual dimensions, more or less tension, and the absence of those more severe local conditions which characterise a strangulated bowel. By careful manipulation, the gaseous and fluid contents of an enterocele may be expressed into the contiguous part of the canal, and even solid faecal matter may be felt. During this proceeding the gurgling produced by the flatus and fluid is heard, and the movement of these is sometimes felt as they escape through that portion of the bowel lying within the orifice of the sac.

At first, the constitutional symptoms indicate functional disturbance of the alimentary canal, rather than any morbid state of the tissues of the herniated viscus. There are indications merely of a blocking-up of the tube, and nothing more. Thus, slight pyrexia and nausea, succeeded by vomiting, are the most marked features, which continue even although the large intestine below that portion in the sac has been emptied of its contents after the commencement of the attack. Unless the vomiting be very prolonged, and great depression result therefrom, as occasionally happens in very delicate or old persons, the surgeon will not fail to remark the absence of the urgent symptoms characteristic of strangulated bowel.

In most cases of obstruction the impediment is removed by those means which excite the natural peristaltic movements of the alimentary tube. The local application of warmth and moisture to allay pain; repose; the administration of aperient enemata, or even in some cases, when vomiting has not occurred, of purgatives, by the mouth, are beneficial. Abstaining, in the first instance, from manipulation of the tumour, is a point of great importance. Should the impediment to the passage of the indurated faecal matter depend upon the contraction of the orifice of the sac, the enlargement of this part becomes necessary, either by cutting its tissues or those around it. T

operation must then be performed in the usual manner, and in relation to the special case.

3. *Inflammation of a hernia.*—A hernia in this state shows all the local signs, and excites the constitutional symptoms of inflammation. The combination of those indications which characterise the condition called strangulation do not, however, exist.

Inflammation is usually the result of external violence, and

FIG. 264.



Hernia at the femoral ring of a part only of the walls of the bowel. The patient, fifty-six years old, had been ill nine days, when the exploration of a small swelling in the site of the left femoral ring was made. She was then dying, but survived the operation forty-five hours. After death peritonitis was found, and the intestines in the pelvis were united together. * The cæcal end of the bowel, the tube contracted between it and the hernia adherent to the sac. *a.* Intestine adherent to the sac. *b.* Peritoneal sac. *c.* Lobules of fat outside the peritoneal sac. (Museum Guy's Hospital, Drawing 484²⁰—Preparation 2503²⁰.)

may be produced by the pressure of badly-fitting trusses. A morbid state of the whole alimentary canal, of an inflammatory type, may extend to the hernia, and in that way give rise to trouble.

reducible epiploceles are more liable to be inflamed than kinds of hernia; and a patient the subject of reducible ocele may have the hernia inflamed, from which cause it

becomes permanently irreducible by contracting adhesions to the sac.

The local signs are—pain, increase in the bulk of the tumour, a certain degree of hardness, firmness, and elasticity when it is pressed, with more or less unevenness and irregularity of surface. The margins of the aperture through which the hernia passes do not tightly embrace the protrusion, so that, in a large hernia, the finger can be passed along the pedicle of the tumour for some distance.

The constitutional symptoms are not usually severe. In the first stages more or less pyrexia arises; and when the contents of the tube are arrested in their course, the indications of the condition become marked.

It is unnecessary to describe in detail the local and constitutional measures to be adopted. Reliance must be placed upon those in ordinary use to control and arrest the disease.

There occasionally happens a variety of hernia, first noticed by M. de Littre,* in which only a portion of the wall of the ileum, or a diverticulum from the bowel becomes adherent to a very small hernial sac. By extension of inflammation to the neighbouring viscera, obstruction to the passage of the stercoraceous contents of the tube arises, and death ensues.

Fig. 264 illustrates this variety. It was drawn from a case which was under the care of the writer.

4. *Strangulation*.—A hernia is said to be strangulated when the displaced viscus is subjected to a constriction which impedes or arrests the circulation of the blood through its vessels, the passage of the stercoraceous materials along its canal, and constitutes an impediment to its return into the abdominal cavity. This condition excites constitutional symptoms of the most dangerous and characteristic nature.

Is the condition of a hernia, termed strangulation, solely the result of mechanical constriction produced by the margins of the orifice of the sac; or does it depend upon a morbid action engendered in the tissues of the protruded bowel antecedent to its escape from the abdominal cavity? In several instances of strangulated femoral hernia in women, and in some cases of inguinal hernia in men, the patients themselves voluntarily

* 'Obs. sur une nouvelle espèce de Hernie,' *Mém. de l'Acad. royale des Sciences*, 1700, p. 300.

ted that, for some hours before the hernia caused any inconvenience, or was even in the sac, their bowels had been 'disordered,' 'relaxed,' or that they had been suffering with 'bowel-plaint.' In other cases, there seemed to be a great tendency to a very rapid derangement of the mucous membrane along the whole track of the canal above the hernia, scarcely explicable on the supposition that it had been all excited by the mere existence of a constriction around a small knuckle of intestine. In evidence of this morbid action, I would adduce the fact of a rapid and abundant secretion from the entire mucous face of the small intestines, their great distension and intense vascular congestion, witnessed in some instances.

Again, given any case of reducible hernia, which for months, even years, has readily glided into the hernial sac, and has been returned as easily into the abdominal cavity, why on some particular occasion should it become irreducible when it has passed through the same orifice it has been in the habit of re-aversing, and in the tissues of which no appreciable changes have occurred? For it is in vain we seek any marked alteration in the structures around or composing the orifice of the hernial sac itself.

In giving an affirmative reply to the above inquiry, we are justified by facts in attributing the strangulated state of a hernia to a predisposing cause, commencing in a morbid state of the alimentary canal generally; at least, in some cases. Patients often relate how they had observed that the hernial tumour was 'larger than usual' before they suffered much inconvenience; and this circumstance is commonly attributed by them to a greater bulk of the protruding viscus. But we have no proof that this is the correct explanation of the increase of size of the tumour in all cases; in some doubtless it may be, but it is due to a distended state of the bowel rather than to its quantity.

Let us examine a case of reducible inguino-scrotal enterocoele. At one moment all that the surgeon can detect is a slight fullness produced by an empty hernial sac. At another, a small knuckle of intestine, having descended in an empty condition, becomes more or less filled with the stercoraceous contents of the alimentary canal from above it, and an elastic swelling results, which is produced by the bowel and its contents, fluid and gaseous. Assisted by the peristaltic movement of the viscus itself, or by gentle pressure, these contents of the tube

these phenomena the peristaltic action of the canal, the contraction of the tissues of the hernia by congestion of its vessels, and the irritation of its nerves? During these progressive stages the protrusion has attained a size dependent upon more or less distension, caused by accumulation of its contents, and the swelling of its tissues, induced by the effusion of inflammatory products, reacting against the narrow orifice of the sac, and the more or less retardation or arrestation of the sanguiferous current in the blood-vessels of the rupture, and its strangulation, the result. The increased bulk of the protrusion prevents the retrograde passage through the small aperture by which it escaped from the abdomen. The body of the sac being stretched, may mechanically contract the dimensions of its orifice in some measure; whilst the arrangement of the contents of the tumour, in relation to the mouth of the sac, may likewise constitute an impediment to its reduction.

The filaments of the nerves of the injured bowel play an important part in transmitting thence to the nervous system the local disturbance of their sentient extremities, and consequently the sympathies are excited. The heart contracts more frequently and with greater force; the pulse is full, and more than normal; the cheeks are flushed; the surface of the body is warmer than usual, often moist, though sometimes dry; the tongue is covered with a white fur; nausea and intolerance of food are complained of; tenesmus occurs, and flatus and a small quantity of feces may be expelled, if the large intestine

cate the commencement and progress of strangulation of intestine, we must now systematically describe—

First, the condition of the hernia when strangulated.

Secondly, the changes taking place in the tumour.

Thirdly, the constitutional symptoms excited by the local disease; and,

Fourthly, the morbid conditions developed within the abdominal cavity.

The structural changes taking place in the hernia.—When the bowel is in the condition implied by the word ‘strangulation,’ the circulation of the blood within the vessels of the part is, at first, impeded. Congestion of the capillary vessels is the consequence, and the tissues of the viscus become swollen. When exposed to view, in this first stage, the serous membrane is of a deep-red tint, and through a lens the minute blood-vessels may be distinctly seen, with their outlines well defined. To the touch the hernia feels firm, and resists slight pressure; but the elasticity and resiliency of its tissues still continue unimpaired.

Next, the circulation of the blood is arrested. The tissues of the bowel, about which the constriction is tight, become now more swollen; they have palpably a solid, leathery consistence; the colour of the serous surface is a dark purple; it is dull, lacks its usual lustre, and sometimes, varying as regards shade or depth of colour, it is mottled with a red and purple tint. Patches of extravasated blood appear in the subserous connective tissue, and an adhesiveness to the fingers is very striking.

When the bowel has been strangulated many hours, its tissues are more swollen and soft; they no longer possess their characteristic elasticity, but they remain in the same position in which the pressure of the finger places them. The serous surface has lost all its characteristics; it is black, ash-coloured, flocculent, and adhesive. In this stage all the tissues of the bowel have undergone more or less disorganisation; blood is extravasated in the connective tissues between the different coats of the viscus; the mucous membrane is soft, flocculent, and easily detached from the fibrous walls. Probably in some part of the bowel, at its convexity, or where it is in immediate relation with the mouth of the sac, its tissues have sloughed, or ulceration is concealed by shreds or patches of adherent lymph.

The intestine is also usually firmly fixed to the mouth of the sac by inflammatory adhesions.

The last stage is that in which the entire mass of the protruding bowel has become gangrenous, or has passed into a condition termed sphacelus.

In the diseased conditions just described we may recognize morbid changes similar to those which are observed in strangulation on the outside of the body. If a string be tied around the neck of a bladder near the prepuce, the latter organ becomes first swollen, secondly inflamed; thirdly, it suppurates; and lastly, it mortifies. Ulceration also takes place at the part with which the ligature is in contact.

The coats of the bowel are often ulcerated by the pressure of the constriction to which they are subjected, but the various tissues of which the viscus is composed are endowed with different powers of resistance to those morbid processes. It may be seen, on cutting open the intestine, in cases in which the bowel has been deeply sulcated or grooved on its external surface, without any trace of abrasion thereon, that the mucous membrane is ulcerated in a line corresponding to the groove immediately embraced by the medium of constriction. Occasionally it happens that the line of ulceration forms on the free end of that end of the strangulated bowel directly continuous with the upper and distended portion of the alimentary tube, where there is none at the other end. This is probably caused by the pressure of the distended bowel within the abdomen, and it may be seen, particularly in some cases where the knuckle of intestine lies in close relation with Gimbernat's ligament. We have even observed blood in the fecal evacuations of patients who have died with the bowel thus diseased.

Mr. Bryant states* that ulceration at the line of stricture is most frequent in inguinal hernia, although the sulcation of the bowel is as common in femoral as in inguinal hernia, and that fecal extravasation, if not produced by rupture of the bowel from taxis, generally follows ulceration at the line of stricture.

Small, recently-developed herniæ are more frequently strangulated than large ones, and those which have existed some time. Sir A. Cooper writes: 'A small hernia is more easily strangulated than a large one, the pressure on the contents being

* *Guy's Hospital Reports*, 1856.

iolent, and the symptoms are much more urgent, as the stricture acts with much more effect upon a single knuckle in topping its circulation, than when the contents of a hernia are large and voluminous.*

It is a matter of great importance to distinguish between the two classes of hernia—'RECENT' and 'OLD.' Mr. Pott observes: 'Recent hernias are in general more liable to stricture than old ones, for reasons which are obvious from what has already been said; but when old ones get into the same circumstances, the symptoms are much the same; though I think in general they are not altogether so pressing, and the latter generally admit of more time to attempt reduction in.' †

And very lately Mr. Bryant has demonstrated, by the cases admitted into Guy's Hospital, that in recent hernia strangulation frequently occurs, and that the risk attending it is very great, even although the bowel be speedily liberated.‡

Morbid states of the hernia induced by violent manual pressure.—Let it not, however, be assumed, that the morbid states of a hernia, above described, are always inevitably due to the compression which the orifice of the sac has exerted around it. Such is not the fact. The tissues of the herniated viscus are rarely so firmly constricted as to produce in a short space of time complete mortification, absolute death, of the whole of the knuckle of bowel in the sac; yet this state of the hernia is often met with. In these cases the protruded viscus has been subject to another source of damage and destruction. Under the mild expression of 'the use of the taxis,' the hernia has been forcibly compressed by manipulation; its tissues have been contused and irreparably damaged; blood has been extravasated in profusion between the different layers of the tissues composing the viscus, and complete disorganisation of its structures is the result of the injury inflicted. I have been for many years interested in ascertaining, from the observation of the state of the bowel after death, the comparative amount of injury inflicted on the viscus by the natural constriction of the orifice of the sac, with the tissues around it, and that produced by violent manipulation of the tumour, with the hope of reducing

* *The Anatomy and Surgical Treatment of Abdominal Hernia*, part i. chap. vii., the last paragraph.

† *The Chirurgical Works of P. Pott*, ed. t. 1808, vol. ii. p. 63.

‡ *Guy's Hospital Reports*, 1861.

not often happen that mortification of the bowel is observed in those cases in which the order happens not to have been recognised when the manipulation of the tumour has not

I may represent the danger which is associated with attempts to reduce the hernia, by stating, in more irreparable damage may be inflicted in a few minutes by coarse, careless, impetuous manipulation than the natural means of constriction could produce.

Need I add, that in the attempts to reduce a hernia, the employment of such violence as is necessary to produce these results is reprehensible in itself, and justly condemned as not only opposed to every principle of surgery but because it is in violation of all the principles of surgery?

Artificial anus.—Two conditions of the intestine are necessary for the formation of an artificial anus. One, in which the wall of the intestine sloughs, leaving a sinus, as if a piece had been punched out, which maintains the continuity of the alimentary canal, and allows a portion of its contents to escape. This forms at about the centre of the convex knuckle of intestine, and at a point farthest from the orifice of the sac.

Another variety is due to ulceration of all the coats of the bowel, even to the mesentery. It usually occurs at the mouth of the sac: and, in con-

the wound of the integuments heals perfectly, and complete recovery ensues.

After the second, however, the prognosis is not so favourable; for, commonly, an artificial anus is permanently established, and more or less of the contents of the alimentary tube are discharged hereat.

As the result of long-continued constriction by the mouth of the sac, and the pressure made upon the two pieces of intes-

FIG. 265.



Artificial anus after strangulated ileum, at the convexity of the tube. The patient, seventy-two years old, was the subject of strangulated femoral hernia for 'two days,' but probably more, when the protrusion was returned by operation. The bowels acted freely by the anus, but on the fifth day from the operation fecal matter was discharged from the wound (*b*). Fæces continued to pass through it until death, which took place seventeen days after the operation. The perforated bowel was the ileum, about a foot from the cæcum, which is seen in the wood-cut (*a*). It is raised from the mouth of the femoral aperture to show it more distinctly. Fæcal extravasation into the peritoneal cavity was prevented by adhesions between the margins of the perforation, neighbouring coils of bowel and omentum (*c*, *d*). The patches or marks on the coil above and behind are the remains of those adhesions. (Drawings, Museum Guy's Hospital, 486^b.)

time lying therein, their walls become adherent at the points of contact, ulceration takes place, and the continuity of the tube is thus established within the abdomen, by the processes of nature alone. This condition is very rarely met with, and I may therefore refer the reader to a published case.*

* *Transactions of the Pathological Society*, vol. x. p. 128; the preparation is preserved in the Museum at Guy's Hospital, No. 2492¹⁰.

the hernia is strangulated, differs in a remarkable manner in different individuals. In some persons the tumour

FIG. 266



Illustrates that variety of artificial anus in which the continuity of the rectum is destroyed in consequence of destruction of the entire calibre of the bowels. This is a case of femoral hernia in a woman sixty-seven years old. The bowels were strangulated fifty-five hours when herniotomy was performed, and the hernia was reduced into the abdomen. Four days afterwards faeces escaped from the wound. Sloughing took place around the wound, and the woman lived through. The symphysis pubis is *b*. Fistula leading from the intestine *d*. The low ileum *f* ending in the caecum *e*. *c*. Part of abdominal walls. (Drawn from a dissection at Guy's Hospital, No. 486¹².)

quickly sensitive, and even intensely painful, so that the patient is intolerant of the most gentle manipulation, especially in the region of the orifice and neck of the sac. Sometimes

rapidly; its surface becomes more uniform and regular; the integuments tense, smooth, shining, red, with increase of temperature; and the general outline of the whole swelling assumes a more pyriform shape, or is more locally circumscribed, according to the variety of the hernia. Fluctuation may also be detected in some cases. It must, however, be remembered, that the displacement of gas by pressure upon the tumour causes a sensation very closely resembling that which indicates the presence of fluid, and that the difference between one and the other is not always perceptible even by surgeons of great experience in delicate manipulation.

The sudden enlargement of the tumour is explained by the fact of a rapid secretion and accumulation of serum within the hernial sac, in numerous instances; by the distension of the enterocele with gases, in others.

Characters of the serum in the sac.—The serum varies in character according to the duration of the strangulation. When the sac containing a strangulated hernia is cut open, some fluid usually escapes. In colour, consistence, and odour, this serum varies considerably in different cases; and these diversities may be regarded as important indications of the condition of the tissues of the strangulated bowel.

It is pale yellow, clear and bright, when strangulation has existed a few hours only; and, under these circumstances, the intestine is simply deep-red, from vascular congestion, and its tissues elastic.

It is dark-brown, but clear, when strangulation has existed many hours; and the intestine is then of a purple tint, but its tissues elastic.

Its colour resembles that of a strong infusion of coffee; it is turbid; blood and small coagula are mingled with it, when the protrusion has been long strangulated, or protracted and violent taxis employed; now the intestine is dark purple, inclining to black; its tissues are leathery, not resilient, infiltrated with blood, and often flakes of lymph are adherent to its surface.

When turbid, dull brownish-yellow, and containing blood, coagula, pus, flakes of lymph, and even faecoid matter, in which state the odour of faeces is perceptible, the intestine is usually approaching a gangrenous condition, if it have not ready passed into sphacelus.

The escape of gas through serum, of the nature last described,

producing bubbles, is an indication that the herniated bowel is probably ruptured, and that a communication exists with the interior of the alimentary tube.

If the progress of the disease be not arrested by the liberation of the bowel, its tissues mortify, the coverings of the sac become inflamed, infiltrated with serum, and at last with pus. Crepitation may be felt, which arises either from the development of gases in the decomposing tissues, or depends upon their escape from the alimentary tube; and finally gangrene of the skin ensues, and through the opening caused by the separation of the slough the contents of the intestinal tube escape, and an artificial anus is established.*

In cases of entero-epiplocele, after the reduction of the intestine by the taxis, the irreducible omentum sometimes becomes inflamed, and, together with the sac and its coverings, suppurates, sloughs, and becomes detached in the usual manner; the patient subsequently making a good recovery.

Constitutional symptoms induced by the morbid state of the alimentary canal above the hernia.—Many of the general or constitutional symptoms of strangulated hernia are excited by the morbid condition of the part of the alimentary canal above the portion herniated, more than by the state of the hernia itself. The simple obstruction to the passage of the stercoraceous contents of the tube induces severe constitutional disturbance; an illustration of which is afforded by cases where the canal is crossed by fibrous bands, or directly compressed from any other local cause.

There are so many functions disturbed by the retention of the hernia in the sac, that it is very difficult to assign the constitutional symptoms to any single circumstance, such as the constriction of the hernia. Yet it is almost impossible to deny

* John Hunter writes: 'It is very curious to observe in hernias, that while the gut is in the sac and alive, no inflammation takes place on the sac or intestines; but the moment the gut becomes mortified or dead, the stimulus of a extraneous body takes place immediately; an outlet is then endeavouring to be made by the inflammation and suppuration of the sac forming an abscess in which matter, with the contents of the gut, is brought to the skin. While this is going on, the sound gut within the abdomen, where it passes into the ring, adheres to those rings all round; so that when the abscess is formed, burst opened, and the mortified parts sloughed off, these ends of the gut open into abscess, and not into the cavity of the belly.'—*Hunterian Ms. Descrip. Cat. the Path. Specimens in the Mus. Roy. Coll. of Surgeons of England*, vol. iii. p. 117.

that some very marked influence arises from this single cause alone; for what change is more marked or striking than the cessation of vomiting immediately ensuing upon the liberation of the bowel? The hernia is indisputably the primary and exciting cause; but much of the constitutional disturbance must be referred to those morbid phenomena which progressively arise as the result of mere obstruction of the alimentary canal.

We may consider these effects of obstruction under two heads. First, those produced in the portion of the viscera within the abdomen; secondly, those excited in the nervous, respiratory, and vascular systems.

Effects of obstruction.—That part of the alimentary canal which is between the stomach and the hernia becomes by degrees distended with flatus and fluid of a dark-brown colour. The mucous membrane is deeply injected, sometimes of the colour of blood. The serous membrane is also red, from vascular turgescence; and often patches of a deep-red tint produce mottling of its surface. After long protracted distension, the contractile tissues of the small intestine seem to lose their function, which in some cases they never regain. When obstruction occurs to a part of the colon, the patient may die from perforation of the coats of that bowel, or even of the cæcum. The portion between the hernia and the anus, whether small intestine or large, is contracted; the mucous and serous membranes are generally pale. The bowel often contains a little tenacious mucus.

Results of peritonitis are seen after death in the form of shreds of plastic lymph adherent to and lying between the coils of the bowels; or a large quantity of sero-purulent effusion occupies the peritoneal cavity.

Constitutional symptoms of strangulated intestine.—The constitutional symptoms, which are regarded as indicative of strangulated intestine, when associated with a local tumour in any site of a hernia, become marked and highly characteristic after vomiting has once commenced.

That train of symptoms seems to be mainly due, first, to vious irritation starting from the herniated viscus, and afterwards to the interruption of the passage of the stercoraceous matter; for any circumstance producing mechanical obstruction causes very similar results. Thus, if any part of the ali-

strangulated, especially if certain local indica-
to mere swelling, also existed.

Usually, the first symptom the patient com-
ing. Upon careful inquiry, however, it is
especially if the case be one of old hernia, that
pain had been experienced at the site of the
for some hours before, it was observed to be
Nevertheless, so little trouble is sometimes
tumour, in the first instance, that patients of
protrusion of the hernia to the effects of the
act of vomiting. Occasionally, too, soon after
in the morning, the patient complains of
rapidly succeeded by vomiting, without being
of the existence of any hernial tumour. This
history of females, who perhaps have been to
the day before. The same likewise happens
knowing themselves to be ruptured, and ha-
truss, designedly conceal the fact. So little
is sometimes caused by the hernia, that the at-
and medical attendant being entirely engrossed
vomiting alone, both remain in ignorance of
the hernia.* Upon enquiry it may be elicited

* *Importance of searching for a hernia in certain cases.*
fore, of the fatal results which almost inevitably ensue if it
be not speedily liberated, it is the paramount duty of every
called upon to afford relief in cases of continued vomit-

possessing ordinary intelligence, that before the nausea commenced, a sudden sensation of uneasiness was felt in the abdomen, especially about the hypogastric region; that this was followed by a repeated desire to go to stool, with tenesmus, and that nothing but a little flatus escaped.

Nature of the vomit.—The composition of the vomit varies at different stages of the illness. At first, it usually consists of substances last swallowed; next, principally of yellow bilious fluid, which after a time changes to a greenish hue; until at last the colour changes to brown, and resembles more or less exactly, especially in odour, a mixture of feces and water. It is then termed *feculent*, or *stercoraceous*.

During the active stage of vomiting the sufferer is unable to retain anything in the stomach for any length of time. But as the extreme state of prostration of the nervous system becomes more and more imminent, and also if the narcotic effects of opium be induced, the irritable condition of the stomach subsides, and may cease altogether. The surgeon must not, however, be thrown off his guard by this apparent calm. His anxieties must not cease in consequence of this composed state of the patient. On the contrary, it betokens a most dangerous depression of the vital powers, a degree of exhaustion from which some persons never rally.

The stage of prostration is now established. The pulse, which during the earlier hours of the attack was more full, and beat more frequently than normal, and which, as the vomiting continued, became weaker and more rapid, now beats slowly and with little force. The surface of the body is chilled; the hands and feet are blue, their integuments shrivelled; the aspect of the countenance is one of anxiety and distress; a peculiar expression as of suddenly increased age is very characteristic; the muscular system has lost its tone; the tongue is dry, furred, and frequently brown; and the secretion of urine falls below the normal standard of quantity. At this stage the hernia is sometimes reducible. The most gentle manipulation suffices. This occurs more commonly in inguinal hernia than femoral. It probably depends upon the relaxed condition of the abdominal muscles, and the removal of the constriction they once exerted around the mouth of the sac; in the same manner as chloroform exerts a beneficial influence. Death occasionally takes place at this period of the attack. This fatal issue is

attributable to the extreme prostration of the nervous system, induced by protracted vomiting; for in after-death examination of such cases we do not discover any organic lesion to cause it.*

The next stage, characterised by distinct symptoms, is after the development of peritonitis. But this does not always occur over so large an extent of surface as to excite grave apprehensions for the result. Peritonitis is often purely local, and confined to a small area around the region in which the hernia is situated. Perhaps the local inflammation is only just sufficient to produce adhesions between the orifice of the sac and the protruded intestine. However, when general peritonitis is developed, the usual symptoms of the disease are well marked. Hiccough is especially regarded as an unfavourable indication. Peritonitis, in the most severe form, arises when the walls of the bowel have been perforated by ulceration, thus allowing their contents to escape into the peritoneal cavity. Intense abdominal pain immediately follows this accident, collapse supervenes, and death speedily releases the sufferer.

The last stage is collapse: a condition in which the vital powers scarcely seem to exist any longer; the sufferer, although not physically dead, yet seems to linger on the threshold of that state. The patient may continue some hours pulseless; the extremities cold, and the breath also; the facial aspect deathlike; the power of speech but slightly impaired; until a sudden change comes over the features, and life has departed.

We have before described the pain felt in the hernial tumour. Besides this there is a peculiar sensation of tightness or constriction referred to the umbilicus. This is a highly characteristic feature of a strangulated hernia. The patient describes the suffering as resembling the effect which would be produced by encircling the mesogastrium with a tightly drawn cord. To

* Mr. Obré showed a dissection of a case of entero-epiplocele illustrating this fact, at the Pathological Society (*Path. Trans.* vol. vii. p. 219). A woman, fifty-three years old, died after four days' strangulated bowel, without receiving medical advice. After death a small portion of intestine was found perfectly strangulated in the left femoral ring; it was intensely blue, but its tissues unaltered. The small intestines were preternaturally red, but not inflamed.

M. Malgaigne also reports a case of the same kind. Strangulation of the intestine had existed eight days, when the woman died. There was neither gangrene, ulceration, nor peritonitis. He then observes that we see that a strangulated hernia excites such an extreme depression of the vital powers that patients may succumb without any anatomical lesions appearing to produce that event.—*L'Union méd.* 1854, p. 248.

ometimes added a dragging sensation, extending from the epigastrium to the hypogastric region.

Regarded as a symptom of peritonitis, is an uncertain indication.

This disease frequently exists without much pain being experienced, and even pressure on the abdomen is without causing complaint. Pain is more particularly marked in those cases where peritonitis arises without ruptured bowel. If, however, fecal extravasation be the cause of the inflammation, then the pain immediately becomes intense, often being, indeed, pathognomonic of that occurrence.

Prognosis of strangulated hernia.—Hospital experience must be regarded as a fair criterion by which to judge of the results induced by the effects of strangulated hernia alone. The high rate of mortality observed in this class of patients must be considered as the result of the nature of the cases admitted into hospitals for treatment. In a large proportion of cases the prospects of saving life are passed before the patients even leave their homes. Men and women labouring under the most adverse circumstances, irrespective of the particular disease under observation, are, for the most part, the sufferers submitted to treatment. Their constitutional strength has been previously and irreparably reduced by habits of dissipation, places of residence, persistent recklessness, and usually of voluntary submission to almost every condition the reverse of conducive to health,—such are the cases hopelessly beyond cure. These facts alone are sufficient to account for the high mortality which occurs in our hospitals in cases of strangulated hernia requiring a cutting operation to liberate the bowel. Undoubtedly it is to the length of time which the bowel is allowed to remain strangulated, and to the delay in the performance of the operation for its liberation, that the high mortality must be generally attributed. This circumstance must be too strongly impressed upon the mind of the patient; the medical attendant must act too promptly in the employment of the only treatment which can avert such untoward results.

Mr. James of Exeter was evidently surprised at the frequency of strangulated hernia occurring in the hospitals of London when he observed that 'There can be little doubt that the majority of these cases (of hernia) were the victims of time; . . . this, I believe, is an error little imputable to the distinguished
IV. Y Y

surgeons by whom these operations were performed generally the unfortunate patient who is the cause of disaster.*

This statement very truthfully represents the state of the case, in some instances, without doubt,—in those, for instance, in which the patient, being conscious of the existence of the rupture, conceals that knowledge from his medical attendant, or refuses all interference beyond external or internal medical agents. It does not remove the imputation of culpability from him who, when the case is under his medical care, leaves his patient for several hours in suffering; nor can it justify the delay which is caused by a surgeon who, duly appreciating the nature of the illness, refrains from liberating the contents of the bowel, or asking others to do so, after appropriate measures have been unsuccessfully employed to return it by the taxis.

We have never known an instance of a patient dying as a consequence of the bowel being liberated at too early a period. We have had to operate upon many whose chances of recovery have been absolutely sacrificed by the inexcusable delay which has preceded the operation. We make this assertion after a large experience, extending over many years.


Why is it that the mortality attending strangulated hernia is so large? How does it happen that every writer on the disease and every hospital surgeon has to deplore the high mortality of patients subjected to the operation for the liberation of the bowel? The average annual mortality in this metropolis from hernia, as published by the Registrar-General, amounts to 148.† This average is calculated from the total number of deaths from this cause registered during the last thirty years. The highest number recorded is for the year 1862, 214; the lowest in 1850, 128. We may assume that the increase of the population, and greater accuracy on the part of those who register the causes of death, may account for the difference between these extremes,—42. Now, judging from the conditions of the cases which have come into hospital under our own hands, we believe we are correct in stating that the mortality may be reduced at least by two-thirds if the liberation of the bowel is effected immediately after the failure of judicious at-

* *Practical Obs. on the Operations for Strangulated Hernia*, p. 79.

† *Summary of Weekly Returns of Births and Causes of Death in London*, published by the authority of the Registrar-General, 1863.

place the hernia by the taxis, assisted by proper means, or soon after the vomiting of fluids regurgitant from the intestinal tube. I do not hesitate to pronounce this judgment after carefully ascertaining the facts of the cases which have fallen under our treatment. For example, in inguinal hernia: of eight cases, three were incurable in consequence of the condition of the bowel, irreparably diseased by long-continued constriction; two others were in a similar state from the bowel having been injured by violent compression employed to reduce it. Cases of femoral hernia are even still more fatal from the causes above mentioned. In twenty out of twenty-eight, the immediate cause of death was attributable to the disease of the

FIG. 267.



This woodcut represents the injured tissues of a knuckle of small intestine which had been violently contused during efforts made to reduce it. The dark shade between the serous and mucous membranes represents blood extravasated in the walls. The whole of the knuckle was gangrenous although it had been strangulated only a few hours. The lines of ulceration in the mucous membrane are distinctly seen. (Copied from a coloured drawing and preparation in Guy's Hospital Museum.)

bowel which had formed the hernia; this morbid state depending upon injury, over which the mere liberation of the bowel exerted but slight influence towards repair. For it should be remembered that the simple replacement of the viscus in the abdominal cavity does not ensure the restoration of its functions; the surgeon only replaces it in that situation where the reparative powers of nature are most likely to exert their influence with activity; and that if the tissues of the hernia when reduced are incapable of performing their functions, those of the intestine cannot be performed, and therefore death must ensue. In several of these cases the disease of the bowel was rendered still more surely irreparable by the pressure to which it had been subjected in the taxis; and in two instances the

damage inflicted by this means was so great as to have absolutely killed the entire coil of intestine which was in the sac, fig. 2.

But all the causes which produce the morbid conditions of the bowel above mentioned, and more fully detailed in preceding pages, can be prevented. They may not be always under the control of the medical attendant, nor amenable to surgical skill, but we confidently promulgate the doctrine that the salvation of life in cases of strangulated hernia entirely depends upon the liberation of the bowel at the earliest moment practicable, and the converse may be predicated, that the destruction of life will inevitably ensue in proportion to the length of time the constriction has existed, with its constitutional consequences and the prolonged and forcible taxis employed.

The vital importance of liberating the bowel from constriction at the earliest moment cannot be over-estimated. As upon the speedy accomplishment of this the salvation of life depends, little precipitate action may even be forgiven, so hazardous is the position of a patient with the bowel strangulated. But what is the risk attending the operation of exposing the hernial sac, dividing the impediment to the reduction of the hernia, and reducing it, even should the peritoneal sac require to be opened? Practically, none. In comparison with that of leaving the bowel strangulated, it is harmless.

To what cause, then, may we attribute the culpable negligence of those who leave the poorer classes of the community in this most perilous moment—in a condition replete with jeopardy, fraught with results hazardous in the extreme? It would seem that the embarrassment and the delay are caused not by surgeons being timid and slow to propose a remedy, but from their really being ignorant of the amount of mischief certain to arise by allowing the constriction of the bowel to continue.

We once heard a physician relate the following circumstance: Being asked to see a poor woman who had been vomiting for several hours, he discovered a hernial swelling. He suggested that a surgeon should immediately see the patient, intimating that an operation was urgently needed. The gentleman in attendance replied, 'But will it not be desirable to wait until the vomiting has ceased before the performance of the operation?'

Can any principle be deduced from the facts detailed, to serve as a guide in determining the moment at which the

Attempts to reduce the hernia by the taxis should be given up, only as hopeless of good results, but fruitful of pernicious results, and a cutting operation be urged as the only safe means to liberate the bowel from constriction? We may certainly assert this without fearing the accusation of too great precipitation: when the period has arrived at which it is certain, from the nature of the fluids vomited, that regurgitation of the contents of the duodenum and jejunum has taken place, delay in effecting the reduction of the bowel is certain to be attended with progressively increasing evils. If there be evidence of regurgitation from the ileum, the condition of the patient is still more hazardous. Even assuming that the case was first seen for the first time at this period, and that attempts had not been made to reduce the protrusion, delay is inadmissible. The administration of chloroform should be recommended, with an understanding between the patient and the surgeon that the operation should be immediately performed when its anæsthetic effects are fully developed, he fails to reduce the hernia by manipulation.

Desault, fully appreciating the injury too often inflicted by the hernia by the injudicious and violent employment of the taxis, wrote: 'Always think favourably of a case of strangulated hernia when the taxis has not been used.'

Not two years since I operated upon a case of femoral hernia in a woman who had been labouring under symptoms of strangulated intestine for fourteen days. This was carefully ascertained. The bowel was thickened and like a piece of leather. Taxis had never been employed, and this patient recovered perfectly.

Treatment of irreducible hernia. Reduction of the hernia by the taxis.

—The endeavour to reduce the hernia, which consists in replacing it from its abnormal position, and passing it through the orifice of the sac into the peritoneal cavity, by dexterous manipulation, is termed the use of the taxis. Of course, the desire upon the part of the surgeon is to accomplish this. But all attempts to replace the bowel in the abdominal cavity should be relinquished until the circumstances attending the special case under observation have been carefully ascertained and considered.

The principal circumstances to which attention should be directed are as follows:

1. The kind and variety of the hernia regarded in its anatomical relations.

which have taken place in it during this stage ; which it gives rise, if merely local or extending into men, with or without manipulation ; the condition of the organs ; its probable contents, so far as may be conjectured from the evidence, assisted by touch and sight.

5. The treatment already adopted by the patient, or other persons, before the observation of the surgeon.

These are all subjects upon which profound and serious meditation are urgently demanded.

Injurious effects produced by the taxis.—It is necessary once more that irreparable injury is frequently inflicted on the herniated bowel by violence, used at all stages of the disease. The danger of any mischief occurring by the taxis increases in proportion to the length of time the bowel has been strangulated. This necessarily arises from the viscus becoming less and less able to resist the pressure of the hand in consequence of a morbid state due to the stricture.

The principles by which the surgeon should be guided when he is desirous to reduce a hernia without the knife, or by the taxis, are these :

1. Before vomiting commences. It is evidence of bad judgment to abstain from manipulation of the tumour at this stage, and until other remedial means have been tried, with the object of forming a correct diagnosis of the nature of the case. The taxis, unassisted, often fails to replace the bowel.

ould be enjoined to lie on the back, the pelvis being elevated, and the knees flexed. Or, in other words, the abdominal muscles must be relaxed, by posture, as much as possible. Warm fomentations are to be disposed over the region of the mouth and neck of the sac, especially in children; and the tumour, if it be a scrotal hernia, supported, or never allowed to pendent, thus dragging on the neck and mouth of the hernial sac by its own weight. The lower bowel may be induced to empty itself by administering an enema of warm water with a long flexible tube, and tincture of opium should be given in doses suitable to the age of the patient. It is desirable, however, that a full dose be given to adults: and with this view thirty drops diluted with camphor-mixture is not too large a quantity. A few hours may be allowed to pass away in order to afford time for this treatment to take effect, unless urgent symptoms arise.

Some patients cannot maintain the posture just described, it might be highly injudicious to enforce it. The principle of relaxing the abdominal muscles may then be carried out by allowing the patient to lie on the side of the trunk; but even in this position the hernial tumour must be carefully supported.

Under the influence of this treatment the patient very often falls asleep, and on awaking finds that the protrusion has retracted into the abdominal cavity. Now, should that happy result not have occurred, it will be desirable to manipulate the tumour, gently, which perhaps has become soft and flaccid. This must be done upon some fixed principle. A successful result will not accrue unless the pressure employed be directed with intelligence. The anatomical relations of the protrusion to the mouth of the sac and the apertures in the abdominal wall must be carefully considered; and the course or direction which the hernia takes in its descent should be strictly regarded. All efforts undertaken with a view to its reduction. The patient must be disposed in a posture to relax all the abdominal muscles which can in the slightest degree contract around the mouth of the sac. This part of the sac and its neck should be held, as far as practicable, with the fingers of one hand, whilst the fundus of the tumour is held in the palm of the other. This proceeding can only be adopted when the tumour is large.

The object desired by the manipulator is twofold: first, to relax the mouth of the sac; and, secondly, to diminish the bulk of the protrusion. The area of the orifice of the sac may be

the operator is enabled to exert a more direct influence on the hernia.

2. During the stage of vomiting. It behoves the surgeon to be always on his guard in the use of the taxis, which is coincident with a hernial protrusion. The longer, indeed, the vomiting has lasted, the greater the iritation and inflammation of the tumour. If before the commencement of the operation delicacy of handling be important, how much more so is the necessity for gentleness when the state of the bowels, as indicated by the symptoms, can be foretold! When vomiting has occurred, the tissues of the bowel were certainly not healthy, and the damage inflicted by pressure during the operation is in a great measure increased. After vomiting, especially if it has been prolonged for several hours, and it has become fæculent, the tissues have become morbid by the constriction to which they have been subjected. Ill-prepared, then, are they now to resist the effect of violence which will surely be to bruise and even to rupture them.

The measures which were employed in times past by the surgeon to apply the taxis with more effect, are now generally prescribed at this stage of a case of hernia, and need more than simple mention here. An exception is made in favour of opium. There is one period at which the administration of opium is attended with decided advantage. As we have been watching cases of strangulated hernia, we believe we have recently noticed that there is a short interval of time in which the bowels continue to function, and the patient continues until regurgitation takes place from the stomach. This occurs when the stomach is not yet

We believe the statement to be correct, that all other modes of treatment have been abandoned in favour of the administration of chloroform. Why? it may be asked. An analysis of the effects of those therapeutic agents formerly in vogue shows that they exerted an influence over the constitutional and local causes preventing the reduction of the hernia, similar to that which is now accomplished by chloroform more speedily, more certainly, with less risk to life, and under circumstances much more within control. The constitutional remedies were bleeding, the warm or hot bath, and tobacco-enemata; all of which exert a powerful but uncertain influence over the vascular, nervous, and muscular systems; on which account they cannot be employed without some risk arising from effects which are often uncontrollable and persistent for a longer period than is desirable. But the anæsthetic qualities of chloroform, together with the complete suspension of all voluntary muscular action induced by the inhalation of its vapour, renders this therapeutic agent of inestimable value, especially in those cases in which the abdominal apertures are under the influence of muscles controlled by the will, as well as stimulated to involuntary contraction by local irritation.

It is especially beneficial in all kinds of inguinal hernia. Its good effects are most striking in cases of hernia into the vaginal process of the peritonæum occurring in infants and youths, as also in those varieties developed slowly in adult life, and which have existed many years. In fact, the operation for the liberation of a strangulated inguinal hernia in middle age and elderly adults, is now rarely required in comparison with the frequency of its performance in past years. As soon as voluntary muscular contraction ceases, then the tumour gradually becomes softer or less elastic, under gentle and well-preconcerted pressure, smaller and of different shape, until at last the hernia escapes from the embrace of the mouth of the sac with a sudden jerk.

The sensation the surgeon experiences when it quits the sac and enters the peritoneal cavity is very peculiar and striking. A characteristic is this of the complete freedom of the bowel from constriction, that in its absence the expert surgeon immediately suspects that all is not quite right.

Usually, vomiting ceases after the reduction of the hernia at once. Occasionally, however, it may continue from the effect of the chloroform alone; but the pain in the abdomen is imme-

diately less, and quickly ceases altogether, especially that characteristic, dragging pain across the mesogastrium.

The patient should maintain the recumbent posture for a short time, with the abdominal muscles relaxed. We do not consider it necessary to place a pad and bandage over the mouth of the sac. Diet of a bland nature and semi-fluid consistence may be allowed in small quantities, and all stimulation of the alimentary canal by aperient medicine must be avoided. As soon as the injured viscus has recovered its functions, the bowels will act spontaneously. Stimulants should be administered according to the constitutional depression existing at the moment.

But great caution as well as discretion are necessary qualifications on the part of the surgeon when he manipulates the tumour. During the insensibility of the patient, there is considerable risk of bursting the intestine or lacerating the hernial sac. For twelve hours, or even perhaps for twenty-four in old protrusions, after the first vomit, the danger of using the taxis with sufficient violence to burst the bowel is not very great; but so much damage may be inflicted on its tissues as to preclude all hope of the repair of the mischief. When twenty-four hours have expired, during which time the bowel has been strangulated, it may be very easily burst by forcible pressure, especially if during the latter part of that time the patient has been vomiting fluids which have regurgitated into the stomach from the small intestines. If there be hiccough, the taxis is inadmissible.

The indications of burst bowel are very characteristic. If an enterocele, the hernia glides away from the pressure of the finger, and consequently the tumour disappears. This act is not, however, accompanied with that sudden and peculiar sensation which the replacement of an unburst bowel within the peritoneal cavity produces. The patient immediately complains of severe pain in the abdominal region; vomiting ceases, but retching and hiccough may arise instead; collapse rapidly supervenes, and death closes the scene in a few hours. In some instances inflammation and suppuration have taken place in the sac, and extending to its coverings, an intestinal abscess has been developed. A fistulous opening thus becomes established, which communicates with the interior of the alimentary tube. (See preceding remarks on Artificial Anus, p. 680.)

There is a period, then, 'when the symptoms of the rupture

ne very far, that it is imprudent to reduce it, even if ; but as it is impossible, perhaps, to tell when the ation of the gut is gone too far for reduction, it will, in be attempted while life exists, with the hopes of a Upon the other hand it may be asserted, or supposed, it is not reduced, the person must also die ; but this o certain as the other ; for the mortification of a gut does not kill—it only kills from its consequences ; and a material difference between a mortified gut out of r, and one within. The consequence of one within is death ; but the one without in general endeavours to y producing inflammation and suppuration of the parts, producing a fistulous orifice, or artificial anus.*

applications.—Considerable advantage attends the appli- f cold over the tumour. But when either ice or freezing s are used, their action must be widely extended over th, neck, and body of the sac. The local effects of cold iminish the bulk of the protrusion, by exciting contrac- its blood-vessels ; to retard inflammatory processes ; reducing nervous sensibility, to permit an advantageous ation of the tumour. Frigorific applications are valu- nts when there may be unavoidable delay in obtaining rm, or in liberating the bowel from constriction. The for their employment is after vomiting has commenced, consider they are more suitable to cases occurring in t than in the earlier periods of life. The advantages g their employment, however, are so trifling in com- with the injury certain to be inflicted on the bowel by d constriction, that it is safer to proceed at once to its n by operation, than to allow any great length of time e in the expectation of an advantage, which is in any btful. In practice, cold applications can only be re- n the light of very useful prophylactics. After marked ns of strangulated bowel have existed twenty-four old as a local therapeutic agent is scarcely admissible, , on account of the delay which must necessarily occur mportant moment.

distressing thirst, an accompaniment of continued , is greatly allayed by taking small pieces of ice into th.

The introduction of every kind of purgative medicine into the stomach must be scrupulously avoided in all stages of strangulated bowel. When swallowed, they are usually speedily vomited; and therefore it is sometimes suggested that under these circumstances they cannot produce a bad effect. This, however, is not always the case. We have seen fatal results ensue from diarrhœa quickly supervening after the bowel had been reduced, the exciting cause of which was referable to repeatedly swallowing purgative medicines during the stage of vomiting.

A purgative enema seems to exert a beneficial effect occasionally. But after one has been used in the early period of the attack, a repetition of this treatment is perhaps hurtful in the majority of cases.

Many other local and constitutional remedies have been employed with the intention of assisting the reduction of a hernia, and with occasional benefit. But the uncertain result which attends their employment; the disease, which is always progressive in the herniated viscus; and the risk to life, a certain accompaniment of continued strangulation of the hernia, and its attendant consequences—deter the experienced surgeon from persisting in entertaining the hope of reducing the protrusion by the taxis, after chloroform has been fully administered, and the hernia remains unreduced.

Herniæ have been replaced whilst completely reversing the ordinary position of the trunk, by keeping the head nearest the ground and the pelvis upwards. A patient may be placed in this posture (head downwards) by hanging over the back of a man, or over the side of a high bedstead or sofa, whilst the knees are at the same time flexed. Another method consists in encircling the mesogastrium with a folded sheet or round-towel, and at the same time drawing the contents of the pelvic region from below upwards, whilst the patient lies in a recumbent posture.

When the hernia is but recently strangulated, and it is an object to reduce it as speedily as possible, without a cutting operation, any method suggested which has been once successfully employed is perhaps worthy of a trial.

Injuries inflicted and accidents occurring in the use of the taxis.—These affect the following structures, separately or in combination; but we may consider independently—

those affecting the hernia; those affecting the sac; and those involving the tissues covering the sac.

The hernia, whether consisting of a hollow or a solid viscus, may be bruised, and its blood-vessels ruptured. Under these conditions blood is extravasated into the tissues composing it, producing either patches of ecchymosis, or, when an enterocele exists, layers of effused blood appear between the different coats of the viscus.

The tissues of the bowel being most delicate, they are more commonly subjected to irreparable injury by violent taxis than the omentum. The presence of this structure, perhaps, often tends to prevent the serious mischief just described. The degree of irreparable injury inflicted on the bowel cannot be accurately estimated by the length of time the symptoms of strangulation have existed. Much rather does it depend upon the violence and prolongation of the attempts employed in the taxis. At any moment during the first twenty-four hours following the first act of vomiting, the entire portion or coil of bowel in the sac may be so disorganised by pressure, that its vitality is entirely destroyed, and so literally killed, that nature throws off the part by the processes usually attending the separation of living from dead tissues. Under these circumstances, the entire calibre of the canal is divided, and an artificial anus is formed: or death takes place in consequence of extravasation of the stercoraceous material into the peritoneal cavity. Sometimes the tissues of the bowel are cut by pressure against the orifice of the sac, or the structures surrounding it; as in femoral hernia, against the free edge of Spigel's ligament. (See Fig. 267.)

After the expiration of twenty-four hours, the constriction to which the bowel has been exposed having led to inflammatory action and softening of its textures, there is great risk of bursting the tube. This rent usually occurs at the convex border of the gut, at the farthest point from the mesentery, and corresponds with the direction of the circular contractile fibres. Another effect of violent compression is to cause the mixture of blood and coagula with the serum in the sac, and flakes of lymph or puriform effusion arise from the same injurious interference. The local and constitutional indications of these injuries have been described in a preceding page.

Injuries of hernial tumours by accidental violence.—We may

here briefly describe the treatment to happen to hernial tumours from during the pursuit of the ordinary afflicted with reducible as well as reader will derive more information possible to introduce here, by the paper, entitled 'On the Proceeding Injured Intestine from a Blow on a H Aston Key.*

A simple contusion is the mildest the signs of more severe mischief then be returned. It is important abstain from bodily exertion, and th in a state of repose. If the violence sufficient to cause inflammation and slightest indication of any such s watched with the most anxious ca ment seized when an incision into escape of the contents of the alimen ceeding avoid the risk of extravasatio

But when there are plain indicat of the violence has been to lacera chance was contained in the sac at inflicted, time should not be was measures; but the hernial sac m thus a ready means of escape be r fluids. The symptoms indicating are described in Vol. II. p. 670
INTESTINE.

Another variety of injury of a h patient has an irreducible epiplocele. to the mouth of the sac, and entire whilst the body of the sac remains cavity. A contusion of such a tu and suppuration within the sac. C may be experienced in precisely di ture of the disease.

Injuries of the hernial sac.—There which produce important effects

* *Guy's Hospital Reports*, 184

hernia. Both of them render a case of strangulated hernia exceedingly complicated and embarrassing. The violence causing the mischief may be applied to the tumour either by the sufferer or another person; but considerable force is requisite to rend or tear the tissues composing the sac, or to separate it from its surrounding connections. These injuries more frequently happen in association with long-continued pressure than after short trials of the taxis. The two varieties are as follows :

A. By displacement.—The sac is detached, to a greater or less extent, from the surrounding structures. Those parts of the sac called the orifice and neck are more frequently affected by this injury than any other portion. They become detached from the inner surface of the internal abdominal fascia. Together with these, that portion of the parietal peritonæum immediately circumjacent to the sac's mouth is also severed from its attachments, and in this manner a pouch is formed within the fascial membrane of the abdomen, into which the hernia may be forced; and being retained therein, is lost to touch and sight. Sir Charles Bell has recorded a case which illustrates this injury.*

It is stated that the entire hernial tumour may be pushed within the abdominal walls, in a mass, whilst the hernia is still strangulated by the orifice of its sac. This injury was first described by French writers, under the appellation of '*réduction en bloc.*' However, more recently, examinations made after death indicate other lesions to be the probable causes of the disappearance of the hernia, in some cases; and certainly afford conclusive evidence that such an accidental disposition of the sac, when strictly examined in an anatomical point of view, must still be regarded as a very rare occurrence indeed. We refer the reader to Mr. Luke's paper for further information.†

In offering the following explanation of many of these cases, we court the inquiry of future observers, to support its correctness or to prove it to be erroneous. The tissues of the scrotum are very loose, and readily change their position. Every observer must have noticed the variable length of the spermatic cord, between the external abdominal ring and the testis, at different times. The hernial sac, attached to the

* *Lond. Med. Gaz.* 1828, vol. i. p. 485.

† *Med.-Chir. Trans.* 1843, vol. xxvi. p. 159.

anterior surface of the spermatic cord in a like manner. When the hernia occurs it extends lower than when it is empty, perhaps only just emerges from the inguinal canal. Now let its mouth at the internal abdominal ring, and the walls by the margins of the orifice be pushed outwards. The fundus of the sac attached to the scrotum is not on this account severed, but merely ascends towards the inguinal ring, and tending to contract, it is gradually within it with its walls in contact. The walls, rather thin, are not very recognisable. The course of the hernial sac has been often followed during the progress of an operation for the removal of the herniated gut, is attested by the reports of the reporters.

B. By laceration or bursting.—As long-sustained compression of the hernial sac, the serous membrane of the sac is rent, but the gut makes its escape through the aperture of the serous tissue. Its course outside the inguinal ring, by continued pressure; and detached from the neighbouring peritonæum, it forms a separate sac, that serous membrane and the internal ring.

The part of the sac a little below the mouth, usually styled the neck, within the inguinal canal, is most frequently burst apart. This injury is more commonly produced in inguinal hernia, the sac of which is a peritoneal process of the peritonæum, whether attached to the testis or not. It is inflicted when the patient is fully under the influence of chloroform, especially if he be also in a state of insensibility. The rupture of the hernial sac happens to be in the direction of the opening in its posterior wall, and pursues its course in that direction. The course which the hernia would pursue.

The indications of the accident are as follows: the tumour becomes flaccid, and the colour which alteration in its features is produced.

* *Med.-Chir. Trans.* 1859, vol. xlii.; see

ch the sac contained being squeezed through the rent into connective tissue around the sac; the bulk of the tumour diminishes as the pressure is continued, until at last very little, if anything, can be felt, but the surgeon has failed to experience that sudden jerk so characteristic of the escape of hernia from the gripe of the mouth of the sac, as it enters abdominal cavity; and if he have had much experience of reduction of hernial protrusions, doubts will arise in his mind as to the probable direction this one has taken. After the effects of the chloroform have passed away, all the symptoms of strangulated bowel recur, and perhaps with increased force. In the tumour itself may reappear, and recede on the application of slight pressure.

Now all these circumstances are highly characteristic of the accident that has occurred; and there remains but one proceeding to be adopted immediately. It is this. The hernial sac must be exposed and opened. Perhaps it may appear to be very tight; and even the finger passed upwards and along the internal division of the sac enters a cavity through a well-defined aperture, in which intestine is felt. This abnormal aperture may be mistaken for the internal ring and the cavity into which it leads, that of the abdomen. Acting upon this belief, however, would lead to the commission of a fatal error; one which, if detected at the moment, will surely compromise the life of the patient. An effort must now be made to draw the bowel out, if it does not come forth spontaneously; and when this is accomplished, the true mouth of the sac will be discovered by passing the finger upwards along the anterior surface of the entry. By the orifice of the hernial sac the protrusion is usually constricted. The constricting tissues, therefore, require to be cut; after which operation, the exercise of great care and attention is needed to prevent the entrance of the hernia once more into the abnormal space outside the peritoneal cavity. The salvation of life depends upon the return of the protrusion through the natural orifice of the sac, considerable freedom in the use of the knife is justifiable, if the attainment of the red end is thus facilitated, and the risk of failure in doing so thereby removed. All cases of this description not only give rise to great embarrassment and difficulty, but they are associated with very unfavourable consequences and results.

Injuries to the coverings of the hernial sac.—A very short
 . IV. Z Z

description will suffice to indicate the injuries inflicted on the tissues by the unjustifiable violence in attempting taxis. We have seen the result of the rupture of the integuments; extravasation of blood; infiltration into the subcutaneous tissue; inflammation of all the structures of the scrotum; suppuration, sloughing, and phlegmonous inflammation of the neighbouring regions. Those local measures and ordinary topical measures suited

The disappearance of the hernia from one sac into another ;—intra-abdominal. As to those rare cases of inguinal hernia with an intra-parietal sac. An intra-parietal sac in these cases consists of a peritoneal sac which passes along the inguinal canal and is lodged in the walls of the canal. In some of these curiously developed sacs the peritoneum is so paper-thin that the hernia is strangulated by the ventral or the scrotal division, it may be reduced, and thus, the tumour disappears. The operation is not employed with the utmost care and thus, the tumour disappears. The constitution of the patient indicates that this is not the case of the abdominal muscles, or in the case of the abdominal muscles, the hernia will sometimes reappear after the liberation of the bowel is accomplished. The operation to accomplish this end is to pass the bowel from one sac into the abdominal cavity.

The operation for the removal of the bowel, or the liberation of the bowel. The point in the treatment of every case is reached when the impediment to the passage of the bowel must be removed by a cutting operation after the failure to reduce the hernia by taxis.

The part of the practical surgeon is to apply it; and, however important the patient, or creative of alarm the

It is his paramount duty to urge concurrence in this step, as one offers the surest prospect of the salvation of life.

For too many medical men the cutting operation seems to be reserved in the light of a last resource, and one which it is proper to use until the danger to the life of the patient appears imminent. But the sooner that this fatal error is eradicated, the sooner shall we discover upon what fallacious views it has been based. We believe that death resulting from a strangulated bowel would be a rare occurrence, if it were possible to return every strangulated enterocele within twelve hours after the commencement of the symptoms, even assuming that the cutting operation be required to accomplish that end in every case. For let this conviction be firmly impressed upon the mind, that death results not from the operation, but because the operation was not performed at the right moment.

Every medical attendant upon a case of strangulated hernia should reflect on these indisputable facts, that so long as the patient remains in that condition the life of the patient is slowly passing; that to his judgment and foresight has been committed the safety of the sufferer; and that upon his discretion, firmness, and resolution hangs the fatal issue. For it is no exaggeration to say, that each minute as it elapses carries with it the chances of recovery further and further away.

We hope that the mortality arising from strangulated hernia is not now so large as it once was.* Nevertheless, at the present it is excessive; and when we know, and all professing surgery should know, how surely the liberation of the bowel tends to save the life of the patient, why is it not done the moment all other measures have failed to replace the gut in the abdominal cavity? Some medical men do not seem to be sensible of, or to appreciate the vast amount of injury which is certain to accrue from the persistent vomiting, nor to value the complications derivable from the character of the vomit. They do not assiduously attempt to check the vomiting by administering medicines; but they utterly reject, at the suitable moment, the only means by which it is to be arrested.†

* Mr. Hey states, that when he entered upon the profession of surgery, now 30 years since, 'the operation for the strangulated hernia had not been performed by any of the surgeons in Leeds;' and he adds, that he 'lost three patients in five upon whom the operation was performed.'—*Practical Obs. in Surgery*, 3rd edit. 1814, p. 129. Are the results of the treatment of strangulated hernia more successful at the present day? We fear not.

† We may have expressed our views a little strongly, and thus laid ourselves

same time including the walls of the sufficiently large to admit a grooved director, the sharp edge of which must not enter the contents of the sac, but laterally. The tumor should be placed nearly flat on the surface of the instrument, all risk of injury removed. A grooved director having been introduced, the opening and firmly held against the inside of the tumor, indeed stretching its tissues over it, the director is then moved upwards and downwards, or towards its or

open to criticism and animadversion; for we cannot help being influenced whilst writing these lines by an occurrence of the kind at the time of doing so. A fine woman for her age, who had been known to be the subject of strangulated femoral hernia, had persisted for between ten and eleven days. At the end of the hospital 'for the operation,' which necessitated a journey, she arrived cold and pulseless, though intelligent; the operation was performed, she was in fact moribund. As soon as practicable the operation was effected. The escape of purulent fluid from the tumor was characteristic of the condition of the peritoneal cavity. The fluid applied to the surface of the body and stimulants administered expired in a few hours after admission. Yet, during the operation medicines had been prescribed to arrest the vomiting, and employed, and the marvel was, as a friend who accompanied her, that she reached the hospital alive. What other result could have been anticipated? *Ten days'* vomiting, and starvation, surely enough to destroy life at the age of seventy-one. She had a perforating ulcer at the point of junction of

sually the escape of more or less serum is a sufficient indication that the hernial sac has been opened. But the operator should not be thrown off his guard by an absence of serum, when the sac has been cut, any more than by seeing a flow of blood when it has not been even reached. The latter somewhat unusual occurrence is due to the development of a cyst upon the peritoneal tumour, and is at once explained by passing the finger into the circumscribed cavity. For the variable conditions of the disease, see page 683.

When an enterocele is under treatment, the intestine is brought into view as soon as the sac is opened; but in an enterocele it often happens that the omentum only is exposed. The surgeon then carefully raises this structure, for the bowel lies underneath it in the majority of cases. It is wholly enveloped by it in some instances, when that arrangement termed an omental sac exists. In this case the omentum requires to be fully unravelled. We employ the last word in order to direct the use of much care and caution in the division of its vessels.*

The condition of the bowel should be now carefully ascertained, for by the morbid state of its tissues the surgeon is reminded upon the propriety of placing it within the peritoneal cavity, or allowing it to remain outside of the abdomen. Multiple adhesions of the bowel to the body of the sac may be easily broken down; but the greatest caution is required in the treatment of the hernia where it lies within the orifice of the sac, as by rough manipulation its tissues may be torn, when extravasation of the contents of the tube into the peritoneal cavity will take place. If the bowel is ruptured, it must be sutured to the edges of the wound by sutures.

To reach the impediment to the reduction of the hernia, the tip of the index finger is now directed upon the anterior surface of the mesentery, towards the mouth of the sac. Should the end of the finger pass freely through the ventral orifice of the sac into the peritoneal cavity, when introduced with great gentleness, an attempt may be made to reduce the hernia without cutting this part of the sac. In the manipulation of the bowel great delicacy is required. The surgeon gently

A very interesting paper on these 'omental sacs,' written by Mr. Prescott Pett, is published in the *Med.-Chir. Trans.* 1844, vol. xxvii. p. 282; and another, on a case of scrotal hernia, with compound omental sac, in *Path. Trans.* vol. ii. p. 98.

the anterior surface of the hernia, its d the index finger of the left hand. It s the abdomen sufficiently deep to be e striction formed by the tissues around further. Careful examination is nec ascertain that the bowel does not ove director, nor lie between it and the ante and orifice of the sac. To avoid all risl when that viscus is quite clear of the g instrument should be firmly pressed ag are about to be cut. A bistoury, specia operation, is gently glided along the , and a slight resistance is usually en through the constriction, which, howev edge reaches, and in its passage divid cause the impediment to the reduction .

When those tissues are firm and rig or a peculiar sensation is felt, on divi the cutting of thick leather. The op index finger to examine the opening he sufficiently large to admit the passage attempts its reduction ; if it prove st bistoury may be passed along the fing creased. The impediment to the reduct thus entirely removed, its replacement cavity is easily effected.

But we must revert to a few subjects merely alluded to in the preceding description of the operation.

Of opening the hernial sac.—When the anatomical relations of the hernial sac were carefully examined, it was discovered that the impediment to the reduction of the hernia was not in instances caused by the narrow passage of the ventral orifice of the sac alone. Careful dissections showed that those structures which lie in close connection with the outside of its mouth and neck offered considerable obstruction to the retrograde passage of the hernia through those parts of the sac.

For the last hundred years surgeons of eminence have applied these anatomical facts to treatment, and many distinguished operators have dwelt upon the advantages to be derived from reducing the hernia in the abdomen without cutting open the peritoneal sac. On the grounds above alluded to, they have devised that mode of performing the operation for a strangulated enterocele called ‘the division of the stricture without opening the sac,’ or ‘external to the hernial sac.’

The advocates of the operation believe its most favourable points to be as follows: the peritoneal cavity is not opened; the peritonæum at the mouth of the sac is not cut; the diseased intestine is not exposed to the influence of the atmospheric air, nor to the direct contact of the fingers of the operator; nor is there risk of hæmorrhage into the peritoneal cavity so great, nor would a large vessel be accidentally cut. They consider, then, that the operation itself is not attended with the same amount of danger as when the peritoneal cavity is opened, its parietal reflection cut, and the inflamed bowel exposed.

The simplicity of the proceeding commends the operation highly; for the wound inflicted is very little more than an incision through the integuments, and its advantages must have seemed paramount at a time when the dangers of the operation were exaggerated, and it was assumed that patients with strangulated hernia died from the effects of the operation itself.

Bearing in mind, however, that death occurs from the constitutional and local effects of long-continued strangulation of the bowel and its injury by forcible taxis, and because the bowel is not liberated by the operation early enough, the advantages of any particular mode of operating, especially of this one, are perhaps not quite so great as at one period many surgeons were inclined to admit. The method of performing

guard the risk to be great of reducing regulated by adhesions, omentum, or even hernial sac itself.

Next, pointing to statistical facts, and of the two operations as performed in hospitals,—in those where the preference the sac in all cases, and in those in which pursued,—they maintain that the greater at those institutions where the sac is inci

It is, however, quite clear that we are facts upon which to institute a safe comparison the success of the one practice or the results of the cases under treatment, in themselves worthless, because the incidents in any two never precisely alike, and accuracy, as absolutely required to arrive at the truth

We believe the dictum to be equally surgeon says that in every case the sac another directs that it must not, except proper practice consists in making a judgment cases : viz. those in which the operator fixed principle ; those in which he does ing to his judgment and experience, then do it.

In many of the metropolitan hospitals to the sac is performed in suitable cases.

curs of frequently operating on cases of strangulated hernia, will be guided by his own judgment as to the propriety of opening the hernial sac in every case, or of leaving it intact in selected cases. But if we may be permitted to enunciate the principle upon which inexperienced surgeons may act with safety, it would be this. In all those cases in which the surgeon would deem it safe to return the hernia by the taxis, if it were practicable, he may do the same thing after he has removed the impediment to the reduction of the hernia, by cutting those tissues lying outside the sac which are the cause of that impediment. But in those cases in which the symptoms are suggestive of an aggravated morbid condition of the bowel, and on account of which it would be improper, nay hazardous, to attempt to reduce it by the taxis—in those the sac must be opened.

In making a suitable selection of the cases for the first plan, great discrimination is therefore required. We regard those cases to be favourable for the operation in which the symptoms of strangulation have existed but a few hours, and when they have not been very severe: when the vomiting is not stercoraceous, or the patient very prostrate; when the tumour is a simple enterocele, and when it has escaped forcible attempts at reduction. Of course no cutting operation of any kind will be attempted until after the failure of well-applied taxis to reduce the hernia.

On the contrary, the indications for incising the sac are: a long continuance of the strangulation of the hernia; inability to empty the sac completely; the persistence of stercoraceous vomiting; the prostration of the patient; the compound confirmation of the tumour (it being an entero-epiplocele); and after repeated, protracted, or forcible taxis has been used.

Guided by these facts, the number of cases in which the operation of division of the impediment to reduction of the hernia external to the sac may be regarded as preferable to any other, becomes very limited. But we cannot refrain from expressing our conviction, founded upon the results obtained in those cases in which this operation has been carried out, that it is easy of performance, free from the risks attributed to it, unaccompanied by some of those accidents which occasionally occur when the sac is opened, and most favourable to the rapid recovery of the patient. It must, however, be remembered that these are select cases, and belong to a class most favourable for recovery; and we cannot deny but that they would probably be successful, even should the peritoneal sac be cut open.

4 A grooved director, specially adapted for inserting the tissues forming the impediment to the reduction of the hernia.

5. A bistoury of peculiar construction, with which to divide those tissues.

6. Retractors.

The *director* (4) and the bistoury (5) are the only instruments requiring special notice.

The *director* has been constructed after a great variety of forms—straight and curved; long and short; narrow with wings or lateral projections to protect the back of the knife; with and without a handle affixed; with a point or end or beak, with the groove terminated, near the stop, or without such a stop. That form of the *director* invented by the late Mr. Aston Key has been employed with our observation many years with advantage; and we therefore, to recommend its use. It is usually known as the hernia director.*

The *bistoury* used to divide the tissues which constitute the impediment to the reduction of the hernia, wherever it is situated, has been made in a great variety of shapes and breadths, and usually with some contrivance, fixed, to prevent its end and cutting edge injuring the tissues. It being impossible to describe every variety of the *bistoury*, perhaps it will suffice to state the characteristics of

culars the two instruments resemble each other. The metallic part extends about $3\frac{1}{2}$ inches beyond the handle. Its sides are $\frac{3}{8}$ of an inch broad where it unites with the handle, and they gradually taper towards the end, where the breadth is $\frac{1}{2}$ of an inch. The blade terminates in a button or blunt end. At the terminal $\frac{3}{8}$ of an inch, the sides of the metal are bevelled off, and one edge is made to cut. This sharp edge should be slightly concave. All the other edges and surfaces of the metallic shaft are carefully rounded off and polished. Sir A. Cooper's hernia-knife has a cutting edge $\frac{6}{8}$ or $\frac{7}{8}$ of an inch long, beyond which extends a blunt terminal portion $\frac{3}{8}$ of an inch long. The objection we entertain to this knife relates to this long blunt beak, which must be passed into the abdomen before the cutting edge reaches the tissues which require division.

When the index finger, of either hand most convenient to the operator, is employed as a director, its point is passed to the orifice of the sac, outside or inside; the bistoury is passed upon the anterior surface of its first phalanx, and the side of the instrument being pressed against it, lies in a depression on its soft parts. Thus the sharp edge of the knife is prevented from injuring the bowel. The end is then dexterously insinuated beneath the tissues to be cut; and as they slowly yield before it, the finger follows the direction of the knife, and the operator judges of the size of the opening he is making by the facility with which his finger passes into the abdomen.

When a metallic director is used, the bistoury is gently carried along the groove of the instrument.

Of the treatment of the strangulated intestine.—The primary object of every surgeon who has a case of strangulated enterocele under his charge, should be the return of the intestine into the abdominal cavity as quickly as possible. With this object in view, 'the operation,' as it is termed, is undertaken after the failure of all other means.

But there are certain morbid states of the bowel which preclude the operator from doing this, after the liberation of the viscus from the constriction to which it was subject. The morbid conditions having been already described (p. 677), we need only briefly refer to them here, in order to direct their treatment.

Commencing, then, with the worst condition, in which the entire knuckle of the intestine in the sac is dead, mortified, or

sphacelated. Before the operation, adhesions will have formed between the coils of bowel in the abdomen, as well as between them and the parietal peritonæum in the vicinity of the mouth of the sac. It would be very injudicious to interfere with the processes of nature to break down these adhesions. They are the result of peritonitis; and this form may be termed protective, for its effects, especially the adhesions, may possibly prevent the extension of the peritonitis to the mischief which would certainly accrue if the contents of the alimentary canal escaped into the peritonæum.

It is sometimes difficult to find the end of the upper part of the alimentary tube, from which the stercoraceous matter escapes; meddling with the processes of nature is not desirable, and therefore, unless the perforated viscus is easily reached, it is better to leave the parts as they are, exposed. The edge of the wound should be left free and open, in order to allow a ready escape for the contents of the bowel. In this manner an artificial anus is established. (See Fig. 266.) Should the open end of the gut be accessible, one, two, or more sutures may be inserted through all the structures constituting it, and they may be thus fastened to the integuments.

If there be a small perforation in any part of the herniated viscus the result of disease, there are generally adhesions to the wall of the abdomen, which prevent the extravasation of its contents into that cavity. When, therefore, the opening does not admit the whole calibre of the canal, the herniated viscus may be placed just within the abdomen; for sometimes the aperture will entirely close, even after stercoraceous matter has flowed through it for several days. Even a ligature may be tied around a small hole in the intestine with the best result. This treatment was successful in a case reported by Dr. Watson of Edinburgh.* (See Fig. 265.)

Should the intestine be accidentally wounded, the cut edges may be carefully drawn together by using the glover's suture, and returned just inside the mouth of the sac.

When the tissues of the herniated bowel are not perforated but appear to be in such a morbid condition as to induce the operator to fear that their restoration to a healthy state is not probable, although just possible, the hernia may be returned, if the adhesions at the neck of the sac are not strong. The la-

* *Edinb. Med. Journ.*; July, 1860.

Aston Key advocated this plan of treatment. He did this upon the principle that the abdominal cavity was the place in which the injured bowel was more likely to be repaired than in any other. 'The danger of abdominal extravasation will not be increased by replacing the injured bowel at the neck of the sac; for, should sloughing of its coats ensue, the slough may be walled-in by adhesion of the surrounding peritonæum, and fecal extravasation be prevented.'* When adhesions prevent this treatment it is better to divide the orifice of the sac slightly, and to leave the bowel in the sac. Under such unfavourable circumstances, patients occasionally recover without a fecal fistula. An interesting case of this kind is published by Dr. Watson, in the *Edinburgh Medical Journal*, March 1870.

In some cases the operator encounters great difficulty in reducing the hernia, even after the mouth of the sac has been freely enlarged. This depends upon the following causes:

1. *Its bulk*.—The hernia sometimes consists of a very large quantity of intestine, which is distended with flatus; and although a portion is reducible, the reduction of the entire mass is impracticable. After the failure of these attempts, a plan may be pursued which was adopted by Mr. Tatum with complete success.†

The bowel may be punctured, and the flatus allowed to escape. Mr. Tatum used a grooved needle for the purpose; we would suggest the employment of the filiform trocar and canula. This instrument may be obliquely inserted through the tissues of the bowel, thus forming a valvular wound.

2. *Adhesions*.—These may be *recent* and *chronic*. The latter are often formed long antecedent to the conditions for which the operation was undertaken. In the recent state the adherent surfaces are usually easily separated, whether they unite the coils of the bowel to one another, to the sac, or to the omentum. When this morbid condition happens to the viscus *in* the body of the sac, there is not much risk of the interference making mischief; but when the bowel is adherent to the mouth of the sac, the condition usually indicates an advanced morbid state of the viscus. It is an indication that nature has made preparations to prevent extravasation into the peritoneal cavity, in the event of the bowel bursting from the distension caused by

* *Guy's Hospital Reports*, 1842, vol. vii. p. 264.

† See INJURIES OF THE ABDOMEN, Vol. II. p. 666.

accumulations in that part of the digestive tube above the hernia. Meddling with such adhesions is attended with great risk of tearing the bowel. The safest procedure is to reduce the protrusion, if possible, after having enlarged the mouth of the sac; but even allowing it to remain in the sac, relieved from constriction, would be preferable to lacerating the bowel in the attempt to separate the adhesions, and so establishing an artificial anus. Recent adhesions to other structures may be separated with ease. Flakes of adherent lymph had better be left where found. By detaching them, the coats of the bowel may be torn; and they frequently overlies small sloughs.

Enterocèles of long standing sometimes become permanently united to a part of the sac, although very rarely. In such cases the orifice of the sac is usually large, and the chance of the bowel becoming strangulated proportionally remote. The peristaltic action of the canal proceeds as usual in the hernia, with occasional inconvenience arising from accumulations therein causing obstruction. Should an operation be required, two proceedings may be adopted: either the orifice of the sac may be enlarged, and the intestine left in the sac; or the adhesions of the viscera may be destroyed, and the bowel returned. The choice of these two plans might be decided by the condition of the viscus and the extent of the adhesions.*

3. *The depth and inaccessibility of the mouth of the sac.*—That part of the sac which lies in close relation with the general peritoneal cavity is sometimes so deeply seated, that considerable embarrassment arises not only in making its orifice larger, but in returning the hernia into the peritoneal cavity when that has been done. By employing an assistant, who carefully holds the opened hernial sac by its cut edges, to draw it forwards and outwards, or from the abdomen, its mouth is more fixed and slightly stretched. This manœuvre greatly facilitates the reduction of the protrusion.

Treatment of the omentum.—Omental protrusions or epiploceles rarely require the removal of the impediment of their reduction by a cutting operation. But after the return of the intestine, in cases of entero-epiplocele, it often becomes a question, what is to be done with the omentum.

The treatment is as follows: it may be reduced; it may be

* Relating to these cases, the reader may consult *A Treatise on Obstructed and Inflamed Hernia*, by Henry Stephens.

in the sac; it may be cut off; or a ligature may be tied round it.

The first proceeding would be generally chosen when practicable. In some instances, however, the great bulk of omentum, in its diseased condition, necessitates a choice between either leaving it in the sac or removing it.

There are some disadvantages in leaving the omentum in the sac: the fat becomes inflamed; it suppurates, and sometimes ruptures; the healing of the wound is consequently delayed, and the after-treatment of the case thereby protracted. Usually, however, the omentum shrinks up and contracts, and eventually the wound heals, but with more or less of a tumour at the abdominal aperture. This swelling may interfere with the successful application of the pad of a truss. Some surgeons, however, believe the remains of the omentum to be useful in picking up or plugging the ventral aperture.

The removal of either the whole or a part of the protruding omentum may be effected by cutting it away at once, or by applying a ligature around it near the orifice of the sac, and leaving the portion below the ligature still attached, or cutting off. Besides, there is the alternative of leaving the truncated end of omentum in the sac, or of replacing it in the abdomen. With the exercise of due care, omental adhesions to the sac may be safely separated, whether recent or chronic.

If the omentum be cut off, there is great risk of bleeding from its cut vessels when it is replaced in the peritoneal cavity. Even after every precaution had been taken to secure the blood-vessels, cases in which omentum has been excised have proved fatal from internal hæmorrhage alone.

It has been assumed by some writers to be injudicious to tie a ligature around the omentum, on account of the untoward results which may ensue. But the reader should refer to Vol. I. p. 663, where he will find the good results of this plan of treatment related by Mr. Pollock, as well as the mode of applying the ligature.

By the kindness of the editor, Mr. Holmes, we are able to give the result of the practice of removing diseased or superfluous omentum, after ligature, as performed at St. George's hospital. Of twenty cases of hernia in which the omentum was securely tied, a few died; but the notes of the after-death examination of them show the cause of death to have been, in all cases, independent of the ligature placed around the omen-

was removed, than in those in which it was

Cases of double hernia.—In patients suffering from protrusion on both sides of the body, or in difficult cases, it is sometimes difficult to ascertain which hernia is the most recent. The surgeon must, of course, employ every means to arrive at a correct diagnosis before commencing the operation. If he should be unfortunately have arrived at a premature conclusion, and having opened the sac of one side, find it a impediment to its reduction, nor indication of a second hernia, he is justified in cutting open the sac of the other side. And, as a precedent, we may refer to a case reported by Dupuytren performed a double operation, and cured the patient.

A man, aged forty, was the subject of double inguinal hernia. With much difficulty reduced both tumours. Well-marked hernial sac existed, although there was no tumour at the inguinal ring. Dupuytren opened the right inguinal canal first, because there was the most pain when pressed on that side. The sac was entered through the left inguinal canal and found a hernial sac containing a small portion of the small intestine. The orifice of the sac was very deep; it was cut; the bowel was reduced, and the patient recovered.*

Treatment of the case after the operation.—The effect of the reduction of the hernia is usually followed by vomiting, especially if the bowel had not been in the sac very long time, and neither peritonitis nor hi-

The peritoneal cavity is the rapid relief from pain across the mesogastrium.

It is manifest that no single plan of treatment, to the exclusion of every other one, can be urged as applicable to all cases of hernia, after the liberation of the bowel. As the disease occurs at every age, in every variety of constitution, and under individual peculiarities of the most diverse kind, so strangulation of the hernia may happen with the idiosyncrasies and opposite conditions of youth and senility, strength and decrepitude, health and disease, temperance and intemperance. It must, however, be carefully borne in mind, when treating a case of this kind, that we have to deal locally with an injured intestine, and a penetrating wound of the abdominal cavity; and that the constitutional powers of the patient are greatly reduced by continual vomiting, starvation, suffering, and alarm.

In some youthful, robust patients the tendency to inflammation becomes, at an early period, strongly developed, whilst, on the contrary, in the aged and decrepit little dis-ease may appear demanding interference.

The after-treatment has been conducted upon the most opposite principles. Surgeons of the highest eminence may be found amongst the advocates of the two methods, which may briefly stated as follows:

The first class are those who administer purgative medicines soonest immediately after the operation, and persist in their continuance 'until the canal is completely unloaded.' Calomel, extract of colocynth, sulphate of magnesia, and castor-oil, given by the mouth, or aperient enemata injected into the rectum.

The second class, relying almost entirely upon the restorative powers of nature and the influence which repose exerts on the restoration of injured tissues, depends upon diet and, when required, opium.

In selecting one of these plans of treatment for adoption, we give the preference to the last, since it seems to be most suitable to the majority of the cases occurring in hospital practice. In many cases we have allowed the patient to recover without giving any medicine whatever; great care, however, being taken that a mild and farinaceous diet, with milk, be only sparingly taken.

This necessity for repose, writes Mr. Key, in an injured state of the bowel it should be well to bear in mind, after the operation for strangulated hernia.

operation.

The bowels are often relieved spontaneously by the reduction of the hernia ; but when the patient is active, the question arises of the use of medicines. We have embraced abundantly the advantages to be gained by abstaining entirely from the use of the former, and by giving the preference to the latter plan, but when the patient is loaded, and any discomfort arises from the use of an enema of warm water, or gruel with castor-oil, or even sweet-oil, mixed with gruel, it results. Cases under our treatment have been cured, and even the wound has healed, before the patient is relieved, and not the slightest trace of inflammation from the constipation.

When there is much tenderness on the abdomen, in the neighbourhood of the hernia, the application of leeches is of great service, as often as required, but in relation to the patient.

If acute peritonitis be developed, it must be treated in accordance with the ordinary rules for the management of this disease.

Stimulants are often required soon after the operation, and should be given in small quantities. It is also required that all aliment should be enforced, that all aliment

disturbed by the cutting operation should be placed in relation to each other, and the divided edges of the integuments brought together by sutures. The number required may be left to the discretion of the operator, but no more need be used than sufficient to keep the upper two-thirds of the wound united. It is always advantageous to leave an opening at the lowest point to allow of the escape of blood and discharges. A piece of wet or dry lint may be laid over the incision, and a pad of wet lint is adjusted over it with a bandage by some surgeons. However, a bandage is not required in every case, nor is it on any account essential.

If, during the subsequent progress of the case, the connective tissue in the wound, the omentum, or the hernial sac should become adherent, the appearance of the integument soon betokens that the union by adhesion is hopeless. In that case the lowermost vessels, or all of them, must be removed, and the treatment to be adopted need not differ from that commonly employed in every inflamed, suppurating, or sloughing wound. The grand point is to keep a channel open for the free escape of the discharges.

The prognosis of a case of strangulated hernia.—Reflection on the facts already described will enable the surgeon to form a very nearly accurate prognosis of a case of strangulated hernia. But in order to concentrate the attention of surgeons on those incidents which more especially cause a fatal termination, we must briefly recapitulate the most important morbid conditions which call for special observation.

1. The condition of the tissues of the herniated viscera, and especially of the intestine. Their morbid state is too frequently produced by violence in the use of the taxis, by long-continued obstruction, or very tight strangulation of short duration. In such cases, then, the prognosis is usually unfavourable on account of the destruction of the portion of the bowel injured, and of its inability to repair the damage inflicted upon it.

2. The morbid state of the peritonæum, visceral and parietal, and of the tissues of that portion of the track of the alimentary canal between the stomach and the hernia. The existence of peritonitis antecedent to the liberation of the bowel cannot be regarded in any other light than a symptom of grave significance. Associated with it, the alimentary canal above the hernia is usually greatly distended, and its contractile fibres

prostration, therefore, precedes the reduction, and must be regarded as a most unfavourable sign.

4. Neither should we overlook the condition of the viscera of the thorax and abdomen. A hernia, so commonly existing in the chest, may enter into our hospitals. Over the influence of this, however, we can exert but little control. It is an important element in prognostication.

5. All these conditions are also influenced by the patient's social habits, general cachexia, and idiosyncrasies—features in a case to which no special remedy is given.

Now the lesson that we learn from these cases is, that states which bring about a fatal termination, may be avoided by the reduction of the herniated viscus into the abdomen. We must, therefore, to accomplish this with the utmost discretion, and be consistent with the requirements of the patient. We must carefully between those cases in which the reduction is liberating the bowel without the loss of the bowel, and in which some means are justifiably resorted to. By acting thus energetically upon the case until the risk of danger is averted, the surgeon may hope to rescue a large majority of the cases of strangulated hernia.

form his part. All matters relating to the operation only perhaps, reached perfection; but it would really appear, the condition in which patients are sent into the hospitals operated upon, that the greatest amount of ignorance is of the principles upon which an irreducible hernia is to be treated.*

PART II.

SPECIAL KINDS OF HERNIA.

The following classification of abdominal ruptures is arranged according to the anatomical divisions of the abdomen in which they appear:

the EPIGASTRIUM	{	1. Diaphragmatic.
		2. Epigastric.
the MESOGASTRIUM	{	1. Ventral (also in other regions).
		2. Umbilical.
		3. Lumbar.

Many patients who died after the operation had been performed by the Guy's Hospital survived the operation as follows:

1 only 17 hours.

5 survived 24 hours and less than 48 hours.

4 " 48 " " 72 "

1 " 72 " " 96 "

1 " 96 " " 120 "

3 " 144 " " 168 "

1 " 168 " " 192 "

3 survived longer periods, but with artificial anus.

1 died of bronchitis.

Causes of death were—

Artificial anus, &c., in . . . 3	Perforation of bowel in . . . 1
Strangulation in 5	Bronchitis and cachexia in . . 2
Peritonitis in 8	Neglect to seek surgical aid in 1

There is no little doubt that the lives of at least two-thirds of these patients were sacrificed by the delay in having recourse to the operation for the relief of the bowel; which, however, was performed as soon as possible after admission.

III. In the HYPOGASTRIUM

- | | | |
|---|-----------------------|------------------------|
| { | A | 1. Above Poupart's lig |
| | | inguino-scrotal; c |
| | B | 2. Below Poupart's lig |
| | | femoral. |
| | C | 3. Through the aper |
| | | the pelvis; in fr |
| | | neath the ramus |
| | pubes, obturator. | |
| | 4. In front—beneath t | |
| | of the pubes, peri | |
| | 5. Pudental. | |
| | 6. Vaginal. | |
| | 7. Behind—through t | |
| | atic notch, ischia | |

HERNIÆ IN THE EPIGASTRIC REGION.

Diaphragmatic hernia.—Of these cases there are three first, that in which the diaphragm muscle becomes str consequence of a loss of tone, and is pressed up into the cavity by the contents of the abdomen; secondly, that i in consequence of congenital defect, some of the visc through the aperture into the thorax; and, thirdl some of the abdominal viscera pass into the chest the natural apertures in the diaphragm, which have stretched.

Epigastric hernia escapes in the angle bounded by tl lages of the first false ribs, the apex of which correspon appendix of the xiphoid cartilage. M. Malgaigne has infant with a tumour in this region which swelled i time any effort was made with the abdominal muscl sac of this hernia is generally more movable than th umbilical, and the mouth through which the hernia en usually large. The reduction of the protrusion is t easily effected. A suitable bandage is required to pre escape of the hernia, which sometimes consists of a p the stomach, or arch of the colon and omentum.

* Exclusive of cases in which the abdominal viscera pass into through a laceration in the diaphragm. Such cases are treatd INJURIES OF THE ABDOMEN, Vol. II. p. 637.

HERNIA IN THE MESOGASTRIC REGION.

Ventral hernia.—Of this hernia there are two kinds: first, those which occur in the linea alba; and, secondly, those which pass through other parts of the parietes, except the so-called rings.

Those of the first kind usually escape in the linea alba between the umbilicus and the pubes. They are produced as the consequence of general relaxation and stretching of the fibrous tissue in the middle line of the abdominal walls after they have been greatly distended during pregnancy or ascites. We have seen hernia consecutive to an abscess in which the pus had made its way through at the linea alba. This kind of hernia is more common than the second, which escapes through any part of the anterior or lateral walls.

The usual indications of a rupture accompany these kinds also; the chief risk of an error of diagnosis arises where circumscribed collections of pus exist in the parietes without much local pain or any constitutional disturbance. If the hernia be reducible, the surgeon would not be likely to fall into error; and if not, the history of the case, with a careful manipulative examination, will enable him to form an accurate diagnosis.

Umbilical hernia.—By this term we understand that kind of rupture which escapes from the abdominal cavity, either through the umbilical ring in the foetus, or by an aperture formed at a later period of life in consequence of a separation of the fibres of the linea alba in the umbilical region. (See Fig. 272, p. 766, in which a small one is delineated.)

The synonyms are, exomphalos, omphalocele, or, in common language, ruptured navel.

It is developed at all periods of life, and in both sexes. Most commonly observed in infancy, it forms a protrusion either before or soon after the separation of the umbilical cord. In youth it is rarely developed, provided that the umbilical aperture has been once well closed. As years advance, it is more frequent, especially in females, and particularly so in persons exposed to obesity.

The hernial sac is always an acquired formation in this rupture; that is to say, there is not any peritoneal extension at the

closure of the ring is effected, and may
tissue of the cord itself, before that
We have seen such a case in which th
the sac sloughed. Rather later, after th
the hernial sac may be protruded in c
lical aperture remaining imperfectly cl
only by the integuments. In youth, a r
by escaping through a partially closed r
pressure, it dilates, unless precautions
And, in adult life, the fibres of the line
by stretching or yielding to the pressu
rupture escapes at the site of the on
immediate vicinity. The coverings of
rally very thin, and often inseparably
consist of the integuments; some fat,
often in proportion to the thickness of
regions of the abdomen; and the deli
fascia. The body of the sac is usuall
rather stronger near and at its orifice,
tissues outside of it form a very firm, re
The mouth of the sac is often large in
the protrusion. The relation which
when the protrusion is large should b
the bulk of the hernia increases, it
around the ventral aperture upon the
abdominal walls but it descends towa

Various organs of the abdomen are found in the sac of an umbilical hernia; very frequently the omentum, the stomach,* portions of the small intestines and of the large. When the disease has been neglected, we have seen nearly the whole of the intestinal canal in the sac enveloped by omentum. The last-mentioned organ sometimes becomes firmly united to the sac; and when bands or septa are thus produced within it, the intestines may become entangled by them. The omentum likewise becomes hard and contracted, forming masses within the sac which sometimes give rise to difficulty in diagnosis.

This rupture begins by forming a small soft projecting ovoid tumour at the navel. Gentle pressure with the finger pushes something into the abdomen, and a small hole is then felt, with very sharp, rigid edges. Directly the finger is removed, the skin either remains relaxed or flabby in the fossa of the navel, or it is slowly projected forwards, and reappears in the same bold relief. If the progress of the disease be not arrested, the protruding viscus descends lower and lower, so that the broadest part of the tumour lies below the mouth of the sac. A curious example of congenital umbilical hernia, coupled with malformation, was shown by Dr. Gibb at the Pathological Society, and in the same volume of the *Transactions* there is a description of a case of triple umbilical hernia which contained portions of the stomach, duodenum, jejunum, ileum, omentum, and ductus communis choledochus.†

Treatment when reducible.—Should there be any disposition to a protrusion at birth, or soon after, the simple application of a bandage is not merely palliative, but materially tends to the cure of the case. In infancy, after the separation of the cord, the protrusion may be prevented by fixing a piece of cork over the umbilical ring. It should be circular, about one inch in diameter, slightly convex on both surfaces, and covered with soft leather. With adhesive plaster to retain it in its proper place, and a bandage encircling the abdomen but not violently compressing it, this plan is sure to be attended with advantage. In the majority of cases the complaint will be cured. When the protrusion occurs in an adult, a suitable truss or bandage must be employed, with a view to prevent the dilatation of the

* A very interesting case is related by Mr. Moore, in the *Med.-Chir. Trans.* vol. xlv. p. 181.

† *Trans. of the Pathological Society of London*, vol. vii. pp. 216, 220.

sac and the increase of the tumour. If a large irreducible hernia has to be treated, a suitable appliance is required.

The radical cure of this hernia has been successfully effected by an operation performed by Mr. Barwell, on three patients at the respective ages of six months, fifteen months, and five years.* It is, however, open to further inquiry, how far a cutting operation is justifiable in infancy, as so large a proportion of cases are curable by bandages or trusses. Mr. Wood states that he has cured cases by inserting needles through the sac, after carefully reducing its contents, and keeping the faces in close contact by twisting a ligature over them. Wood makes use of rectangular pins and the wire suture.

Mr. C. Heath, having been called on to operate in a strangulated umbilical entero-epiplocele, after having reduced the bowel, dissected the sac from its attachments; and then passed a ligature through its base and the omentum, cutting it off. The patient recovered; and when the report was given to the Pathological Society there was not then any hernial protrusion.† But we are at a loss to understand how the opening in the linea alba was closed by these means; and therefore it can be shown that this has been accomplished, we should recommend the adoption of this measure.

Cases of umbilical hernia offer remarkably well-marked examples of that condition termed obstruction from accumulation of stercoraceous matters, and patients are constantly subject to severe constitutional disturbance arising from this cause, and not from positive strangulation of the bowel. We must not, therefore, be in a great hurry to operate in these cases, but after a free administration of aperient enemata, the contents of the sac become unloaded, and evacuations from the alimentary canal are obtained.

The operation.—An incision should be made in the line of the body, commencing about two or three inches from the upper border of the mouth of the sac, and continuing upwards upon the sac as far as may be requisite. The small opening by which the surgeon can remove the impediment to the reduction of the hernia, or liberate the bowel from compression, the better; and when it can be accomplished without opening the peritoneal sac, it is desirable to do so.

* *The Lancet*, 1861, vol. ii. pp. 419, &c.

† *Transactions of the Pathological Society*, vol. x. p. 131.

on is necessary whilst cutting the different coverings of the and should it be necessary to cut the tissues forming the incision should extend upwards in the same line the first.

hernia in the loins.—Lumbar hernia has been described by several surgeons. These protrusions arise after contusions of the region, or perhaps after lacerations of the abdominal muscles in the part, as well as from relaxation of the tissues. In the third *Bulletin des Travaux de la Société de Médecine de Marseille*, Dr. Chapplain* relates the case of a man, aged sixty, after being squeezed between a wall and a carriage, found a tumour between the crest of the ilium and the last rib. It appeared at first like a chronic abscess, but the presence of intestine was easily ascertained. Mr. Kingdon has seen a case of this kind. The bowel protruded just above the crest of the ilium at its highest point, about three inches from the spine, where the quadratus lumborum and abdominal muscles meet. The man was fifty-four years old, tall and thin. He died with hæmoptysis and emphysema of the lungs.

HERNIE IN THE HYPOGASTRIC REGION.

Inguinal hernia is the most common kind. The statistics of the Society of London Truss Society prove this; for rather more than two-thirds of the total number of applicants of both sexes professes to support every description of hernia, were afflicted with inguinal ruptures of one kind or the other.†

* *Union médicale*, 1862, vol. i. p. 157.

† TABLE B.

Showing the proportions between inguinal and femoral hernia in both sexes at different ages. (Constructed from Kingdon's tables, 1860–61.)

Ages.	Inguinal.		Femoral.	
	Male.	Female.	Male.	Female.
1 to 20	2435	249	23	38
20 „ 40	2954	322	139	432
40 and upwards	2154	128	144	278
	7543	699	306	748

the table shows—

that inguinal hernia is most common.


that in females femoral and inguinal hernia occur in about equal propor-

tion, but that femoral hernia in the male is the least common of these kinds.

Of inguinal hernia the following varieties are described: the oblique or external, and the direct or internal. The terms external and internal have reference to the course taken by the internal epigastric artery in relation to the orifice of the hernial sac. In the oblique kind, the mouth of the sac is situated to the outer or external side of this artery; in the direct, this important part of the sac is placed to the inner side of the same vessel.

There are also varieties of the oblique kind named according to the situation of the protrusion; thus the tumour is called a

FIG. 208.



The annexed woodcut represents double inguinal hernia. On the man's right side the protrusion has passed through the inguinal canal, and is just entering the scrotum. On the left side it is called 'bubonocoele.' The outline of Poupart's ligament, below the swelling produced by the protrusion, is distinct.

'bubonocoele,' when the hernia has passed the internal inguinal ring, but has not protruded at the external, when, in fact, it is retained in any part of the inguinal canal. It is called 'scrotal hernia or oscheocoele,' when it has passed through the external abdominal ring, and is therefore in the scrotum.

We propose the following anatomical classification of the varieties of oblique inguinal hernia.

1. Into the vaginal process of the peritonæum (the congenital hernia of Haller, and subsequent writers).

2. Into the funicular portion of the vaginal process of the peritonæum.

3. Scrotal, the sac being thrust into the tissues of the scrotum.

Of inguinal hernia in the male.—This species escapes from the abdomen, *above* Poupart's ligament, through the internal and external inguinal apertures or abdominal rings. There are two varieties, the *oblique* and the *direct*.

Oblique inguinal hernia, having passed through the internal inguinal ring, lies in the inguinal canal. Pursuing the oblique direction of this canal, it emerges at the external inguinal ring, and enters the scrotum, into which it descends. The mouth of the hernial sac is situated to the outer side of the internal epigastric artery, whilst its neck and body are usually in front of the structures composing the spermatic cord. But in rare cases these organs are divided; sometimes the blood-vessels pass over the tumour, the vas deferens behind it, and *vice versâ*; or they are attached to the sides of the tumour. The relative positions of the hernial tumour and testicle differ. The variable site of this organ depends upon congenital defect, and hence in some cases the testis cannot be distinguished from the tumour produced by the hernia. However, in the majority of cases the testicle is situated at the posterior and inferior region of the scrotum; more rarely, it may be detected at the front of the fundus of the tumour. An endeavour should always be made to ascertain the site of this organ, in every case of inguinal hernia, and under all circumstances.

The coverings of the sac of an oblique inguinal hernia are formed by the tissues of the region in which it lies. To expose the sac of a bubonocoele, it is necessary to divide the integuments of the groin, the aponeurosis of the external abdominal oblique muscle, and the internal spermatic fascia. Some of the fibres of the free edges of the internal oblique and transversalis muscles are observable along its upper border, and some fibres of the cremaster muscle skirt its lower edge. To reach the sac of a scrotal hernia, we must divide the integuments of the scrotum, including the dartos, the fibres of the cremaster muscle, often strongly developed, and the internal spermatic fascia. This last-named fascia is the delicate membranous extension of the internal abdominal fascia along the spermatic cord.

The mode of development of the sac of an oblique inguinal

its development is slow; its progressive stages manifest, and, although a prolongation of the peritonæum, its due to a morbid action, which allows displacement of this serous membrane.

Hernia into the vaginal process of the peritonæum
development of these two very different kinds of sac already described (see p. 654), in considering the hernia; and at the same place the reader will find which induce the author to prefer the designation into the vaginal process' and 'into the funicular those of 'congenital' and 'infantile,' formerly in also the natural varieties found in the condition of are described.

Under abnormal conditions, the following states of are found :

1. The whole length of the canal remains open.
2. The entire canal may remain open, and an a striction of its walls take place between the external ring and testis.
3. The tunica vaginalis propria testis is perfect now a closed sac, but the canal continues to exist at testis.
4. The abdominal orifice is completely closed, but below this exists.


The diagrams are intended to illustrate the conditions of the vaginal process of the peritonæum.

following works : Camper, *Icones Herniarum*, tab. x. fig. 3 ; Hunter (John), in *Medical Comm.* pt. i. chap. ix. ; and in Palmer's edit. of the *Works of J. Hunter*, plates xxv. xxvi. For numerous references to books on this subject, refer to *A Treatise on Ruptures*, by W. Lawrence, F.R.S., 5th edit. 1838, p. 564.

The variety of hernia developed in consequence of this imperfection is the hernia congenita of Haller and subsequent authors.

Fig. 270, A. In this figure the vaginal process of the peritoneum is open, as in the last imperfection ; but a contraction has taken place at the point above the testis.

FIG. 269.



This imperfection is demonstrated by a preparation in the Museum at Guy's Hospital, No. 2368 ; and a drawing from a case reported by Sir Charles Bell, in the *London Medical Gazette*, 1828, vol. i. p. 485.


Examples of hernia into a vaginal process of this kind are noticed by Scarpa ; see Wishart's translation, *Memoir*, ii. § 10. A drawing of a similar case may be seen in the work by Sir Astley Cooper on *Abdominal Hernia*, the 2nd edition by Mr. Key, pl. ii. fig. 2. Another is related by Mr. Lawrence in his work on *Ruptures*, as before, p. 574 ; and I have myself operated upon cases of this kind in young men. A large hernial sac with this remarkable condition may be examined in the Museum of the Royal College of Surgeons of England, No. 1943.

Fig. 269, B. This diagram shows the division of the vaginal

process of the peritonæum, into the inferior testicular vaginal process; and the superior funicular vaginal process, at the ventral end of which a communication with the abdomen remains, as on the other side.

This imperfection is delineated by Camper. In the work *Icones Herniarum*, folio, 1801, plate x., he contrasts it with that one represented in fig. 269, A. The drawing was made from the dissection of an infant, in 1759. Seiler also gives a figure to illustrate a rather more contracted state of this part of the vaginal process, in a work entitled *Obs. de Testiculorum ex abdomine in scrotum descensu*, &c. More recently, M. Malgaigne has described the frequency of hernia in association with this imperfection, *Leçons clin. sur les Hernies*, Paris, 1841; and in *L'Union médicale*, 1854.

FIG. 270.



Examples of hernia into the funicular portion of the vaginal process of the peritonæum occur very commonly in children, and a preparation of a hernial sac, originating with this defect, is preserved in the Museum of the Royal College of Surgeons, No. 1328.

Fig. 270, B. In this diagram the vaginal process is represented as a tube passing down in front of the spermatic cord and testis. The ventral orifice has been closed above.

This is the condition of the vaginal process of the peritonæum described by Mr. Hey, in which he 'found that the tunica vaginalis was continued up to the abdominal ring.'—*Practical Obs. in Surgery*, 3rd edit. 8vo, 1814, p. 227.

I suspect that it is this imperfection of which preparations exist in museums, described as 'a congenital hernial sac with

'mouth obliterated,' when unaccompanied with a history of the man from whom it was removed had ever been rup-

associated with congenital persistence of the vaginal process of the peritonæum, the testicle of the same side as that on which the defect occurs frequently occupies an abnormal situa-

tion in its normal site at the lower part of the scrotum, it may always be distinguished from the hernial protrusion in that it occupies the same serous sac as this organ and overlies it; but although this region should contain a hernial tumour, the testis may not have reached the scrotum at all.

The situations in which it is then found are as follows:

(1) Within the abdomen; the vaginal process extending into the vaginal canal, but not reaching further than just through the external abdominal ring, or into the upper part of the scrotum.

(2) Fixed in the inguinal canal out of the reach of manipulation; whilst the serous canal, passing into the scrotum, forms a sac for the reception of a hernia.

(3) Immediately outside the external abdominal ring at the lower part of the scrotum; in which state, when a hernia descends, it passes in front of this organ into the scrotum, even as its fundus.

A defective development of one or even both sides of the vaginal process is associated with the cases of deformity above described, and may lead the surgeon to discover an abnormal position of the testis; a fact of great importance in the treatment of these cases of hernia.

Clear view of the abnormal conditions of the vaginal process of the peritonæum, deviations from the normal situation of the testis, and the relative position of the hernia.

Hernia into the vaginal process of the peritonæum:

The vaginal process continuing open and common to the testis and epididymis. (Fig. 269, A.)

The testis may be situated—

- a, in its normal site at the fundus of the scrotum;
- b, just outside the external abdominal ring, or between its pillars;
- c, within the inguinal canal;
- d, within the abdomen.

The testis in the scrotum.

N.B. The hernia is never in conta

APPENDIX.—*Additional sacs, or prolong
the vaginal process within the a*

Concurrent with these instances of m
of those complicated cases of hernia wh
in the configuration and anatomical di
sac. In a majority of the cases in which
any very unusual direction, the hernia
same sheath as the testis, or a port
sufficient proof of the precise nature of
cause of its development, if any be requi
recorded in which a sort of second sac or
process of the peritonæum extended l
composing the abdominal walls. He
parietal,' 'ascending, or intermuscular,'
been applied to this variety of hernial s
this category are the cases termed 'herr
writers. These cases form two classes :

1st. Those in which the sac extends
dominal walls.

2nd. Those in which it extends into ti

In the first class, the sac extends mo

neurosis of the external abdominal oblique muscle, lying between it and the integuments; and when the hernia protrudes, it forms a tumour in the groin above and parallel with Cart's ligament. An example of this rare variety is cited by Scarpa,* and another case has been recorded by Fano.†

In the second class, the sac extends into the iliac fossa and upon the iliacus muscle, between the internal abdominal wall and peritonæum; or directing itself inwards, it passes behind the horizontal ramus of the pubes, and reaches the side front of the urinary bladder.‡

Hernia into the funicular portion of the vaginal process of the peritonæum.—When the surgeons of the last century discovered that a hernia could pass into the vaginal process of the peritonæum, and there be in contact with the testicle, they appear to have been content with this fact, and, without further research, to have assumed this variety to be the only form of hernia dependent for its origin upon non-closure of the ventral orifice of this canal, or defective obliteration of the upper part of the vaginal process of the peritonæum. Thus many writers acknowledge the fact that an infant may be the subject of scrotal hernia, and, after detailing such a case, they add, 'but not congenital;' implying by that expression simply the anatomical fact that the hernia is in a distinct sac, and thereby separated from the testis, in the same manner as it occurs in the adult. The term suggested in this essay precludes an erroneous view of those varieties of hernia which originate in congenital defect of the coverings of the testis and spermatic cord, and renders the disease in the infant and in youth much more easy of comprehension.

The reader should recall to mind the previous statements relating to the descent of the testicle, and the development of the tunica vaginalis propria testis out of the inferior termination of the vaginal process of the peritonæum. For surely there cannot be a tunica vaginalis propria testis developed so long as a canal exists along which a hernia may descend and touch the testis.

Treatise on Hernia, translated by Wishart, 1814, 8vo, p. 171.

L'Union médicale, December 1861.

Cases by Dr. Parise, in *Mém. de la Soc. de Chir. de Paris*, 1851; *Mém. sur les variétés nouvelles de Hernies*.

Herein consists the deficiency; in this anatomical fact defect. But when the testis is enveloped by the two layers of the serous membrane, the visceral and parietal, when the cavity of the tunica vaginalis testis is formed, in its own vaginal sheath, and is entirely shut off from every other viscus, there may yet remain the upper portion of the process of the peritoneum communicating with the abdominal cavity through the internal abdominal ring. Into this canal, termed the vaginalis propria funiculi, or the funicular portion of the process of the peritoneum, a hernia may protrude, and the canal thus becomes converted into a hernial sac; quite 'congenital,' too, as the variety ordinarily characterised by this term, for it is dependent upon a congenital defect, viz. the non-closure of the abdominal orifice of the canal, and its position in front of the spermatic cord as far as the testis is concerned. This variety is now distinguished as the funicular portion of the vaginal process of the peritoneum.

In the new-born infant, or at a later period, some months after birth, it is not at all uncommon to see a hernial protrusion occupying the inguinal canal, and the scrotum, but entirely separated from the testis, which is situated, at the fundus of the scrotum.

M. Malgaigne was, I believe, the first surgeon to point out this variety of hernia, and to point out its origin, and its relations, and distinctive features. It is exceedingly common. Carefully instituted examinations lead us to conclude that it is nearly as frequently met with as the ordinary variety described; but, from the fact of its having been confused with the scrotal hernia of the adult, very little attention has been bestowed upon it. The diminutive size of the testis has led to its relative situation, as regards the hernia, being overlooked and neglected; for cases which we have often examined have generally been regarded as of the ordinary variety, termed 'congenital,' and not worthy of further consideration. In the earlier periods of life, the distinction is not, perhaps, of primary importance; but the recognition of this variety in youth and adult life becomes a necessity of great moment in some instances, not only in reference to correct diagnosis, but also with respect to the judicious treatment of the case.

In these cases the hernia being enveloped by a peritoneal sac, the pathologist must admit one of two explanations

velopment: either that the parietal peritoneum of the abdomen was suddenly pushed downwards before the hernia to form its sac; or that a serous canal existed continuous with the peritoneal cavity, which became converted into the hernial sac. The last explanation, being strictly in accordance with anatomical facts, appears to me to be the one which the morbid anatomist will adopt.

The congenital patulence of the funicular portion only of the vaginal process of the peritonæum is capable of proof by anatomical examination, and there is no more reason to question its remaining so during the later periods of life, than to dispute the fact of the occasional unobliterated state of the entire length of the vaginal process to the fundus of the scrotum in adult life. We are able to adduce abundant facts to prove the development of hernia in such cases, if space permitted.

In advancing adult life the orifice of the hernial sac still maintains its original relations to the internal abdominal ring, and especially its depth. Under these circumstances, a hernia having existed many years, the abdominal rings are not approximated, but remain widely separated. A man, forty-five years old, was operated upon by myself on account of strangulated oblique inguino-scrotal hernia, with which he had been afflicted since boyhood. This was the first time he had found any difficulty in returning the protrusion. The orifice and neck of the sac were deeply seated; that is, the index finger was passed along the inguinal canal to reach the abdominal orifice of the sac. It did not require division, but was sufficiently large to allow of the reduction of the hernia after the sac had been opened and a large quantity of serum had escaped. His case terminated successfully. The testis was at the fundus of the scrotum, and separated from the contents of the hernial sac.

The next question relates to the frequency of these varieties of hernia into the vaginal process of the peritonæum.

After making inquiry of patients afflicted with inguino-scrotal hernia for many years, we were much surprised to find that so many of them stated that the protrusion occurred suddenly, and was first observed when the dimensions of the scrotum were increased by the presence of the hernia. Another circumstance also attracted attention. Robust, healthy, and well-developed men, of middle age, asserted that the rupture first descended

by those patients who seem to possess a fair average and might be presumed to give reliable information the subject of inquiry. During the two years 1854 and 1855 forty-four patients, suffering more or less from inguinal hernia, were admitted into the wards of Guy's Hospital. There were twenty-six cases of hernia into the vaginal or peritonæum; and of these, twelve had the testis in contact (the hernia congenita of Haller), whilst in the remaining fourteen the hernia occupied the funicular or canal, and was separate from the testis. In seven cases the hernia was developed suddenly, and reached the surface the first time it was observed, even in adult males being robust and well-developed. The respective patients, at the time they came under examination, were between fourteen months and forty-seven years of age.

Pott wrote, nearly a hundred years since: 'Rupture with that particular circumstance which brings the description mentioned in the title, are said to be very different from what I have observed, both in the living and the dead. I am inclined to believe that they happen much more frequently in adults than they are suspected to do.' *

To demonstrate, perhaps even more satisfactorily, the large proportion of the cases of inguinal hernia which occur at a very early period of life, we may refer to the Reports of the City of London Truss Society, so industriously directed by Mr. Kingdon.

The total number of males applying at the Society's Dispensary during the year 1854 was 1,234.

33, or 383 more than half, had reached only thirty years of

Thus it appears certain that inguinal hernia is developed frequently before thirty years of age.

This statement, which may be accepted as correct, is fully supported by reference to the table I have arranged from the report of the City of London Truss Society (see Table, p. 645).

Thus it appears that inguinal hernia occurs very frequently during the first twelve months after birth. This happens in consequence of the congenital persistence of the canal of the vaginal process of the peritoneum. Its frequency then diminishes to puberty, after which period the cases rapidly increase to the completion of thirty years of age.

Taking periods of ten years, or decades, the frequency of inguinal hernia at different ages is well seen (see Table, p. 731).

It is remarkable that the first decade contains the largest number of cases, and next the third or that between twenty and thirty years of age. From this period of life the cases decrease in a rapid ratio.

Hour-glass-shaped contraction of the sac of oblique inguinal hernia.—An uncommon peculiarity in the conformation of the sac of an oblique inguinal hernia occurs in consequence of a throwing of its walls at a point corresponding with the site of the external abdominal ring, or a little below that point in the rotum between the testis and that aperture. When a hernia is the sac, it may become strangulated by this constriction. The outline of the hernial tumour, as well as that of the sac, exactly resembles the shape of the ordinary hour-glass; hence the name given to it. It is always associated with a congenitally defective state of the vaginal process of the peritoneum. The result of the imperfect closure of this sheath at that part where the junction of its walls normally takes place, is to give rise to the formation of a ring or circular constriction within this process of the peritoneum at the point where the tunica vaginalis propria testis and tunica vaginalis propria funiculi meet together. Instead of the upper division of this canal being closed, and the two divisions, by this means, rendered distinct from each other, they communicate freely. The result is, that a hernia passes through the abdominal orifice of the funicular portion of the canal, traverses it, and then passes through the constriction into the cavity of the tunica vaginalis propria testis, which gland it touches. Instances of this constricted condition of the sac of an oblique inguinal hernia are related by

between the internal ring and the upper part of which were red, tense, and shining. The whole superior division and an inferior. This last, which unaltered by manipulation, was firm, and resisted elastic, and was very painful when compressed. and the swelling entirely disappeared when slight was reproduced by the contraction of the abdomen the abdomen above the internal ring. All the intestine being strongly marked, and considering this was imperative, I administered chloroform; and fully under its influence, that I was still unable determined to operate at once. An incision of long, was made, the direction of which was parallel to the tumour, and the centre corresponded with the superior ring. This aperture, although seen, was not sharp to my finger upwards on the outside of the hernial trace its boundary. A very distinct constriction of the tumour, producing the form of an ordinary hernia, corresponded with the ramus of the pubes over the superior division of the hernial sac above this constriction only. I next endeavoured to extricate the superior division of the sac; but I was unable to release it. I carefully introduced into the inferior sac, and moved it along the groove. Still, even when all the serous membrane was made on the bowel from above, it was impossible to reduce the bowel by the taxis. Having been divided, the strangulated bowel was removed from the abdominal cavity without any enlargement of the abdominal cavity required. The left testis was visible at the external ring, evidence that the hernia had descended into the scrotum. This man recovered.

of oblique inguinal hernia, which Mr. Kingdon selected for me, in order to illustrate quite a different subject, I find that he observed the characteristic stricture of the scrotal tumour in four men in whom the hernia was developed at the respective ages of seventeen, twenty-nine, and thirty years, and one in boyhood.

But what explanation can be given of the constriction of the bowel by the stricture at such an unusual part? Let us seek for it by an examination of preparations and a reference to the physiological changes taking place in the final process of the peritonæum before, or soon after, birth. Before doing so, let us distinctly isolate and exclude from the class of cases about which we are writing all those in which the impediment to the reduction of the hernia, or the cause of constriction of the bowel, depends upon the development of adventitious bands crossing the cavity of the sac without any definite specific arrangement. Hernial sacs, showing such peculiarities as last mentioned, are preserved in the London museums; but they are quite foreign to our immediate purpose, and belong to a perfectly distinct category as regards their development and relations to the sac and its contents.

In the cases we are now describing the hernial protrusion and testis are in contact, otherwise the case would not belong to this class.

In the Museum of the Royal College of Surgeons there is a Hunterian preparation, No. 1343, thus described in the catalogue: 'The sac of a large congenital hernia. The exterior of the sac is uneven and sacculated, through the unequal yielding of different parts of its walls: the testicle is situated at the widest part.' This account is not sufficiently explicit or descriptive of the structure of the hernial sac and its relations to the testis. A careful examination of the interior of the sac shows a large superior division and a small inferior. In the latter the testis may be recognised; but between the divisions

an imperfect septum, or perforated diaphragm, existed before the vertical division of the entire sac was made, and by means of this foramen a communication was established between them. The sharp, well-defined, free edges of this septum are seen stretching from the sac-wall to the testis, as well as its strong attachment to the sac itself. Now assume the cut edges of this sac

to be joined together, and the superior and inferior sacculi would be complete, though any foreign body might be passed from the one into the other. A hernia, for example, filling the upper sacculus, might pass from it through the aperture in the diaphragm into the lower, and then become constricted by the margins of this annular contraction.

But a preparation in the Museum at Guy's Hospital, No. 2368, at once affords a satisfactory clue to the solution of these somewhat remarkable cases. It is the vaginal process of the peritonæum taken from a child. It shows a portion of the parietal peritonæum of the abdomen with the whole process extending downwards from it. This process has been opened along the whole extent of its anterior surface. At the superior extremity the canal communicates with the abdomen, and at the inferior extremity the testis may be seen; and between these two points, about midway, the cavity is contracted. When this tube was uncut, there must have been an opening or narrow passage near its centre, through which a protrusion, coming by the abdominal orifice, and traversing the funicular portion of the vaginal process, must have passed to reach the testis. A constriction resembling this formed the impediment to the reduction of the hernia in the cases above described.

Sir Charles Bell describes a hernial sac of this kind which came under his observation in connection with an obscure case of hernia, which he has recorded in the *London Medical Gazette* of 1828.* 'There was another remarkable circumstance observed, which related to the sac lodged in the scrotum. At the lowest part there was a hole of communication between that sac and the cavity which is between the coats of the testicle. This orifice was so large that the finger could be passed through it; and its margins were so dense as to resist dilatation.' In the explanation of the woodcut, he writes: 'A bougie has been passed from the hernial sac into the cavity formed by the tunics of the testicle, through an opening which forms a communication between them.'

A careful examination of the second figure of the 2nd plate of Key's edition of Sir Astley Cooper's work on Hernia will satisfy any one that it delineates a case of the kind we are describing. It was a 'congenital hernia firmly constricted by an annular contraction of the sac itself,' although described as 'a band of membrane adhering to the sides of the tunica vaginalis, through an aperture in which the intestine has protruded and become strangulated.' The hernia is seen both above and below this aperture; the portion below 'discoloured by the effects of strangulation' produced by its margins; the portion above, 'below the internal ring and above the strangulating band, free from pressure and retaining its natural appearance.' Now in this description we have but a repetition of the appearances I observed in the case under my own treatment, related at p. 744. In chap. xvii. on *Congenital Hernia*, Sir Astley Cooper states that in this variety 'the stricture is much oftener found to be *within the sac*, than in the ordinary kinds of hernia;' and he relates a case, upon the authority of Mr. Hodgson, of Lewes, in which the intestine 'was found strangulated within an aperture of the tunica vaginalis. The intestine appeared to have suffered more from this stricture than from the ring.'

Lastly, Mr. Lawrence,† when describing cases of this kind, records one in which, 'about half-way between the testis and groin, the hernial sac was so contracted that a probe only would pass into the stricture.' In this case, too, the abdominal orifice of the sac 'would not admit the smallest portion of the tip of the finger,' and it also formed an impediment to the reduction of the hernia; in this respect differing from my case.

By these quotations, to which numerous instances might be added, especially those recorded by Mr. Pott, with dissections, case 14,‡ it is capable of demonstration that this class of cases must be confined to those forms of inguinal hernia dependent upon congenital defect or persistence of the vaginal process of the peritonæum throughout life.

But a glance at a plate in Camper's work § will more easily explain the nature of these cases. Figure 2 shows the cavity of the right vaginal process of the peritonæum opened vertically

* Vol. i. p. 484; also a woodcut.

† *On Ruptures*, 5th edit. p. 574.

‡ His *Chirurgical Works*, edit. 1808, vol. ii. p. 184.

§ *Icones Herniarum*, tab. x. 1801.

from the abdomen to its inferior termination; at which point is seen the testis. Figure 3 shows the left side, in which, just above the testis, the entire canal is divided into two portions by the internal surfaces of the serous membrane united together. Should it, however, happen that this septum is not complete, a hernial contraction takes place, and a ring is thus formed, keeping up a communication between the persistent funicular portion of the vaginal process, which is above, and the cavity of the imperfect tunica vaginalis testis below. By the constant pressure around a hernia, the tissues composing this annular contraction of the hernial sac become more and more dense, until it forms a firm, fibrous, callous ring at the situation where the tunica vaginalis ordinarily terminates just above the testis; its variable distance from that gland depending, in different cases, upon the larger or smaller size of the protrusion, and the consequent distension of the tunica vaginalis testis.

The differential diagnosis between these cases, of hernia into the vaginal process of the peritonæum, and the ordinary form of inguino-scrotal hernia, is formed by the manner of their development, the age at which they first appear, and their configuration. Whilst the common form is developed slowly, these appear suddenly, and often pass at once into the scrotum without resting in the inguinal canal. They are noticed in the earliest infancy, throughout childhood, and during early adult life. Their form or outline is highly characteristic. The tumour projects in a remarkable manner from the external outline of the inguinal canal; directly the rupture escapes from the embrace of the external abdominal ring, the form is more globular and rounded in comparison with the pyriform outline of the hernia of slow formation in adult life. The testis can or cannot be felt; when it is easily detected, it is evident that the fund and rupture are in separate sacs, and *vice versâ*.

In the hour-glass-shaped variety the configuration alone is sufficient to attract notice and lead to its diagnostication.

When the hernia is reducible, the depth of the internal abdominal ring and the length of the inguinal canal are discoverable by tactile examination.

The importance of recognising these cases.—It is highly important to recognise the cases of hernia into the vaginal process of the peritonæum, in a practical point of view, especially in relation to their treatment. Much more severe constitutional symptoms usually accompany strangulation in these cases, and this con-

generally useless if the bowel has been strangled for hours, and persistence in such attempts is erroneous. Under any conditions, the surgeon must remain patient until distinct and separate contractions exist, through which the contents must have passed; one of these being situated in the hernial sac, the other being its true ventral orifice. If no contractions, or both of them, may offer insuperable obstacles to the reduction of the hernia, unless their treatment is resorted to.


If we scrutinise any large number of cases admitted into the hospitals, a majority of them require the liberation of the bowel by a cutting operation. At St. George's Hospital, 30 years of age. At St. George's Hospital, 11 had passed that age.* At 57 cutting operations performed in order to liberate the bowel in inguinal hernia, 33 belonged to the chronic variety, 24 were of the old slowly-forming variety. In order to demonstrate with how much greater facility an inguino-scrotal hernia is reduced by the operation of descending into the vaginal process of the peritoneum, I refer to 129 cases of oblique inguinal hernia treated at Guy's Hospital. Of these, 59 had traversed the vaginal process of the peritonæum, 70 belonged to the inguino-scrotal variety. Or, tabulated thus:

Of inguino-scrotal hernia, requiring a cutting operation.	Of oblique inguinal hernia, requiring a cutting operation.
30	30
11	11
57	57
33	33
24	24
59	59
70	70
129	129

the taxis. In some of those cases in which the hernia, when strangulated by the orifice of the sac, was supposed to have returned into the abdomen in a mass, more careful examination after death showed that the inguinal portion of the sac had burst, and that the bowel had been pushed through the perforation, and was lying outside the peritonæum, between it and the internal abdominal fascia. This accident is more fully described at page 704. In these cases, also, the mouth of the sac is sometimes detached from its connections. When the patient is youthful, and the anæsthetic effects of chloroform are produced, great caution is, therefore, necessary in using the taxis, on account of the slight and delicate connections between the peritonæum to the neighbouring parts at this early age. Difficulties also attend the operations on these cases, in consequence of the depth of the orifice of the sac, the variable position of the testis, and the varieties which may be encountered in the disposition of the hernial sac, all of which have been before described.

Infantile hernia.—The infantile hernia of Hey, and the cysted hernia of the tunica vaginalis of Astley Cooper, are synonymous terms for a variety of the oblique inguinal hernia, depending likewise upon an abnormal condition of the tunica vaginalis peritonei. The state of that sheath which precedes the development of this kind of hernia depends upon the ventral foramen being closed, but the canal persisting from that point to the testis. The hernia slowly pushes before it the parietal peritonæum of the abdomen into this sheath, and when the parts are dissected, it is seen that ‘the tunica vaginalis is continued to the abdominal ring, and encloses the hernial sac,’ as Mr. Cooper describes. In this manner ‘the protruded parts, together with the sac, are contained in the tunica vaginalis testis’ (Lawrence, *A Treatise on Ruptures*, p. 576).

The name given by Hey to this variety of hernia leads the reader to infer that it is always developed in infancy. This, however, is not the fact. Hey’s case was an infant fifteen months old. Forster’s case, related by Cooper, was thirty-one years old at the time of the operation. The duration of the rupture is not specially stated, but the description induces us to believe that it had not existed long. In Lucas’s case the rupture was developed at about seventeen years of age. The rupture in Forster’s second case had been ruptured all his life,



he has incised the first serous sac. The
the testis attached to its walls, falls out, and
suspended by its attachment to the margin of
abdominal ring.

Inguino-scrotal hernia of slow formation.
oblique inguino-scrotal hernia which occurs
late adult life, and forms for itself its own
parietal peritonæum before it, causes at first
at the internal abdominal ring, slowly trans-
canal, and at last occupies more or less of
the 'pointing' of the hernia at the internal
M. Malgaigne aptly terms it, the attention
sometimes drawn. Upon inspection a slight
integuments over the internal inguinal ring
becomes more prominent when the patient
coughs or sneezes, especially if standing
posture, and under similar influences, if the
over the ring, the hernia is projected against
tion thus induced is termed the impulse of the

At this period a truss should be constant
to fit comfortably, it will prevent the further
viscera; and by counteracting their propulsion
the parietal peritonæum, the development of
arrested

in and the aponeurosis of the external abdominal oblique muscle. Along its superior border it is overlapped by the free edge of the internal oblique and transversalis muscles; along its inferior by the cremaster muscle. It lies upon the internal abdominal fascia as it emerges at the internal inguinal ring, afterwards upon the conjoined tendons of the internal oblique and transversalis muscles, just before it reaches the external inguinal ring. Under these dispositions of the fleshy and tendinous parts of the abdominal muscles, the hernial sac with its contents is pushed in a very especial manner under the influence of their contractions; more especially in the region of its mouth and neck: it is a frequent impediment to the reduction of a hernia. The spermatic cord usually lies behind the tumour at the internal inguinal ring, and is attached to the posterior surface of the sac as it traverses the inguinal canal.

As the development of the sac advances, it is pushed through the external inguinal ring, over the os pubis, and into the scrotum. As the hernia points at the external inguinal ring, it forms a somewhat globular swelling; but as its bulk increases in its progress towards the fundus of the scrotum, the shape of the tumour is generally pyriform, when the scrotum is distended by the hernia. In the body of the tumour contractions or pulsations, generally following an oblique or transverse direction, are occasionally observable. Such appearances are due to the more or less yielding tissues of the scrotum.

The testis is usually below the tumour, sometimes behind it. The elementary structures of the spermatic cord may be traced along its posterior or outer boundary, generally close together, side by side, but sometimes, though very rarely, separated from each other.

When the hernia is reduced, the forefinger may be passed with facility into the abdominal cavity through the external inguinal ring, the inguinal canal, much shortened, and the inter-inguinal ring. This is practicable on account of the approximation of those rings taking place from the pressure of the hernia and the relaxation of the surrounding structures. The mouth of the sac and the inguinal apertures are indeed sometimes so stretched that the finger may be freely passed to the abdominal surface of the symphysis pubis; and the pulsation of the epigastric artery, or of the external iliac, may be felt in the usual situations of those vessels.

A tabular arrangement of the characteristics of the three varieties of inguinal hernia in males.

A. Into the VAGINAL PROCESS of the PERITONÆUM. (<i>Hernia congenita Halleri.</i>)	B. Into the FUNICULAR PORTION of the vaginal process of the peritonæum.	C. INGUINO-SCROTAL tissues of the
<p>1. Is developed most commonly in infancy; occasionally in youth; rarely in adult life.</p> <p>2. Is suddenly produced.</p> <p>3. At once descends along the inguinal canal into the scrotum.</p> <p>4. May rest in the inguinal canal when the testis is <i>not</i> in the scrotum.</p> <p>5. Envelopes the testis, and lies in contact with that organ.</p> <p>6. The orifice of the hernial sac is contracted, and corresponds with the site of the internal abdominal ring.</p> <p>7. Neck of the sack long and tubular; it lies in the inguinal canal, between the rings, which are <i>not</i> approximated, even in the adult.</p> <p>8. Hernial sac a direct, congenital prolongation of the peritonæum; tubular when not distended with a hernia. It reaches the testis in the scrotum, and envelops it. Thus there is a congenital serous canal to receive the hernia.</p>	<p>1. Is developed frequently in infancy; often in youth and in adult life.</p> <p>2. The same.</p> <p>3. The same.</p> <p>4. May rest in the inguinal canal when the testis <i>is</i> in the scrotum.</p> <p>5. The testis, at the fundus of the scrotum, occupies its own serous sac, which separates it from the hernia. It usually produces a very distinct prominence at this part.</p> <p>6. The same.</p> <p>7. The same.</p> <p>8. Hernial sac, a direct congenital prolongation of the peritonæum; tubular when not distended with a hernia. It does not extend into the scrotum so low as the testis. A congenital serous canal exists to receive the hernia.</p>	<p>1. Occurs in infancy, and is, I believe, exclusive of the adult.</p> <p>2. Is slowly produced.</p> <p>3. By slow degrees, it turns the inguinal canal into the scrotum.</p> <p>4. May remain in the inguinal canal for a long period, the testis being in the scrotum.</p> <p>5. Is separate from the testis, which is at the fundus of the scrotum, and is not in contact with it.</p> <p>6. Orifice of the hernial sac is contracted, and near the external abdominal ring.</p> <p>7. Neck of the sack is short and broad; the rings are approximated.</p> <p>8. Hernial sac a direct congenital prolongation, and also developed by the pressure of the abdominal viscera on the parietal abdominal peritonæum, which is before the hernia is formed into a sac. Here the hernia is not in its own sac by the peritonæum before the tubular character is wanting.</p>

diagnostication.—Every inguinal hernia escapes from the inguinal cavity above Poupart's ligament. If, therefore, the hernia is formed by an inguinal protrusion be examined whilst in the inguinal canal, Poupart's ligament is traceable along its superior border, and the opening where the tumour seems to escape from the abdomen is above the same fibrous band.

It may be always distinguished, then, from crural hernia; the latter kind generally escapes through the crural ring, which is situated behind and below Poupart's ligament, and that ligament may be always traced along the superior border of the tumour. Figs. 268, 272 should be compared.

In an inguinal hernia reaches the scrotum through the external inguinal ring, the outer pillar of which is formed by the fibrous attachment of Poupart's ligament to the spinous process of the pubes. Now place the tip of the finger upon the external process of bone, and if the neck of the tumour is on its inner side, or between the finger and the symphysis pubis, the tumour must be formed by a protrusion which has passed through the external inguinal ring; a fact sufficiently illustrative of an inguinal hernia. Should the tumour be to the outer side of the finger, the probability is that the hernia has passed through the crural ring.

Differential diagnostication.—The surgeon is often required to distinguish between inguino-scrotal hernia and some other tumors developed in the inguinal region and scrotum.

These are—1st, the different kinds of hydrocele; 2d, the testicular spermatocele connected with the epididymis; 3d, varicose veins of the spermatic veins; 4th, inflammation of an old inguinal sac, and the results of such inflammation; 5th, inflammatory affections and other diseases of the testis, the cord, and its coverings; 6th, hæmatocele; 7th, malpositions of the testis; 8th, inflammatory and other diseases of the inguinal lymphatic glands; 9th, growths of fat in the connective tissue of the inguinal canal and upon the spermatic cord; 10th, diseases of the integuments of the scrotum, especially growths, phantiasis.

The nature of these diseases being described in other parts of the work, the observations we have to make in relation to the differential diagnosis between them and inguino-scrotal ruptures may be condensed into a tabular form. This plan was suggested by the perusal of a chapter written by M. Vidal (de Cassis),

divisions :

I. The reducible. (*See table on next p*

II. The irreducible. (*See table on p. 72*

In the first division there are :

1. Inguinal hernia.

2. Hydrocele of the vaginal process of

3. Hydrocele of the funicular portion

4. Varicocele of the spermatic veins.

The tumours in the second division are of fluids only, sometimes almost exclusively of both solids and fluids in variable

Those composed of fluids are—1, hydrocele of the tunica alba testis; 2, hæmatocele in the tunica alba testis; 3, encysted spermatocele connected with the epididymis; 4, hydrocele of the spermatic cord.

Those formed by solids, or of solids and fluids, are—1, diseases of the testis, a, of inflammatory or degenerative growths; 2, hæmatocele of some standing; 3, diseases of the spermatic cord; 4, diseases of the spermatic cord; 5, diseases of the tissues of the scrotum.

Between the local signs of an inflamed spermatic cord, especially if its orifice be plugged by omentum, and those of an inguinal hernia, there is a close resemblance. But

either in their local or constitutional manifestations. It is more probable that a hernial protrusion might be considered to be some affection of those organs, than that they be mistaken for a hernia. The history of the case usually settles all doubt as to its nature.

Characteristics of the Reducible Tumours compared with Hernia.

	Their entrance or return into the abdomen.		Their passage from the abdomen.
	Characters in common.	Special characters : when without complications.	Special characters.
HERNIAL.	<i>All</i> return into the abdomen most easily when the patient lies down on the back and when the abdominal muscles are relaxed.	1. <i>Hernia</i> enters most readily. When once commenced, passes in quickly and suddenly. Entrance complete. Opaque and thick neck of tumour. Testis may or may not be perceptible until reduced. <i>No</i> vibration.	1. Is developed from above, descends when the patient rises or exerts the abdominal muscles, and more quickly than others. The finger pressed over the ring prevents its descent.
HYDROCELE OF PERITONÆUM.	—	2. <i>Hydrocele</i> of vaginal process of peritonæum enters slowly, and never suddenly. Entrance complete. Translucent and small neck of tumour. Testis imperceptible until the fluid has entered the abdomen. Vibration.	2. Seems to be developed from below upwards. The serum sometimes remains in spite of the horizontal position.
HYDROCELE OF FUNICULAR DIVISION OF VAGINAL PROCESS OF PERITONÆUM.	—	3. <i>Hydrocele</i> of funicular division of vaginal process of peritonæum enters like No. 2. Entrance complete. Translucent. Neck of tumour may pass into inguinal canal. Testis perceptible at fundus of tumour. Vibration.	3. Similar to No. 2.
VARICOCELE.	—	4. <i>Varicocele</i> enters very slowly. Entrance not complete, the bulk of tumour only diminished. <i>No</i> vibration.	4. The tumour increases like hernia when the patient rises; but it increases also if pressure be made over the course of the spermatic veins in the inguinal canal, or if the blood be retarded in its passage along them in any way.

Characteristics of the Irreducible Tumours compared with Hernia.

Disease.	Weight.	Translucency.	Fluctuation and vibration.	Relation of testis to tumour.	Figure and development.	Size.	Consistence.	Pain.
HERNIA	Lighter than either the fluid or solid tumours.	Very rarely so; generally opaque.	Only when fluid co-exists with the hernia.	Position variable, but the testis usually discoverable.	Pyriform, but with thick neck. Occasionally globular or ovoid. Begins at neck of scrotum and descends.	Variable; at times very large.	Soft and yielding, except it be omental.	Painless unless dis-eased.
FLUID TUMOURS.	Hæmatocele rather heavier than hydrocele.	Hydrocele particularly so. In rare cases opaque. Hæmatocele opaque.	Distinct; vibration very characteristic, as in hydrocele.	Perceptible in spermatocele; not in hydrocele of tunica vaginalis propria testis, usually.	Pyriform, but with very thin neck. Outline very regular. Globular as hydrocele of spermatic cord. Oblong as in spermatocele, and nearly transverse to vertical axis of scrotum. Begin near fundus of scrotum and ascend.	Rarely very large.	Yielding and elastic. Exceedingly incompressible as hydrocele of cord.	Painless, unless testis be squeezed, usually.
SOLID TUMOURS.	Generally heavier than fluid.	Opaque.	Absent.	Often involved and imperceptible though its site may be discoverable by pressure.	Outline of testis often preserved. Sometimes surface irregular.	Large; steadily increasing.	Resisting, firm and rarely hard.	Painful, but variable in degree.
Mixed TUMOURS.	Heavy.	Sometimes in parts of	In some parts, not in	Involved.	Irregular outline.	Large; at times rapidly increasing.	Resisting in parts, soft in	Variable.

Treatment.—The palliative treatment of reducible oblique inguinal hernia consists in maintaining perfect and uninterfering retention of the hernia; for if this be done carefully and with method, a permanent cure may be effected, especially in children. The cases of hernia into the vaginal process, if carefully treated immediately after the first appearance of the tumour, and before the walls of that canal have been stretched to any length of time, are sometimes cured by the employment of a well-adjusted truss. But in the use of this instrument before puberty, great attention must be given to the site of the tumour; for the pad of the truss might press injuriously upon the organ. The spring of the truss must not be too powerful; and if the pressure of the pad upon the walls of the abdomen is very strong, their tissues are absorbed where the pad presses, and they become, in consequence, seriously weakened. The truss should prevent the escape of the hernia by its accurate adjustment, rather than by the force with which it presses against those apertures through which the hernia passes. On this account, each individual requires to have a truss adapted to the configuration of the body. The pad should be applied over the internal inguinal ring, pressing gently upon the inguinal canal to afford support to the tissues of that part, and not upon the external inguinal ring and pubes, as we have often seen it applied. The patient should be enjoined to wear the truss uninterruptedly, except only when lying in bed; to adjust it carefully before leaving bed in the morning, and on no account to permit the hernia to remain for a moment in the sac.

When the hernia has acquired large dimensions, or has become irreducible, a particular form of appliance, termed the bag-truss, becomes indispensable.

The radical cure of reducible inguinal hernia.—The mode of performing the operation as practised by Wutzer is as follows (a special instrument is required, which consists of two pieces of wood and a flexible needle):

The patient being placed in the most favourable position, and the rupture carefully reduced, the forefinger of the operator is inserted to invaginate the integuments and to push the fundus of the sac into its orifice. The cylindrical wooden plug is next inserted in place of the finger, and the needle thrust through the invaginated skin, sac, anterior wall of the inguinal canal, and skin of the groin. The grooved piece of wood is then laid over the inguinal canal, held in position by the needle, and

hernial sac and the entire circumference of its entire obliteration of the other parts of the inflammation. Unfortunately, however, the results do not uphold the theory.

Mr. Wood * has contrived an operation by which the chances of failure above mentioned are avoided. The operator requires a special needle, of wood, glass, or porcelain, and some strong force for the contraction of the abdominal muscles shown to be controlled by the influence of chloroform.

After an incision has been made through the skin of the scrotum, the forefinger of the operator is pushed into the hernial sac as far as possible into the inguinal canal in front of the spermatic cord, at the same time separating the tissues. The needle is next carried by the forefinger and passed through the conjoined tendons of the external oblique and transversalis muscles, the internal inguinal ring, and the integuments. Great care is taken to avoid including the spermatic cord. A thread is then passed through the eye of the needle, and the needle is then withdrawn, leaving one end of the thread in the puncture made in the scrotum, still threaded, is next passed through the external abdominal oblique muscle near to the internal inguinal ring, and out through the opening in the integuments.

and the inner pillar of the ring. Its point is now brought through the first puncture for the third time, and the needle is then withdrawn from the thread.

A loop and two ends of thread pass through one opening in the skin. 'Two portions of thread are thus placed across the hernial canal, invaginated fascia and sac, closely embracing, but not including, the spermatic cord, and connecting the posterior or deep wall with the anterior and superficial, perforating aponeurosis of the external abdominal oblique muscle in two places; but escaping by the same aperture in the skin' (Fig. 10). The compress is next applied over the canal obliquely, and the two ends of the thread passed under the loop and tied in such a manner as to give equable adjustment to the cure.

The after-treatment of the case is the same as that adopted in similar operations.

Mr. Wood describes several modifications of this operation in which wire was used with good results.

By taxis.—When a surgeon has a case of irreducible inguinal hernia to treat, his attention should be particularly devoted to a careful examination of the tumour, in order to diagnose the variety under observation. He should ascertain with accuracy the anatomical relations of the tumour to the surrounding parts; the position of the testis, especially; the length of time the patient has been ruptured; the mode of development of the hernia, whether suddenly or slowly; the treatment pursued to prevent the protrusion from taking place; and the progress of the present state of the rupture.

Having determined the variety, if the state of the tumour permits, he may employ gentle pressure in order to reduce the hernia, remembering the direction or course it has taken, and the circumstances which probably cause the impediment to reduction. The abdominal muscles must be relaxed as much as possible whilst the patient lies on the back.

It is only necessary to describe, in this place, the manipulative proceedings, as the constitutional measures employed to render them more effective have been detailed at page 696. A much larger proportion of the cases of inguinal hernia are now reduced by taxis than formerly. This is due to the use of the trochanter, the effects of which in producing total annihilation of the contractile function of muscular tissue are so important. In all inguino-scrotal protrusions are not reduced with the

same facility as large ones. When such a protrusion occurs in a young person, and has been only recently developed, after a fair trial of the taxis, when the patient is fully under the influence of chloroform, and the efforts to reduce it have failed, any delay in removing the impediment to the replacement of the hernia is reprehensible. To resort to any other measure which necessitates the occupation of time, is only to abandon the sufferer to additional risk of the loss of life.

If the protrusion be compound, that is to say, when it consists of a large mass of omentum and a very small knuckle of bowel, the reduction of the latter is effected with difficulty. The same thing occurs when the hernia is one of irreducible omentum of long standing, from old adhesions, and a small piece of bowel. Also, when the sac is distended and rendered very tense by a considerable effusion of serum. The efforts of the surgeon to reduce the bowel are thereby rendered abortive for if the serum cannot be made to pass through the mouth of the sac, into the abdomen, the hernia never will.

The operation.—The patient should lie on the back, with the pelvic region slightly elevated, unless some firm, resisting body, as a mattress, cushion, or table, be underneath. The anatomical points which the surgeon particularly wishes to reach and display in the cutting operation are, the external inguinal ring, the aponeurosis of the external abdominal oblique muscle, the internal inguinal ring, and the mouth of the hernial sac. The various structures which constitute the coverings of the hernial sac need not be displayed with rigorous anatomical precision. They differ exceedingly in density and development in different cases. The length of the incision should not be proportioned to the size of the hernia. Its length must be just sufficient to expose freely the anatomical points, alluded to above, and more. In ordinary cases, the point of the scalpel should be inserted through the integuments at about one to two inches above the assumed centre of the external inguinal ring, and carried downwards upon the anterior surface of the tumour, to terminate about two or three inches below the same mark. The margins of the external inguinal ring are next carefully exposed, and their relations to the tumour examined. They should not be cut unless it be imperatively required to do so. The deeper coverings of the hernial sac are next carefully incised, with or without the assistance of a grooved director, until that important structure is reached.

The operator, pressing his finger upon the sac, next innuates it through the external inguinal ring, in order to ascertain if there be any structures which firmly encircle the neck and orifice of the sac outside. Should he discover any, a grooved director may be guided by the finger underneath them, and they may be cut. After slight pressure made upon the sac, its contents are sometimes returned into the abdomen at this stage of the operation. If their reduction be not practicable, the peritoneal sac must next be opened. It must be done after the manner described at p. 708, and with great care. Probably some serum will escape when the sac is cut, but this is not always the case. The opening in the sac requires to be sufficiently large to allow the operator to reach its orifice easily, and to examine its contents. The index finger is now passed along the anterior surface of the protrusion upwards towards the mouth of the sac, when an impediment to its further passage is encountered. The tissues which bound this narrow opening constitute the impediment to the reduction of the hernia. The operator next passes the hernia bistoury along a grooved director, or upon his finger, through the mouth of the sac, and divides the structures in contact with the knife sufficiently to allow the ungual phalanx to be passed freely into the abdominal cavity. The direction of this incision should be parallel with the linea alba. A few small arteries are occasionally cut during the operation, which may be twisted immediately before further proceeding.

Oblique inguinal hernia in the female sex.—This variety occurs at very early periods of life. In fact, with the exception of umbilical hernia, it is the only kind developed before five years of age. Even until the age of puberty it is more common than any other variety. Its development in infancy is due to the patulous state of the vaginal process of the peritonæum, otherwise known by the name of the canal of Nuck. Doubtless such a condition of this process remains throughout early life, into which a hernia occasionally and suddenly descends. In the adult female this hernia generally forms slowly after thirty years of age. But I have met with double inguinal hernia in a well-formed woman aged only twenty-four years, who had never been pregnant; she had worked hard, however.

Its anatomical relations are alike in both sexes, merely sub-

that it is not uncommon to feel the ovary in inguinal hernia of an infant.

The tumour formed by an inguino-lab attains the size so often reached by inguino-lab have seen one which descended more than half middle of the thigh, and the coverings of which attenuated that the convolutions of the intestine were distinctly visible. When of moderate size, it has a more globular figure, and it has a longer neck, or more narrow and cylindrical neck, according to sex.

In forming a correct diagnosis of the female, the same anatomical points must be observed which have been already described when the male was under consideration. The surgeon should particularly remember that a hydrocele of the neck of the canal of Nuck, is occasionally developed, and may lead to an error in diagnosis.

The palliative measures described as being the treatment of cases of inguinal hernia in the male are applicable in these also; but if the hernia becomes irreducible the surgeon must regard the accidents arising from it, such as taxis, and delay in liberating the bowel with pain and danger, and act with promptitude and decision. We have seen, under these circumstances, the

nial sac as in the male. And he should remember that, occasionally although very rarely, the bowel may be constricted by a contraction of the sac itself near the external inguinal ring, and not by its ventral orifice only.

Direct inguinal hernia.—The variety of inguinal hernia characterised by the term 'direct' occurs in both sexes. It belongs to that class in which the hernial sac is formed slowly, as an accidental or acquired formation. The only exception to the slow mode of formation is in those rare cases when the

FIG. 271.



Direct inguinal hernia.

ectures immediately behind the external inguinal ring are created by violence and a hernia protrudes. It is never the result of a congenital imperfection. This variety is sometimes called the internal inguinal, from the fact that the mouth of the sac, where it is continuous with the parietal abdominal peritoneum, is placed to the inside of the internal epigastric artery. It is comparatively rare.

At its commencement it forms a prominence where it points into the external inguinal ring, through which it passes into the upper part of the scrotum. The outline of the tumour

in direct inguinal hernia than in oblique. When viewed from a short distance it will be seen passing through the vertical or long axis of the tumour with the linea alba, and there does not occur any inclining or curving of the neck of the tumour towards the crest of the ilium, as is observed in oblique inguinal.

In its passage from the abdomen it directly traverses that small portion of the inguinal canal immediately behind the external inguinal ring, and the fibres which form that part of the floor of that canal before the hernia, or they are lacerated when it escapes through the opening so formed. The conjoint tendons of the internal oblique muscles and the pubic portion of the internal oblique. The spermatic cord and round ligament are not in the hernial sac until it has reached the external ring. When it has passed that point, they lie to its sides and are usually less identified with its tissues than in the variety.

The distinction between this variety of hernia and oblique must be formed under the guidance of the facts before described.

The palliative treatment requires a particular support which must be so constructed as to give support

are often very attenuated. Considerable care is therefore required in the operation of incising them.

Inguinal hernia sometimes pass through abnormal apertures in the aponeurosis of the external abdominal oblique muscle.—It would be an oversight not to allude to the fact, that an inguinal hernia sometimes passes through an opening or division of the fibres in the aponeurosis of the external abdominal oblique muscle close to the ring, and not through the true external inguinal ring, which is then traversed by the spermatic cord alone. Such anatomical varieties are rare, and would only require some slight modification of the operation, to meet their peculiarities.

Femoral hernia, also called *crural* or *merocoele*, was not accurately distinguished from some forms of inguinal until the middle of the 17th century. Its designation in a measure indicates the region of the body where it forms a tumour, which, however, may be stated, with greater precision, to be the upper and inner part of the thigh. Its anatomical relations were not accurately defined for many years after it was known to pursue a different course to inguinal; and it is curious to follow the gradual development of our knowledge of those structures immediately interested in, or associated with, this variety of aperture.


The principal anatomical points which should be examined, from the outside of the thigh as well as from within the abdomen, are the following: the ligament of Fallopius or of Poupart; Gimbernat's ligament; the crural canal or ring; the fascia transversalis and iliaca, or the internal abdominal fascia of the iliac fossa; the falciform process of the fascia lata and the apphenous opening; and, when a hernial sac exists, the fascia propria of Sir Astley Cooper.*

Relations of the mouth of the hernial sac to the internal epigastric artery and vein.—The orifice of the peritoneal sac holds im-

* The anatomy of these various parts must be assumed to be known by the reader. We may, however, allude here to the importance of noticing the structure last named, viz., the fascia propria, first described by Sir Astley Cooper from a dissection made in 1800, and which is still preserved in the Museum of Guy's Hospital (No. 2503). It is an envelope which is found on the outside of the peritoneal sac, usually separated from it by a layer of adipose tissue.

portant relations to the internal epigastric vessels. 8 pathologists have established three varieties of the ordinary kind of crural hernia on this basis. The first, in which the orifice of the sac is situated to the outer side of the epigastric vessels, is called external crural hernia; the second, where it lies to the inner side of the same vessels, the middle crural hernia, the most common; the third, in which it is placed to the inner side of the renal vessels, is called internal crural hernia.

Fig. 272.



The illustration represents double femoral hernia. The outline of Poupart's ligament being remarkably distinct above the tumour formed by it.

of the obliterated umbilical artery. The first and third varieties are very rare.

Here we may also state that the obturator artery is occasionally given off from the internal epigastric or femoral, that when the former vessel lies to the outside of the orifice of the hernial sac, the obturator artery may cross closely its neck, and dip down by its inner side to enter the obturator foramen.

A very interesting case, with a well-drawn illustration recorded by Mr. Spence in the *Edinburgh Medical Journal* July 1855. The pulsation of a large vessel was felt with the finger before its division.

varieties in the course pursued by the sac of a crural hernia.—
 In a memoir published by Dr. Le Gendre, he describes four
 varieties of crural hernia.* They may indeed be regarded
 as curiosities, on account of their rarity.

1. The hernia, as soon as it traverses the crural ring, passes
 entirely internal to and behind the femoral vessels, and rests on
 the pectineus muscle, the aponeurosis of which sometimes forms
 an envelope to it; this he calls the pectineal crural hernia, on
 account of its situation, or the hernia of Cloquet, in honour of
 the surgeon who was the first to describe it. Callisen, Vidal
 (Cassis), M. Richet, Dr. Le Gendre, and Mr. J. Adams,† have
 recorded similar cases.

2. This variety, although placed internal to the femoral
 vessels, is, however, rather far from them; it passes through
 the resisting fibrous structure which bounds the crural canal
 sheath internally; that is to say, the ligament of Gimbernat.
 Laugier, who was the first to notice it, calls it the *crural hernia*
through Gimbernat's ligament, or the hernia of Laugier. Cru-
 hier, Demeaux, Nuhn, and Le Gendre have also dissected
 examples of this kind.

3. This comprehends the variety of crural hernia which
 Hesselbach has so well described and figured.‡ The hernia in
 this case traverses several openings in the fascia cribriformis,
 and then presents several distinct lobes, which give it a cha-
 racteristic appearance; it is the *hernia with a diverticulum*
through the cribriform fascia, or the hernia of Hesselbach.
 Le Gendre and Malgaigne have dissected examples of this
 variety of rupture, which the former believes to be not so very
 rare.


4. Lastly, a variety in which the hernia, after having escaped
 beneath Poupart's ligament and traversed the cribriform fascia,
 sends one or more prolongations through the superficial fascia.
 This variety may be termed the *crural hernia with a diverticulum*
through the superficial fascia, or the hernia of Astley Cooper.§
 Dr. Le Gendre describes a dissection of a case of this kind, and
 publishes a drawing of it in his work.

Mém. sur quelques variétés rares de la Hernie crurale; avec 6 planches. 8vo.
 Paris, 1858.

Med.-Chir. Trans. 1860, p. 127.

De ortu et progressu Herniarum, 4to, 1816.

Anatomical and Surgical Treatment of Abdominal Hernia, chap. xx.; case
 No. 12. Sheffield.



parietal peritonæum down upon the thigh or more below Poupart's ligament. The most favourable for the study of the crural hernia. By degrees such a diaphragm at the crural ring becomes into a sac by the repeated and continued which forms the hernia. The sac extends thigh beneath Poupart's ligament; but lower and lower towards the knee, its direction towards the crest of the ilium, so that the tumour lies parallel with Poupart's ligament and fascia lata. When large, the tumour is prominent, and seems to occupy the region of the canal. But in every case of crural hernia, by a protrusion which has escaped from below and behind Poupart's ligament; it is able to trace that structure running along.

Crural hernia rarely increases to the size of the fistula, occasionally, after an operation for the hernia, the related bowel, when the tissues around it have been weakened by cutting them, and then it is to employ the support afforded by a truss. The protrusion of the alimentary canal may protrude. The neck of the sac become so remarkably attenuated that the movement of the intestinal convolutions is impeded.

that region. The great size of the tumour is not a feature of serious importance, for a very small sac, having a contracted neck, more readily entangles the hernia and prevents its reduction.

Diagnosis; first, from other kinds of hernia. Those in which a crural may be confounded are inguinal above Poupart's ligament, and obturator below the horizontal ramus

FIG. 273.




piece of the ileum about three feet from the cæcum removed from a patient, aged fifty-seven, who had been operated upon for strangulated femoral hernia of seventy-three hours' duration. She survived the operation forty hours, and the first symptoms of the attack 118 hours. She had been in bad health previously, was extremely emaciated, and sank from exhaustion. Large and hard masses of scybala were found in the ileum as shown in the part of the bowel cut open. The small piece of bowel which had been in the sac, and was but little diseased, is indicated by the dotted line. The contracted and opened bowel below this contained mucus only.

The figure in the centre represents the sac. Through its mouth a piece of glass rod is inserted; and although the sac is very small, it was remarkable for the layer of fat, *b*, between the peritonæum, *c*, which is not cut, and the fascia propria, *a*. (Museum Guy's Hospital; No. 477².)

the pubes. It would seem to be impossible, however, to mistake a crural hernia for an inguinal, or *vice versâ*, if the observer will but carefully trace the outline of Poupart's ligament. But he must remember that the fundus of the sac of a crural hernia sometimes takes a course upwards, and overlies

that structure. Nevertheless, the course of Poupart's ligament can be always ascertained with more or less precision; whether an inguinal hernia overlies it from above—a very rare occurrence—or a crural from below, which is very common. The fingers of the examiner can still trace it from the ilium to the pubes—of course, in one case more distinctly than in another. If the whole of the tumour lies below the ligament, and

FIG. 274.



Demonstrates the appearances produced by a small inguinal hernia just pointing to the external inguinal ring, and a medium-sized femoral hernia, both on the same side, and the outline of Poupart's ligament between those tumours. On the left side of the scrotum a tumour is produced by hydrocele of the tunica vaginalis. (From nature, and copied from a drawing in Guy's Hospital Museum.)

neck can be traced continuous with the crural aperture, its contents have escaped beneath Poupart's ligament, and it is a crural rupture; but if the contrary, it is an inguinal, especially if the neck be traceable to the internal inguinal ring. Crural is distinguishable from inguino-labial hernia by placing the finger on the spine of the pubes, or upon the attachment of

adon of the adductor longus into that point of bone ; for even very fat persons the pubic spine can be always felt. If the tumour be to its outer side, the protrusion has escaped below Poupart's ligament, in which case it is crural rupture ; if, on the contrary, it has passed through the external inguinal ring, it must be an inguinal. The same point of bone will aid in distinguishing between a pudendal hernia and a crural.

A crural hernia is much more superficially seated than an obturator, and in consequence the tumour which the latter produces is neither so prominent nor so well defined as the former. The latter escapes from the pelvis behind the horizontal ramus of the pubes, and therefore Gimbernat's ligament and the crural aperture can be felt. The elevation of the integuments produced by an obturator hernia is certainly in the locality of the crural aperture ; but the depth of the tumour is its striking feature, and the facility with which the crural aperture can be felt is pathognomonic of the relations of the protrusion.

Crural hernia requires to be distinguished from other diseases which occur in the same region.

1. From *psoas abscess*.—The history of the case, the locality of the swelling, and the results of a palpable examination of the tumour, are quite different from hernia. Psoas abscess is generally preceded by pain in the back, constitutional disturbance, and inability to extend the hip-joint completely without pain. These symptoms do not accompany a crural hernia. Psoas abscess does not burrow down below Poupart's ligament and the crural ring, but to the outer side of the sheath of the femoral vessels. But an infallible test that the tumour is formed by a circumscribed collection of fluid is obtained by placing the patient flat on the back, and then gently applying the right hand above Poupart's ligament and the left below it, *vice versâ*. When pressure is made upon the swelling with the hand only, the other is elevated ; and when the action is reversed, the opposite effect takes place. With this result on manipulation of the swelling, there can be no doubt of the nature of the disease.

2. *Enlargement of the lymphatic glands*.—The correct history of the development of the swelling will aid in forming the diagnosis, especially its permanence and progressive increase. If there be a suspicion that the tumour is of glandular origin, it is as well to institute a careful examination of those somewhat

secluded regions in which certain specific ulcerations nately occur, giving rise to angioleucitis.

3. *Dilated and varicose veins.*—We have seen a dil and a tumour formed of varicose veins occupying the femoral hernia; but the diagnosis was easily made b the patient in the recumbent posture, when the swelli peared. Digital pressure being made at the crural ring the vein, the swelling was at once reproduced.

4. *Cysts in the superficial fascia.* Cysts are sc although very rarely, developed in the region of the cr in some cases associated with hernia, but sometimes The permanency of the swelling, its persistent and i size, the fluctuation discoverable on pressure, and the the case, are aids to correct diagnostication. When present, and there are any symptoms suggestive of str bowel, the propriety of accurately ascertaining the co the tumour by an exploration of its interior cannot be

Palliative treatment.—The treatment of crural hern use of trusses is most important. The prevention of ti tion of the sac is the primary point to which attentio be directed; and this may be accomplished by giving in the region of the crural ring, especially if it be maintained.

We gladly avail ourselves of the large experienc Kingdon in offering a few observations regarding t trusses in this variety of hernia. In a truss to pr escape of a crural hernia, the spring should fall s suddenly from the point where it passes around the hi along Poupart's ligament. The pad should be rather s convex. The cross-strap should fasten high up on the of the spring, in order to keep the pad well down in t The thigh-strap should start from near the pad, and after encircling the thigh, to the pad itself. When l ligament, or rather the whole crural arch, is lax, an backwards and forwards with the varying size of the s the pad should press upon that ligament; for then th aperture is made smaller, and the rupture more efficien tained. For that purpose a larger and flatter pad is But when the crural arch is strong and steady, the sm is both more effective and, by reason of its smallne convenient and less inconvenient. Also, it is less displacement by the movements of the hip-joint.

When the rupture is large, or where it comes down under the ala, it is necessary to use a thigh-belt, with a triangular projecting on the inner surface, and forming a soft continuation of the pad, to fill the triangular space where the form fascia occupies the saphenous opening of the fascia

sometimes after an operation, in which Gimbernat's ligament has been freely divided, a cross tongue in addition, to be to the free end of the truss, is needed.

Reducible crural rupture of course requires a hollow pad, for epiplocele purely, or enterocele.

Attempts have been made to effect a radical cure of crural hernia by a cutting operation. The patients recovered from effects; but we have no data by which to arrive at any conclusion as regards the advantages obtained.

Crural hernia very rarely occurs before puberty. I have seen it in a girl ten years old.

Crural hernia is much more common in the female sex than in the male. But the general belief, that it is much more common in women than inguinal, does not appear to be founded on facts.

The following table demonstrates the proportions in which they occur. The comparative numerical equality is expressed by the circumstance, that before puberty inguinal hernia is common, whilst crural is extremely rare. In 193 girls under 15 years of age, Mr. Kingdon met with 184 cases of inguinal rupture, and only 9 of crural. Even to the age of 20 years, the cases of inguinal hernia are much more common than crural, as the subjoined table shows. In a total of 1,442 married females, at all ages from birth upwards, the majority of cases of crural hernia over inguinal was only 54.

After twenty years of age, crural hernia is much more commonly developed than inguinal.

Decades.	Inguinal.	Femoral.
Birth to 10 years . . .	146 . . .	1
11 to 20 " . . .	103 . . .	37
21 to 30 " . . .	153 . . .	180
31 to 40 " . . .	164 . . .	252
41 to 50 " . . .	76 . . .	158
51 to 60 " . . .	33 . . .	84
61 and upwards . . .	19 . . .	36
Total . . .	604 +	748 = 1,442 cases.*

This table was made from Mr. Kingdon's reports of the City of London

Total cases of femoral	748
Deduct cases between 20 and 40 years	432
At all other ages	316
Deduct cases before 20 years of age	38
After 40 years of age	278

This table demonstrates a fact for which the pathologist be scarcely prepared, viz., that the largest number of crural hernia is developed during those years usually to the prime of life, *i.e.* in women between twenty years old and forty.

But it must be observed that this happens to be that period of life when parturition is most frequent, and when consequently the peritonæum and the tissues of the abdominal wall become much stretched by the development of the gravid uterus. Their power to resist the weight of the abdominal viscera becomes diminished; hence a cause of their protrusion.

To ascertain whether there was any relation between the development of hernia and parturition, Mr. Kingdon made the following table of 680 ruptured females, and found the proportion of mothers to be as follows :*

	Inguinal.	Femoral.
Infants and girls under 16 years of age	87	3
Single women	50	61
Mothers	178	262
Married women who had not borne children	19	20
Total	334	346

And next, with a view to ascertain the influence of repeated pregnancies upon the development of hernia, he examined women who had been mothers before discovering the hernia. Of these, 180 women were afflicted with inguinal, 262 with femoral; which gives a majority of 82 in favour of femoral. The influence of the first pregnancy is very remarkable. Of 180 women who had been mothers before discovering the hernia :

Truss Society for the years 1860 and 1861. It shows the ages of the individuals at the time of first discovering the hernia.

M. Malgaigne states that inguinal hernia is even more common in females than crural. *L'Union méd.* 1854, p. 154.

* *Report*, 1860, pp. 10, 11.

		Inguinal.		Femoral.
and borne one child . . .		27	. . .	40
„ two children . . .		26	. . .	25
„ three „ . . .		22	. . .	31
„ four „ . . .		23	. . .	32
„ five „ . . .		15	. . .	25
„ six „ . . .		8	. . .	17
„ seven „ . . .		15	. . .	19
„ eight „ . . .		13	. . .	17
„ nine „ . . .		7	. . .	18
„ ten „ . . .		10	. . .	9
„ eleven „ . . .		5	. . .	11
„ twelve „ . . .		4	. . .	4
„ thirteen „ . . .		2	. . .	2
„ fourteen „ . . .		2	. . .	8
„ fifteen „ . . .		—	. . .	3
„ eighteen „ . . .		1	. . .	—
„ nineteen „ . . .		—	. . .	1
		180	+	262 = 442

preceding table the fact was demonstrated that crural hernia is most commonly developed between twenty years of age and forty. Now as we know that there is a much larger number of females alive between twenty and forty years of age than in any other age, we might conclude that the number of cases of crural hernia is in proportion to the numbers of the population in each respective age. Under twenty years of age this is certainly the fact, for the population at those early ages of life is the largest, whilst cases of crural hernia are exceedingly rare. The cause of the frequency of inguinal hernia in the early life of females has been before explained.

It is highly interesting to compare the frequency of crural hernia at various ages with the numbers of the female population in at the same periods of life; and in this inquiry we have the use of the census-table of 1851,* and Mr. Kingdon's table before alluded to. From those data the subjoined table is arranged:

		Crural hernia.
Cases under 20 years . . .	493,260	38
from 20 to 40 years . . .	453,809	432
„ 40 upwards . . .	308,609	278
	1,255,678	748

* Census, 1851, vol. i. p. 193, table ii.

crural hernia under certain conditions. lity is probably due to the natural con- coupled with its attendant circumstan- of the class of persons which would a as well as, perhaps, to configuration, causes, or hereditary influences.

Morbid states of the crural hernia. A plocle is very often seen ; an enterocele rare. With a view to ascertain whet the bowel was in any degree influence or absence of omentum, we made an a of strangulated crural hernia, which our own care for operation. The prop being thirty-one pure enteroceles, thir

A reducible crural epiplocele may at e give rise to local pain, and increase in tion of cold, and absolute rest, are the to prevent adhesion between the ruptu dition of an omental rupture in the gro for angioleucitis. But the history of t evidence derived from the absence of lymphatic irritation should suffice to diagnosis.

Both Pott and Astley Cooper allude are of vital importance in the treatm

strangulated crural hernia, only 38 were reducible by the taxis. This, of course, is explicable by remembering that the majority of such cases is not sent to the hospital until the further use of the taxis is evidently unjustifiable. Ten of these cases were of recent development, that is, they were strangulated on their first recognised descent. This is an important feature in crural hernia, bearing especially on the results of treatment in these cases. It is cases of this kind which are so frequently overlooked by the medical attendant. He suspects that the symptoms exhibited by the patient are referable to a rupture, and he asks the question whether there is any tumour to be felt in the groin or elsewhere. Receiving a negative reply, he is content therewith. But that should not satisfy him. The surgeon should carefully examine the crural rings, and the other regions in which a hernial tumour may exist.

The mortality arising in cases of strangulated crural hernia of all kinds is very large; but in those where the bowel becomes strangulated on the first descent, the death-rate is the largest. The experience of a large number of these cases teaches that the bowel should be liberated as soon as possible, and, if the taxis be not successful when the patient is fully under the influence of chloroform, that the cutting operation should not be delayed a moment.

The youngest girl we have heard of suffering with strangulated crural hernia was between eight and nine years old.

Application of the taxis.—The patient should lie on the back, with the hip-joint more or less flexed. The surgeon, taking a position most convenient for the purpose, gently compresses the whole swelling, at first endeavours to diminish its size by pressing any serous effusion that may happen to be in the sac into the abdomen, and next to empty the bowel of its contents. Then recalling to mind the site of the crural ring, he presses the bowel backwards, inwards, and upwards.

The operation of liberating the strangulated bowel by incision.—The tissues to be divided to expose the sac are, the integuments, the superficial fascia, the fascia propria, and often a layer or covering of fat. The points to be particularly observed as guides for the safe and certain manipulation of the sac are, first, Poupart's ligament in front and above it; and, secondly, Gimbernat's ligament at the pubic border of its neck.

Slight variation in the line, direction, or figure of the incision through the integuments is admissible, according to the incli-

nation of the operator; but for all practical purposes we make a linear incision of from two to three inches in length, which should cross over the pubic side of the neck of the tumour, and extend on to the abdomen about one inch above Poupart's ligament. Its form being slightly concave, the neck of the tumour is embraced by the concavity. The integuments and superficial fascia being divided, a well-defined, membranous sac is usually seen, and the operator should next display clearly Poupart's ligament above the tumour, and feel Gimbernat's ligament on the pubic side of its neck. This covering is the fascia propria. Next, he may pass a hernia bistoury, with great care, between the last-named ligament and neck of the tumour, outside the fascia propria, and, directing its sharp edge upwards, cut through some of the fibres of Gimbernat's ligament, or of those which form the falciform process of the fascia lata to it (Hey's ligament), and thus enlarge the crural ring. This done, the reduction of the rupture may sometimes be effected by the application of gentle pressure upon the contents of the sac. This process constitutes the minor operation. It is simple, safe, and well adapted to cases of recent enterocoeles which have been surgically regulated a very few hours, and when there is reason to believe that the bowel has escaped injury by the taxis. Obviously, therefore, it is only justifiable in select cases.

Failing to reduce the contents of the sac, the fascia propria is next carefully divided upon a grooved director, and a layer of fat is very frequently exposed. Inexperienced operators, mistaking this fat for the omentum, are puzzled when they cannot find the bowel, or they mistake the peritoneal sac for intestine when that is seen upon clearing away the fat. The hernial sac, being exposed, it should be traced upwards to the crural ring, and the fascia surrounding it in the crural canal divided. The division of the structures forming the crural canal has not been reserved for this special moment, must now be undertaken as before described. But the hernia remains irreducible. With great care, and *secundum artem* (p. 708), the sac must now be incised. Through the first puncture a stream of serum usually flows. Empty the sac of the fluid by enlarging the opening sufficiently to reach its orifice. If nothing but omentum is seen, carefully raise it from below or gently unravel it, and towards the mouth of the sac, secreted behind that part of the omentum which has just escaped through its mouth, the bowel will be seen. Next, to enlarge the orifice of the sac, intro-

point of the finger between the protrusion and Gimbernat's ment; but if that cannot be done, the hernia bistoury must be insinuated in front of the omentum, directing its cutting edge forwards and holding the blade parallel with the linea alba. Having then liberated the bowel from its constriction, it is easily returned into the abdominal cavity. It is not always necessary to cut the orifice of the sac, after having divided the tissues on one side of it, as before described. Its tissues frequently yield to the gentle introduction of the finger.

Arrange the wound in such a manner that an opening exists at the lowermost angle for the escape of discharge, and flex the joint by placing a pillow beneath the popliteal region for the limb to rest upon. Any further special detail for the treatment of the case is here unnecessary.



Mr. Kingdon has informed us that the extent to which Gimbernat's ligament is divided is a subject of grave importance regarding the future comfort of the patient. He states that Gimbernat's ligament should rather be cut than Gimbernat's, and that when the latter is at all extensively divided, it is almost impossible to retain the hernia after the thickening, subsequent to the operation, has passed off. Certainly the most numerous crural hernial tumours we have seen occurred in women in whom an operation had been performed for strangulated hernia.

Arteries, cut during the progress of the operation, should be immediately twisted.

Wounds of the obturator artery.—This blood-vessel, when it passes to the obturator canal, after dividing from the internal iliac or gastric or femoral, lies close to the neck of the sac, and it has been occasionally cut at the same moment as its orifice. Before using the knife, it might be practicable to feel the pulsations of this artery with the tip of the finger. When cut, an attempt should be made to arrest the bleeding by twisting the ends of the wounded vessel. See Mr. Spence's case, p. 766.

Omental sacs are most commonly met with in cases of crural hernia. An account of them has already been given at p. 709.

Obturator hernia.—The obturator canal is situated at the upper and outer part of the obturator or thyroid foramen, and is bounded above by the horizontal ramus of the pubes, and in the lower parts by the obturator membrane or ligament and the obturator muscles, which are attached to its internal and



THIS rare variety is described as hernia
tor canal, the foramen ovale, the thyroid
and sub-pubic femoral. In the early pa
M. Garengeot * called the attention of su
hernia, which, after having escaped from
the obturator canal, forms a swelling
muscles and in the pubic region of the th
sac lies behind the horizontal ramus of th
obturator canal, and makes its way so
uppermost fibres of the external obtur
times above them. The fundus and body
by the fascia of that muscle. By the
cases, the obturator vessels and nerve
ferently placed as regards the tumour. V
found the artery six times to the outer sid
to the inner side, and three times behind
tion of this artery to the sac probably bea
origin. In Mr. Stanley's case,† both a
above the hernial sac; the former to its
towards its outer.

Both sexes are liable to the formation
large majority of the cases on record occu
anatomical reasons for this are obvious.

There are cases on record in which th
and in which the viscus was strangulat

a part was found, are marked illustrations. In Mr. Oubr 's
 *, 'the eye detected a slight degree of fulness in Scarpa's
 angle,' and 'a distinct hardness could be felt, slight in its
 extent.' In the only instance we have seen, the hernia gave
 rise to a very distinct, well-defined tumour, deeply seated among
 the adductor muscles, to the pubic side of the femoral vessels.
 In Mr. Stanley's case,† a crural epiplocele existed on the same
 side as the obturator entero-epiplocele. The latter was, how-
 ever, extremely small, and produced 'no swelling on the inside
 of the thigh.' The hernia 'was found with its peritoneal sac
 beneath the obturator externus muscle, between the muscle and
 the obturator fascia.'

The foregoing observations illustrate the difficulties which
 attend the detection of protrusions through the obturator
 foramen.

We have collected twenty-five of the most recently recorded
 cases of this variety of hernia, and the remarkable features they
 present are well worthy the attention of the reader. They may
 be divided into two classes: the first embraces a large majority,
 and consists of those cases in which the hernial tumour was not
 discovered during life; the second, those in which it was dis-
 covered by palpable examination, either immediately at the
 inner side of the thigh, or, as, in one instance, only by an
 examination per vaginam.

In several of the cases of the first class, the tumour produced
 by the protrusion was so small, that the sac, with its contents,
 being placed deeply among the adductor muscles and beneath
 the pectineus muscle, could not give rise to any local external
 swelling. In two cases the tumour was found between the ob-
 turator ligament and obturator externus muscle. The contents
 of the sac in six cases consisted of only a portion of the calibre
 of the canal of the intestine. This had, however, become con-
 verted into a diverticulum, in some instances of considerable
 length. We meet with this condition of the hernia occasion-
 ally in other regions, but the cases are not common; and we
 therefore conclude that, in proportion to the rarity of
 obturator hernia, this may be considered to be one of its remark-
 able features. In one case, the Fallopian tube and ovary formed
 the contents of the sac; in another, a portion of the urinary
 bladder.

* *M d.-Chir. Trans.* xxxiv. p. 233.

† *Trans. of the Path. Society*, vol. iii. p. 94.

the obturator nerve, described by the feature of the case, and pressure over the aperture of the obturator canal gave rise of great severity. It is due to the state, that he seems to have been the first particularly upon this fact, which some strongly insisted upon by Romberg.† T seen with this variety of hernia complain sudden pain at the upper and inner part which extended thence down the inside of front of the leg, foot, and great toe.

The cases of the second class are those was discovered during life, either in color or by a careful examination of the region developed.

Dr. Roeser‡ proposed that advantage should be of the male, and the vaginal of the female, to instigation of the obturator region within the pelvis. Lorinser, in 1857, discovered an obturator hernia.¶ in the same year, in a case in which not even fulness perceptible, was induced, by the severity of the strangulated bowel, and the pain caused by local pressure obturator region, to explore that part; and his judgment by the discovery of a very small hernial tumor were returned into the abdominal cavity by gentle pressure the sac.|| Before this Mr. Obré had successful

strangulated obturator hernia, in 1851, in which 'the eye detected a slight ree of fulness in Scarpa's triangle, on the right side;' this triangle of the osite limb being well-marked with a hollow or depression passing down its tre, but which was lost on the affected side. Also, on pressing firmly 'a le below the saphenous opening, a distinct hardness could be felt.' Lastly, the case successfully operated upon by Mr. B. B. Cooper, in 1853, the nan being very thin, there was both ocular and palpable evidence of an isputable nature. At first sight, the swelling in Scarpa's triangle seemed e identical with that which might be produced by a protrusion through the ral canal; although it had not the appearance of being in close relation with apart's ligament, for it formed a somewhat globular fulness, rather than a umscribed tumour. Then the point of the finger could be placed in the ral ring, which of course was conclusive evidence that the hernia had not aped through it. Really, therefore, there were no practical difficulties to vent the formation of a correct diognostication in this case. We may here te that these observations are made from notes of the case taken personally the bedside of the patient.

Complications.—But we must allude to a class of cases, of ich there have been several instances, in which the existence an obturator hernia was complicated with the development other kinds, especially those in the inguino-femoral regions. one case there was a reducible crural hernia on either side; three cases, a crural hernia on the same side as the obturator; one case, on the opposite side; and in another case, an inguinal hernia on the same side as the obturator.

The treatment adopted in three of these cases was that which ost surgeons would approve. The symptoms of strangulated testine being clearly marked, an exploration was made of the eruinal sacs, which were visible and tangible, in the expectation f finding a small knuckle of bowel retained within them. But othing of the kind being found, the examination was not pro- cuted further; although, in two of the cases, an obturator rnia, seated immediately beneath the site of the operation, s proved, by after-death examination, to have been the cause eath.

As a possible complication, we must here allude to the fact at in some of the cases, in the same individual, a sac was and to pass through both right and left obturator canals.

In fourteen of the cases the obturator hernia was not discovered until after th. In one case the symptoms indicated the existence of this variety of nia, and the patient recovered without any interference on the part of the geon. In ten of the other cases the protrusion through the obturator canal s discovered during life; and the treatment adopted and its results we may efly describe. In one case the protrusion was reduced after the application he taxis by Dr. Roeser, of Bartenstein. This was in a female thirty years

an obturator hernia, the integuments over which were gangrenous. He made an incision; the contents of the sac were turned out, and the patient died next day.§ The accuracy of the diagnosis was proved by death.

Dr. Heiberg operated upon a woman, fifty years of age, who had an elastic swelling in the left adductor region. He opened the hernial sac, stretched the obturator foramen, and turned the bowel. The patient, however, died.|| Dr. Heiberg operated upon a woman seventy-five years old. The swelling was very perceptible; the pain on pressure and down the thigh was very severe. The sac was opened, its orifice cut, and a piece of the sac was removed. The aged patient survived the operation thirty-six days.¶ Dr. Nutt operated upon a woman, aged seventy-five years, suffering with a strangulated intestine, but without any tumour or redness. He pressed in one particular spot to diagnosticate the hernia, and there was not even 'fulness' perceptible, over the spot. He cut over this painful spot, found a small tumour, and reduced its contents by gentle pressure. The patient died three days. After death, the bowel which had been strangulated was found in repair. An obturator hernial sac was seen on dissection.

We have now to describe the cases which have been recorded in the late Mr. Obré, of London, was the first surgeon to operate for an obturator hernia. This was in 1851. In a woman, aged seventy-five years, and stout, he diagnosticated this hernia. He cut over the swelling, and cised the sac and its orifice slightly, and returned the bowel.††

In 1853 I assisted Mr. Bransby Cooper in an operation for an obturator hernia.

a woman, forty-nine years old. She was exceedingly emaciated, with twelve children, and was affected with bronchitis. She was being ruptured. For sixty-two hours before coming to the hospital, suffering with strangulated intestine. Her illness began with a swelling at the upper and inner part of the right thigh, and extended from the inside of the thigh, knee, front of leg, and to the great toe. Swelling of the region below Gimbernat's ligament, a deeply seated swelling, was felt. Mr. Cooper made an incision over the tumour, divided the muscle, and reached the hernial sac. Gentle pressure was made and the contents passed readily into the abdomen. She slowly recovered, the wound entirely healed; but before she left the hospital she died with acute bronchitis, which caused her death.

Mr. Cooper, in 1857, discovered an obturator hernia by an examination. The age of the patient is not stated. On the eleventh day after the commencement of the attack, he cut down upon the tumour, opened the sac, and the contents in a state of gangrene. A fæcal fistula was established, frequently closed, and the woman survived the operation eleven

months. We have now given a brief but instructive summary of the facts from which useful practical lessons may be derived. There are many cases published, which have been discovered after death, and there is no doubt that the fact of the existence of the disease.

Mr. Cooper believes he has met with five cases of obturator hernia in men. Two of them were in men; in both the femoral artery was in swelling, and pushed forwards at each forced expiration. The men were of middle years of age, thin and gaunt. He does not recollect whether the artery was pushed forward in the female, whose case he otherwise remembers, but he distinctly recollects that he was able to feel the bony upper edge of the aperture through which he had returned the hernia. The woman was emaciated to the last degree of voluntary locomotion.

We propose next to select the most prominent facts from the number of cases, with an especial view to the diagnosis of those cases in which there may not be any palpable or evidence of a rupture.

Diagnosis of an obturator hernia,—from its position. Emerging along the obturator canal, it emerges upon the lower part of the horizontal ramus of the pubes, to the inner side of the femoral artery and vein; and to the inner side of the adductor longus. The tumour formed by the hernia is covered by the pectineus muscle. From its position, therefore, it may be distinguished by observing the positions of the horizontal ramus of the pubes and of the femoral artery. Those structures occupy, in fact, a position between these two kinds of hernia. In obturator hernia they

are in front of the tumour ; in crural hernia they are. In the former, then, they are easily felt ; in the latter not be without difficulty—not, perhaps, until the reduced. In those cases in which either a fulness, slowness, tumefaction, or swelling exists, coupled with weak indications of obstruction or strangulation in some alimentary tube, the difficulty of diagnosis is not so great but how much embarrassment arises when those signs which betoken strangulated bowel exist, and a hernia nowhere to be felt, let the numerous cases on record which the rupture has only been found after death.

Assuming, for the sake of illustration, that we have a patient suffering with well-marked constitutional indications of strangulated bowel, and that, after the most careful examination of all those apertures through which ruptures commonly escape, we have failed to detect any palpable signs to assist in explaining them, how are we to discover an obstruction, protrusion, or obtain sufficient evidence of the probability of one to justify an exploration of the obturator region with the scalpel? We reply, By pain during the development of the hernia ; by pain at the commencement of the present attack ; by pain of a peculiar character during the progress of the disease ; by pain referred to the course of the cutaneous branch of the obturator nerve and the plexus formed with its internal cutaneous and its distributions ; by pain on certain definite movements of the hip-joint ; by pain on local pressure carefully applied ; and by pain when attention of the pelvic orifice of the obturator canal is directed to the vaginam.

We shall now describe these important indications with a little more precision. During the formation of the hernia the patients had suffered at irregular intervals with vomiting, nausea, and pain resembling colic in the pelvis, and one woman related how upon repeated occasions she suffered thus, and felt relieved directly after examination as if something slipped back into the abdominal lower part. In Mr. Hilton's case, some months before the attack from hernia on the left side, the patient had with symptoms of bowel-obstruction and pain referred to the right side. After death a right obturator hernial sac was found ' more distinct than on the left side, large enough to be freely the forefinger.'

The patient narrated, in a very striking manner, how she had suddenly seized with a violent pain at the inner and upper part of the thigh. Soon afterwards, nausea, followed by vomiting and all the other indications of strangulated bowel, existed. In several of the cases there appears to have been something characteristic in the nature of the pain. Admitting the possibility of meeting with any two or more persons who would describe the pain so closely resembling each other's sensations as to be an infallible guide, yet that described during the progress of the illness seems to afford some clue in these cases. It is described as spasmodic contraction of the abdominal muscles, not as pain within the abdomen. For the explanation of this phenomenon, we may refer to the association existing between the cutaneous filaments of the obturator nerve, which is irritated by the pressure of the tumour in the obturator foramen, and the muscular filaments distributed in the abdominal muscles; for all are branches of the lumbar plexus.

The pain described by several patients, in both instances, whether a hernial tumour was palpable, as well as when it was wholly concealed, could not have failed to attract the notice of the surgeon to the distribution of the cutaneous filaments of the obturator nerve, and the more internal of the internal cutaneous. In this fact we have a most valuable aid to the diagnosis of some abnormal pressure upon its trunk, as it traverses the obturator canal. Certain definite movements of the hip-joint subservise very usefully in diagnostication, as, for example, when there may be some doubt on which side the protrusion exists, or when it chances to be so small as to be under the influence of the obturator muscles, when only in a violently contracted state. We may therefore employ the actions of these muscles of the femur outwards, to compress the tumour, and thus excite pain; and by comparing the influence of these muscles on both sides of the pelvis, valuable aid might be obtained. It has been demonstrated after death, that in some cases the protrusion passes between the fibres of the external obturator; and therefore we may suppose that they could readily compress the neck of the tumour. Want of space prevents further detail; we must therefore leave the adoption of this method, and the method of employing it, to the intelligence of the surgeon.

In several cases, the effect of local pressure over the external orifice of the obturator canal has been to cause acute pain, and

abduction and adduction.

The pelvic aperture of the obturator foramen from the vagina in the female, and the urethra in the male. In cases of difficulty and doubt, a diagnosis may be gained by taking advantage of the fact that these passages afford for the necessary examination.

Accuracy in diagnosis being of prime importance, cannot conclude this part of the subject without pointing out the great advantage to be derived from the use of Scarpa's triangle on both sides of the pelvis, impressing upon those who may happen to be examining a suspected obturator hernia under examination, the importance of seriously reflecting upon all the features of the case, to nervous derangements, as aids to the diagnosis in perplexing and difficult cases.

Treatment.—Should a hernia be detected, the pressure to the tumour may succeed. The pressure should be directed in such a manner as to push the hernia from the ramus of the pubes, and into the pelvis.

Failing to do this, or even with a weak effort, the surgeon should explore the hernia. The incision through the ligament should be either parallel with the trunk of the femur, or the fibres of the adductor longus muscle.

original incision, or its fibres being separated with the handle of the scalpel, some fat and the fascia of the obturator muscle are reached. As yet, however, the tumour has not been felt; nevertheless it may still be there, under cover of the fibres of the obturator muscle. These must be separated; for it is not until the finger can be placed upon the outlet of the obturator canal that the search for the hernia should be abandoned as hopeless. Care must be taken to avoid cutting the filaments of the obturator nerve.

How far the surgeon would be justified in making an examination of the opposite side, after having failed to find the hernia on the one first explored, is a proposition we must leave to the judgment of those who may happen to be placed in such an unpleasant dilemma.

We conclude with this observation of Camper: 'Forsan frequentiores sunt, quam quidem creditur; in ilio igitur ad huius generis herniarum possibilitatem necessario attendendum


Perineal hernia.—The sac of this hernia is formed of the perineum, which in the male lies between the prostate gland and the urethra, and in the female between the vagina and rectum. In both sexes, when the hernia fills the sac, there is a tumour in the perineum. This kind of hernia is most common in women, and usually consists of a portion of the alimentary canal, of the rectum, or of the urinary bladder.

The sac of a perineal hernia escapes between the anterior fibres of the levatores ani, and is covered by the internal abdominal fascia of that region of the pelvis.

The protrusion of the hernia may be prevented by using a T-shaped bandage.

Pudendal hernia.—Pudendal hernia forms a small elastic tumour in the labium pudendi. It lies in the posterior and inferior part of that organ, and forms a somewhat elongated projection on the side of the vagina. The neck of the sac is placed between the ascending ramus of the ischium on the outside, and the lesser sciatic foramen on the inside.

This hernia may be developed at the early age of 22 years (A. Cooper): and in this case it had existed some months. The tumours in the labium, which may be confused with this, are those formed by inguinal protrusions and collections of fluid. Of the last are cystic formations and hydrocele of the



FROM CERVICAL DYSPLASIA TO IS GROWING
tumour being placed entirely to the
ischium, and the muscles attached th

A suitable bandage is required to

Vaginal hernia.—This hernia is u
who have borne several children.
peritonæum, and that portion of the
protrudes. The orifice of the sac is
yields to pressure. Consequently it
the exception of the discomfort and a
attended with urgent symptoms. Th
however, become so much compress
sometimes suffering, may arise from

The protrusion requires to be repl
necessary support to maintain it th
use of a suitable bandage.

When the urinary bladder forms
sionally happens, great distress ari
to micturate. The urine becomes a
so irritable, that a few drops of urine,
intolerable inconvenience. Treatme
quired to improve the condition of t
the bladder, whilst the rupture m
supported.

the hernia is in the sac, it forms a tumour of variable tense, yielding, reducible on pressure, and causing less pain.

the hernia become strangulated, and the operator be to enlarge the mouth of the sac, Sir A. Cooper advises incision be carried directly forwards.

JOHN BIRKETT.

IN the following essay on Diseases treated the subjects in the order of importance: Hæmorrhoids; Prolapsure; Cancer; Polypus; Pruritus; has been said in reference to malformation of the rectum, as this subject is considered in the following chapter.

HÆMORRHOIDS

It has been the custom of surgeons to divide hæmorrhoids into two classes, external and internal. This is a classification perhaps not well to deviate; for hæmorrhoids are seen where it is impossible to divide them into internal or external,—and, pathological hæmorrhoidal tumours, wherever situated, consist of the same morbid elements, viz. an inflammation of the rectum, and an infiltration of the surrounding parts. In still circumstances which exist, or which may exist, with this disorder so modify the pathological process that in numerous instances there is a wide difference between those hæmorrhoidal tumours which are external and those involving the gut within the sphincter.

I shall therefore adhere to the classification of hæmorrhoids into two classes, external and internal.

however, they become attacked with inflammation, the tumour increases much in size, the blood in the vein becomes coagulated, and not unfrequently the vessel gives way, allowing the coagulated blood to escape into the surrounding cellular tissue, where it will form a distinct sheath for itself. In the course of time the inflammatory action subsides, the blood is absorbed, and the tumour diminishes in size, or wholly disappears. If, however, the same irritating causes recur, and no effectual treatment be adopted, the tumour again increases, the skin becomes thickened, and the cellular tissue more extensively infiltrated; and in this way distinct and permanent tumours are formed around the anus, which sometimes reach a large size, consisting mainly of thickened integument and cellular tissue enclosing veins which are at times capable of distension and repletion. In their quiet state these tumours are distinctly external; but when increased in size they may encroach upon the cavity of the rectum and be covered with mucous membrane and thus be partly internal. Coexistent with this state there is not unfrequently a very œdematous condition of the cellular tissue and mucous membrane at the verge of the anus.

With regard to the actual structure of these hæmorrhoidal tumours, it will be found on examination that they are composed of thickened integument, infiltrated cellular tissue, and in most cases of one or more dilated veins; if the part is at perfect rest and has not been lately inflamed, there may be distinguished nothing beyond infiltrated tissue and thickened skin; but on passing into an external pile which has been somewhat irritated, and is increasing in size, there will be found either a vein considerably dilated and containing semi-coagulated blood, or the blood will have escaped from the vessel, and have become either extravasated into the surrounding cellular tissue, or have formed for itself distinct cellular sheaths. This coagulation of the blood

is the reason why an incision into an external hæmorrhoidal tumour, after having become inflamed and swollen, is scarcely ever attended with bleeding, and it is one of the main pathological features in this form of tumour, and far different from what obtains in instances of the internal affection.

These external hæmorrhoids vary much in size, the swellings being sometimes not larger than peas, while at other times they are the size of a walnut. Their presence is accompanied with unpleasant symptoms: as irritation; pain when at the closet; and a sense of bearing down, even when they are in a quiescent



state : but the main source of suffering is their disposition to become suddenly enlarged and inflamed ; which occurrence will arise from allowing the bowels to become constipated, from the straining attendant upon a stricture of the urethra, from excess in food and drink, or from exposure to damp and cold. Under these circumstances, the tumour, which has been hitherto small and flaccid, becomes much swollen, distended, and livid in colour. From the peculiarly sensitive character of the skin at the anus, this distension is accompanied with the most exquisite pain ; and it is remarkable to witness the complete prostration with which the most powerful man is overcome when suffering from acute inflammation of external piles, and especially when the symptoms have been allowed to go on for some days without the proper means of relief being afforded, either from obstinacy on the part of the patient, or from insufficient pathological knowledge on the part of the medical attendant.

With regard to the causes which produce this form of hæmorrhoids, there is every reason to believe that the same circumstances which tend to the production of external, induce the internal affection as well ; and therefore this description will apply to both. It appears that there is in some persons an hereditary disposition to hæmorrhoidal affections ; and we shall every now and then see father and son, or mother and daughter, suffering one after the other in the same way. Possibly, however, this may be owing to similarity in habits, certain of which undoubtedly much more than others induce these affections. Thus, those who have to sit continually at the desk and take little walking exercise are very liable to them ; those too who have to stand long in certain positions, as dentists and hairdressers, are remarkably prone to hæmorrhoids. The great source of hæmorrhoidal affections, however, is anything which prevents the healthy return of the blood from the vessels of the rectum. And thus it is that congestion of the liver, or other obstructive disease of the same viscus, is frequently associated with these affections. A constipated state of the bowels, both from the mechanical effect produced upon the vessels, and from the straining efforts necessary, is found to be the cause in a vast number of cases ; the pressure also of the pregnant womb, and of ovarian tumours, produces hæmorrhoids. In other cases, the irritation caused by the frequent taking of aperient medicines is reasonably considered to be productive of the first symptoms of the disease. Violent horse-exercise, indulgence in the use

Highly-seasoned dishes and other indigestible food and strong passions, together with immoderate sexual intercourse, which deranges the blood more freely to the pelvic region, are each the sources of hæmorrhoidal affections. And it is highly necessary, before any treatment is commenced, to inquire carefully into the peculiar habits of the patient.

The treatment which should be adopted for the removal of the affection must be conducted upon the ordinary principles of surgery. In the more simple cases, little beyond a strict attention to ablution, to the regular action of the bowels, and the avoidance of those causes which are known to produce the affection will be necessary. If the bowels are inactive, a draught of cold water before breakfast, or the use of brown sugar, with a moderate amount of walking exercise, will in many cases beget a healthy tone in the intestinal canal; if, however, aperient medicines are needed, they should be of the mildest description. The compound rhubarb-pill in doses of ten grains, taken occasionally before dinner or before going to bed, is a simple and unirritating aperient. A teaspoonful of a confection of senna is also a useful and efficient aperient. Where there be much irritation about the anus, an occasional dose of calomel should be taken, either before or in conjunction with the aperient medicines. At the same time that great care is taken to provide a healthy action of the bowels, local remedies should be made use of. The ordinary lead-lotion, or one made of one or two grains of sulphate of zinc to an ounce of water, should be applied to the parts morning or night; or if a more powerful astringent application is required, the patient should use the compound gall ointment, which is an admirable agent. By these means, and by careful attention to diet, most of the ordinary cases of external piles presented to our notice may be cured, or so relieved that they will hardly excite attention.

If, however, one or more of these tumours become enlarged and inflamed, a much more energetic treatment is required; for there is in such cases very great suffering both local and general. Where there is much swelling, and the parts are exquisitely sensitive, the patient must be confined to bed, leeches should be applied to the part, and the bleeding should be encouraged by warm fomentations; and subsequently poultices made either of warm bran or of bread, into which half a drachm of opium is dropped, should be applied, and changed from time to time. Opium should be given internally; and as soon as



round mass on one side of the anus the circumscribed form, of a blue colour, and tension. In such an instance, the sufficient accumulation and coagulation of the treatment is to puncture the swelling; there is an escape of coagulated or semi-immediate relief to the painful symptom; application of cold water, or the lotion to the parts freely, will cause an almost entire relief. If, however, there is much loose and soft site of the swelling, it should be removed after the part has been punctured.

After repeated attacks of this nature, surrounded with distinct tumours, more liable to become swollen and inflamed; a surgical operation is required. It is simple and consists in the removal of these tumours with curved scissors. As the patient kneels upon his side, the surgeon lays hold of the tumour with hooked forceps, and excises it, with the skin. Chloroform is rarely necessary. If the patient is exceedingly timid, the parts may be anæsthetized, and thus much pain may be escaped. There is very little bleeding, especially if great care is taken of the mucous membrane.

at : therefore, unless there is an absolute necessity for the this membrane should not be cut, and only the external hemorrhoids, with portions of the redundant integument, should be excised.

Internal hæmorrhoids are more frequent, or at all events are often presented to the notice of the surgeon, because they are productive of much more distress, and more serious consequences are liable to result from them than from the affection when external : and here it will be as well to mention the symptoms which are produced by them, and which are local and general.

The first local symptom which attracts the notice of a patient suffering from internal piles is, in many cases, a more or less severe attack of hæmorrhage, which may not recur for some days or months, but which may persist ; more or less weight and uneasiness are felt at the seat ; and in course of time there is considerable pain when the bowels are being evacuated. As the swelling or swellings increase in size, the evacuation of the contents of the rectum will be more difficult and more painful ; straining efforts are necessary ; hence the hæmorrhoidal tumours become protruded on each visit to the closet. At the earlier periods of the disease they may be with facility returned ; as time wears on, the pain attending defæcation becomes more severe, and the process of returning the piles becomes more difficult. Not only, however, do they protrude at these times ; but if the patient neglects advice, the tumours come down below the sphincter whenever he takes walking exercise, and the constriction caused by the muscle produces congestion in the piles and extreme pain, which is only relieved by their reduction or by a spontaneous flow of blood, which, however, occurs at most inopportune periods. In addition to these symptoms, there is pain and uneasiness felt in the loins and even the thighs, more especially in females, who very often suffer most acutely, and not unfrequently have their sufferings referred to that prolific storehouse of morbid phenomena, the womb. There is, moreover, a considerable discharge of mucus, or muco-purulent fluid, from the anus ; the bladder is rendered sometimes very irritable, and retention of urine not unfrequently takes place.

Patients who suffer from internal hæmorrhoids are liable to have them inflamed from some exciting cause, such as an excess

at the table, or great irritation of the bowels, and then the symptoms are extremely severe; the tumours protrude beyond the anus, and become constricted by the sphincter. Violent pains are experienced in the pelvic region, and there is a high state of constitutional disturbance, denoted by flushed face, furred tongue, rapid and wiry pulse, and extreme restlessness. If these symptoms are not relieved either by the accidental induction of bleeding, or by surgical assistance, the congestion and inflammation increase, and to such an extent that mortification of the hæmorrhoidal masses ensues, and thus is produced a natural cure; but, on the other hand, it is not desirable to encourage this rude attempt at cure, for death may occur from the violent action set up. Dr. Bushe mentions having seen such a case occur.

When internal hæmorrhoids have existed for a length of time the general health becomes much influenced; the patient complains of indigestion, flatulence, an inability to follow ordinary occupation or amusement; moreover, if, as is frequently the case, the disease be attended with periodical bleedings, the face becomes blanched, the pulse weak and rapid, and other well-known symptoms of loss of blood ensue. This is the most serious condition connected with hæmorrhoids of long standing and hence the reason why it is most important to adopt proper treatment at an early period of these affections.

Internal hæmorrhoids present various appearances. On making an examination of a patient who suffers from the mild form of the affection, the veins of the lower extremity of the rectum just within the anus will be found enlarged and distended, forming small fusiform tumours of a deep-blue colour covered by a somewhat thickened mucous membrane. In other instances, and especially where the patient complains of bleeding and sense of weight, with scarcely any protrusion, the inferior extremity of the rectum, for an inch or more, will be found highly congested and vascular, the mucous membrane having here and there distinct patches of morbid vascularity from which, through a speculum, which it is necessary to use in such cases, blood of an arterial colour will be seen to issue. This is the condition which the late Dr. Houston, of Dublin, likened to the diseased lining membrane of the palpebræ in cases of chronic conjunctivitis. In the majority of instances, however, of internal hæmorrhoids, one or more distinct tumours of a rounded or oblong form will be seen to fill up, as it were, the

e of the anus. In some cases their character and size can be ascertained by an ordinary inspection; but it is always best, in order to arrive at a proper diagnosis, to throw up an injection of warm water, and allow it to be discharged before the examination is made. By this means the tumours are brought fairly out. There are frequently two or three distinct tumours, varying from the size of a cherry to that of a walnut. In one case the diseased part presents a bright red appearance, easily

FIG. 275.



Internal Hæmorrhoids surrounded by external hæmorrhoids.

When touched, is sessile and not very raised; in another case the tumour is large, prominent, of a deep blue or reddish-brown colour, having a large broad base, or being attached by a narrower peduncle, and does not bleed. In these cases the vessels appear to be largely dilated, the mucous membrane lining them being shining and tense, or thick, granular, and frequently ulcerated. Besides these appearances, portions of the mucous membrane, highly vascular and thickened, may be protruded at one or more points, as a consequence of the mechanical weight of the internal tumours. In by far the majority of

cases of long-standing piles, the integument surrounding the anus is in an unhealthy condition, being much thickened and now and then forming a distinct ring or long pendulous flaps.

There is one point of importance connected with the seat of internal hæmorrhoids which should not be overlooked, but which, as far as I am aware, has not been mentioned by any writer on this subject. The circumstance I refer to is this, that occasionally instances are met with where the hæmorrhoidal tumours are placed, as it were, in separate rows, so that two or three distinct masses exist near the anus; and about half an inch or more above, other tumours of a similar nature are disposed just in the same way. There are one or two specimens indicating this in the Museum of the Royal College of Surgeons. This is a condition of practical importance; for it shows how necessary it is to make a most thorough examination of a person suffering from internal piles. Cases every now and then occur where the ligature has been applied to one or more internal tumours presenting themselves at the anus; and as the operator is thinking his proceedings are satisfactorily terminated, the patient makes some violent straining effort, and another tumour, or series of tumours, which have escaped notice hitherto, are forced into view. These are formed higher up in the bowel and do not generally protrude; but if a satisfactory cure is expected, they must not be left alone.

As regards the structure of internal hæmorrhoids,—when first forming, they are composed in many instances simply of dilated veins, in others of dilated veins and arteries too. As the diseased condition increases, the cellular tissue in connection with the vessels becomes thickened and infiltrated. In a more or less circumscribed space, the mucous membrane also becomes thickened, and is bulged out by the increase in size of the vessels, and thus distinct tumours are formed. The surface of the mucous membrane becomes also exceedingly vascular. On making a section of the lower part of the rectum in some cases of old-standing piles, the veins will be found to be greatly dilated, sometimes partially and irregularly, so that there will be the appearance of distinct cysts; in other instances, the dilated vessels will be found to be filled with coagulated blood and fibrin. In those cases where the hæmorrhoids are of a very bright red colour and sessile, not unlike a strawberry in appearance, and easily bleeding, the structure consists mainly of a series of small arterial ramifications; but where the tumour

of a darker colour, and like a mulberry, they are composed of a large extent, although no doubt the arteries enter into their formation; for when the mucous covering is removed or incised, the blood which flows is of an arterial hue. In those cases of long standing, where the tumour has become large, and has been submitted to great irritation, a section reveals scarcely anything beyond a mass of highly-condensed thickened cellular tissue, with some vessels penetrating the interior of the tumour.

The *treatment* of internal hæmorrhoids requires more consideration than that which is adopted for the disease when treated externally. In the cases where the piles have not stood long, are not large, and give only temporary annoyance, much may be done by the patient paying simple attention to his habits, and avoiding those exciting causes which engender the disease. If it is ascertained that a sedentary life has produced the affection, by determining the blood to the rectum, the patient should take as much walking exercise as possible; if the bowels are sluggish, their action should be encouraged by a compound rhubarb-pill, or by a teaspoonful of the confection of senna; and a quarter or half a pint of cold water, or of infusion of quassia, should be thrown up the rectum daily. Dietetic measures must be strictly attended to; for many patients, especially those who are robust and whose circulation is sluggish, will tell us that they feel much more annoyance from piles after they have been dining-out, or have taken larger quantities of wine than usual. Hence the necessity for those who suffer from internal hæmorrhoids to abstain as much as possible from the pleasures of the table. Women in an advanced state of pregnancy suffering from the irritation of piles should be very careful about the condition of their bowels, and should keep the horizontal posture as much as possible.

When internal hæmorrhoids increase to such an extent as to protrude at the closet, and produce considerable pain and bleeding, greater precautions and more decided treatment are needed. The bowels should never be allowed to become costive, as to necessitate straining efforts; the protruded parts should be carefully sponged with cold water, or with a strong infusion of quassia or of decoction of oak-bark and alum in the proportion of half a drachm of the salt to twelve ounces of the decoction, and should be carefully returned by the patient; or, instead of these lotions, the gall-ointment may be smeared over

the piles with great benefit. The bleeding, which annoying, may be checked by an injection of sulphur and water, in the proportion of one to two grains to an ounce of the latter; or, if necessary, a lotion in the proportion of eight grains to an ounce, may be used. It must be borne in mind that a moderate amount of bleeding is beneficial to persons who live high, and whose vascular system is healthy; it is beneficial rather than otherwise, and should not be checked; the popular notion as to bleeding from the rectum being salutary is by no means incorrect, when applied to persons in whom it is salutary. When, however, the hæmorrhage arises from a pathological change in the tumour, such as ulceration, or excessive vascularity of the mucous membrane, and when it is continuous and goes on to such an extent as to endanger the patient's health, producing a pallid face, a weak and irritable heart, it should be put a stop to.

A very common internal remedy for piles is that of black pepper, in the dose of a drachm twice a day. It may be given by itself, or, as I often use it, mixed with a small part of confection of senna. It is difficult to say how the remedy acts; it certainly does good not only in this disease, but it is highly serviceable in other affections of the rectum, especially in those cases where the wounds become healed after the operations for fistula, or for internal hæmorrhoids.

From the close connection between the neck of the bladder and the rectum, it follows that the affections of the bladder and viscus, together with those of the prostate gland, will influence the rectum much; and thus in middle-aged and elderly persons, special inquiry should be directed to the state of the bladder, for not unfrequently hæmorrhoids and prolapse of the rectum will be found to be much aggravated, if not caused, by violent straining efforts made in the difficult attendance of the water. If stricture exists, the urethra must be dilated, for there can be any hope of curing the piles; and even when there is not any stricture, and yet there be a loss of the power of the bladder from debility or old age, this may be artificially emptied by the catheter.

When internal piles become inflamed and protrude beyond the sphincter, the patient will suffer much both locally and constitutionally. He must be confined to bed, and, if possible, should be carefully returned by the surgeon.

be a work of great difficulty from swelling and congestion, leeches should be applied, and subsequently warm fomentations and poultices. Ice locally applied in a bladder is a valuable agent to diminish inflammation and pain. Opium should also be given in full doses. An operation which may be considered feasible should not be put in force whilst the hæmorrhoids are in a state of active inflammation. Sometimes, as I have before mentioned, the constriction of the sphincter produces sloughing, and a spontaneous cure takes place; if this is occurring, the process should be expedited by the liberal use of warm bathing and fomenting, and pain must be conquered by the administration of opium.

By the adoption and right application of these remedial measures, a large proportion of cases of internal hæmorrhoids may be cured, or relieved to so great a degree as to prevent annoyance; but many of the cases which come under the notice of the surgeon have existed so long, have reached such a size, and are productive of such troublesome and even serious symptoms, that some active surgical interference is required, in order to bring about a cure or render efficient relief. Originally the usual remedy in aggravated cases was the excision of the diseased part, and it was a remedy accompanied with little pain and difficulty; but the danger of hæmorrhage proved to be so great, that after the sacrifice of several lives the practice was abandoned. It is necessary, even when the excision of external hæmorrhoids is being performed, to take care that the mucous membrane is not too freely clipped, otherwise dangerous bleeding may result. I saw a gentleman nearly lose his life from the laceration of a portion of mucous membrane in the blades of the scissors during an operation for external hæmorrhoids: the operation was done at 2 P.M.; and at 6 I was sent for, and found that he had been bleeding profusely.

Until within the last few years the radical cure of internal hæmorrhoids was effected by strangulating the tumour or tumours by means of one or more ligatures so tightly applied that in the course of a period extending from five to ten days the tumours sloughed, and together with the ligatures separated from the bowel. Indeed even now amongst the senior members of the profession this operation is practised, and there cannot be a doubt that, in the absence of a less severe method of treatment, that by the ligature was the best that could be desired, as it was unaccompanied with the dangers which attended

upon the operation of removing internal hæmorrhoids with knife or scissors, and was much more efficacious than the of treatment by strong nitric acid.

The operation is performed in the following manner. patient, who has previously had an enema of warm water, thoroughly to bring down the hæmorrhoidal tumours, kneels on an arm-chair, or lies on his side upon a bed on An assistant separates the buttocks, whereupon the surgeon lays hold, with a long pair of forceps or vulsellum, of the tumour which is to be operated on. The assistant makes traction with the instrument, so as to expose and isolate the tumour as much as possible. The operator then, by means of a well-curved needle set in a strong handle, passes a thread of strong silk or twine through the base of the tumour, and having cut the thread and removed the needle, ties the two halves of the tumour as tightly as possible. The ends of the ligatures are cut off close to the knot, the protruded part is returned within the bowel, and the operation is finished. It is better to notch the circumference of the tumour at the point where the ligature will fall before it is tied, as the separation will take place earlier: for this hint I am indebted to Mr. Curling. If there are two or three distinct hæmorrhoidal tumours, each of them must be operated upon in the manner I have described.

If there is much loose integument about the anus, its redundancy should be removed with the scissors. Too much ought not to be taken away, as inconvenient contraction might take place; but the surgeon who understands his business will not be likely to fall into an error of this kind. If the flaps of thickened integument which are so frequently seen in connection with hæmorrhoids are not taken away by the surgeon, there is a probability that the cure will not be perfect. Several cases have lately been under my care where the ligatures had been applied by various surgeons, and where the disease had returned. In some of these cases the external folds of integument had not been removed, and I cannot help believing that this neglect had been the cause of the failure of the operation. The proceedings above detailed are finished in a few minutes. If the patient is courageous and determined, I would not operate without giving him chloroform; but if he is highly sensitive and timid, it would be better that he should inhale ether, as the pain in such instances may be very severe.

The patient must keep his bed for some days after the operation. A full dose of opium should be given the first night, so that pain may be prevented and the bowels be confined; if there be much pain about the seat of operation, ice applied in a bladder continuously will give great relief. Retention of urine is to be avoided after this operation, and when present must be relieved by the catheter. It is desirable to keep the bowels quiet for three or four days, if possible, and then to obtain an evacuation by a dose of castor-oil. Very likely the ligatures will separate on the first action of the bowels; at all events they generally come away on the fifth or sixth day. Some pain is felt in the part a few days afterwards, and during this time the patient should keep quiet: convalescence may be expected in a fortnight after the time of the operation.

This is the general course of things when the operation is successful and unattended with any evil results of any kind; but it is well known that the use of the ligature is occasionally followed by very serious results: and, indeed, several instances have been made public where death has ensued after this operation, the cause of death being effusion on the brain, pneumonia, and pyæmia, the latter disorder most frequently. Several fatal cases have occurred in the practice of my own personal friends, and I have had the misfortune to lose two patients after the use of the ligature. One of these patients was a gentleman in the prime of life and very fair health, who died with the well-marked symptoms of pyæmia; and in the other case, a fatal event was due to the loss of blood during and after the operation.

But besides the risk of death from some of these causes after the operation, it is known that other serious results occasionally follow and give much trouble.

Thus I met with a case of a lady who had lost much blood prior to the operation. I had to employ several ligatures, so that the operation was of necessity severe and prolonged, and it was followed by the most intense pain, so that, notwithstanding the employment of large doses of opium, frequently repeated, I was fearful that she would actually die from the effects of the severe pain she experienced, and which did not lessen until the ligatures separated, at the end of a week. After the subsidence of this, other symptoms manifested themselves, and caused great alarm, culminating in fact in a severe attack of lumbago of the left leg and thigh, so that she was not enabled to leave the house to go out for a drive until a period of six weeks had elapsed.

In another instance which occurred to me not long after this, I employed ligatures for removing a mass of hæmorrhoidal tumours. The patient was a healthy middle-aged man, of temperate habits, living in the country; the

of long-continuing and painful ulceration more of the ligatures, for a period varying Everyone who has had much to do with have met with such cases. These ulcerat to heal, and give great annoyance both geons.

To sum up briefly then, it may be st there are the following objections to the u The supervention of more or less pain; a t occasionally death from pyæmia, tetanus, tional disturbances, and long-continued a at the seat of ligature.

It is the duty of the surgeon to adopt s has them in his power, by which these res and by which the disease can be cured ; enabled to effect this by employing a mod is now being pretty extensively used, and under the name of the operation by the clæ which I will now proceed to speak of ; t that modification of the operation which v profession by myself, I must briefly refer t mode of treatment, which probably will alm the employment of the ligature.

As far as I am aware, this plan of tr tumours was originally practised by the

There was another kind of instrument, shaped somewhat like a pair of scissors and which was much more suited for the purpose.

In the year 1863 I first began to treat cases of internal hemorrhoids and prolapsus by the clamp, employing, after division, the strongest nitric acid for the purpose of obviating bleeding. I soon became convinced of the value of this mode of treating these disorders, but in a short time I found out that very great improvements in the construction of the clamp were needed for, and, with the assistance of the late Mr. Matthews of Portugal Street, I introduced to the notice of the profession the instrument which bears my name and which is now generally used. It is shaped somewhat like the ordinary clamp used by Mr. Curling for the purpose of seizing and applying nitric acid to hæmorrhoidal tumours; but the blades are so constructed that they meet accurately at their edges, and are

FIG. 276.



ly fitted into one another by means of a raised surface on one edge and a groove on the other edge. In this way, and if the blades are made perfectly parallel, the most perfect and accurate compression is made upon a tumour or portion of mucous membrane. A more important feature, however, in the mechanism of this instrument, is the adaptation of a light yet powerful spring to the shank of the instrument, just above the handles. The object of this is twofold—in the first place to compress the tumour, and in the next to take off the compression after the actual cautery has been applied; and by the slightest turn of the handle, this may be effected so gradually, and yet so thoroughly, that should any vessel be left uncauterized, its situation can be at once seen by the issue or jet of blood, and the cautery can be reapplied.

The latest, and a very important addition to the instrument, consists in the adaptation of a layer of ivory to the outer surfaces and external edges of the clamp, so that the heat from the cautery cannot be transmitted to the surrounding parts, and thus much suffering is prevented.

I will now proceed to describe the mode of application of this instrument, and this description will apply both to cases of hæmorrhoids and prolapsus of the rectum.

In the first place, the part to be operated upon is made to descend either by means of an enema or by the patient sitting over hot water: the tumour is then seized by a pair of hook forceps, and firmly held by an assistant. The clamp is not carefully applied to the base of the tumour, and the blades are closed by means of the screw. After having seen that this is done effectually, the surgeon takes a sharp pair of scissors and cuts off the whole of the tumour; the cut surface is wiped, and carefully and repeatedly touched by the heated iron until it is considered that the entire surface is thoroughly cauterized.

The screw is now gently turned, the grasp of the clamp is thereby somewhat relaxed, and if any vessel has not been sealed up it will emit blood, and the cautery can be again applied. Sometimes this process may be needful two or three times before the bleeding is quite stopped. As soon as it is ascertained that the surface has been thoroughly cauterized the clamp is removed, and if more than one tumour exists each of them is successively dealt with in a similar manner. The finger, oiled, is then introduced as far up as possible into the bowel, so as to push the parts well up and cause the sphincter to contract; a suppositorium of two grains of extract of opium is introduced, and the operation is complete. If only one tumour requires removal, the proceeding may be finished in a minute or two; but if there are several tumours, the operation cannot be safely or satisfactorily concluded under ten or fifteen minutes.

There are one or two points connected with the operation which demand attention. In the first place, the facility of the whole business depends greatly upon the extent to which the parts are exposed. If the tumours are well brought down on one side the anus, the clamp may be applied very readily; but when the tumours are not thoroughly protruded, or are very small, it is very difficult to apply the clamp. In some instances when the skin of the anus and the mucous covering of the hæmorrhoids

rhoids run into each other, I apply the scissors at the point of junction before applying the clamp. It is well not to cut the tumour too close to the upper surface of the clamp, but to leave about the eighth of an inch of the tissue, which can thus be more readily acted upon by the cautery. On no account should an endeavour be made to embrace more than one of the tumours within the blades of the clamp at the same time. I generally apply the cautery either at a black or dull red heat; if it is too vivid, the agency is more destructive than is needful. We should err rather on the side of a too liberal application of the hot iron than otherwise.

With regard to the employment of chloroform, I am guided by the following considerations. If the patient resolutely demands it, of course he must use it; if at the same time that the operation on the internal piles or prolapsus is being done, any extensive removal of skin be required, I recommend chloroform; or if the patient objects to that, I use the anæsthetic ether, which, when thoroughly applied, answers very well in mitigating, if it do not altogether prevent pain.

For the purpose of locking up the bowels, the same precautions as are needful in other operations on the rectum are taken. At the end of the third or fourth day a dose of castor-oil is given, and the bowels are generally easily moved; a little blood which has collected in the bowel either during or after the operation may come away. If the patient has not been much weakened by long-continued hæmorrhages, he or she may be allowed to get up; and in by far the majority of cases they do so; but if the health has been much shaken, or the operation has been extensive and has involved the removal of much skin, it will be advisable to keep the patient in bed or on the sofa until the bowels have been acted upon a second time, two days afterwards, when it will be found that the patient is able to move about or attend to his ordinary business.

I have now performed this operation in upwards of 200 cases, and I can with truth state that, if it be properly done, there is hardly an element of danger in it. I have operated at all ages, from puberty up to nearly eighty, and in instances of the utmost severity, in cases where the tumours have been immense, and when the patients have been so thoroughly blanched by long-continued bleeding, that one would not dare to perform any operation, unless convinced of its absolute safety. I have not met with a single instance where secondary hæmorrhage has

but I had removed freely some external the use of a bougie for some time soon tion.

In two instances death followed the ference to the details will show at once tl each was a post hoc, but not propter hoc—

In the one instance the patient was a lady ætat dreadfully from hæmorrhage for many years, and I health that for some time before I saw her she could and in this state the prolapsus came down, and returned. I was summoned into the country to see in such a feeble state, I, in conjunction with her me best to relieve her of this annoying cause of irritati clamp and cautery on January 26, 1868. The opera with loss of blood. On the fourth day, the bowels and so far as the actual operation went, it was, to use attendant, 'very successful,' but she never rallied fr been suffering, and died from exhaustion on the thirt

In the second case, a lady between fifty and sixty in October 1868. She also had been suffering for yea bleeding from internal piles, and when I was cons wretched state from bloodlessness. The operation wa loss of blood, and she was able to get on to the so this day a favourite daughter suddenly came to see l the most violent state of hysteria, which lapsed i coma, in which she died on the thirteenth day, wit local mischief about the wound, which was nearly hea ascertained that a favourite son of this lady had sudden that then she had a violent attack of the same nature

case the patients were advanced in life; and the operations attended with as little disturbance as in perfectly healthy individuals.

In one instance I refused to employ the clamp, and this was in the case of an aged gentleman, to whom I was called by Mr. Cross of Petersburg, who had a very large prolapsus. The patient was otherwise seriously debilitated, and some indications of cerebral mischief; and although the disorder had evidently aggravated his condition very much, I quite agreed with Mr. Cross that it would not be wise to operate even with the clamp, and the wisdom of our determination was shown in a few days by the sinking and death of the patient.

The advantages which I claim for this proceeding, which I always adopt in cases of hæmorrhoids or prolapsus, are, first, there is considerably less risk to life than when the clamp is used; that in the majority of cases the suffering both during and following the operation is much less, and that the period of convalescence is rendered much shorter. It is true that the operation itself is complicated and somewhat tedious, as compared with that by the ligature, but these considerations ought not to weigh with the surgeon when the advantages of any particular proceeding are great and palpable. Nitric acid was strongly recommended by the late Dr. Keane of Dublin as a means of getting rid of some forms of internal hæmorrhoids; and it is somewhat curious that, although the paper was published in the *Dublin Quarterly Journal* more than fifteen years since, and Sir W. Fergusson, in his *Practical Surgery*, soon afterwards drew especial attention to the subject, yet the use of the agent is still comparatively unknown, or at all events has not had that attention paid to it which it deserves. I have been in the habit of using the strong nitric acid in some cases of hæmorrhoidal affections for many years. Mr. Lee has incorporated with his *Surgical and Pathological Essays* an admirable paper on this subject, and has therein highly of the plan of treatment which I am now about to consider.

I have stated, whilst describing the nature of internal hæmorrhoidal diseases, that the condition in which they are found varies. I have endeavoured to point out those cases to which an operation is applicable, and have stated that this mode is justifiable in those instances where the tumours are chiefly composed of tissue in which the veins predominate, and have become indurated.

There are, however, other instances where the hæmorrhoidal

prove most destructive to the health, and a great deal of blood. Now, in such cases it undoubtedly be as effective as in the cases described; but this proceeding is not so of the nitric acid is so eminently suitable, which one single application of the acid is remarkable: and an excellent cure may be effected on the diseased texture be subjected to

About this particular kind of case I have in the mind of any surgeon who has seen the operation performed in the proper manner. There is, however, a case where the remedy is an uncertain one, where the surgeon is justified in trying it, and where it succeeded when I little expected it. In a case where there is a hæmorrhoidal mass, or a tumour mainly composed of venous matter, of a dark colour, with one or more presenting the appearance of a sessile pile; or one portion of the tumour presenting the dark blue appearance and another portion of it may be brightly red, the mucous covering granular or slightly fissured. In a case of mixed case I do not hesitate to try it, and it is particularly desirous; but I make it impossible to depend upon any case, although in some instances the

Houston has compared this condition of the rectum not to that of the thickened conjunctiva after long-conjunctivitis ophthalmia. The application of the acid to the diseased part from which the bleeding proceeds will soon remedy all the symptoms.

The following is the mode of applying nitric acid:—The patient having been well evacuated some hours previously, the part of the rectum to which the application is going to be made should be well exposed by making the patient sit over hot water for a few minutes; or if this is not sufficient, an enema of oil should be thrown up the rectum, and the hæmorrhoidal tumour will be brought well into view. The part to which the acid is to be applied should then be carefully wiped with a piece of lint. The surgeon then dips the extremity of a small piece of wood into the nitric acid, and touches the diseased part carefully with it. The part touched, and the neighbouring mucous membrane, is well smeared with oil, and the patient is returned within the orifice.

I can truly say that it has never occurred to me to witness any thing like a fatal, or even a dangerous result, after having had a large experience of this remedy. In one case, of a patient who was most anxious to be cured by one operation, I used the nitric acid much more freely than usual, and produced great suffering for two or three days, with the effect, however, of making a good cure. In another instance, I heard, but was not a witness of the fact, that copious bleeding followed the use of the acid. In a third instance, which occurred very recently, a young lady was treated with the nitric acid for a florid hæmorrhoid. Severe and unaccountable suffering was produced for several days. On making a careful examination at the end of the period I discovered a small ulcer situated at the posterior part of the anus, and exquisitely painful; suitable remedies relieved the pain in a few hours. On enquiring more minutely into the particulars of this case, I have reasons for believing that this ulcer existed before I applied the acid, and that some other caustic came into contact with the sore and produced the severe suffering.

I have now and then met with cases where retention of urine and some bleeding have occurred after the free use of the nitric acid, but never sufficient to cause me anxiety; and they are symptoms which are easily met.

It is not necessary to confine patients to their bed after the


surgeons in this country. The object of the instrument is the more or less rapid removal of the hæmorrhage. That this can be done in a number of instances there is not a doubt. If it were likely to follow, this practice would be adopted in many of those cases where it is not adopted notwithstanding that there is a great objection upon the process. One very serious objection to the plan of treatment has been met with in Paris: this is the occurrence of stricture after the wound has become cicatrised. This objection to the employment of the instrument is counterbalanced by its real or supposed advantages. It would not be wise entirely to discard it, for it must be rare, and in those few instances where it is required, the surgeon would do well to avoid the occurrence of contraction.

PROLAPSUS OF THE RECTUM

During the time that the rectum is protruded in a natural and normal manner, more or less of the bowel occurs; but this is only momentary. When the action is finished, the mucous membrane is drawn within the anal orifice and no

placed in the situation of the prolapse, the protruded part is found to form a tumour of an oblong and cylindrical form, presenting externally the smooth surface of the mucous membrane, which is generally of

FIG. 277.




Prolapsus in Child.

more or less bright red colour, and covered with mucus; at the extremity of the tumour is the orifice or cavity of the bowel, at the anus there is no deep furrow between it and the protruded part, as there is in the intussusception of the rectum.

In the adult, when the prolapse is large and of recent occurrence, this mucous membrane may be as unchanged in appearance and texture as when it occurs in the child, but the

only somewhat thickened and more vascular than natural; should the bowel have been habitually prolapsed for some years considerable changes take place, and on examining an old case the following will be the appearances. Externally there will be a ring of thickened integument; within this the flap of mucous membrane hang down, their lower portions being more thickened, having lost the peculiar character of mucous membrane, and being assimilated to integument; this change is taken place because the most dependent portion is that which either habitually remains protruded altogether, or is protruded for a longer time, and more exposed. On separating the flap of the prolapsus, the upper part of the membrane is found either but little altered from its natural character, being red, smooth and vascular, or superficial ulceration may have taken place

FIG. 278.



Prolapsus in Adult.

where the two portions have been in contact. There is more or less mucous discharge produced, but in pure prolapsus there is little hæmorrhage.

Sometimes, in persons advanced in life, the protruded part consists of a tumour as big as the fist, which has habitually protruded for a long time. In such a case, a very large proportion of the tumour consists of membrane more like leather than the natural tissue. In these old-standing cases the sphincter becomes extremely relaxed, and the anus very capacious; there is generally a redundancy of loose and thickened skin around it, sometimes it hangs down in long pendulous flaps—this state of the parts adds materially to the facility of the occurrence of prolapsus.

In many cases the prolapsus of the rectum is complicated with distinct hæmorrhoidal tumours, which in fact are mainly, if not entirely, the originators of the affection; for when one or more internal tumours exist, they themselves, each time they act, become protruded, and drag portions of the mucous membrane down with them; so that not unfrequently a patient presents himself with one or more folds of prolapsed membrane, at the same time with distinct hæmorrhoidal tumours.

The inconvenience and suffering which prolapsus causes is considerable, for although at the onset of the affection the protruded part may pretty readily return within the sphincter after an evacuation, as time wears on it becomes necessary for the patient himself to return the part, which not rarely is a matter of difficulty, and attended with pain; moreover, from the proximity of the rectum to the neck of the bladder and urethra, there is often great distress of these parts,—constant irritability, and even retention of urine, being an accompaniment of the affection; pain and uneasiness is felt in the loins and in the thighs: the intestinal canal and stomach also sympathise, the patient being troubled with flatulence, loss of appetite, and low spirits.

When the prolapsus cannot be returned by the patient, most of the symptoms occur—extreme pain in the part, and retention of urine; and if unsuccessful attempts are made to reduce the swelling, which is in all probability tightly constricted by the sphincter, violent inflammation of the part, attended with general constitutional suffering, occurs; and in some instances strangling of the protruded bowel takes place, by which means death is brought about, but the mischief may be such as to lead to death. In cases where a prolapsus occurs in children to a great extent, and has been allowed to remain down for two or three days, the local and constitutional changes are not so severe; the prolapsed membrane, however, becomes exceedingly congested.

The causes of prolapsus are constitutional and local; thus, the disease is very frequently met with in individuals who have suffered from general debility and laxity of fibre. In children especially the affection is met with in instances where the strength has been much reduced by insufficient nutriment, bad management, and want of proper attention. An adult or old person who suffers much from prolapsus, usually has a weak pulse, a flabby tongue, and impaired digestion; and in the child there is an

In considering the treatment of prolapsus, shall first refer to that which is necessary in affection as it is met with in young child; it is necessary to seek for its cause; and be made with reference to the urinary apparatus. frequently happens that it is the irritation of the bladder which produces the extrusion of the stone. If there be not stone in the rectum, the ascarides in the rectum may originate the destruction of these parasites by a few doses of calomel, together with the daily injection of a strong infusion of quassia, will prevent the

In by far the larger proportion of cases, the general health will be found at fault, and should be attended to before the prolapsus can be got into place, it is necessary to return the protruded bowels sometimes a work of difficulty, because the child violently and cries. The protruded bowels, when firmly grasped by the right hand, well oiled, and as to empty the vessels, should be employed, and returned within the sphincter. When the child is large, however, and the child is very violent, it will fall again; and in this case the best plan is to keep the child under the influence of chloroform.

may be bathed with a solution of sulphate of iron, gr. ss . of water; or an injection of tincture of sesquichloride of iron, ss . to ʒvj . of water, may be thrown up every morning before the evacuation of the bowels, and after the protrusion has been returned. In some cases the prolapsus will recur when the child evacuates. This accident, however, may be prevented by so managing that the child should be in a kneeling posture during the act. Another plan which sometimes succeeds in preventing the protrusion, consists in an attendant pulling on one side the skin of the anus with some force during the time the bowel is being emptied; by this means a certain degree of temporary contraction is produced, which prevents descent of the gut. In adults, a considerable number of the cases which are not severe, and which have not been of long standing, may be cured by careful attention to the removal of the causes which have produced the disorder. Thus, if it has resulted from violent straining and constipation of the bowels, a mild aperient should be exhibited occasionally, which will prevent accumulation in the bowel, and render the contents more fluid. The compound rhubarb-pill at night will have this effect; or, what is perhaps better, one or two teaspoonfuls of infusion of senna should be taken. The patient should not eat heartily, and should especially avoid vegetables in any quantity. He should take exercise, and be careful to use plenty of cold water to the parts after the action of the bowels. Occasionally a little cold water, or a few ounces of the decoction of oak bark, may be thrown up the rectum; and if there be the least inflammation left after the evacuation of the bowels, the gut, after being first well sponged, should be carefully returned. Attention to these various measures a prolapsus of small size may either be entirely cured, or be prevented from increasing or proving troublesome. And therefore it is of the greatest importance to place reliance upon medical treatment in the early cases. In by far the majority of cases, however, which are presented to the notice of the surgeon, the prolapsus is either very large, or has existed so long a time that medical treatment would be of no use whatever. And then some strictly surgical means must be adopted, if a cure, or even if palliation of the disease, is looked for. If the case is of only recent occurrence, and yet the prolapsus be very voluminous and incapable of being returned, thereby causing much alarm and suffering, it is the duty of the surgeon at once to reduce the prolapsed

return of the prolapsus, the whole or a portion of the mucous membrane should be smeared over with silver, previous to its being returned by the remarkable case of immense size occurring, he adopted this plan, with the result of obtaining a bandage, which he continually had worn.

In order to bring about an effectual cure in severe cases of prolapsus, more decided measures are required. We have seen that the disease essentially consists in a relaxed and thickened condition of the mucous membrane, and that the muscular contraction, as it were, of this from the mucous membrane is involved also, there ensues a weakness of the whole of the thickness of the bowel from the rectum to the anus. The object to be obtained is to restore the normal contraction or relaxation of the mucous membrane between the several tissues composing the wall of the rectum up to the anus and the sphincter. The first was the first to propose a proceeding which would produce this result; and this consisted in removing the redundant flaps of skin which existed around the anus. This was the case alluded to by him in his *Practical Surgery*, p. 443. In some cases, where the rectum is relaxed, and the flaps of integument large, it may be brought about by the removal of a portion of the prolapsus is very large, and a considerable

necessary; the operation is painful, but is soon accomplished; the wounds heal, contraction takes place, the aperture of the anus becomes diminished and braced up, and the prolapsus no longer occurs.

It is important to bear in mind that, in very severe cases, not only is it necessary to remove the relaxed integument, but portions of the mucous membrane, which in instances of long standing has become converted into a tissue more like leather than anything else, must also be taken away: if this step be resorted to, a disappointment will ensue as regards a complete remedy of the prolapsus. Hence the surgeon must think of the possibility of somewhat severe hæmorrhage, which will occasionally occur. After a portion of the mucous membrane, however small, has been snipped away, I have seen it occur to a very great extent, and when it was least expected.

Hæmorrhage to a large extent does occur after a surgical operation on the rectum, the patient will in a few hours complain of tenesmus, and express a desire to go to the closet; he then evacuate a large quantity of blood, and become faint. In such a case it will be necessary to clear away any coagula which may be in the gut, to elevate the pelvis, and introduce ice into the bowel. Should this not stop the bleeding, a digital examination should be made with a speculum, and the bleeding orifice be looked for and tied. Sometimes it will be difficult, or almost impossible, to effect this; and then the anus must be carefully plugged by portions of sponge or lint, through which a thread should be tied, that the compress may be readily withdrawn when the bleeding has ceased.

It may here be stated, that the risk of severe hæmorrhage after the mucous membrane has been excised may be in a great measure obviated by the surgeon taking care to introduce, through the edges of each incision, one or more fine sutures before the patient is left.

Another mode of curing prolapsus consists in the application of the ligature to portions of the prolapsed membrane. This is especially adapted to those cases where there is great redundancy of the mucous membrane, and where the surrounding integument is not much involved; also to those cases where the prolapsus is associated with one or more hæmorrhoidal tumours. This operation was originally proposed for prolapsus by the late Mr. Copeland, who found it to answer his expectations most admirably. It is easily done, by pinching up, with

the disappearance of the tumour.

In either of the operations just described, a certain amount of danger, and they come to bed and to the house for a week, and there is any other agent by which the prolapsus without producing either the danger to the patient to bed, it should be adopted. The solution locally applied to some forms of hæmorrhoids, so well, has lately been used by me in the most standing cases of prolapsus with considerable success; however, only in certain forms of the disease, it will act beneficially. In the cases of long standing, and of very long standing, where the membrane has become very much thickened, and characterized by its appearance, the acid will do little or nothing. In cases of simple prolapsus of the bowel, where the more large folds of mucous membrane, the membrane is extremely vascular, presenting the appearance of a raw surface, or is perhaps superficially ulcerated, a strong nitric acid, applied carefully to a small portion of the diseased membrane, will cure it. It should be used in the same manner and with the same cautions as when employed in instances of hæmorrhoids. The entire surface of the prolapsed membrane, if it is not too large, one application alone will suffice to


mucous membrane, a very weakened condition of the sphincter, an abundance of loose and thickened integument around anal orifice; consequently the use of nitric acid alone will suffice to remedy the disorder, even if the mucous membrane not much altered in its character. In such cases I combine use of the nitric acid with the removal of two, three, or slips of the thickened integument from around the margin of the anus. The acid should first be thoroughly applied to the mucous membrane on one or more occasions; and after the lapse of a few days the subsequent part of the treatment should be adopted by using the scissors in the manner before described. Beneficial results will follow from this practice, and it may be undertaken without fear of danger in cases where the ligation would not be justifiable.

Although either of these methods of treatment may be adopted with success, I now almost invariably remove a protrusion in the adult by the operation with the clamp and cautery, in instances of hæmorrhoids, and in precisely the same manner. If the tumour be small, it is only necessary to remove one or two folds of mucous membrane; but if, as in some cases, it be very voluminous, it is needful to apply the clamp to several portions of the protruded membrane. If there be great distention of the anus with redundancy of skin, two or three slips of the latter should be removed with the scissors, as in the operation of Dupuytren. I find this treatment is just as successful as when it is practised for the removal of hæmorrhoids. In those cases where any operative measure is not advisable, great relief may be obtained by the use of a pessary or spring.

FISTULA IN ANO.

The artificial communication between the cavity of the rectum and the neighbouring textures denominated fistula in ano constitutes an affection of very great interest and importance, as there is a difference of opinion regarding the exact pathology; and for the most part an operation more or less severe is required for the cure of the disease.

Very opposite opinions have been propagated by eminent surgeons regarding the formation of fistula. Thus, it has been confidently asserted by Sir B. Brodie that the origin of the disease was the existence of an ulcer in the mucous membrane, and that the suppuration in the neighbouring textures was



ulceration of the mucous coat of the
taneously and as the result of injury pr
ulceration having occurred, some of th
escape into the surrounding cellular t
tion.

In the majority of cases of fistula ir
formation of the disease is the existe
cellular tissue surrounding the bowel
the various textures are involved, th
pressed upon, the mucous and muscu
a greater or less extent, and at last
penetrated.

It has been asserted that the correc
pathology of fistula is of little conse
impossible to mention a surgical disea
ful to know the steps of its formatio
the malady, which must ever be looke
tance than its cure, depends almost en
of the fact that in a great number o
side the bowel is the cause, and not th

When suppuration has taken pla
around the rectum, and the matter
cavity of the abscess commonly does
probably of the periodical disturbanc

are kind, where there has been extensive burrowing of matter, there are two or three openings situated at a greater distance from the anus—as, for instance, over the ischium or in the perineum—and the matter may even extend upwards, and open into the groin.

The external opening may present itself as a simple minute aperture, and when situated close to the anus may be so hidden between the folds of the integument as readily to escape observation, unless it be carefully looked for; the only thing which betrays its presence being a slight moisture escaping from one when pressure is made. In other instances of more extensive and old-standing disease, the opening or openings present themselves as somewhat prominent apertures, not unlike those seen in connexion with dead bone, and surrounded by more or less erythema of the skin, so that there is not the great difficulty in finding them.

Although for the most part it will be found that the external opening of a fistula is by the side of and some little distance from the anus, it is well to know the fact that sometimes it is so close to the circumference of the anus that it is necessary for the patient to protrude the parts well before this aperture can be discerned.

An external opening does not always exist, and then is constituted what the older surgeons named a blind or incomplete fistula. This form of the disease is somewhat rare, and it is very likely to be overlooked or misunderstood.

The internal opening of a fistula nearly always exists; and indeed, according to that view which looks upon the formation of the disease by preliminary ulceration of the mucous membrane, an opening must necessarily obtain; but there is no doubt that there are cases of incomplete fistula where there is no opening into the bowel; an examination on the living body might not be a sufficient proof of this, for the opening may exist, and not be ascertained; but no such source of deception can occur after death. There are three specimens in St. George's Hospital Museum, as well as two preparations in the Museum of St. Bartholomew's Hospital, which show this. On careful examination, however, it will be noticed that the mucous membrane of the rectum at the spot where the opening would have existed is very much thinned.

Formerly it was considered that the internal opening was situated much higher up than is really the case. Subsequent

cases are occasionally met with where distinct apertures in the gut. And how this occurs; for in some of them near the rectum the matter extends a rating it from the cellular tissue to the inches; and it is but likely that they become thinned, and ultimately perforate the matter at more points than one.

The course which the sinus takes is subcutaneous. Very frequently it runs to the external sphincter, or it may be distant from the levator ani. In one very specimen at St. George's Hospital Museum the sinus may be running closely beneath the fibres of the levator ani.

Sometimes the result of extensive disease of the lower part of the rectum is the formation of either complete or incomplete; that is, a fistulous tract on either side, having an external aperture; or a sinus may enter the bowel, and yet there be only an open aspect; or there are two sinuses, one entering into the bowel and on the skin, which has an internal aperture. There is occasionally a condition, where the fistulous sinuses surround the rectum, and the matter issues from several points.

the causes which produce fistula are various, and, understanding its pathology, we must look to those circumstances which are liable to excite inflammation and suppuration about the anus. In some rare cases the disease is of a traumatic origin.

Thus a fish-bone may have adhered to the mucous membrane of the rectum, and produced ulceration and subsequent abscess, which has degenerated into fistula; or the ulceration may have primarily taken place in the mucous membrane in persons who have suffered from severe dysentery. In the greater number of cases, however, when abscess has been the primary step, the morbid action has been idiopathic, occurring in persons who have been out of health, or who have been ill with some specific disease. Those who have lived long in tropical climates, and who have suffered from disease of the rectum, are doubtless more liable than others to get fistula. A violent blow upon the perineum or ischio-rectal region will produce abscess which will terminate in fistula. I have recently treated upon a very healthy young man where the disease could be traced to this source.

When an inner opening exists without any external aperture, diagnosis is somewhat difficult. The patient will complain of having suffered from more or less pain about the rectum; this gradually becomes more severe, until there was perhaps some alleviation, accompanied with a discharge of purulent matter from the anus. On examination, the surgeon will detect a distinct and circumscribed induration by the side of the rectum, very painful to the touch; and on pressing the part towards the anus purulent fluid will be observed to escape in considerable quantity. And when the integument has become thickened, fluctuation may be discovered; but not in all cases.

It should not be taken for granted that, because a sinus is close to the anus, it must be connected with the rectum; abscess and fistulous openings may exist in connexion with the rectum in other parts. Thus sometimes an ordinary pelvic abscess may discharge itself close to the anal aperture, or an abscess in communication with the hip-joint will open into the ischio-rectal fossa. An abundant discharge from a cavity connected with the prostate gland took place in this situation in a recent case under my own care. Necrosis of the tuberosity of the ischium or of the extremity of the sacrum may exist, and an aperture in connexion with the disease may be present in the same situation.

and a very common, and one which is recovered.

In order to bring about the cure necessary in the great majority of cases, operation, which fortunately, from the disease by Pott and others, is a matter now than it was in former days, are used in selecting the cases, is almost success.

Before, however, proceeding to the treatment for the disease when the fistula has been formed, it is necessary to make the importance of dealing promptly with the suppurative process which forms the fistula. When a patient is suffering from the abscess near the rectum, he should lie on his left side, and the bowels should be thoroughly purged with calomel, followed by castor-oil; hot fomentations should be assiduously employed, and by the early adoption of these measures the inflammation may be arrested, and all the symptoms subsided. It is a good reason to believe that matter will be necessary, in order to prevent the necessity of an incision into the centre of this swelling. In some cases, where the

In many cases, where the inflammation of the cellular tissue has not existed long, and where the general health is not much impaired, an abscess close to the rectum will completely heal by the prompt evacuation of its contents, and thus the fistula will be prevented. Hence is shown the vast importance of properly understanding the pathology of this disease, and thereby being able to arrest it in its first stage.

In those cases, however, where there has been extensive misadventure, and the lower part of the rectum has been much denuded, where the general health has been weakened from this or from other cause, even this prompt surgical treatment will not prevent the abscess from degenerating into a fistula; and in other numerous cases, the abscess has been allowed to burst, and a stultous communication between the gut and the external surface has been formed.

In some cases where the sinus is short and free from induration, it may be made to heal by the employment of irritating substances thrown into the canal. Those which are likely to be of most service are injections composed of the pure tincture of cantharides, or of the tincture of iodine undiluted, which should be used daily. Another method of healing these fistulae consists in irritating the sinus by passing a silver probe covered at its extremity with fused nitrate of silver. Each of these cures is occasionally successful, especially when, at the same time, the general system is improved by medical treatment and by change of air. Therefore, in those cases where there is no cause for hurry, or where the patient is unwilling to undergo an operation, these means of cure should be tried persistently, and both patient and surgeon will sometimes be rewarded with success.

Fistula in ano may be treated successfully by ligature, and it is a method which has of late years been warmly recommended by Mr. Luke. The operation is effected by passing a common thread, by means of an eyed probe, through the external aperture into the bowel; then adapting the two ends of the thread to a small screw, which can be tightened from time to time as the ligature is cutting its way out from the fistula. The process of separation is generally completed in a week or ten days; and as the ligature ulcerates away, the cavity behind it comes gradually filled up. The mode adopted and the instruments used by Mr. Luke are described in the *Lancet* for 1845, p. 221.

There are some advantages in the ligature. Thus, it may be employed without fear of producing hæmorrhage; and therefore in those rare instances where the inner opening of the fistula is situated high up, and where serious bleeding may be expected to arise after a cutting operation, this mode of treatment should be adopted. Again, there are patients every now and then to be met with who are so nervous about a cutting operation, that even with the promise of chloroform they will not submit to the knife; in such the treatment by ligature can be advantageously substituted. It is not a method adapted to cases where the sinuses are very tortuous and extend in various directions; but in those instances where the fistula is simple, there is no doubt that the application of the ligature will be followed by success, although the process is a somewhat tedious one, and sometimes it has to be abandoned in consequence of the pain and annoyance caused.

In by far the majority of cases of fistula in ano, an operation by the knife—that commonly denominated *the operation for fistula*—is needed, if a satisfactory and efficient cure be looked for. This consists in the division of the structures situated between the sinus and the cavity of the intestine, and, in fact, the laying open of the entire fistula from one orifice to the other. The proceeding is rendered necessary in many cases of obstinate sinuses situated in other parts of the body; and it is easy to understand how imperatively it is called for when a sinus implicate a part periodically acted upon by a powerful sphincter muscle, whose contractions alone will suffice to prevent the healing of the fistula, even were there no other causes at work to impede this process. The object, therefore, of the surgeon in performing the operation is not only to lay the sinus open and thus place it in a more favourable condition to heal, but by dividing the sphincter muscle to paralyse its action for a time and thus to keep the wound at rest—a proceeding found to be essential for a cure, not only here, but in the treatment of wounds, sores, or injuries in other parts of the body.

The operation, as performed now, is, in the majority of cases, a simple one, compared with the same before, or even after, the days of Pott, whose writings tended so powerfully to diffuse correct views regarding the pathology and treatment of fistula in ano. Instead of excising the fistulous tract, or even of making very free incisions high up in the gut, the surgeon only finds it needful to divide those structures which are limited by

Two orifices of the sinus; and it is now pretty clearly ascertained that, in the majority of instances of the disorder, the opening is met with at a point not higher than an inch, and, the more, from the anus; and it is not necessary to carry the incision higher into the rectum, even though the sinus may extend for some distance by the side of the bowel above the opening.

The operation is performed in the following simple manner: The patient, who has had the bowels well cleared out previously both by castor-oil and by an enema, and who has been rendered insensible by chloroform, lies upon his side; an assistant separates the buttocks. The operator introduces his left forefinger, well oiled, into the rectum, and then passes a curved, double-bladed bistoury with a blunt point through the external opening, and carries it along the sinus until the point is made to enter the bowel through the internal opening and to come in contact with the forefinger, the bulb of which is turned towards the orifice of the sinus. This being effected, the surgeon, by a sawing motion with the hand holding the knife, and aided by the left forefinger, pushing the instrument downwards, divides the whole of the structures between the sinus and the anus, bringing out both knife and forefinger together. In those cases where the sinus is single, this one incision is sufficient; but where there are one or more tracks branching off, the bistoury must be carried along them, so that the undermined integument may be fully opened and the wound brought to one.

It has been ascertained that an internal opening does not exist, the knife must be forced through the mucous membrane into the cavity of the bowel at that spot where the tissue is found to be thinnest, and the operation should be then completed. It is not necessary to carry the knife to the very extremity of the sinus, if this extends to a considerable length, whether there be an inner aperture of the fistula or not.

After the operation is completed, a small strip of oiled lint should be placed lightly between the edges of the wound, but there should be no plugging or thrusting in of large pieces; a band and bandage may then be lightly yet firmly placed over the wound; the patient should be kept quiet in bed, and should have a mixture of chalk-mixture and laudanum, in order that the bowels may be confined; his diet should be such as not to cause much fecal accumulation. At the end of three days a dose of castor-

case is the confection of black pepper, i night; or it may be needful to send the of air, when the wound, which has hithe will often commence healing again rapid

Before undertaking the operation fo should take especial care to examine locally, but as to the general state of he not unfrequently complicated with other may either render any cutting operatio induce delay in order that some prepar adopted. Thus it is well known that unfrequently associated with pulmonary found to exist, it certainly would not be operation, unless the suffering from the be very great, and the mischief in the If the fistula be cut when the pati phthisis pulmonalis, the wound, in the not heal up, even though life may be spa time.

There are other cases also where : persons who have had their health br residence in hot countries, and have s and disease of the liver; or cases are seen or kidneys have become structurally alte in exultant spirits. An operation in such

It may have been divided. In this case, either this wound will not heal, or its cicatrix will put on a scirrhus character.



I was called to see a middle-aged woman, who had been operated upon on several occasions for fistula, unsuccessfully, by a surgeon of large experience. On examination, I found that the wound had only partly healed, and had taken on a cancerous character. On examining the rectum, I found that there was a firm stricture, about an inch up, and there was much hardness about the opening together. I gave an opinion at the time, that it was a question whether the disease was not of a malignant character, although the features were not decided.

At any rate, the ill success attending the operations for the fistula led to the existence of the disease, which perhaps at the time was over-looked. I ascertained from a medical friend that this woman died six months afterwards, and that the disease had manifested itself as cancer.

If a fistula be complicated by a simple stricture of the rectum, this latter be overlooked, the operation will in all probability be unsuccessful. When the stricture is recognised, preliminary treatment should be employed before the knife is used. The contracted part should be dilated by the bougie, and subsequently the sinus is laid open; during the after treatment, too, this use of the bougie must be strictly adhered to, otherwise contraction will ensue, and the sinus will not close. When the inner opening of the fistula is situated above the stricture, this latter may be divided at the same time that the sinus is laid open; but it is not even in such a case to employ the bougie prior to the operation, if there be not any decided objection to this proceeding.

Very now and then a weakness or entire loss of power of the rectum occurs after its division, more especially in those cases where two or three operations have been rendered necessary. In three instances which have lately presented themselves to my notice the operation had been repeated thrice of the patients, and twice in the other. In two out of three the loss of power was complete; when it is but partial, recovery may take place after the parts are thoroughly soundly cicatrised.

In some rare cases there will be an external opening on the outer side of the rectum, with only one inner aperture. It then becomes a question as to what should be done. The operation performed on that side where the inner opening is, together with slightly enlarging the mouth of the sinus on the other, will sometimes suffice; but if this is found to be insufficient, both the fistulous tracks must be freely incised. The objection to this double operation consists in the circumstance



tended with some striking phenomena an excessive amount of suffering, and judiciously treated are easily remedied of surface which are found involving brane or the skin itself at the extremity differing in regard to situation and size feature, viz. acute sensibility. There are of morbid action so limited, producing and discomfort as painful ulcer of the obvious—it is the periodical action which causes this great sensibility, independent of pain experienced from the presence of the outlets of the cavities of the body and urethra.

The simplest form in which these ulcers occur is that where there is a breach of surface around the anus, not extending within the case the entire extent of the mischief ; if the patient be told to extrude the parafinger into the rectum, the mucous membrane is healthy ; the patient complains of a smarting sensation when the stools are passed, and the uneasy sensation or pain goes off entirely : it is at this point very carefully in the examination by this means shall we be able to deter-

the treatment necessary in such cases is, firstly, strict attention to cleanliness; the patient should be desired to wash with warm water to the parts two or three times a day thoroughly, by means of a sponge. The solid nitrate of silver should be used once or twice to the ulcerated surfaces, and frequently the application of the red precipitate ointment or the lotio nigra will bring about a speedy cure. It will be necessary to attend to the state of the bowels; and the surgeon should not forget to inquire if there be any vaginal discharge; should there be any, to put a stop to it by suitable injections.

The variety of ulcer next to be considered, and that which is frequently seen by the surgeon, in consequence of the great distress which accompanies it, is that where the ulcer is situated partly without the anus and partly within the rectum. The existence of this form of ulcer is not unfrequently overlooked, because, although the symptoms are peculiar and striking, the pathological change causing them is not readily detected, unless by one accustomed to look for it. The patient is suffering from this form of the affection will complain of what he terms the piles; on questioning him closely, it may be ascertained that there is not any protrusion, but that he feels more or less acute pain in passing the contents of the bowels. This pain, instead of diminishing, increases in severity during the act, and lasts for a considerable time, varying from a quarter of an hour to four or five hours; it then ceases, and there is no suffering until the bowels are again moved, when a repetition occurs; and as time goes on the symptoms become more severe. There is also in some cases a discharge of blood, and in all a more or less purulent discharge. On examination there will be nothing visible externally, with the exception, perhaps, of a small excrescence or pile about the size of a pea; and this is found in a large proportion of cases of ulcer of the anus and rectum, and is a sign of great importance. In requesting the patient to protrude the parts as much as possible, and separating the sides of the anus very carefully, the ulcer will be seen situated at the base of the little tumour, which in nine cases out of ten is at the posterior border of the tumour. In one case the ulcer may be only the eighth of an inch in length, and its extent may be defined by the eye alone; in another instance it may be more than half an inch in length, extending beyond the verge of the anus, implicate the

was a peculiarity regarding this: it
the external excrescence posterior
from one another, and an ulcer of
from the base of either of these tun
narrowing as it encroached upon
forming a complete triangle, with it
Instead of being so large as this—v
not be bigger than a large pin's-he
surface is either smooth, the mucou
verge of the anus is the seat, bein
somewhat rough and excavated w
implicated, and the appearance of
red or ashy-grey colour. In some
separated more or less by a raised p
to become nearly distinct.

It is extremely important to empl
investigation of these cases; for a
overlooked, in consequence of th
enveloping the part and hiding
therefore, when this disease is susp
and to take care that the examin
bowels have been well evacuated ar
With regard to the pathology of th
come to any correct conclusion. In
of surface, at first slight, is produce

The writers, as Mr. Copeland and Mr. Syme, to constitute a peculiar affection; but it is doubtful whether the effect in reason is not rather one of the prominent symptoms of an inflammation of the orifice of the anus. There is good reason to believe that in all or most of these cases of spasmodic constriction of the rectum, a diligent search would bring to light some slight lesion or excoriation, on which the symptoms depend. One of the most recent authors of repute, Mr. Quain, holds the view that the sphincter is but secondarily or sympathetically affected, and that the painful spasm is not a disease in itself.

I think, perhaps, that I have more frequently met with this affection in women than in men; and most certainly they are more liable to other morbid conditions which produce much the same kind of symptoms as those afforded by the painful ulcer. Amongst these I may mention the displacement of the womb, the pressing upon the rectum gives rise to a difficulty and pain in defæcation; that painful condition of the coccyx which is the result of an injury, produces also very similar symptoms. I recently a very marked case of this kind, where an eminent physician-accoucheur considered that an ulcer of the rectum existed, whereas, on examination, no such lesion was found, and appropriate remedies applied over the coccyx soon brought about a removal of the symptoms complained of.

The treatment of this affection consists either in the employment of local agents, or in the use of the bistoury. In those cases where the ulceration is only of slight extent, is not situated far within the sphincter ani, may be comprehended by the naked eye, and the symptoms have not long existed, a cure may be brought about by careful attention to the bowels, by the employment of rigid cleanliness, and by an occasional application to the ulcerated spot of the solid nitrate of silver or the phosphate of copper. After the sore has been touched once or twice, the effect will be, in those cases which are benefited by this treatment, a considerable mitigation of the severe pain which has troubled the patient when at the closet and afterwards; the sore, instead of presenting the greyish appearance which is usually observed, puts on healthy granulations, and slowly contracts in size. In order to assist this process, a lotion of the sulphate of copper or sulphate of zinc, one grain to an ounce of water, should be applied two or three times a day; or if these should not succeed, the ordinary black wash may be used with effect. The daily introduction of a full-sized bougie, made of

wax or of yellow soap, will sometimes act beneficially by distending the sphincter, and producing such an irritation as will set up a healing process in the sore.

In the more severe cases, however, this local treatment fails to produce a cure; and there are some persons who have been brought to such a state of suffering, local and general, that it will be unwise to trust to measures which either may not succeed, or which may be slow in giving relief. A mode of the operation which was originated by the French surgeon Boyer, and which consisted in the division of the sphincter and the tissues around, is now generally undertaken by surgeons, and is found to be successful in remedying the disease. The French surgeon and those who had adopted the operation were induced to make a complete division of the sphincter and ani, under the impression that the muscular contraction was the cause of the ulcer, and that the pathological tissue could not be removed unless the action of these fibres was temporarily destroyed. This severe operation doubtless was successful in bringing about a cure; but it was reserved for the English surgeon Copeland to show that it is only necessary to make a small incision through the ulcer or fissure itself without dividing the sphincter. Sir B. Brodie, Mr. Syme, and Mr. Quain have all advocated in their writings this limited operation; and it is almost universally acknowledged, that if the ulcer is divided in its long axis, the same relief is produced as if the sphincter was divided, and the operation itself is much less severe and painful. The relief which is given to the patient is most remarkable. In a few hours the aspect of suffering is so well marked in most of these cases, is almost entirely removed; and even on the first evacuation of the bowels after the operation, the patient feels scarcely any pain being produced by the raw surface of the wound.

The operation itself is simple and easy of execution. The bowels having been well cleared out previously by the use of castor-oil or an injection, the surgeon should then take a narrow-bladed, *probe-pointed* knife, and having passed his forefinger into the rectum beyond the extremity of the ulcer, should introduce the point of the knife very carefully, commencing his incision above the ulcer, carry the cut straight fairly and quickly through the centre of the sore or fissure. It is absolutely necessary to include the whole length of the ulcer, and not to cut to one side; and therefore it is well

stant should open the orifice of the anus as far as possible, under that a good view of the greater part of the ulcer should be obtained by the operator. There is very frequently a small anal pile or thickened fold of integument at the base of the ulcer; this should also be removed, as well as any other pendulous flaps of skin which may encircle the anus. Scarcely any dressing is required after this operation; a small strip of lint dipped in oil may be introduced, and kept in for a day or two, but this is not necessary. It is as well to give an opiate after the operation: in the first place, to lessen pain; and secondly, to prevent the action of the bowels, which should not be allowed to act two days or more; and then this action should be produced artificially by a dose of castor-oil. The wound should be daily dressed with a dossil of lint dipped in a weak solution of sulphate of zinc and water. This dressing is not absolutely necessary, but there are some cases in which the wound becomes gangrenous, and then a slight stimulation is desirable. The use of the confectio piperis nigri will assist the healing of the wound after this operation, as after that for rectal fistula.

Certain precautions and modifications with reference to this operation have been suggested by surgical authors: thus Sir B. Brodie has cautioned surgeons not to make the incision towards the vagina in the case of a female, for fear of the occurrence of loss of power over the sphincter vaginæ; others have suggested that the incision should be made on either side of the anus, in order that the sphincter muscle should be effectually divided: but now it is not found necessary to do more than simply incise the muscle, and in the most severe cases only to divide some of the fibres of the sphincter, the surgeon need not obey either of these directions, but be content with making a fair incision through the diseased part, wherever it be situated. The posterior part of the anus is found to be the seat of the affection in by far the majority of cases; sometimes the ulcer is found on one side, and very rarely it is seen on the anterior aspect; thus, out of twelve cases which I have recently met with, the ulcer existed in front only in one instance, whereas in all the others the disease was situated behind. It is very necessary to make a careful examination of the rectum at the time of operating, for it is very possible that a polypoid growth will be found to exist in conjunction with and above the ulcer, and which may easily escape observation. I have quite accidentally discovered these growths more than once lately after I had completed this operation and

had introduced the finger high up into the bowel for the purpose of passing a suppository. If one or more of these growths were left behind, it is probable that the cure would not be complete, or at all events their presence might produce a return of the ulceration; for I believe that not unfrequently it is the presence of this polypoid growth that originates the fissure or ulcer.

It has lately been recommended in France, that an operation effecting the same purpose as division of the ulcer should be performed. This consists in the forcible tearing open of the sphincter by the introduction of the two fingers into the anus, and using extension. There has been sufficient evidence placed before the profession to prove that this mode of dealing with fissure of the rectum is attended with success; but it is a proceeding so rude and unsurgical, that we trust the surgeons of this country will not forsake the simple and effectual means we have described when any operative measure is required.

There is a form in which *fissure of the anus*, in the true sense of the term, occurs, and which should not be considered as the same affection we have before described, although they are by most writers classed together under the same head. Thus the ulcers of the lower part of the rectum and anus are called by several surgical authorities *fissure of the anus*, whereas the term 'ulcer of the rectum' is much more applicable; for although to a careless observer the disease may have only the appearance of a fissure, on a more careful examination there is found to be a well-defined breach of surface, varying from an eighth to a quarter of an inch in breadth; and the true painful ulcer, independent of syphilitic or gonorrhœal infection, always involves the mucous membrane of the rectum itself. In that form of the disease to which the term 'fissure of the anus' may well be applied, the affection consists in one or more cracks or streaks of excoriation, or superficial ulceration, situated between the folds of the muco-cutaneous surface immediately at the anus itself, involving the rectum but slightly. They produce a considerable amount of uneasiness, smarting, and itching; and, from the periodical action of the anus, the cracks or fissures become deeper and longer, and thus are troublesome to heal. Sometimes excruciating suffering, as is noticed in the true painful ulcer, exists; and on the most scrupulous examination nothing but a long crack or streak of excoriation of the kind mentioned can be seen, and it is reasonable to infer that the

ring is produced by this very slight disorder; it is not difficult to account for it, considering how extremely susceptible persons—especially females, in whom this condition of the rectum is mostly observed—are of painful impressions.

The treatment to be adopted in such cases is, in the first place, to endeavour to procure regular and healthy evacuations of the bowels, and to use scrupulous cleanliness after the motions. A considerable number of persons who suffer from this affection are those who sit a great deal. This should be avoided as far as possible, and plenty of exercise should be taken. The application of the solid nitrate of silver on one or two occasions sometimes cures the disease; or a lotion composed of from ten to ten grains of the same to an ounce of water may be applied to the part carefully by means of a camel-hair brush every other morning. The ointment of oxide of zinc, or one containing chloroform, will sometimes be equally useful in allaying the irritation and healing the part.

It may, however, as will sometimes happen, none of these measures succeed in giving relief, and the patient suffers from continued distress after the evacuation of the bowels, an incision should be made through the fissure; the part will subsequently heal, and the troublesome symptoms disappear.

STRICTURE OF THE RECTUM.

Like the other mucous canals, the rectum is liable to a contraction of its cavity; and although this disease may not be considered as one of very frequent occurrence, it is by no means rarely met with, several cases generally presenting themselves annually in the various public institutions.

The disease consists essentially, in most cases, of an adventitious deposit thrown out upon or around the coats of the rectum. In the most simple form of stricture there is a more or less prominent ring of apparently hypertrophied mucous membrane, either entirely or only partially surrounding the cavity of the rectum. A careful examination on the living body with the finger, or an investigation of the diseased part after death, shows that the thickening is produced in the areolar tissue underneath the mucous membrane; although there are preparations to be met with which lead to the belief that the encroachment upon the cavity of the gut is caused merely by a prominent fold or ring of the mucous membrane itself. In the more marked cases of

the disease, not only is the deposit found in the submucous cellular tissue, but there is a thickening of the muscular coat as well, produced, not by hypertrophy of this investment, but by an infiltration of the fibrous exudation through the meshes of the muscular texture. In the more severe cases, where the disease has lasted for a long period, this fibrous deposit becomes more extensive and dense. In most cases there is, together with a very narrow contraction, a large amount of thickening of the coats of the bowel; but every now and then a considerable amount of contraction is found, without any, or with scarcely any, consolidation of the surrounding tissues. In one very interesting preparation in the Museum of St. Bartholomew's Hospital, there is a tight contraction of the gut about three inches from the anus, the cavity being diminished to about one quarter of an inch in diameter for the extent of an inch; but there is not any thickening of the tissues whatever, and the contraction is supposed to be owing to muscular action. In some rare instances, the stricture is caused not by any deposit in the walls of the gut, but by fibrous bands running across the cavity of the bowel; thus, in King's College Museum there is a curious specimen, which shows a stricture about one inch and a half from the anus, consisting of a cup-shaped septum formed of thick bands running across, in which are three or four distinct openings which would each admit a small quill.

In a few instances the fibrous deposit producing the thickening—which is the essential feature of the disease—involves only a portion of the circumference of the bowel, as was found in a case of stricture of the rectum lately brought to me for consultation. In by far the greater number, however, the whole circumference of the gut is implicated in the condensation which has taken place. It may be much more decided on one side than another.

The extent of bowel affected lengthwise also varies much. The disease may involve only one or two lines of the gut; much more frequently the deposit extends from half an inch to an inch and a half in length; sometimes the consolidation implicates three or four inches of the intestine; and, in rare cases, there is a general thickening of the tunics, involving nearly the whole of the rectum, and producing a considerable diminution in the cavity of the bowel.



The pathological changes which occur as the result of stricture of the rectum affect the bowel in the immediate locality and

And it; important results are also witnessed in the tissues surrounding or in connection with the bowel.

The intestine immediately above the stricture is, in the majority of cases, more or less dilated; in one instance the being widened only to a small extent, whilst in another, the stricture has been very tight and of long duration, the intestine is distended in a great degree; thus in one specimen in the Museum of Guy's Hospital, the rectum is five inches in diameter. There is also a thickening of its tunics above and immediately diseased, in a great measure owing to an increase in the development of the muscular coat; the law being observed here, that if there be any obstruction to the passage of contents of a muscular organ, the increased energy necessary will be marked by an increase in the size and strength of the muscular fibres. The mucous membrane above the stricture is generally in a diseased condition, being either preternaturally enlarged and thickened, or, in long-standing cases, superficially and deeply ulcerated, and sending forth a copious unhealthy discharge. Below the seat of stricture there is not much alteration in the parts; the mucous membrane is, however, sometimes found to be inflamed and thickened, or even ulcerated; and there may be thickening and induration of the coats of the intestine, the result of inflammation and suppuration extending to the immediate locality of the stricture.

In addition to these morbid phenomena, which are usual, very serious changes are occasionally seen in connection with the stricture of the rectum. Thus the irritation of the disease may give rise to inflammation and suppuration in the tissues surrounding the bowel; one or more abscesses may form, and communicate not only with the bowel at separate points, but open upon the surface near the anus, or in the front of the perineum. Sometimes a communication exists between the rectum and the vagina or the urethra, and more rarely an aperture forms above the stricture, between the bowel and the cavity of the peritoneum.

The connection between stricture of the rectum and fistula is one of its results is an important one; the two diseases are so means unfrequently met with together. It is supposed by some authorities that the fistulous opening in the bowel is usually seated beyond the stricture, and that the morbid change is owing to the obstruction and subsequent mischief induced by the stricture; whilst others, among whom is



bowel above the obstruction, and or
ture. In another specimen, a fistula
side of the gut for half an inch :
patient lately seen by me, where ti
ciated, I found, on examination, tha
into the bowel just above the strictu
there can be little doubt that ulce
brane is the first step in the form
others suppuration in the surroundi
sequent formation of the sinuses, is

With regard to the seat of strictu
that one particular locality is more d
another. Thus in by far the major
traction is met with at the lower pa
inch or an inch and a half from t
the living body proves this; and th
of the morbid preparations in the va
that, in very nearly one-half of all t
was found to exist at a point abou
orifice. Next in frequency to this p
at a point varying from two to thre
is occasionally met with at four or f
the contraction involves that part o
junction with the sigmoid flexure of

The symptoms which are produ

necessitating the use of aperient medicine; diarrhœa has probably supervened, the motions being tinged with blood; and this has been followed by constipation, preventing almost entirely the natural action of the bowels.

As the disease advances, the constipation increases, and there is a sense of obstruction about the rectum, which causes the patient to strain violently whilst at the closet; the fæces are passed in small quantities at a time, and are much diminished in size, as the bowel becomes more contracted; the general health begins to suffer from the retention of fæcal matter in the intestines; the abdomen becomes distended; there is marked dyspepsia, with a loaded tongue and general lassitude; and the patient loses his natural appearance of robustness. The local symptoms also become more severe: there is a continual uneasiness about the rectum; considerable pain attending the passage of the fæces, which is effected two or three times in the twenty-four hours, in a liquid form; the bladder becomes irritable; pain is felt in the loins and down the thighs. Occasionally, a large accumulation of fæcal matter having taken place above the stricture, there will occur suddenly from time to time a violent attack of diarrhœa, followed by the most obstinate constipation. In advanced cases, one very troublesome symptom consists in the excoriation and ulceration which is produced around the anus by the acrid discharge which takes place from the seat of the stricture. In some cases this discharge is very abundant at the time the patient is at the closet, and is one of the most important diagnostic symptoms.

Although the general health in many cases of stricture of the rectum does not suffer much at first, the disease will, if unchecked, destroy life gradually by the local irritation and suffering, and by the impairment to nutrition almost surely following from constant and long-continued obstruction to the passage of excrementitious matters. In other cases the patient dies suddenly, from symptoms of acute obstruction and inflammation of the bowels, the strictured canal becoming blocked up either by an accumulation of hardened fæces, or by some foreign body. In one remarkably interesting preparation in the Museum of the College of Surgeons, a stricture of the rectum which had existed for years had suddenly become entirely closed, lymph being produced by the irritation of a fishbone which had been swallowed, and which had become arrested at the contracted part.

dense and hard obstruction, or it can only be ,
through the diseased part. When withdrawn, the
rally covered with a muco-purulent secretion, if t
advanced.

If the symptoms are well marked, and yet th
reach the obstructed point, it will be necessary
the patient strain violently, or to examine him
upright posture ; in this way a stricture wh
beyond the reach of the finger in an ordinary ex
be discovered. Should this fail, a wax or gum
about as large as the adult forefinger, and well
carefully introduced up the rectum. This meas
rarely necessary for the mere purposes of diagn
to several sources of fallacy.

In making an examination of the rectum, th
bear in mind that it may be pressed upon by bo
the bowel. An abscess of the prostate gland, e
chronic nature, will so press upon the rectum as
cavity and prevent the passage of the contents c

Latterly I had a patient at the Westminster Genera
twenty-five, who applied with symptoms of obstruction of t
had passed for a week, and he was in continual suffering, an
much reduced in health. On examining the rectum with
that the cavity of this gut was almost closed by a large elas
in front. The patient had not had any gonorrhœa, but abc
catheter had been introduced, and had caused severe pain an

rectum as nearly to obliterate its cavity; the uterus may be inverted, or a solid tumour growing from that organ may so press on the bowel as to cause the symptoms of stricture. We have seen that the essential pathological feature in stricture of the rectum is a fibrous deposit in or between the tissues of the wall; and there can be no doubt that the chief instrument in the formation of this product is inflammation of the coats of the intestine of a more or less chronic character. The causes of inflammation leading to the formation of stricture are three: spontaneous inflammation of the mucous membrane may arise; or the habitual presence of hard fecal matter, and frequent attempts to evacuate, will produce it; foreign bodies lodging in a portion of the rectum for a continuous time will bring about the same result. Muscular contraction may cause stricture. I have alluded to one specimen where, with very good reason, is supposed to have been the cause. One of the most severe instances of stricture of the lower part of the rectum, or, more properly speaking, of the anus itself, was produced by the cicatrization resulting from wounds made with the scissors in the removal of external piles, especially if no precaution be not taken to excise only the superabundant tissue.

Sometimes, in these operations, large portions of skin, as well as circular pieces of mucous membrane, are cut off; the consequence is, the formation of a contraction at the anus. I met with an instance of this kind in the person of a body who had been operated upon for external piles by one who did not understand his business. Such a tight and unyielding contraction occurred as proved fatal; and the only manner in which life could be rendered at all comfortable was by passing a bougie daily, for the patient would not permit to divide the stricture.

By no means infrequent cause of stricture of the rectum is the contraction of the mucous membrane of the bowel, terminating in cicatrization and subsequent contraction of the cavity of the intestine. This is occasionally witnessed in those persons who have lived a long time in India, and have suffered from dysentery. Mechanical injuries to the bowel will produce stricture, such as the operation for fistula, or the infliction of a wound by means of a probe or an enema-pipe.

There is one cause, however, of stricture of the rectum which is almost entirely overlooked by the majority of writers, in this country at least. I allude to the venereal poison. As a direct consequence of the application of the poison to the part by

syphilitic stricture of the rectum. Mr. Partridge that he had a case in King's College Hospital could be no doubt that the disease was produced intercourse. Bushe, in his valuable work, says ulceration of the rectum may arise from direct the venereal poison; or it may be consecutive the genital organs, and then co-exists with other symptoms.'

The treatment of stricture of the rectum must upon the same principles as obtain in the other remedial measures for stricture of the urethra. In the first place it must be stated that the surgeon cures, in the full sense of the word, an organic stricture of the rectum, but he can remedy it in a great measure, increase, and thus ward off those secondary ills which necessarily ensue if the original disease be neglected.

The great object the surgeon has in view is to prevent contraction, and to restore the calibre of the rectum as far as possible to its natural condition. This can in a great measure be effected by the employment of dilatation in those cases which are situated within the reach of the finger, and where the primary and secondary morbid changes have not proceeded to a serious state. The agent employed is the bougie, or some other mechanical contrivance for dilating the stricture as a dilator. The bougie which is

employ in cases where considerable pain is experienced on the introduction of the finger, or where, from the existence of much muco-purulent or sanguineous discharge, ulceration of the mucous membrane is suspected. It is important to be very particular in cleaning bougies after use. The gum-elastic-bougie is the one perhaps, which is most generally useful when well made. It is sufficiently firm to compress the stricture well, and yet elastic enough to yield to any obstruction through which it is not intended to pass. The metallic bougie, which is a short cylinder attached at pleasure to a handle, is especially useful in narrow strictures attended with a large amount of induration and situated close to the anus.

Before any attempt is made to dilate a stricture, it is necessary to make a very careful examination, so as to ascertain the exact position and the nature of the stricture. In this manner the surgeon is enabled to ascertain pretty accurately the size of the bougie which should be employed—a step of importance, as thereby the patient is saved much unnecessary probing, and the bougie is passed into the stricture with more facility. If the point of the index-finger cannot penetrate the obstruction, a bougie smaller in size than this finger should be well oiled and gently passed along to the face of the obstruction. A very slight amount of pressure will generally cause the bougie to penetrate the contraction. If it is held tightly, the instrument should be left for half an hour; if, however, the stricture readily yields, a size larger should be passed.

The effect of this operation must be watched, and if no undue irritation be produced, a bougie of somewhat larger size may be introduced on the second or third day, and be repeated at the same intervals, until as large an instrument as the anus can admit is passed without difficulty. In those instances of stricture where there is but a slight amount of thickening of the coats, the rectum may be dilated almost to its natural calibre in a short period; but when there is great induration of the tissues around, and when the diseased part is exceedingly sensitive, the progress of the case will be slow, and attended with delay and difficulty. The patient may complain of great pain when the bougie is passed through the stricture; and its
may be followed by local and constitutional suffering. If stricture is very sensitive, the wax bougie should be used, the size increased very slowly. When the instrument is withdrawn, a suppository of opium should be passed into the

gut beyond the stricture if possible ; and the patient should sit in a hip-bath, and remain perfectly quiet for some hours afterwards.

By taking these precautions, a case of bad stricture of the rectum, which at first sight would appear to be very refractory to treatment, will be brought into subjection much more readily than is imagined. This will especially be the case in those instances where the patients can remain perfectly quiet. In some cases where the act of dilatation is attended with and followed by much pain, it will be desirable to confine the patient to bed ; but in those cases where there is not much suffering produced, this step is by no means necessary.

It is a very important matter, during the treatment in question, to attend to the state of the bowels, and prevent accumulation above the stricture. In fact, it is well, before the use of the bougie is commenced, to clear the intestines out by the exhibition of castor-oil internally, and by the use of injections of warm water ; and whilst the dilatation is being carried on the action of the bowels must be brought into as healthy condition as possible by the occasional use of these remedial measures.

When the stricture is seated beyond the reach of the finger the use of the bougie is attended with more difficulty, as there is no certain guide to the exact position of the stricture, or the size of the instrument which it will admit. With very great care, however, the surgeon may dilate a stricture when it is situated several inches beyond the anus with safety ; but it must be borne in mind, that the folds of the bowel or the prominence of the sacrum present themselves as obstacles, which may be mistaken for stricture. And the greatest possible care should be taken not to do mischief ; it is a well-known fact that in several cases the walls of the gut have been penetrated by the bougie even where there has not been any stricture in existence.

It has been stated that, in certain instances of stricture of the rectum, there has been a venereal origin ; and when this can be pretty clearly made out, the local treatment should be assisted by the exhibition of small doses of mercury and iodide of potassium ; the general health will be improved, and we may expect a more speedy absorption of the morbid product constituting the stricture. It will be well also, instead of using oil, to smear the bougie with the strong mercurial ointment.

division of the strictured portion has been practised by some persons; and, as an adjunct to dilatation, it is, in certain cases, a useful means. It is not, however, free from danger. I came under my observation a short time since, in which ensued upon the division of a stricture with the knife, several similar cases have been recorded. This procedure, therefore, should not be adopted unless there is a necessity

division is necessary in those cases of stricture, rarely met with which are of traumatic origin, and situated at the verge of the anus, or within a very short distance; there is generally much cicatrix and much induration around, so that dilatation with the bougie is painful and difficult. It is also advisable, if actually necessary, in old-standing instances of the disease situated in the usual locality and arising from the usual causes, where, from neglect and continual irritation, the induration has become excessive, and the stricture most unyielding to the bougie. In either case the operation is best done by introducing the left forefinger into the rectum, and guiding it so that a straight, blunt-pointed, narrow bistoury, with which the stricture is to be divided. Deep incisions must not be made, but the stricture should be simply notched at several points of its circumference. A bougie should be introduced immediately for a minute or two; a suppository of opium is to be passed into the rectum; the patient must keep quiet two or three days, when the bougie is to be employed, and continued at regular intervals. In this manner a stricture which previously had defied all attempts to dilate it, will be so reduced that the patient may obtain great comfort.

A stricture of the rectum when once fully formed, is seldom ever thoroughly cured, even under the most assiduous care. There is the same tendency to re-contraction here as in instances of stricture of the urethra; therefore, when the disease has been brought under subjection, it will be absolutely necessary to keep up periodical dilatation; this may, however, be easily effected in the majority of cases by the patient himself. In some cases once a week will suffice; where, however, the disease has been traumatic, it will be needful to pass a bougie every day, or every other day, to prevent re-contraction.

If sudden obstruction of the bowels should happen to a patient suffering from stricture of the rectum, it will in probability depend upon some great contraction having

orifice, it must be dislodged ; and this may be effected by the introduction of the finger, or by the use of a probe, which are to be carefully passed through the stricture.

When sudden obstruction of the bowels takes place, as an instance of stricture of the rectum situated high out of reach of the finger, the patient is placed in the most dangerous condition, for it is impossible to ascertain the extent of the obstruction. I have referred to one preparation in the Annals of the College of Surgeons, showing a stricture of the rectum situated near the colon, which had become suddenly obstructed by an effusion of lymph, the result of a fish-bone stricture.

In such cases the long tube must be carefully introduced, and the obstruction may be overcome ; a large quantity of water should then be thrown up. If, however, it cannot be passed, and the use of powerful agents is of no avail, death will be imminent ; and the only method of obviating it will consist in making an opening in the rectum, either by the method adopted by Littre in the operation of Trepan, or by the method suggested by Amussat. The latter operation is usually selected ; there are a few cases on record in which it has been preserved, and there are instances in which it has been undoubtedly justified in resorting to this expedient.

CHAP. IV. OF THE RECTUM.



which render the local disease much more formidable than simple stricture. The features of distinction between these particular kinds of the affection are generally pretty broadly marked, and, indeed, cannot well be confounded, when the one we are considering has become fully developed.

Cancer may attack any part of the rectum, but it is perhaps most generally met with in the lower portion; so that in a large number of cases the disease is within the reach of the surgeon's finger, and its character may be readily appreciated. In the early stage of the disease it appears to commence as a hard deposit between the coats of the intestine; the deposit in some instances consisting of distinct nodules scattered here and there over the bowel; in other cases there will be a complete circle formed by the disease. The actual seat of the deposit is in most cases the areolar tissue, between the mucous and muscular coats. As the disease advances, this is more distinctly ascertained by the circumstance that the mucous membrane, in a more or less healthy condition, is raised up by the deposit underneath it; in course of time, however, this membrane becomes ulcerated, generally in the centre of the deposit.

In more rare instances the malignant deposit is situated between the muscular and serous coat of the bowel, and the morbid growths penetrate into and between the fibres of the muscular tissue. A general thickening and induration of the whole structure of the bowel takes place, so that when a section is made it is impossible to distinguish one texture from another.

This means the cavity of the bowel becomes encroached upon, much in the same way as when the simple fibrous thickening takes place.

As time advances and the cancerous growth increases, the cavity of the intestine becomes encroached upon to a greater extent by the formation of one or more tumours, or nodules, which project into the bowel; by-and-by ulceration of their surface takes place, and in some instances there will be found enclosed together a large hard nodule of cancer, and a distinct fungous growth with a bleeding ulcerated surface. The morbid growths increase upwards and downwards, and involve the rectum to the extent of four or six inches, and spread towards the office of the bowel, and appear there as warty growths or masses of fungoid disease. It not unfrequently happens, however, that there is a small portion of the bowel perfectly healthy between the anus and the deposit.



action, or subsequent disorganisatic
quent result of the mischief in the r
artificial opening in the base of the l
escape into and blocking up of this c
some instances the ulceration penetr
and opens into the cavity of the per
there may be a communication betw
more distant parts, as for instanc
examined a preparation of scirrhu
about two inches from the orifice ; t
ing above the stricture, communic
abscess of the hip-joint. In some c
takes place as the result of the exten
marked ; whilst in other instances the
an enormous amount of thickening
together of the various parts around.
the pelvis is found to be almost en
mass of malignant disease, originatin
cases the peritoneum is thickened wi
the omentum also may be loaded w
have malignant tumours scattered th

All the different varieties of cance
rectum ; of these, however, the true l
frequently met with ; next to this, pe

bloody mucus, and this latter symptom becomes more and more prominent. Thus at first the symptoms are those which attend upon ordinary stricture of the rectum, and the more especially as in some instances there is very little disturbance of general health. As the disease advances, however, there is on a greater difficulty in voiding the fæces, the act is accompanied with considerable pain, and at each time the fæces are moved there is a discharge of blood. As in ordinary stricture, there is great distension felt, which is now and then relieved by a profuse diarrhoea.

The general health at first is but slightly affected; but as the disease advances, its effects on the constitution become marked; the patient complains of indigestion and flatulence, and of an indescribable depression. His face becomes anxious—takes the peculiar dull aspect of one suffering from malignant disease; and emaciation takes place. The patient is deprived of sleep; and indeed there can be few more pitiable objects than those of those who are suffering under cancerous disease of the rectum.

In the course of time, the gut becoming more narrowed, and the texture of the diseased textures getting into an ulcerated state, it ensues of necessity a greater hindrance to the passage of fæces; painful attempts are constantly made to pass them, and these efforts are accompanied with an acrid sanious discharge, mixed with fæculent and muco-purulent fluids. And eventually this action comes on without the patient having the control over himself. As the disease involves the surrounding tissues, abscess forms by the side of the bowel, and degenerates into fistula. And when the deposit encroaches upon the bladder, there is either great irritability of that organ, or daily retention of urine; this latter symptom may be produced by impaction of fæcal matter in the urethra. I have lately seen a preparation where a fatal retention of urine proceeded from this cause.

Death generally follows this disease within three or four years after its symptoms are marked. And it is produced either by a gradual sinking of the powers of life, or by the patient coming on of an attack of obstruction of the bowels. When the disease has existed for some time, and is not situated high up in the bowel, the surgeon can have little difficulty in forming a correct diagnosis; and in many cases there is more or less scirrhus induration at the orifice of the



Cancerous growth, protruding at the anus,

also is of a more offensive and san-
the earlier stages of the simple st-
be difficult notwithstanding to ascer-
malady at its outset; therefore a
given.

I saw, a few months since, a patient, about
upon twice for fistula, neither operation succe-
the last had failed, I made a careful examina-
the lower part of the rectum. I am not aw-

on of life. We have seen that, in instances of simple stricture of the rectum, the affection may be cured, or at all events kept under control, by the careful employment of proper medical means and appliances; because the deposit forming the stricture is merely the result of inflammation, and thus may be absorbed or much diminished by the use of pressure; moreover, the general constitutional health is not affected when the disease is of a cancerous nature. Relief, however, is given to those unfortunate persons who labour under the complaint; and it is important to enquire what are the means best calculated to produce that relief.

There is a difference of opinion amongst surgeons as to whether the employment of dilatation is proper. As a rule it is stated that bougies should not be used in malignant stricture of the rectum; but it is unwise to discard them entirely. When the disease is met with in its early stage, has not ulcerated within reach of the finger, and is producing much constriction of the calibre of the bowel, a wax bougie well oiled may be passed gently through the contracted part; and its use may be repeated once or twice in the week, if pain be not produced. In a case of extensive cancer of the rectum which had ulcerated, I employed the bougie in this manner for a long time with the effect of giving relief. It is not to be expected that any absorption of the diseased tissue should take place, and that which is deposited in connection with cancerous stricture is as the mere result of inflammatory action. When, however, the disease has advanced to the ulcerative stage, it would be highly imprudent to use bougies.

The main object in the treatment is the alleviation of pain; this is best effected by the use of anodynes exhibited in the form of suppositories made of *pil. saponis comp.* or of morphia mixed with extract of hyoscyamus, in the proportion of half a grain of the salt to ten grains of the extract. This should be introduced each night into the rectum, and in the course of the treatment the strength of the anodynes must be increased. I have given very great relief given to the patient for many weeks by the use of this suppository. As the disease extends, and involves other parts, the suffering is much more severe, and it is then found necessary to give opium by the mouth as well as by the rectum, in order to produce a mitigation of pain. The bowels are to be unloaded by the occasional use of cathartics, and, when there is not much irritation caused, by the

exhibition of enemata of warm water. In order to soothe the ulcerated and painful parts, injections of tepid oil or of decoction of poppies must be used from time to time : the surgeon must not forget to take great care in the introduction of the enema-tube, as serious mischief may soon be effected.

During the time the malady is running its course, the strength of the patient, which will gradually decline from day to day, must be supported by those articles of diet which contain the most nourishment in the smallest bulk, and which leave but little fæculent matter in the intestines ; and it will be needful to avoid as much as possible vegetables and bread. The profuse sanious and foul discharge which sometimes accompanies this disease may be checked by an injection composed of sulphate of copper and opium, or a very diluted lotion of solution of chloride of zinc may be employed with benefit.

If sudden and complete obstruction of the bowels should occur during the course of this disease, it will be a question as to how far the surgeon is justified in making an artificial anus. If this accident happens in the last stage of the disease, when the patient is much exhausted by previous suffering, I do not think that the operation should be performed ; but if the obstruction, which must be fatal, should occur during the earlier course of the affection and before the constitution of the patient has been much impaired, the surgeon is only doing his duty if he recommends a proceeding which, although in itself surrounded with difficulty and danger, may yet prolong life for several months at least, and allow the patient to die at last in comparative ease. Some surgeons were, a few years since, in the habit of performing excision of the lower part of the rectum when affected with cancer ; but this proceeding must be looked upon both as barbarous and unscientific, and it is now, happily, exploded from the catalogue of surgical operations.

POLYPUS OF THE RECTUM.

Polypus of the rectum, although a somewhat rare disease, is not unfrequently seen by those who have large opportunities of studying the affections of this portion of the intestinal canal. It is met with in various forms, and it is important to be acquainted with the disease, inasmuch as it often produces most severe symptoms, and at the same time is easily remedied.

The most simple form in which a polypus is seen is where

are one or more short processes, as it were, of the mucous membrane standing out prominent in the cavity of the rectum. Bodies, if they may be so termed, produce little uneasiness, and are generally only accidentally discovered when the surgeon is making an examination of the bowel with the speculum for another disease. They are usually situated above the sphincter, and cause very little inconvenience, and therefore do not demand the attention of the surgeon.

A soft gelatinous polypus, like that growing from the mucous membrane of the nostril, is very rarely indeed met with; it is not an infrequent occurrence to observe a polypus growing from the mucous membrane of the rectum about half an inch from the orifice. It varies in length and thickness, from half an inch to two inches in length, and as thick as a quill or as the little finger. This form of tumour is of a reddish-white colour, firm in texture, and attached to the mucous membrane sometimes by two distinct stalks; or it may have only one attachment and a bifurcated extremity. On section this form of polypus is found to be fibro-cellular in structure, and very slightly vascular. In one specimen of this kind of polypus, which I lately removed from the mucous membrane of the gut just within the verge of the anus, some curious scopical appearances were observed. The structure was partly fibrous, but the mucous covering was arranged so as to form a number of very distinct villous processes. It looked like a piece of small intestine; lacteals could be seen in the villous processes, and their trunks could be followed for some distance into the tissue of which the tumour was composed.

When seated high up, this polypus does not produce much uneasiness; but most generally it is attached to the bowel so close to the anus, that when the patient is at the closet the weight of the tumour protrudes, and perhaps becomes compressed by the sphincter, and in this manner very annoying and painful symptoms are produced. Sometimes even the tumour protrudes when the patient walks about; great irritation exists; there is a discharge of mucus, sometimes of a little blood; and the patient is obliged to wear a support, which gives

He is conscious of some foreign body at the anus, but the symptoms are usually attributed to piles.

The only efficient treatment for this kind of polypus is to remove it; and as there is so little vascularity, it may safely be removed by the scissors, especially when the tumour is very

near to the anus; for if there be any hæmorrhage, it is easy to stop it. When the polypus is situate an inch or more above the anus, and the base is at all broad, the tumour should be well brought down by an injection previously given, and a thread can be tied tightly around the base before the growth is snipped away. In several cases, however, where I have cut these fibrous polypi away without first tying them, there has only been a very slight amount of bleeding subsequently.

FIG. 280.



Warty polypi at the anus.

A form of polypus very rarely met with is where the growth is warty.

I lately saw a gentleman of middle age, who consulted me for prolapsus; on coming to examine him, after making him protrude the parts well, I found attached to the base of the prolapsed portion, about an inch from the verge of the anus, a curious-looking mass, composed of an aggregation of small growths arranged upon a peduncle, just like a bunch of grapes. This polypus, if it be so termed, was an inch in length, of a reddish-white colour, and it protruded at the centre of the anus. I removed this growth; and on microscopic examination its texture was chiefly or entirely epithelial, in fact it might be considered as a warty growth.

A *vascular* polypus is, perhaps, of most frequent occurrence; chiefly met with in young children. The tumour varies in form from a cherry to a pea; is of a bright red colour; its structure is fibro-cellular, and eminently vascular. It is either seated within a short distance of the anus, or it may have its attachment one or two inches up the gut, by means of a long narrow stalk. The symptoms which this form of the disease produces is the prolapse of the tumour on the child evacuating the bowels, and hæmorrhage of a more or less profuse character.

Bleeding not only takes place when the child is at the stool, but it will persist afterwards, when he is running about. The occurrence of this loss of blood in a child should lead to a careful examination of the rectum for polypus; for if the tumour is detached high up, it will recede immediately after the action of the bowels, and escape observation. It is necessary, therefore, to make the inspection immediately after the evacuation of the contents of the rectum either by medicine or an enema.* The treatment which should be adopted is simple, and merely consists in the removal of the polypus, either by twisting it off with a pair of forceps, or placing a ligature around the pedicle, returning the tumour within the bowel and allowing it to fall off. Excision should not be resorted to in these cases, there might be a considerable amount of bleeding, which would be especially prejudicial in a young child.

It is very important to make a thorough examination of the rectum immediately after the bowels have been acted upon, when there is a suspicion of a polypus, otherwise the disease may escape notice. A remarkable illustration of this occurred in a lady to whom I was called by a medical practitioner, who had been in constant attendance upon the patient, whose symptoms were referred to external piles. These were removed, but with scarcely any relief to the pain experienced after the action of the bowels, and to the sense of pressure. I made a careful examination, after the bowels had been well cleared by an enema; but nothing was observable beyond a slight crack at the posterior part of the anus. It was thought right to divide this, in the hope that the severe pain would be removed by the operation, but this failed in its object; they were vainly endeavouring to ascertain the cause of her suffering, until one occasion we found the patient had just got into bed after having had the bowels freely acted upon. I took the opportunity of examining her intimately, and to my great surprise found a large fleshy polypus attached high to the posterior wall of the gut by a narrow peduncle. Without further delay, I, with some difficulty, placed a double ligature around this tumour, and fortunately removed a disease which, from being long undetected, had caused great suffering. It was strange that this polypus had not shown itself before. I examined her on a previous occasion, after the action of the enema.

A rare form of polypus of the rectum is occasional in adults, when the tumour reaches a size as large as it has been not inaptly termed the *villous* tumour, it mainly consists of elongated processes or villi, excruciating. This form of tumour produces most severe effects as a foreign body in the bowel, and is the cause of hæmorrhage to a large extent, by which the health is very much broken down. It is a question with regard to whether this form of polypus is malignant or not.

Mr. Quain has made some investigations into the nature of this tumour, and he has found that it was comp

FIG. 281.



Villous tumour of the rectum. From a preparation in the Museum of St. George's Hospital.

processes slightly held together, each process supporting villi. And as the mass is very vascular, he has found that the disease is not of a malignant nature, according to Rokitansky.

A very interesting specimen of this rare form of tumour of the rectum was preserved in the Museum of St. George's Hospital, and on examination it exactly the features described by Mr. Quain. It was situated about two inches from the verge of the anus, in the anterior wall of the rectum, springing from the submucous cellular tissue. It is round in form and prominent, and about the size of an orange. The tufts or villi are remarkably well developed. The tumour so obstructed the cavity of the rectum that almost complete retention of the feces resulted, and the gut

distended. The bladder was much pressed upon by the diseased mass. It was impossible to say how long the disease had been growing, but it was not until about five years before death, when the occurrence of hæmorrhage and the symptoms before spoken of led to its discovery. Several attempts were made to destroy the mass by ligature and other means, but it was not able to effect more than a partial removal of the more prominent portions,—an operation which was performed no less than thirty-three times before his death, and on each occasion with temporary benefit. The patient was seventy years of age when he died, and no other disease was found on a careful examination of the

case. I have only once met with an instance of a growth of this nature, but it was of a very interesting character, inasmuch as it had resisted treatment and had reduced the patient to a very low state. It was that of a married lady, sent to me by Dr. Rooke of Cheltenham. She had suffered much from profuse hæmorrhage from the bowel without any protrusion whatever, and when I first saw her it was extremely difficult to ascertain where the bleeding came from, but after a very careful examination with the speculum, I found a well-defined vascular growth, about the size of a crown piece, two inches from the anus. From the situation of the patient, and the appearance of the diseased parts, I was afraid it might be of a malignant character, and this idea was somewhat confirmed by the continuance of the bleeding, notwithstanding very careful local and general treatment. However, finding milder measures fail, I determined to treat with vigour strong nitric acid to the part. This I did on several occasions, and was much pleased to find that the bleeding became checked, and after about a dozen applications the hæmorrhage ceased, and the patient's health was fully restored. I saw this lady lately, some three years after the treatment, and she was in remarkably good health.

PRURITUS ANI.

This affection is very common, and very productive of suffering; and although certain pathological changes occur at the margin of the anus in connection with it, itching and irritation of this part must be regarded rather as a symptom than as a special disorder. Thus it is frequently associated with an unhealthy state of the secretions of the intestinal canal, or with simple constipation—a fact which is not uncommonly overlooked in the treatment of those who suffer; it is also attendant upon a congested state of the vessels of the rectum, which can hardly be considered hæmorrhoidal, inasmuch as on examination there will be found neither hæmorrhoidal tumours nor prolapsus, but a considerable amount of congestion of the mucous membrane. The form of the complaint is peculiarly prone to occur in those who sit for a long time together at the desk. Pruritus is familiar to us as a symptom of ascarides in the rectum; it is

the secretions of the part is continued, fissures, and in the course of time the integument gets almost into a condition of distressing itching is produced, and the patient lies down in bed—which is

The treatment to be adopted is upon the ordinary principles of medicine, with a very careful examination of the system, both external and internal; and with a view of ascertaining if there is any existence of the pruritus, such as in the large intestines; in either case, and the use of enemata will afford immediate relief. If the unhealthy secretion, exists, a change of diet, and the occasional use of calomel and three or four leeches frequently remove the itching. The use of any local measure such as soap and water. It is, however, a disease, although perhaps not a disease, neglect and irritation, degenerating into a great difficulty. Here, at the same time the

The application should be repeated so long as the eruptions exist. Not unfrequently the itching and inflammation disappear so soon as these heal and the skin returns to its normal state; but there is a tendency for the excoriations to recur, and to prevent this, the patient should be advised to wash the parts well morning and night with a strong solution of potassium permanganate.

If the solution of nitrate of silver fails either to cure the inflammation, or to give relief to the itching when they are present, a treatment composed of one drachm of glycerine to an ounce of water may be used with good effect; and should there be a hemorrhoidal or condylomatous condition of the integument, nothing more useful than the application of an ointment composed of one drachm of calomel and one ounce of lard. As in the case mentioned before by me—should be applied either within or upon the sphincter, the ordinary operation for the disease must be put in force.

In obstinate cases, it will be found that the use of all measures, general and local, will be unattended with success. If this has not already been tried, I am in the habit of recommending the daily introduction of a well-oiled bougie made of india-rubber.

NEURALGIA OF THE RECTUM.

It is impossible to affirm with truth that neuralgia, as in other parts of the body, is ever situated in the rectum; however, now and then cases are met with where extreme pain of a distinctly neuralgic character is experienced for a long time, and where, on the most careful examination of the rectum, external and internal, no appreciable lesion can be detected. We therefore do not hesitate to apply the term neuralgia to this condition of things. The cases are more numerous in the persons of females who have been subjected to some depressing causes; the pain is described as being severe, and continuous, and not particularly aggravated by the motion of the bowels, as is the case when the suffering is due to a painful ulcer of the rectum. It may be of a local character, and be the result of direct exposure

of this affection is extremely difficult, and indeed only a temporary relief can be given. Of course, the treatment applies with the symptoms of neuralgia, the

in our enquiries—colchicum should
remedies best suited for this complain
the cold douche to the anus, and the u
in the proportions of one drachm of tl
lard, either smeared around the anu
rectum by means of a large wax bc
measure be adopted, care should be t
of the belladonna on the system.

DISEASES OF THE URINARY ORGANS.

UNDER the term *urinary organs* will be here comprised the kidneys, ureter, bladder, prostate, and urethra. It is extremely desirable, when dealing with any disease the symptoms of which are referred chiefly or entirely to some portion of the important system defined above as urinary organs, to possess an intelligent acquaintance, not merely with that part which may be especially liable to suffer morbid changes, but with the system itself as a whole. It is absolutely necessary for the formation of a correct diagnosis, to be familiar with the intimate relations which exist between each portion of that system, as well as with the various peculiarities which each is liable to exhibit when influenced by disease. Few things conduce more to error in estimating symptoms than a too exclusive specialism in the observation of disease. The study of stricture and of other forms of urethral obstruction, of vesical diseases, of calculous formations, and of those chronic changes in the reticulating structure of the kidney which are conventionally assigned to the province of the physician, must be pursued whether if the surgeon is properly to appreciate the import of any one of these affections. For example, suppose a surgeon to perform an important operation upon the bladder or urethra (and what operation there is not important?), who is unacquainted with, or is at least indifferent to, the chemical and microscopical characters of his patient's urine; will anyone, who knows what those indications reveal, deny that elements so important to the formation of a correct diagnosis, and, therefore, to an appropriate therapeutical scheme, must be absent from the mind of the surgeon in relation to that case? It may be replied, that the necessary information can be obtained from the physician. Ought the surgeon to be dependent upon an acquaintance with the facts required upon any other server in such a case? Assuredly not; not only is the

to affirm, that no man can deal adeq
of impaired urinary function, whose
the use of the sound or catheter, wh
urinary deposits in all their varieties
renal casts, and who is not acquainte
far as these are known, as well as wi
subjective phenomena which are f
The exploring sound is quite as ess
urinary disease as the stethoscope is
He who is, conventionally speaking,
does) constantly overlook calculus a
detriment of the patient; while the
man will (and does) treat mechanica
only be injured by his manipulation
careful study of all acute and chroni
mended to the student who desires t
for the practice of surgery.

The following is a table of those re
sarily come under the observation of
with the diseases of the bladder, pros
frequently demand his services. T
department of urinary disease in thi
the slightest glance at them; it ap
necessary to make these introductory

Inflammation, (nephritis) { calculous.
 Chronic { from obstruction.
 { diathetic.

Inflammation of pelvis of kidney { from obstruction.
 Pyelitis) acute and chronic { calculous.
 { scrofulous.

Calculous nephralgia.

Abscess of kidney and fistula.

Perinephritic abscess.

Tumours of kidney.

Suppression of urine.

Hæmaturia.

Formations.—There are few deviations from the normal forms which can be regarded as malformations, and these are of little importance to the surgeon. It is only necessary therefore to enumerate the ordinary varieties. (1) There may be only one kidney, which is then usually larger than natural; it may occupy its usual place, or may in reality consist of the two organs united in the median line, and occupy a position in front of the vertebral column. In the last-named condition the form is sometimes that of a horse-shoe with its convexity downwards. The kidney may occupy its ordinary situation, and a tumour may be moveable, forming a tumour recognisable during palpation, or it may be fixed in an unusual situation, generally above or below the natural position, or within the pelvis. (3) The kidney may present an outline more or less lobulated, showing the persistence of a form existing in foetal life, and natural in some of the lower animals.



Nephritis.—Acute and chronic.

Acute nephritis, of a severe form, is common in surgical practice.

It exists also as a sequel to preëxisting chronic inflammation of the kidney, and in Bright's disease.

The causes of the first-named variety, with which almost exclusively we have to deal here, are as follows:

Contusions, bruises, and strains; by extension from the bladder in cases of calculus, stricture, and cystitis, following operations or trauma; retention of urine, causing distension of the ureter and increased fluid pressure on the kidney; renal calculus; entozoa; extension of inflammation from neighbouring parts, as diseased



ating, increased by the uprigate vomiting, even by respiration, and ureter. Occasionally there are a thigh of the side affected, and tender region, sometimes considerable. M urine is mostly high coloured and s often mingled with blood, and ther otherwise; later in the disease it m microscope renal casts may be found corpuscles. Sometimes total sup signs of fever are present, as heat headache, thirst, quick and hard puls also, in cases suddenly occurring fr commonly in cases of extension from

In the latter instances, suppurati kidney is to be dreaded; the sympt there is great depression of the vit often occurs by uræmic poisoning. supervenes on nephritis, there is als tion of the testicle on the affected si

As it is impossible to include in tl renal and allied affections with wh familiar, a few remarks will follow diagnosis of nephritis, so as to illust of those other diseases which might

of primary acute nephritis may sometimes be confounded with the following affections: calculous nephralgia; pyelitis, and calculous; perinephritis; perinephritic abscess; disease, or abscess of the psoas or adjacent parts; lumbar renal neuralgia and hysteria; inflammation of the ear; vesical calculus.

Calculous nephralgia; by which is intended the pain and morbid disturbance occasioned by the descent of a calculus from the kidney to the bladder, and which sometimes resembles closely acute nephritis. Indeed, excepting only the severe symptoms, all those of nephritis are present. There is no pain, generally limited to one side, and shooting along the ureter to the sacrum, with retracted testicle; but the invasion is sudden, and the pain more acute, than in nephritis. There may be vomiting, frequent micturition, and bloody urine; but the constitutional disturbance, although it may sometimes be present in moderate degree, certainly falls short of the fever which always accompanies the inflammatory affection. If the calculus obstructs the pelvis or ureter for any length of time, symptoms of calculous pyelitis may arise. And lastly, all the phenomena may suddenly disappear, which confirms the diagnosis of calculus.

Acute calculous pyelitis may be distinguished by the following characters: the pain in the back is excessively severe and constant, much more so than in nephritis; shooting pains are felt in the groin and inner part of the thigh, radiating to the testicle, which is generally strongly retracted. When the pyelitis is unaccompanied by renal affection, the urine is often scanty, and although it may be scanty at first, is subsequently increased in natural quantity. It may have much pus and albumen mixed with it, and sometimes blood; if so, these elements are found in intimate admixture with the urine. The course of the complaint is less rapid than that of nephritis, and there are frequently considerable intervals of freedom from

Perinephritis; caries of spine.—Inflammation of the tissue surrounding the kidney does not generally make its appearance as an acute attack. The subsequent formation of matter, if it takes place, is accompanied by febrile symptoms; but the gradually increasing swelling, in its later stages presenting fluctuation, while at the same time the urine is healthy, is significant of simple perinephritic abscess, provided the special signs of

paired by the inability to act, from the psoas muscle. Pain on pressure at the front rather than behind, the reverse occurring sometimes in nephritis.

4. *Rheumatic affection* of the do- usually productive of great pain, which necessitates the slightest rest, much more so than in nephritis ; vomiting, where, but vomiting and other constitutional symptoms. Again, the urine is neither suppressed nor increased in character ; it may be more than alkaline.

5. *Renal neuralgia* is met with, but severe pains are complained of by hysterical temperaments, which it is difficult by observation to separate from nephritis. The urine, mostly pale and abundant, the absence of constitutional symptoms, and the existence of renal pain are sufficient to mark these cases. .

6. *Acute cystitis* is sometimes met with, which seem to indicate that the kidneys are not the seat of the disease. There may be present great pain, fever, febrile symptoms of severe character, tenderness of the perineal regions ; and there is often vomiting and suffering in passing water, while

ritis. All the local and much of the constitutional disturbance of the latter complaint may be produced, as their subsidence seems to indicate, from mere temporary irritation of the kidney, occasioned by the presence of calculus in the der. An examination of the previous history and symptoms, the urine itself, and the speedy relief which follows simple medical means, will indicate the real cause, respecting which sound subsequently employed at the proper time will enable to clear up any doubts which may still exist.

Treatment of acute nephritis.—It is of importance at the onset to insure free action of the liver and bowels, which are very obstinate. In addition to active purgatives, enemata are recommended; but distension of the colon with irritating fluids is undesirable, and emollient injections alone are admissible; thus, turpentine should be avoided, lest absorption should irritate the kidney. Perfect rest in bed, and low diet, are to be enjoined. Local bleeding by leeches or by cupping is very serviceable: by the latter mode ten or twelve cups may be taken, and the bleeding may be repeated with advantage if the patient is hale and the attack severe; otherwise, it should be followed by dry cupping, daily or on alternate days. The utility of antimony and salines (generally more or less diuretic) is questionable, and the repeated administration of diuretics is contraindicated; the moderate use of simple diluents, water, linseed-tea, barley-water, or alk-and-water, is preferable. Free action of the skin should be promoted by hot baths or large hot fomentations. For vomiting, when obstinate and distressing, hydrocyanic acid and kreosote fail, though sometimes successful in mild cases; but counter-irritation to the loins and pit of the stomach affords better results; a constant supply of ice in the mouth helps to allay it. When head-symptoms appear, the most efficient remedies are purgation and renewed counter-irritation to the renal region; similar applications should be made to the temples and to the nape of the neck, with hot bottles and leeches to the feet and legs. In applying counter-irritation, tharides should not be employed, on account of its known violent action on the kidney, in some cases even by endermic application; the desired result may be attained equally well by croform, ammonia, mustard, or more slowly by nitrate of silver. If either of the two former are selected, dilute with equal parts of spirit; soak a piece of lint the size desired and lay

on the surface, covering instantly with oiled silk, and making gentle pressure; in a few minutes the full effect will be seen. In later stages of the complaint, internal stimuli and support may be necessary.

Where acute nephritis first shows itself, as is often the case in surgical practice, by extension from the urethra or bladder in a debilitated subject, the general principles enumerated above must guide the treatment, but actual depletion is rarely permissible. All unnecessary irritation of the urinary apparatus must be avoided, and no catheters be passed, if they have been necessary before, unless absolutely demanded by the exigencies of the case, and then with the utmost care and gentleness. If following lithotripsy, all mechanical interference must be suspended, and catheterism used only to relieve an obvious retention. If suppuration take place in the kidney, or the urine be suppressed, little can be hoped for from treatment; active counter-irritation of the renal region and internal stimuli must be perseveringly employed while any hope exists.

Treatment of the attack of calculous nephralgia.—The hot bath; antispasmodics and opium liberally administered; cupping to the renal region, and hot cataplasms afterwards; mild diuretics, particularly the decoction of the couch grass or triticum repens; abundance of diluent drinks. Sometimes the occasional inhalation of chloroform, to relax the muscles for a period of ten to twenty minutes, has been of service.

Treatment of chronic pyelitis.—If occurring from obstruction in the urethra, the most efficient treatment consists in overcoming it, and enabling the urine to pass freely, so as to relieve the dilated and diseased pelvis of the kidney from distension and pressure. At the same time, the purulent discharge is sometimes diminished by full doses of buchu, and small doses of cubebs and the balsams; sometimes by mineral acids. Abundant nutriment is generally necessary. If the complaint depends on calculus in the kidney, and an examination of the urine or the history has indicated pretty clearly the nature of the formation, appropriate remedies can be taken with the view of dissolving it or at least of hindering its increase. Such are alkaline salts, as the citrate of potash in uric acid calculi, and acids, mineral or benzoic, for those which are phosphatic. All sources of local irritation must as much as possible be avoided, as sudden or violent movements of the body, and irregular habits of all kinds. Wherever the signs of general malassimi-

On are present, especially those of the tuberculous diathesis, appropriate constitutional treatment must be enjoined. Profuse pyelitis is, however, rare.

Abscesses may be formed under these circumstances; also as a sequel of nephritis; they should not be opened until the four points and the diagnosis is perfectly clear. Often they are perinephritic, although originated by disease within the kidney itself; occasionally a calculus may be removed through an opening made. If not opened sufficiently early, these abscesses may make their way internally, or along the sheaths of muscles, and open in the groin or elsewhere. They may heal externally, or give rise to renal fistula.

It is necessary to bear in mind that other tumours in the renal region may be confounded with abscess, such as cystic disease of the kidney; simple fluid distension of the pelvis; accumulation of blood in the same situation; encephaloid and circular deposits, and hydatid cysts adjacent to the kidney; and abscesses, which are the effects of spinal caries. Among these, ordinary cystic disease affords no very obvious signs:

The character of the urine may be unaffected; if the cyst is large, dulness on percussion is present. Distension of the pelvis with fluid may be due to retained urine, or merely serous fluid, and may in very rare cases afford the sense of fluctuation externally. If there is a partial communication with the bladder, the amount of water passed will differ greatly on different days. Cancer of the kidney is encephaloid in six cases out of seven; the pain is severe, the growth rapid, the tumour extremely large and abdominal; hæmaturia is present, and the diathesis is generally marked. Tubercle in the kidney rarely produces a tumour appreciable during life, and its symptoms are not much more to the bladder than to the organ affected: Tubercle in the lungs, and elsewhere, usually coexist. Tumours containing blood are almost invariably recognised as originating from direct violence, or from a strain in violent exercise. Tumours containing acephalocysts are very rare; they are usually in connection with the pelvis of the kidney, and may persist for a very long period without rupture, subsequent to which hooklets, &c. are found in the urine. In rare instances, both these and the tumour produced by great distension of the pelvis of the kidney have been successfully punctured.

Chronic nephritis is not always a well-marked disease, except as a sequel of the acute attack, not including under the term

quired by the other affection; never character as a blood disease must here.

The remedial means for simple cl local stimulating applications to t specific renal irritants, before specific stitutional treatment of chronic infla

Renal calculi.—(See the essay on (

Hæmaturia signifies simply the blood; which may exist in any c discoverable only by the microscope constitutes the major proportion o urethra. It is a symptom, merely, c the urinary organs. When blood is p and has been in contact with the uri the latter presents a peculiar brow to the practised eye, and commonly l If in large quantity, under similar c has the appearance of thick or mudd blood in hæmaturia may be any one parts: the kidney, the pelvis of the l prostate, or urethra.



Hæmaturia occurs in acute and

It is always important to determine its source; the following signs will assist in forming a correct diagnosis:

Renal hæmorrhage, not being traumatic, is usually preceded and accompanied by some signs of kidney-disease; which is most frequently acute desquamative nephritis, and renal casts are usually found in the urinary deposit; the blood is always intimately mixed with the urine, which shows the smoky tint. Further, it may be said, as a general rule, that such urine is voided without pain, or any other local symptom whatever, is more likely to derive its blood from the kidney than elsewhere. Hæmaturia from a calculus impacted in the pelvis or ureter is associated with the history and signs of the occurrence (see Calculous Nephralgia). Hæmaturia caused by a blow on the back is mostly from the pelvis or ureter. Malignant disease occasioning hæmaturia presents sooner or later some physical characters associated with the organ which is affected.

Blood derived from the neck of the bladder or prostate generally remains within the cavity, and if in large quantity forms a clot there, perhaps distending the viscus; or the urine has the colour of porter, and a darker sanies comes away at the time of micturition. When altogether from the prostate, a portion usually issues unmixed with urine by the urethra. In malignant disease of the bladder or prostate, disintegrated shreds of tissue come away with the urine, and occasionally cancer-cells (so called), and the blood is apt to appear suddenly in large quantity. In villous tumour, characteristic shreds of soft growth are sometimes detached; in both this and the preceding case the bleeding is large, and its occurrence irregular, but it is much less so in the latter case than in the former. With villous growth the blood is more intimately mixed with the urine, which is constantly more or less stained. The appearance of a drop or two of almost unmixed blood at the end of passing water may be caused by either calculus or chronic prostatitis; the history, the distinctive symptoms, and the examination will determine which.

When the hæmaturia is due to urethral sources, and these are numerous (such as injury to the penis from blows or lacerations, and in falls, with or without fracture of the pelvic bones; chordee; occasionally from rupture in sexual intercourse; from surgical instruments; in some forms of stricture, from severe efforts to pass water; impacted calculus; sometimes in severe gonorrhœa; phagædenic ulceration; and malignant



heat, and the precipitate or albume microscope determines it without fail, often a matter of extreme importance with calculus in its early stage and in the detection of the blood-corpuscle by it is the only certain method.

Treatment.—Renal hæmorrhage in calculus, malignant disease, and vice the recumbent position, and by internal and sulphuric acid, matrico, and acetate of opium; when produced by purpura, iron and other appropriate remedies. The bladder may demand similar general treatment; but local means are here more to be applied to the perineum. The patient maintaining perfect quiet in bed, or iced infusion of matrico, may also be used. Small pieces of ice may be placed on the bladder. It is commended that a catheter should be used to break up the clot, and that efforts be made by applying an exhausted gum-bottle as a stricture; but such interference is unnecessary from retention of urine. The disturbing a clot is liable to open or to recently deposited fibrin, and to pr

in almost all cases it is wiser to allow nature to do the work, than to be officious in rendering assistance. The distress occasioned to the patient by the urgent wants to micturate is instantly relieved by the use of opium either by mouth or rectum. When it is necessary, however, to use an instrument, as when the patient is incapable of passing any urine by his own efforts, a small-sized catheter should be passed, and a syringe, or stomach-pump, or, better still, Clover's exhausting apparatus for lithotritry, adapted to it, by means of which a considerable portion of clot may be gently withdrawn. It should not be forgotten that B. Brodie records a case in which, all other remedies having failed, a dose of Ruspini's styptic, administered internally two or three times in the course of twelve hours, was followed by complete cessation of the bleeding.

Suppression of urine.—This term applies only to a condition in which the kidney fails to perform its function of secreting urine. The bladder is empty, or having been emptied, continues so since it receives no more by the ureters. Meantime the materials of the excretion accumulate in the blood, and the vital powers become oppressed; a comatose state ensues, and finally death, unless the eliminating function of the kidney is resumed, and the urine again excreted. This function is sometimes totally, but only temporarily, suspended; more commonly it is after long impairment by chronic disease gradually extinguished. Advanced chronic disease existing, any sudden shock or operation, however slight, from acute attack, &c., may suddenly induce fatal suppression. The two conditions of suppression and retention have been much confounded. None, however, can be more distinct: in the former the bladder is empty, in the latter it is unnaturally full; in the former no urine is secreted, in the latter the secretion goes on as usual, but some obstacle opposes its removal from the bladder. The symptoms are these: the patient usually complains of languor; fever, vomiting, more or less pain in the back and loins; he is unable to pass urine, or only a very small quantity: he soon becomes drowsy, perhaps delirious, then comatose, and dies in from two to four days if suppression remains complete. There may be anasarca, but often there is none. Catheterism shows that the bladder is empty.

Treatment.—Local cupping; but if the patient is in low condition, without abstraction of blood; hot fomentations and counter-irritation to the renal region. The hot-air bath and

free purgation, to aid in eliminating excreta from the system. I have seen, in one case only, recovery after forty-eight hours of complete suppression: I have seen it not unfrequently after twenty hours. In the later stage of chronic renal disease, little is to be hoped for from treatment.

The Ureters present very little occasion for notice.

Malformations.—The most common deviation from the natural condition is that of increase in number; two or three ureters may issue from one kidney, and may be continued down to the bladder, but much more frequently unite to form one channel. Sometimes one ureter only exists; sometimes the ureter is incomplete, and forms a canal with a closed extremity. None of these conditions possess interest from a surgical point of view. The ureter is liable to inflammation, which extends by it from the bladder to the kidney. It may be the subject either of stricture or of obliteration, which conditions are generally produced by injury caused by the passage or impaction of calculi; occasionally from pressure by a tumour; rarely from external injury. It is often greatly dilated, when obstruction exists at the neck of the bladder, or in the urethra, from hydraulic pressure slowly acting upon it; and sometimes to such an extent as to become a supplementary reservoir to the bladder. It has even been seen prolapsed into the bladder, forming a tumour there. It may be the subject of malignant and tubercular disease by extension downwards from the kidney, or upwards from the bladder. A calculus is sometimes arrested in its passage through the ureter, occasioning great agony (see Calculous Nephralgia).

THE BLADDER.

Malformations.—These are happily not very common, since, of whatever kind, they are the source of great discomfort, and even of misery, to the subjects of them.

1. Absence of the bladder. The organ is sometimes, although very rarely, absent. (a) The ureters have been seen to communicate directly with the urethra; in one case these ducts were somewhat dilated before entering it, in another this condition was not noticed. This abnormality has been only observed in the male sex. (b) Less rarely the ureters have been found entering the rectum, and discharging the renal excretion

ly by that channel. These instances have chiefly, although entirely, been discovered in the male. (c) A few instances on record in which the ureters opened directly into the a.

Two or more bladders, it is said, may co-exist in the same act as a congenital malformation. Two cases only are recorded which have any title to be so regarded: one of these recorded in a man in whom the ureter of each side appeared to its own distinct bladder, the two reservoirs being closely ad to each other at their adjacent surfaces (Blasius); the in a woman who is reported to have had five bladders, five uys, and six ureters (Molinetti). I am satisfied that other reported as belonging to this category, including one tly described with great detail, were evidently examples of lation resulting from long-standing disease. Sometimes rgan appears to be partially divided by a septum, or to be ad of two more or less symmetrical lobes; but although cases have been often regarded as instances of malforma- they are in reality also the effects of disease.

Extroversion of the bladder. This is the least rare of all eces of conformation to which the organ is subject. In these the anterior wall of the bladder is wanting, while frequently, ot invariably, the pubic symphysis is absent also. The rior wall and base of the organ are then pushed forward e abdominal viscera, and form a red and vascular flattened inence, covered by mucous membrane, in the situation of pubes. The extroverted portion is in the adult usually about size of the palm of the hand; the orifices of the ureters may seen just below the centre, and from beneath the lower gin a short imperfect penis projects, which is flattened, as if t in the median line, into the urethra. At the root of the is a rudimentary prostatic structure exists, the ducts of ch can be seen opening into the angle between the bladder pubes. There is usually more or less hernia of the bowels ither side and beneath the extroverted bladder, into a pouch- fold of integument, which is covered with hair, represents scrotum, and contains the testes. The mucous membrane ery sensitive, readily bleeds when touched, and from the ecting orifices of the ureters urine is almost continually dis- ing over its surface. This condition produces much excoria- of the skin on the parts beneath, besides infecting the ent with a continual urinous odour. The malformation

thus described affects in the male. Of eight two only were in the there are some slight but most conform a defect is of a less sex female, as the general in abeyance; the male penis, wholly incapable

Treatment of malformation

be palliated by well reservoir may be employed for adjustment in various

Extroversion of the bladder, the mucous membrane of the ureters are hidden wide penis; the exposed aspect. The large pendulous testis.

by the mechanician. results of past experimental measures to Langenbeck in Germany made attempts to remove the latter being however than extroversion of the 1858, was also partial named condition. In the Hospital, has attempted

ureters into the rectum ; but with results which could be considered successful, although approximating. Other cases of operation have been fatal. In the nature of the operation required is less formidable, such as the indication is rather simply to cover the bladder and alter the course of the ureters. Dr. Daniel Ayres, of New York, U.S., appears to have succeeded in covering the exposed bladder of a young woman with integuments borrowed from the neighbouring parts of the abdomen.* The operation consisted in bringing down a flap of skin from the part above, turning it down over the bladder below, so that the cutaneous surface was in contact with the mucous membrane, and in pulling up the integuments, previously raised by dissection, on each side of the vulva towards the middle line to meet the flaps described. The result obtained was a covering to the bladder, and a small opening for the exit of the urine, which facilitated with ease the application of an efficient apparatus to it. But Mr. Holmes has recently succeeded in covering the bladder by lateral flaps, in a young male child. Incontinence of course as before ; but an instrument may now be easily introduced to catch the urine, and the patient no longer suffers from abrasion by friction. I have examined this patient, and consider the improvement beyond question. (See *Lancet*, 1863, p. 714.)

During this period, Mr. Holmes has developed and improved the method, having several times repeated the operation. By turning in the exposed mucous surface and contriving to leave a small opening only at the lower part for the escape of urine, he has saved the pain of contact between the exposed surface and the patient's clothes, and ensured the free exit of urine without excoriating the skin : very important results and well repaying, when attained, the somewhat tedious and repeated steps of a plastic operation.

Mr. Holmes has also designed a procedure for forming artificially a new artificial cloaca, common to the ends of the ureters and the rectum, by inducing sloughing of the tissues composing the wall of the bladder and the anterior wall of the bowel. For

Congenital Extrophy of the Urinary Bladder. By Daniel Ayres, M.D. New York, 1869. Another successful case is reported by Mr. E. Barker of New York in the *Medico-Chirurgical Transactions*, vol. liii. p. 187.

details in relation to both these methods, it will consult the author's work.*

Mr. John Wood, of King's College Hospital, has carefully studied the subject of plastic operations for stricture, and with considerable success. He has operated (in the summer of 1868) on eight cases, and an interesting account of his modes of proceeding, and of the results obtained, is given in the *Medical and Chirurgical Transactions*.

Injuries of the bladder.—(See vol. ii. pp. 715 et seq.)

Diseases of the bladder.—In entering upon a study of the numerous affections which range themselves under this heading, I shall adopt the following classification, in the hope of perspicuity. We shall study first those diseases which present local anatomical characters, establishing the extent of the lesion which has given rise to the symptoms observed; and, secondly, those more obscure diseases in which similar symptoms occur, but which do not necessarily involve any organic structural changes.

I. *Diseases which involve organic structural changes, the existence of which is generally recognised during life, and is appreciable after death.*

These may be arranged as follows:

Inflammation (cystitis):	Acute . . .	{ due to injuries and
		„ presence of
		(calculi, &c)
		„ prolonged retention
		„ extension from the
		kidney, or from
		chronic cystitis
		„ external cold.
		„ chemical irritation
		of the mouth.
		„ chemical irritation
		of the bladder.

* *The Surgical Treatment of Children's Diseases.* London: 1868. p. 149.

† Vol. lii. pp. 85-131.

ation in) :	Chronic (a with, and b without, catarrh) :	as a sequel of acute.	stricture of urethra. tumours in urethra. bar at the neck of of the bladder. enlarged prostate. calculi or other foreign bodies in the urethra. tumours of the penis itself.
		as the result of obstruc- tion to the outflow of urine from	
		from foreign bodies in the bladder.	
		„ growths in the bladder.	
		„ altered urine.	
		„ paralysis.	
		„ over-distension or atony.	
		„ displacement.	
		„ disease of neighbouring or- gans (uterus, rectum, &c.).	

inflammation of the bladder.

abscess of the bladder.

inflammation, simple and non-malignant, and gangrene.

perforation of the bladder.

atrophy and sacculation of the bladder : dilatation.

hæmaturia. (See Hæmaturia.)

tumours and growths	{	1. Fibrous	{	warty.
		2. Villous or vascular.		polypoid.
		3. Epithelioma.		
		4. Malignant.		

calculus of the bladder.

stricture of the neck of the bladder.

abscess of the bladder.

foreign bodies in the bladder.

hæmaturia. (See the essay on that subject.)

inflammation of the bladder.

stricture of the bladder. (See the section on Stricture.)

conditions which do not necessarily involve organic struc-
tural changes appreciable by known methods of observation, and
are associated with some injury or derangement of a
centre.

Paralysis.

Atony from over-distension
caused by

'Irritable bladder'
quent).

'Spasm of the bladder'
involuntary, and excessive
Perverted sensibility

I shall then consider
tion of micturition produced

Incontinence of urine

Habitual engorgement of the
dium urinæ).

Retention of urine.

Inflammation of the bladder
seat of inflammation
mucous membrane; on the
of the process, the cell
may be involved, and
there is no evidence of
and independent inflammation
the middle and outer coats

The mucous membrane
of the urethra or neck
often most severely, affords
common form of cystitis
urethra, as in gonorrhoea
of the lining membrane
death simply injected :
may have become more
slaty in colour, as happens
siderable period of time
and membrane-like shrivelled
extended layer of this
adhering to the whole
a false membrane has
and necessitated puncture
female it is sometimes t

causes.—The traumatic are numerous. Injuries such as blows, pelvic fractures; operations of lithotomy, lithotrity, and very catheterism; injections; in the female, prolonged and mental labours; mechanical irritation of calculus; the local action of cantharides and of some mineral poisons, and of diuretics; action of the urine itself, retained and decomposed as in stricture and prostatic enlargement; inflammation of adjacent parts, as kidneys, prostate, rectum (?), urethra, finally the last, giving rise to a common but mild variety; chronic pre-existing inflammation; rarely from external and occasionally from gout.

Symptoms; of the graver form, occurring after operations and existing chronic disease. The invasion is ushered in by, the signs of general fever, followed by perspiring skin. Desire to pass water is very frequent, sometimes quite unobtainable; the act is excessively painful, often accompanied by rectal pain and tenesmus. Pain is constant behind the pubic symphysis and in the perineum and in the penis; also in the loins and loins; sometimes in the groin and upper part of the thigh. There is tenderness to supra-pubic pressure, often on the lower half of the abdomen, the patient's legs being raised and bent against its parietes; much pain is produced by rectal examination, still more by catheterism. The urine is at first high-colored, scanty, and clouded; then obviously reddish, like the string of raw meat; soon purulent and loaded with shreds; the blood-tint continues, perhaps deepens. Meantime the patient's condition becomes low; he is often delirious; and if the disease is not checked, he dies in from seven to fourteen days. The attack is frequently fatal.



The symptoms of the milder form, which occurs by extension from gonorrhoeal inflammation of the urethra, are chiefly local. They are, painful irritability of the bladder, sometimes very distressing; supra-pubic pain; pains in the sacrum, perineum, and thighs; loss of appetite, but not always marked fever. The desire to pass water must be complied with instantly, and is extremely painful. The urine is cloudy, and deposits more or less of mucus and pus. It may be fairly supposed that in this milder form of symptoms the mucous membrane of the neck of the bladder is chiefly affected, and not that of the entire viscus; the most common termination is resolution; while in the more serious form described above, the whole of the mucous membrane is affected, and perhaps some of the tissues sub-

be severe. Hot muscad-poultices to the nypog is much pain there. If the irritability of the h half-drachm to one-drachm doses of the tincture every two or three hours should be taken until this failing, opium and hyoscyamus supposi enemata may be administered. The diet shou not lowering. As the acute stage passes off, the may be watched, in one or two ounce doses f and then of small quantities of tincture of cub or of capivi, giving twenty minims of the fo eight of the latter, thrice daily. In the severe ment advised above is to be adopted at the c but if the general fever heightens, and symptom appear or threaten, support and alcoholic stin given freely, and large hot local fomentati maintained, should be substituted for the hip- by the mouth, is usually necessary to tranquil with warm enemata, daily, to unload the bow must be watched for, and the catheter, a soft employed if necessary; but never unless it be If there are signs of renal inflammation, the a treatment must be vigorously applied.

Cystitis resulting from cantharides locall absorbed comes on usually within two to four dose; the symptoms resemble those of gonc

aggravate the symptoms. It is highly important, if a has been the cause, to remove it at once, and to sponge the surface in order to detach every particle of the cantharid which may be adhering to the skin.

Chronic cystitis.—Of all urinary affections, this is one of the common, since it is apt to complicate almost every other at one time or another. Its causes also are numerous and varied. First of all, on the subsidence of the acute form, a chronic condition commonly supervenes, and may continue for a considerable period. Next, all forms of obstruction to the exit of urine produce chronic cystitis, by confining the urine until it becomes slightly decomposed, and consequently acting on the mucous lining of the bladder. This condition is intensified if the contractions of the bladder are also unusually weak and painful. Inability to empty the viscus, from paralysis or atony, when no obstruction is present, acts in a very similar manner. If the urine is unduly acid or alkaline; if loaded with deposits from the kidney; sometimes if it is thin and watery; if it be morbid, as in Bright's disease, or if loaded with irritating matters, such as diuretic salts, &c.,—chronic cystitis may be induced. Foreign bodies in the cavity of the bladder powerfully excite this disease; also tumours of the bladder, simple and malignant. Diseases of the rectum sometimes, such as piles, prolapsus, and cancer, those of the uterus and vagina, by causing pressure, and *à fortiori* when implicating the wall of the bladder, are among well-recognised causes. The bladder is sometimes, although rarely, displaced by adhesion to the neighbouring parts, as to the intestine, usually to a portion of the sigmoid flexure of the large bowel; or it may be forced into a hernial protrusion: in either case its contents are partially retained by pressure, and thus produce more or less of the affection in question. The structure affected is the mucous membrane lining the bladder; rarely is any other implicated. After long-persisting inflammation, it is thick, velvety, and dark in colour, its vessels are loaded, while the underlying fibres are usually hypertrophied. Chronic or subacute cystitis appears in two distinct forms: the first, an ordinary inflammation, in which a more or less active hyperæmia, hypersensibility, and very slight increase of secretion, are the main elements; the second, often a condition succeeding to this or to the acute form, in which a passive congestive hyperæmia from local debility in the capillaries,



there is present subacute or chronic present, capable of being detected in circumstances: and is therefore by import, as it is often taken to be. or near the bladder may give rise to cause being removed, the effects Ordinarily it ceases with rest, hot mild aperients, and sufficient alkal acidity of the urine or nearly so, necessary, the use of buchu, triticum in small doses, may be adopted. The form of the complaint, however, is of treatment becomes wide and im-

Chronic cystitis with increased secretion of the mucous membrane h
Respecting the nature of the discharge of opinion has been manifested by eye characteristics are well known. into which the urine has been perfectly transparent, tenacious, ropy material white of an egg, is soon deposited tents of the vessel, it is seen at first soon it slides down in a mass, and then falls out suddenly and heavily succeeded by some as simple m-

pus fluid, containing numerous pus corpuscles and young alial cells ; but the proportion of the liquor to the corpuscles is much larger than in ordinary pus. The old term, muco-pus, seems it perhaps more correctly than any other.

causes.—An attack of acute cystitis ; permanent obstruction of any kind, causing retention, stagnation, and decomposition of urine ; calculi and other foreign bodies ; tumours in the bladder ; spinal paralysis, not only as a cause of retained urine, but occasioning perverted action of the mucous membrane by increased nervous supply ; and all causes which produce a long-continued irritation of the organ.

treatment.—The inflammation is of chronic and atonic character ; depletion and antiphlogistic remedies are therefore contraindicated. On the contrary, the treatment should be chiefly tonic and astringent. It is also more topical than general, since the remedies taken internally appear to act chiefly, if not entirely, by their local influence, impregnating the urine which is retained in the bladder. It will nevertheless be advantageous to regard it under two heads, as topical and as general treatment.

Topical treatment, strictly so called. One important consideration in the treatment of this affection is to maintain the diseased membrane of the bladder as free from inorganic or other deposit as is possible. The urine itself also, if alkaline and irritating, must not be permitted to remain in constant contact with the diseased membrane. It is therefore desirable to introduce the catheter for the purpose, first, of insuring the complete contraction and evacuation of the bladder, if its own efforts do not accomplish this ; and secondly, to remove morbid deposits by injecting warm water, and washing out the interior of the viscus. This is often productive of great comfort to the patient ; and increased benefit may sometimes be obtained by previously impregnating the water so employed with astringent and adjuvant agents. Of the former kind, one of the best is the acetate of lead, commencing with one-sixth of a grain to the ounce of water ; next to this, the nitrate of silver, commencing with one grain to eight ounces of distilled water ; or nitric acid may be used, in the proportion of ten minims of dilute acid to eight ounces of water. By slow degrees the nitrate may be increased as far as to one grain to the ounce in some obstinate cases, and with much advantage. After washing out the bladder with plain water, at a temperature of about 98° or 100°

(a gum catheter and india-rubber bottle, with nozzle and stop-cock, is the best apparatus), about two or three ounces of the fluid, which should have the same temperature, should be slowly injected, and, after remaining two or three minutes, be allowed to flow out. Two or three drops of carbolic acid to half a pint of water is a useful injection when the urine is fetid.

Anodyne solutions may be used, but are of small value. I have employed the following: Dissolve one drachm of each extract, conium and hyoscyamus, and half a drachm of extract of opium, in two drachms of proof-spirit and fourteen drachms of water; of this solution, add a sixth or a fourth part to three ounces of warm water for an injection, to remain in the bladder five minutes; two-thirds should then be permitted to flow out, and the catheter withdrawn; the rest is retained in the bladder.

All the manipulation connected with these processes must be of the most gentle kind, otherwise more harm will be done by the manual interference than good by the therapeutical element. And where the plan is manifestly not producing some marked benefit after a few trials, and, *à fortiori*, if it increases the symptoms, the application must be discontinued.

It should be borne in mind, however, that on all occasions of washing out the bladder, a small quantity of liquid only, such as two to three ounces, is to be injected. This may often be highly useful, while the employment of double this quantity may be productive of nothing but mischief. The bladder is usually more or less irritated by throwing in so much as five or six ounces; and it is on this account that I have long ceased to employ injections in lithotrity, a plan attended with no little benefit.

Counter-irritation is not very manageable in these cases, although sometimes useful. I have employed a Burgundy-pitch plaster, croton-oil in soap-liniment, and strong tincture of iodine, over the pubes, the latter to the perineum also, with some advantage, after shaving the hair. Cantharides, it is said, must be avoided; but I have found no disadvantage from its application to the perineum in the form of the strong acetum lyttæ; and where the inflammation chiefly affects the neck of the bladder, it has sometimes an excellent effect. No better application to the pubes for general purposes exists than a hot linseed-poultice, the surface of which has been sprinkled with mustard.

The pain and loss of sleep often present demand the use of

es either by mouth or rectum, and generally also such use
erients as the system demands in consequence. Each case
res often a different treatment; some experiencing more
from opium by the mouth, others from suppositories or
lenemata. Opium alone, or with belladonna, or conium, or
pyamus, may be administered often with the best results,
ugh the rectum, and with a most tranquillising effect on
bladder.

Internal remedies.—Among the most popular are the
ions of buchu and uva ursi, and the decoction of pareira
a. The indications for their use appear to be the following:
nic mucous discharge from the bladder in large quantity,
iated with relaxation and debility, no inflammation being
ent, may be controlled by the astringent tonic uva ursi; or
agent may be employed with equal parts of decoction of
ira. Chimaphila is considered by some appropriate in this
plaint. The decoction of senega is perhaps superior. In
state also small doses of copaiba or oil of cubeb diminish
secretion considerably. When micturition is frequent, the
e is cloudy, and inflammation is present, a pint daily of de-
on of triticum repens (two ounces to the pint), or a half-pint
e infusion of buchu during the same period, may afford
t relief.

r. Gross employs a combination, sometimes useful in irri-
e and inflammatory states of the bladder. One ounce and a
of the leaves of uva ursi, and half an ounce of hops, are
ed in two pints of boiling water in a covered vessel for two
s; a wineglass to be taken several times a day.

emulcents are useful by diluting the renal secretion, at the
e time that they furnish some little nutriment to the body,
in some instances, perhaps, exert some special therapeutic
ence. They are also agreeable vehicles for the adminis-
ion of acids, alkalies, and other agents, when these are
ired. Among the most useful are the decoction of marsh
low, or of the common mallow in its absence; the decoction
arrageen, or Irish moss; the infusion of linseed; the in-
on of the bark of slippery elm; the decoction of barley,
er known as barley-water; and a solution of gum arabic in
er.

he alkalies and the acids are valuable remedies when appro-
tely given. The old rule of administering the former when
urine is too acid, and the latter whenever it is alkaline, is

not to be followed. No doubt, when there is an excess of acid, it may be most effectually diminished and nicely controlled by the alkaline bicarbonate, as well as by the tartrates and citrates; but this is rarely the case in chronic cystitis. It is far less easy to neutralise the alkalinity of urine, the condition most constantly present in catarrh. Some benefit is, however, occasionally to be obtained from small doses of the alkali, even when the urine is alkaline from large admixture of mucous secretions and decomposition of urinary salts in the bladder. If the urine is secreted with its full amount of normal acid, or with excess, it is liable to irritate a vesical mucous membrane whose sensibility is already exalted by inflammation. Hence Dr. O. Rees has proposed to give in such cases enough alkali, such as the citrate of potash, to neutralise the acid reaction of the urine, so that this cause of irritation may be removed; and in some cases a certain benefit is thus attained. Care must be taken, however, not to produce alkalinity of the urine, which it is our object to diminish. The mineral acids have little or no influence on the reaction of the urine. Large quantities of nitric or of hydrochloric may be given, without producing any result whatever. The benzoic acid is excreted by the kidneys, but a large quantity of this insoluble drug must be taken to acidify alkaline urine. Lemon-juice in full quantity has a similar effect. Alkaline urine chiefly depends on mechanical obstruction, and local treatment is mainly required.

In almost all instances, the patient should have light and nutritious diet; and sometimes the moderate use of stimulants is called for. It is of great importance to maintain the digestive system in good order, and to avoid obstruction of the abdominal viscera; an occasional dose of blue-pill is sometimes very serviceable to the patient. Rest in the recumbent position, and abstinence from all exertion, especially walking or riding, are in themselves matters of the first importance in treatment.

Suppuration of the bladder is that condition in which the catarrh, already described, consists mainly or entirely of a purulent secretion. It is to be treated chiefly with local mild astringent injections, and has been sufficiently considered with the last subject. The larger the proportion of pus in the fluid of catarrh, the graver is the existing affection.

Abscess of the bladder.—Occasionally deposits of pus occur in the walls of the bladder; these are always the result of pre-

ing severe disease, usually of a chronic character. Their occurrence is rarely to be diagnosed or even suspected during life, and no special treatment is indicated.

Ulceration of the bladder.—This is by no means a common lesion, and seldom occurs except in the last stage of urethral disease, arising from obstruction when, after long-standing inflammation of the mucous membrane, ulceration takes place.

It may result also in advanced tubercular and malignant diseases of the bladder; the ulceration following softening of tubercular deposit, and disintegration of the cancerous masses.

A simple form of ulcer has been observed to occur, permitting a silver catheter to remain with a large portion of the instrument within the cavity of the bladder, upon the mucous membrane of which it has rested or impinged for some time.

Signs of ulceration are, blood in the urine, severe pain at a fixed spot, constitutional depression, usually soon ending in death. The urine contains débris of mucous membrane; and the ulcer increases rapidly, or the tissues become gangrenous, and there is much fœtor and dark sanious matter mixed with the urine. Instrumental explorations are exceedingly painful; there is constant desire to pass water, and the effort produces a severe agony.

Treatment.—Stimulants and support for the general condition; small doses of opiates, by mouth and rectum; small doses of nitrate of silver internally, and diluents; no unnecessary mechanical interference to be permitted.

Vesico-intestinal fistula.—Fistulous openings sometimes form between the bladder and some portion of the intestinal canal. The result is, that some of the liquids from this latter source enter, at first in minute quantity, and give rise usually to much irritability of the bladder, and other symptoms of subacute cystitis. These are sometimes the earliest signs of the existence of an abnormal communication. If the urine is examined at an early period, fragments of vegetable and of animal fibre may be discovered under the microscope. I have myself been enabled to diagnose at an early stage, in otherwise doubtful circumstances, the existence of vesico-intestinal fistula by this means. Usually the symptoms become gradually more severe, and the fœcal odour and colour are communicated to the urine.

At length, considerable fragments of fœcal matter enter the bladder, become partially dissolved there, and pass through the

urethra. I have seen several cases of this distressing condition, both in the male and in the female, but chiefly in the former; and in these, carcinomatous disease in the abdomen has been the most frequent but not the only cause. Thus, adhesion may take place between an ulcerating bowel and the bladder; and the ulceration may extend into the latter, no cancerous disease being present.

Treatment.—Little can be said in relation to this matter. The lesion having been determined, instrumental interference is—except under special necessity, such as for the purpose of removing foreign bodies from the bladder, which are a source of great misery—to be shunned. The use of purgatives is also to be avoided; but the bowels should be maintained, as far as possible, in the natural condition. The food is to be such as will nourish the body without producing a large and coarse residue of fecal matter. In one case under my own care, in which a gentleman passed for several months the whole of the feces by the urethra, a great amount of suffering was occasioned by swallowing indigestible substances, like grape-stones and husks. Subsequently, by a careful selection of fluid nutriment, the most painful features of the case were much ameliorated. Now and then washing out the bladder with small quantities of tepid water gently thrown in contributed to the patient's comfort. In this case the ulceration was not due to cancer.

For some of these cases of non-malignant disease another form of treatment remains. When it can be determined that the fistulous passage connects the bladder with the rectum, or with the sigmoid flexure of the colon, the propriety of opening the descending colon [Amussat's operation] ought to be considered. In one case, recorded in the *Medical and Chirurgical Transactions*, vol. xxxiii., Mr. Pennell of Rio Janeiro performed this operation with success, diverting the passage of feces from the bladder by the artificial anus. More lately Mr. Holmes has achieved a similar success in a case in which the fistula was believed to exist low down in the colon. It is reported in the same *Transactions*, vol. xlix. pp. 66 et seq. The point to be determined, if possible, before operating, is whether the communication exists between the small or the large intestine or rectum, and the bladder. If it is in the descending colon or below, urine will probably issue at some time or another per rectum; if the small intestine is the seat of ulceration and opens into the bladder, no urine can be expected to issue by

well. In favourable cases, then, this proceeding should be adopted.

Hypertrophy and sacculation of the bladder, dilatation, &c.—The muscular walls of the bladder become hypertrophied in most cases of obstruction to the outlet. Increased force being necessary to expel the urine, the organic fibres are augmented in proportion to the action required. Hence the vesical coats may become rugose and columnar, resembling in appearance the left

FIG. 283.



sac projecting from the right side of the bladder. The communicating orifice admits a common quill. Cause, long-standing hypertrophy of the prostate, the thin portions seen to be enlarged below. *a*, prostate gland divided. *b*, cavity of sac laid open.

ventricle of the heart; and they may reach half an inch, and sometimes even an inch in thickness. From the interlacing of fibrous bands which constitute the parietes of the bladder, it often happens, when hypertrophy exists, that the mucous membrane is forced by pressure into the interstices between, and gradually sacculi are formed. These at first are very small, but if the action is continued and powerful, they become larger in size, protruding, in the form of tumours, beyond the neck of the bladder, and forming additional reservoirs for urine. A sac, or sacculus, may thus sometimes attain the size of the

IV. 3 M

although it may be suspected ; no requisite beyond that necessary to the bladder fails to expel its contents, inflammation.

In addition to hypertrophy, the times thickened by an interstitial maturation ; a circumstance which in the muscular action of the viscus. is simply dilated, and even distends the walls being thinner instead of condition occurs, not in stricture, but prostate where, by reason of the simplifying the muscular apparatus at viscus is incapacitated from contraction distension. Hypertrophy does not muscular action is impossible, or not

Vesical hæmaturia. (See Hæmaturia)

Tumours of the bladder.—These are as follows :

1. Fibrous growths, of a warty and
2. Villous or vascular growths.
3. Epithelial growths.

re, protrude, and assume a polypoid form. A single clinical preparation may afford examples of all stages of development between the earliest elevation and the fully formed solid tumour. There is no analogy whatever in origin or nature between them and the uterine growths; nor between them and the polypoid outgrowth, which is frequently found arising from the posterior part of the prostate.

Symptoms are those common to all forms of obstruction and irritation, added to the special signs of foreign body in the bladder. Careful sounding may detect their position and form, and will prove the absence of stone. The complaint is most frequent in children and young people. Unlike the villous and malignant growths, they rarely give rise to blood in the urine.

Treatment.—No special curative treatment is applicable in the case of these growths; much may be done to palliate the symptoms and to support the powers of life, by adopting the means which are fully detailed in considering the effects of obstruction under the heads Stricture, Enlarged Prostate, Chronic Cystitis. In the female, a growth of this kind has been removed by ligature, the urethra easily admitting the same degree of dilatation.

Villous growths.—These have usually been classified as 'benign,' but have no claim, either by their structure, their origin, or by the presence of constitutional cachexia, to be so considered. The villous growth of the bladder is a soft, gelatinous body, about the size of a large marble when fully developed, made up of innumerable villous processes, floating in the urine and branching off in every direction from the base, which is connected solely with the mucous and submucous tissues. These villi are identical in structure with the villous processes of the chorion in its normal state, and are very vascular. Several such tumours may coexist in one bladder; and the whole or a large portion of the mucous lining of the organ may, when examined, be found more or less studded with small nodular processes, similar to those of the tumour itself.

Symptoms have nothing special to distinguish them from those of other tumours, excepting the frequent or almost constant appearance of some blood in the urine without provocation. In the end, death occurs rather from hæmorrhage than from obstruction in these cases, as well as in part from the exhaustion

Treatment.—The indications are
spasmodic action of the bladder,

FIG. 28



Villous growth. From a specimen



internal remedies, and to counteract
by chalybeates and nutritious diet.
carefully introduced into the bladder
acetate of lead or of nitrate of silver

case the symptoms resembled those of stone, only that the slightest of movement produced bleeding, in the later stage. Injections of nitrate of silver, from the half of a grain to one grain to the ounce, controlled this ten-remarkably, and enabled him to walk a mile or two without hæmorrhage. At death the same deposit was found in one kidney. (*Path. Trans.* vol. xviii. 32-4.)

Malignant disease of the bladder is more frequent than the one just described. Sometimes it is primary, commencing independently in the bladder; at other times it is secondary, arising from the prostate, from the uterus, or from the rectum. The species may be scirrhus, but is usually encephaloid, either originating in the bladder; always the latter when arising from the prostate; when from the rectum, it is usually scirrhus; from the uterus, the same, or epithelioma. It is said to have existed, although it must be extremely rare.

The disease generally runs a rapid course, and presents constitutional characters significant of malignant action. In cases of encephaloid a large mass is soon formed, which sometimes almost fills the cavity of the bladder.

Symptoms.—One of the most strongly marked signs of this affection is hæmorrhage into the bladder, generally occurring suddenly and in large quantities, rather than by frequent or continuous oozings from capillary vessels, which latter mode is more characteristic of villous growths. The pain is also very severe and lancinating, and is referred as much to the loins, back, and thighs as to the pubic and perineal regions. All the signs of obstructive disease and chronic cystitis are present. Constitutional cachexia accompanies the local disease in its advanced stage; glandular enlargements are usually distinguishable in the iliac regions; and the patient suffers from loss of sleep and rest. Small portions of disintegrated tissue and purulent sanies are often voided from the bladder; occasionally blood may be found in the urine, which, when examined by the microscope, seem to indicate the specific character of the disease; but the diagnosis is generally tolerably clear without the somewhat questionable support which is to be derived from this source. Without in any respect undervaluing the utility of microscopic examination, I am compelled to conclude that too much stress must not be laid on the indications which cell-growth in the urine afford in respect of the presence of malignant or non-malignant disease in the urinary passages. Even large masses of soft growth however, are frequently



freedom. In all those complaints of the little or nothing remains to be done but support the powers of life, the remedie given with no sparing hand. So severe the patient suffers if left to the act unchecked or only partially checked by thetics, that it must be regarded as c their aid in mitigating his distress, so f available without danger to life; while a holic stimulants are in some cases scarc opium in enabling him to endure the pro

Tubercle of the bladder.—One of the 1 of the bladder is tubercular deposit; p unassociated with tubercular disease in c indeed in almost all—perhaps it may other parts of the urinary system are these are generally the kidneys and pros been found to have followed primary dise disease commences by a deposit in the small tubercles; these become numero sequently the morbid material softens, 1 gives way, disintegration takes place, and a large extent of the inner coat being

ly or middle adult life, will serve to indicate the true character of the disease.

Treatment—That which is adapted for constitutional tuberculosis in relation to the general health. For the local complaint, anodynes, the remedies for chronic cystitis, rest; and mechanical interference when the diagnosis is tolerably clear.

Bar at the neck of the bladder.—The enlargement of the prostate, common in old age, often produces an elevation of the ducts, sometimes amounting to a distinct ridge, which rests at the inferior aspect of the neck of the bladder. A bar may be thus said to be formed there; nevertheless, it has not been usual in this country to apply that term to any product of the enlarged prostate; it has been thought more desirable to reserve it rather to denote any bar which may exist in the spot described, but which is not prostatic in its character. Such bars occasionally exist, and usually in elderly subjects. Mr. Guthrie was the first to point out this fact. He believed the bar to consist in disease of 'the elastic structure,' existing at the neck of the bladder, and as such to be quite distinct from the very common complaint already referred to. After considerable study of this question, I cannot but consider the non-prostatic bar as extremely rare, since almost all permanent obstructions at the vesical neck are due to enlargement of the prostate and or to its effects; nevertheless, in very exceptional instances the bar in question is undoubtedly to be met with.

In all long-continued diseases of the urethra and bladder it is common to find a hypertrophied condition of the uvula and associated muscular fibres, and more or less elevation at the neck of the bladder, or bar, as its consequence; but this is not to be regarded as an example of the disease in question.

The treatment mainly consists in combating the symptoms of chronic cystitis which may arise, and occasionally passing a large catheter, so as to insure a patent condition of the neck of the bladder. Mr. Guthrie and others have proposed to divide the bar by incisions; but the rarity of any true barrier here, apart from prostatic enlargement, will induce extreme caution in affirming positively the presence of this affection during life, and any severe surgical operation for its relief could only be contemplated in presence of very serious symptoms as its undoubted effect. The subject of operations for relief of obstruction at the neck of the bladder is considered at the close of

to be retained in that situation, the bladder is retained in the pelvis, and in part the scrotum. In a hernial protrusion of bowel, or may be this. In the former condition it has been opened by operation for strangulated inguinal hernia, but in the latter it was opened by mistake, as in one of the two cases. The portion of bladder forming the hernia was situated beneath the peritonæum, and consequently covered by that structure. In cases where a truss may be successful in retaining the protrusion of the pelvis; where permanent adhesions have taken place, mechanical support is necessary to prevent further protrusion.

Inversion of the bladder.—This is a rare condition, and is found exclusively among children. Cases only have come to my knowledge which had been seen were from two to four years, probably commenced at a very early age, and gradually prolapsed and protruded through the scrotum, becomes much distended, until the appearance of a pyriform, red, and vascular mass. In some cases narrowly escaped the application of the suppositorium referred to; but on careful examination of the ureters were discovered, and reduced.

abled subsequently to retain some ounces of urine, and slight continence only remained.

Conditions which do not necessarily involve organic structural changes, appreciable by known methods of observation, and which are generally associated with some injury or derangement of a nervous centre.

Paralysis of the bladder.—One of the most common terms used in relation to complaints of the bladder is paralysis. Yet the paralysis of the bladder never occurs without some lesion of a nervous centre; so that the word is often incorrectly employed. For example: the bladder when unable to evacuate its contents from obstruction by stricture, or by prostatic enlargement, or by atony from over-distension, is commonly said to be 'paralysed,' although there is no derangement of the nervous supply; to which condition the use of that term ought, in accordance with our employment of it elsewhere, to be limited. The bladder may possess its normal contractile power, but be unable to overcome an abnormal obstacle in the outlet, and so fail to expel its contents; the complaint in this case is not paralysis of the bladder, but solely obstruction in the channel of exit—two totally different conditions. This distinction must be clearly kept in view in our consideration of this subject.

True paralysis of the bladder is a condition in which the contractile power of its muscular fibres is lost or impaired from failure of nervous supply; thus there may be partial as well as total paralysis. The usual cause of this accident is cerebral or spinal lesion. It happens in some injuries to the brain, in almost all serious injuries of the spine, in softening, in apoplexy and in all serious organic affections involving those parts of the cerebro-spinal system which affect the urinary apparatus. It occurs from spinal debility produced by excesses, especially of the sexual kind; also in temporary functional derangement of the nervous centres; through the agency of reflex action from some adjacent source of irritation, as in hæmorrhoids, and especially after operations for their removal; and after other operations; although in some of these cases the retention is probably due rather to spasm of the constrictor muscles than to paralysis of the bladder. It may also happen after shock to the system from any source, in fevers, &c.

I have met with a single case of permanent loss of power in the bladder attributed to an overdose of belladonna. Certain

regarded as common occurrences, the in almost all cases atony from over-di (see next section).

Treatment of paralysis.—From what is to empty the bladder, by means thrice daily. It should be employed patient has probably lost sensibility, complaining of pain, when injury is in is often in a lax state, and consequer easily made. A full-sized instrument whichever is found to pass most easil signs of chronic cystitis manifest th remedies must be employed, especial which washing out the bladder, follow injections, may be reckoned as the bes



In cases not traumatic, or depend strychnia, steel, cantharides, ergot of a trial. Galvanism, cold douches, ar be employed as local agents.

Atony from over-distension.—Org existed either at the neck of the blad in the course of the urethra (strictur period, and having gradually augmer

dition of complete contraction, are over-stretched, lose their tone, and, even on the obstruction being removed, cannot expel the contents of the viscus. A similar state may be produced when no organic obstruction is present, from long-continued retention of urine exerted voluntarily, or occasioned by temporary spasm, or by some cerebral affection, in fevers and the like. The patient whose bladder has been thus over-distended often finds himself unable to empty it subsequently. Such is the condition termed here atony, and often falsely called paralysis, already noticed in the preceding section.

Treatment.—It is essential, in the first instance, to insure the complete emptying of the bladder for a time, at least once or twice daily. The muscular fibres must be enabled to resume their normal condition of contraction from time to time; and this can never take place unless the bladder is artificially emptied. Sometimes, if the over-distension has not continued long, they gradually regain their tone, or at least recover much of it. When the abnormal state has existed long, it may often greatly benefited, but is rarely completely removed. Without the mechanical aid all medical treatment is absolutely useless, but it may perhaps be useful in combination with the former. The general and local stimulants of nervous action, already advised in paralysis, should be employed in this affection also. The cold douche over the lumbar spine and on the perineum is sometimes particularly useful. Injections of cold water into the bladder itself are the best of all direct applications to the viscus.

Irritable bladder.—This term is often very loosely employed to designate some specific affections of the bladder. Like the term ‘dropsy,’ ‘irritability of the bladder’ denotes nothing more than a symptom; in either case the real cause must be discovered; the terms themselves conveying no meaning beyond the simple fact, in the former case that a collection of fluid is present, in the latter that the patient passes urine too frequently. Irritability of the bladder is a symptom present in almost every affection of the urinary organs, from childhood to old age; and is named here at the head of a section solely on account of the conventional misappropriation of the term, and as affording opportunity of protesting against it. Nothing is more common, even in professional communications, than to be asked what is the best remedy for ‘irritability of bladder.’ It denotes no special pathological state, and always requires to be con-



disease has not been discovered ; but the
causing cause may be predicated as invariably
whatever the cause may be, by general an-
thetics, and antispasmodics, as described
treatment for cystitis, prostatitis, and stricture
it is liable to coexist.

Perverted sensibility of the neck of the
There are a few cases of severe urinary
which neither researches during life nor
have revealed the nature of the affection.
remark that the number of instances ad-
in practice, diminishes in direct propor-
skill in diagnosis which is brought to light.
Usually the patient complains of symptoms
those of stone, but repeated soundings find
any other deviation from the normal state.
urine is healthy, or at most only occasional
undue action of the mucous membrane. It
aggravate the symptoms, sometimes have
air and scene often produce an improvement
only temporary. By careful investigation,
liver, early stages of renal disease, the pro-
commencing cerebral changes late in life
found to account for some cases ; but never
apparently inexhaustible. Such have been

occurs in adult subjects of all ages, when the bladder is already over-distended with urine, a certain quantity, which usually bears a very small proportion to the quantity retained, running off either during sleeping or waking. The following rule almost invariably holds good: in juvenile incontinence there is no retention of urine; in adult incontinence it is present to a serious extent. There is no principle more important to remember in the treatment of diseases of the urinary organs than this, viz. that an *involuntary flow of urine in the adult indicates a distended and an empty bladder.*

1. *Incontinence in childhood and youth.*—This is a common and sometimes a very troublesome infirmity, affecting patients generally under the age of puberty, although occasionally it continues during some years after that period. The single symptom is that, in the middle of the night or towards early morning, the child wets the bed without awaking, and the act may be repeated once or a greater number of times before rising. Sometimes this occurs every night; sometimes there is an interval of one night, rarely of more. The patients so affected do not appear to belong to any particular temperament, nor to be necessarily weak and cachectic. In some the failing is simply a habit resulting from want of management, such as neglecting to take up young children once during the long period which is necessary to them as a night's rest. In such cases careful attention to habits and diet may arrest the infirmity. In many cases, however, no ordinary management succeeds, and medical aid is sought.

Innumerable remedies have been given, and with a very unequal success. Small doses of cantharides and of strychnia, either with or without iron, are sometimes employed. Tincture of iron alone, or the vinum ferri, has sometimes proved successful. Blisters to the sacrum has been a favourite remedy; also cold douches to the spine, and general tonic treatment in most cases. In my own experience, nothing has afforded such good results as belladonna. I have employed it in the form of extract, commencing, according to the age of the patient, with the sixteenth to the eighth of a grain twice a day, and increasing the dose if required as far as it can be safely borne. It is necessary to add, that disappointment has arisen in the use of this remedy from the impurity of the drug; it is absolutely necessary to employ a pure extract. After failure with a pre-

included the removal of worms, and other sour irritation always to be looked for in obstinate may generally be attained sooner or later, although it is occasionally necessary for a long period

2. *Incontinence in the adult*, as has been before in the vast majority of cases, a bladder distension; the cause of which may be stricture, or cystitis, stone, atony from over-distension, or Very rarely does it happen in the male that the urine runs off from the bladder as fast as it arrives there from the ureters. Hence it is a cardinal principle in practice, that when the urine runs off incontinently, a catheter should be passed to ascertain the amount of retention present, and to relieve the bladder. In the female, true incontinence is more frequent, and is usually caused by a traumatic injury to the short urethra, which has impaired or destroyed the sphincter's action, such as from pressure in labour, or from the use of instruments to move the child, from over-distension of the ureters, or from calculus, &c.

In regard to the treatment of incontinence, it is to recapitulate that which is appropriate to the various conditions above mentioned as causes of an unnatural flow from the bladder, since they have been considered under various headings. When that rare affection in the female, true incontinence, maintaining the bladder in a constant empty state, is really present, the cause will be

investigation of the cerebro-spinal system must be instituted, and some other sign of lesion of that system will probably be discovered, in which case the treatment must consist in that which the general affection indicates, together with the employment of some local appliances suited to protect the patient from the disagreeable results of his infirmity, such as a well-adapted urinal. Where incontinence results, as occasionally happens, from mere functional derangement, caused by debility from excesses, the cure will be effected by any means of a tonic character which can restore normal tone to the spinal system. For this object the preparations of steel and zinc, of strychnia, together with quinine and other vegetable bitters, are of value. Nutritious diet, cold sea-bathing, and abstinence from the causes of debility previously acting, form also a part of the treatment for these cases. The local alteration of structure, which does occasionally give rise to the incontinence just described, is a form of enlargement of the prostate, in which the posterior portion, commonly called 'third lobe,' is much hypertrophied, and projects forward in the manner of a wedge between the two lateral lobes, so as to open out the neck of the bladder and render it constantly patent; instead of projecting backwards into the cavity, as almost invariably happens, and forming a species of valve which prevents the outward flow of urine. The former condition permits the bladder to remain constantly empty, and the organ becomes small in capacity. The latter maintains it constantly filled with stagnant urine, unless the catheter be employed to remove it, and produces a dilated and capacious bladder. The treatment of the former condition consists mainly in providing a urinal; nothing in that way can be done to modify the form or condition of the enlarged prostate, beyond that which is effected in preserving good a state of the general health as circumstances admit.

Habitual engorgement of bladder, and overflow of urine.—These terms more correctly describe the conditions met with in the majority of cases in which urine passes involuntarily than that of 'paralysis,' so commonly used, and already referred to at length. Urethral obstruction from enlarged prostate, or stricture, having existed for a certain period of time, the bladder becomes unable to contract with sufficient force or continuance to overcome the obstacle and empty itself; so that a certain portion of urine always remains after every act of micturition. This failure of the bladder to contract fully, when once established,

ureters and kidneys.

The treatment of engorgement and overfilling the bladder from once to three or four times in twenty-four hours, according to the necessities of the case, in using those remedies which the amount of urine voided almost always present to some extent, indicates (see Chronic Cystitis, p. 889).

Retention of urine (see Stricture).

THE PROSTATE.

Malformations.—There are no special malformations noted here, except such as are involved in the condition of ‘extroversion of the bladder,’ which see (p. 881).

Injuries (see vol. ii. p. 735)

Inflammation of the prostate } acute,
(*prostatitis*), } chronic.

Abscess of the prostate.

” periprostatic.

Hypertrophy and simple tumours.

Atrophy.

Malignant disease.

Tubercle.

Cysts.

Acute inflammation of the prostate.—This exists alone, but is generally associated with

the occurrence, and not by extension through continuity of
 This idiopathic inflammation, excepting the cases produced
 by violence, as by instruments, &c., is probably extremely

causes.—These are: the pre-existence of acute inflammation
 of the urethra from any cause, but especially the gonorrhœal;
 inflammation of the urethra in an aggravated form, giving rise to
 inflammation of all the parts posterior to it; the application to
 the prostate of strong injections, cauterisation, the passing of
 catheters, and mechanical violence of various kinds; inflammation
 of the bladder, and calculi. Cold and damp to the perineum
 give rise to it, perhaps most frequently in gouty and rheu-
 matic subjects. Alcoholic drinks may induce prostatic inflam-
 mation, urethritis already existing, but only on this condition.
 Excessive sexual excitement, under the last-named circum-
 stances, is also an undoubted cause.

Symptoms.—At first, a sensation of weight and fulness about
 the rectum and perinæum is experienced, with some pain and
 tenderness at the neck of the bladder. The patient wants to
 urinate more frequently, and does so with much pain,
 especially at the close of the act. These symptoms increase;
 pain becomes severe, shooting and throbbing, and almost
 insupportable; a sense of great tension is experienced, and the
 rectum and perineum are tender when pressed. Pain in the back
 joints, and often in the thighs, is complained of. Movements
 of the body become painful, as does also the sitting position.
 The act of relieving the bowels at stool produces considerable
 distress; and a finger introduced into the rectum encounters
 resistance; the anterior wall of the bowel is prominent, hard,
 hot, and the outline of the prostate may be traced, much
 more than in the natural state. An attack of piles may be
 excited, the close contiguity of the hæmorrhoidal and prostatic
 parts favouring this result. Should a catheter be passed, the
 patient complains of excessive pain when the instrument reaches
 the prostatic part of the urethra. At a later stage, if suppura-
 tion has taken place, the rectal swelling becomes softer. Micturition
 is often difficult, and may even be impossible, total
 retention occurring from the obstruction produced by the swollen
 prostate. General fever accompanies these symptoms, com-
 mencing with rigors, and manifesting itself by all the usual
 signs during the acute stage of the disease. Pus usually
 appears in the urine after a few days, and if there has been

outset than anything else. From ten to
be applied to the perineum and around
to six or eight ounces may be performed
should be taken, followed by a large pill
and the patient be wrapt up warmly in
be repeated with advantage in the course
should not be used for more than ten
rule. If retention of urine occurs, an
once to hot fomentations and a full dose
sized gum elastic catheter should be
relieved. The instrument may generally
and is to be used again if necessary (see

In the course of a few days the severe
subside, often, however, not without the
relapses. These are mostly attributable
the part of the patient, either in the use
or erotic excitement. When convalescent
prostate still remains much enlarged and
of urine is not propelled with the natural

Chronic inflammation.—In this condition
times enlarged, but not invariably; whereas
to the acute attack, perhaps it always
are: diminished force in propelling the
some gleet discharge: the urine is clo

for some weeks, and the surface dressed with simple ointment. Tonics and good diet, with attention to the digestive system, are almost always indicated; these latter being in fair proportion, iron is mostly beneficial; and as the bowels should in all cases not be constipated, the iron should be combined, if necessary, with some mild laxative. When nocturnal emissions are frequent, and especially if they occur without any consciousness whatever on the part of the patient, the application of a solution of nitrate of silver, from ten to thirty grains to the ounce, to the prostatic part of the urethra, is one of the means of treatment. To be successful, an efficient instrument is absolutely necessary, as well as care in injecting the fluid at the right spot: three or four applications may be sufficient. Let it be remarked, that we are by no means to neglect the existence of chronic prostatitis because the patient is not troubled by frequent seminal emissions. These occur frequently from altogether different causes, which, as they have no necessary connection with disease of the prostate, need not be further considered here.

When enlargement only remains after acute inflammation, iodide and the bromide of potassium should be given for a considerable period of time. Conjoined with or following this treatment, a course of tonics, steel and quinine, with sea-bathing and generous diet, often effects a great improvement and completes the cure.

Process of prostate.—Acute inflammation, when neglected, or arising in a naturally weak or delicate constitution, may terminate in suppuration of the organ. This occurrence may be expected when after the first week or two the acute symptoms do not subside, when the pain and difficulty of micturition and retention increase, if rigors occur, and the patient is very anxious and feverish, complaining of tension and pulsation in the perineum or neck of the bladder. Fluctuation may be felt in the rectum; or pressure in the perineum may reveal tenderness and fulness there. The natural course of abscess in the substance of the prostate is generally evacuation by the urethra. Occasionally it issues by the rectum, and this is perhaps as probable, generally speaking, as through the urethra. It may be followed by a troublesome urethro-rectal fistula; but this is rare. Generally the opening of the prostatic abscess into the urethra soon closes; but if this does not take place, the cavity

may long remain open, and become a receptacle for giving rise to fresh abscesses in the vicinity.

In treating a case presenting the symptoms described should watch for any evidence which may exist of a tumor forming in the perineum; and if so, an incision should once made boldly in the median line, so as to evacuate it successfully, the patient will be greatly benefited; but if it is encountered, no harm results, but the relief of pain and tension is probably effected. Occasionally puncture of an abscess has been accomplished in the rectum when fluctuation has been distinctly felt there.

The discharge of matter from an acute abscess of the prostate is sometimes followed by long-continued suppuration. A chronic abscess may arise here spontaneously, though not very frequently; generally resulting from confirmed or neglected stricture of the urethra. When this has occurred, and matter has been discharged into the urethra, it sometimes happens that the cavity enlarges, urine makes its way in, fresh irritation is maintained, and much of the prostate is destroyed; its capsule becoming little more than the surrounding pus-secreting cavity. Sometimes the abscess is limited to one side or lobe only, and sometimes the collections are two or more in number; generally speaking, however, the result found at death is that described. This condition is always one of great import, and its existence is often not suspected during life. There are present the symptoms of chronic cystitis, emaciation, often hectic; and a highly tonic and soothing regimen is usually indicated. Injections of nitrate of silver, of half or one drachm to the ounce, are sometimes useful; but no local interference is desirable which gives much pain, or even temporarily aggravates the symptoms.

Abscesses supposed to be prostatic have not unfrequently turned out to be situated external to the prostate, and not within the envelope of the organ. Such are termed periprostatic. They arise much in the same manner as prostatic abscesses, but are of less serious import than those situated within the capsule of the prostate itself. The treatment requires early incision, and does not differ in other particulars from that already described for prostatic abscess, properly so called.

The organic enlargement of the prostate of advanced age (hypertrophy). This common affection very seldom occurs before sixty years of age; I have never seen or heard of a true exan-

before the age of fifty-four years. Chronic inflammatory enlargements may occur at any age after puberty; but the case now under consideration is wholly distinct, both in time and by causal relations, from such or any other inflammatory state. It has been said to be the common inheritance of old age, but it is, on the contrary, a complaint which a very large majority of elderly men escape. With the view of ascertaining the relative frequency of its occurrence, so far as such a number of observations can be regarded as indicating it, I dissected out with great care the prostate in forty-three elderly individuals, the results of which inquiry are presented in the following summary: Appreciable enlargement existed at the age of 32 per cent.; notable enlargement, causing symptoms of obstructed life, at the rate of 12 per cent. The oldest individuals in the series were among the unaffected; including 1 at ninety, 1 at eighty-five, and 2 at seventy-nine years. Subsequent more extended researches by Dr. Messer and myself have but confirmed these results.

The enlargement consists mainly of an over-development of the glandular tissues which enter into the normal structure of the prostate itself. These tissues consist, first, of unstriated muscular tissue and the connective tissue always associated; and they form at least three-fourths of the prostatic body: secondly, interspersed among this structure are numerous branching glandular tubes and crypts with their accompanying ducts.

Enlargement, or hypertrophy as it is commonly termed, may be determined by an abnormal production of the first-named elements alone, or with some of the glandular tissue interspersed; or this latter may be fully or imperfectly developed. The lateral lobes of the prostate may in this manner be increased four or six times their natural weight and bulk; or an outgrowth may take place from the central part of the organ, in a backward direction, towards the cavity of the bladder, as a more or less rounded or pyriform tumour of prostatic tissue, and constituting what is frequently, but not very correctly, called the enlarged 'third lobe;' and this form is one of those most commonly met with. Generally, however, every part of the prostate partakes more or less in the enlargement.

Again, while the tissues thus described may be generally over-developed throughout the entire prostate, it is very common to find independent, or almost independent, and isolated portions of the same material embedded within the prostatic structure proper. On making a section with a sharp knife of a

prostate so affected, these small rounded bodies are easily divided, and the portions may be removed by the finger-nail. They are usually made up of very compact tissue, like that of the prostate itself, but generally with less of the glandular element, and that, almost always, presenting a defective development. Sometimes they are completely isolated by a limiting fibrous cyst. The presence of these small tumours is extremely common in the enlarged prostate, as well as in the healthy organ in advanced life, and they seem to bear a relation to the containing organ somewhat similar to that which the uterine fibrous tumours bear to the uterus. In both cases they are

FIG. 285.



An example of greatly enlarged prostate, the increase in size being mainly due to the fibrous tumours described; opposite *a*, *b*, and *c*.

composed mainly of unstriped muscular tissue, and exhibit other points of analogy in their nature and history.

The most important result of enlargement is obstruction to the flow of urine. Its first effect on the prostatic urethra is increase, sometimes considerable, of its antero-posterior diameter; there is also diminution of its lateral or transverse diameter, the canal becoming a narrow passage, instead of one which, when distended, is of about equal diameter in every direction. The length of the prostatic urethra is also materially increased, and it is often tortuous also. In some preparations

which I have examined, the urethra has measured four inches from the orifice of the bladder to the membranous portion, instead of one inch and a half, which is the normal length.

The natural direction also deviates: where there is enlargement of the median portion or 'third lobe,' the urethra suddenly bends, producing an angular curvature in the place of a nearly straight line; so that a complete step has sometimes to be mounted at the neck of the bladder before an instrument can enter the cavity. When, with this development of the

FIG. 286.



bladder, symphysis pubis, and enormously enlarged prostate; a portion is removed from the left wall of the bladder, to show the vertical ascent of the urethra as it enters the bladder.

Median portion, there is much enlargement of either lateral lobe, the lateral direction of the canal is also changed. Thus if the right lobe predominates, there will be a lateral curve of the urethra, the convexity of which is presented to the left, and *vice versa*.

The next deviations are those in the form of the internal orifice, or vesico-urethral orifice. When the median portion predominates, this orifice has a crescentic form, the convexity of which is directed upwards. When the right lobe considerably exceeds in size the left, the crescent has its convexity to the left side; and so on. In some preparations, where two or three irregularly-enlarged lobes are combined, the orifice is

much distorted, presenting a tortuous outline. Sometimes overlapped by an outgrowth from the median portion form of a valve. In this case, which is not common, the valvular portion appears to be forced against the neck of the bladder by the effort of micturition, and the obstruction is still more complete.

The size and weight attained by hypertrophied prostate sometimes remarkable. A prostate measuring two inches in transverse direction, and one inch in thickness from before backwards, or which weighs an ounce, must be considered

FIG. 287.





Hypertrophy of the prostate affecting equally the lateral lobes and median portion, a typical example of the usual condition in a moderately advanced case.

trophied. On the other hand, a transverse measurement of three inches is not uncommon. I have seen one exceeding three inches and a quarter; but such a degree of enlargement is extremely rare. The weight has been known to reach six ounces. In malignant disease even these limits are exceeded.

The consistence of the organ varies, being sometimes firm and harder in texture, at others a little looser and softer than in the natural condition. It usually presents the form of a mass of fibrous tissue when numerous fibrous tumours are embedded in the substance.

Symptoms.—In the earliest stage of chronic enlargement of the prostate there is nothing to attract the attention

patient, as it is probable that a considerable period is passed between the actual commencement of enlargement and the occurrence of any marked symptom. One of the earliest signs generally observable is diminution in the force with which the urine is ejected, and the lapse of time which occurs before a stream is established. The size of this is not necessarily much smaller than it was in health, but it cannot be projected so far as the ordinary amount of effort. The desire to pass water becomes more frequent than natural, and the relief afforded by the act of micturition is less complete. On rising in the morning, when the bladder has become distended during sleep, the want occurs again in a few minutes, especially after the first effort; and in course of time sleep is much disturbed by calls to pass water. A sense of weight, fulness, and uneasiness about the perineum and rectum is felt, which the patient soon refers to the neck of the bladder. As the expulsive efforts to pass water become more frequent, irritation of the rectum increases, the contents of the bowel are more frequently passed, an inability on the part of the patient to prevent the act of defecation accompanying that of micturition; and tenesmus, protrusion of the mucous membrane or prolapsus, and hæmorrhoidal swellings, are apt to result. Much stress has been laid by some writers, following J. L. Petit, on the appearance of hardened stools as an indication of enlarged prostate, but certainly without sufficient grounds. In some cases the motions may be passed in the form described; but this is caused by the action of the sphincter ani, and not by the prostate. If the complaint advances unchecked, the symptoms of chronic cystitis appear, and pains also sometimes in the penis and testicles. Occasionally, some muco-purulent discharge from the urethra is observed; this varies with circumstances, sometimes appearing after exposure to cold or damp, or with an attack of retention; while it may happen also that the irritation involves the testicles, which become tender and swollen. Vascular excitement of the penis, producing frequent erections, is also at times concomitant symptom. As the complaint makes progress, relief to the bladder in micturition becomes less complete, as the enlargement gradually closes the neck or urethral outlet, and the bladder is never emptied, a certain portion of its contents only being expelled at each act of micturition. These difficulties, if unrelieved, inevitably increase; the capacity of the bladder yielding to the constantly-augment-



which it extends; it is sometimes found although more frequently to the extent of the fingers' breadth above the pubes.



As the complaint advances, the patient and he is the subject of frequent febrile slight irregularities, or exposure to adduce extreme distress from the fever occasioned. Attacks of complete remission on these occasions, and the functions impaired through the long-continued charge of their secretion. A little hemorrhage, and sometimes relieves congestion, exposure to cold, sexual excitement, which tend to produce a vascular detritus of the viscera.

The characters presented by the urine are the decomposition of some of its constituents, mixed with the products of chronic inflammation of the bladder. More or less of mucous deposit on the lower part of the vessel, and a thin pellicle on the surface, which is sometimes iridescent. The mucus increases in quantity, and becomes tenacious, slimy, adhesive matter so we find with chronic inflammation of the bladder.

Characters are influenced to some extent by the quantity voided, which often varies considerably in the same patient from day to day; the measure being sometimes below, but is generally much above, the natural or healthy standard. As a result of long-continued disorder in the urinary apparatus, and of the changes in the urine itself, which have been described, it is not surprising that the formation of calculus sometimes takes place. Its presence will be suspected where there is much pain about the neck of the bladder at or during the act of micturition; if the pain at the end of the stream is unusually severe, and if the blood and pus are very frequently observed, and especially if fragments of calculous matter have from time to time been passed. But the existence of calculus is often wholly masked by the prostatic disease, because the conformation which the neck of the bladder assumes in this affection tends to prevent the most distinctive symptoms of stone; inasmuch as the foreign body is less liable to be lodged in the vesical neck, but lies back deeply behind the enlarged prostate.

It not unfrequently happens that symptoms of enlargement are long before the real cause is suspected: the frequency of micturition suggesting that there is undue freedom rather than obstruction to the act. The real malady is unsuspected, and the treatment is directed only to those symptoms which have been productive of most discomfort or anxiety to the patient; to some febrile condition, it may be, resulting from a hidden cause. The march of events, however, must ultimately throw suspicion on the state of the bladder; a catheter is passed, and, greatly to the astonishment of the patient, some thirty or forty ounces of urine, or even a much larger quantity, may be drawn off, notwithstanding that the act of micturition has been just performed. And it occasionally happens, during the prevalence of such a state, that some unaccustomed change in diet, or exposure to cold and damp, suddenly produces congestion of the prostate, and complete retention, and so discovers the existence of the affection for the first time.

The last stage of unrelieved disease is indicated more by the signs of a gradual decline of the powers of life than by those of increasing obstruction; although sometimes the final symptoms are those of rapid depression, after sloughing or ulceration of the bladder, and repeated hæmorrhage, or from great discharge



finger in the rectum, more accurate information be obtained than by either explorative or palpation. The patient's knees being drawn up against each other, the finger should be made to pass the sphincter, and when introduced into the rectum, the two phalanges are free to move in the rectum, the prostate may be defined.

If familiar with the normal conditions of the prostate, deviations will easily be recognised by palpation. We may first observe its size and shape, is there any enlargement, is it general or partial? lobes? and to what extent? It is so palpated that the tip of the finger encounters the prostate, enters the rectum, and has to be depressed before it can be carried beneath the prostate, finding the yielding coats of the bladder. When the finger is carried up to its fullest extent, its fullness and firmness may be encountered. Enlargement or outgrowth from the median lobe of the prostate may not be uniform but is usually symmetrical.

Next, we observe whether the tumour is equally so in places; whether there is any fluctuation we can appreciate fluctuation distinct from the prostate in a case of retention of urine.

seen it and the finger in the bowel, and as to the situation and direction of the instrument, &c., in the event of there being difficulty in introducing it.

Having learned relative to these different points all that can be obtained through the rectum, the urethra is to be explored. A small-sized catheter, of the form ordinarily employed by the physician, should be first used, because any phenomena presented arising from those observed when the prostate is healthy are at once made apparent. If by examination through the urethra no variation is found in regard of size, and the urine flows freely when the catheter has traversed not more than the ordinary distance, say from six and a half to eight inches, while the handle of the instrument itself has not required more than the ordinary amount of depression in order that its point may enter the bladder, we may be satisfied that prostatic enlargement does not exist; and we must seek for another cause of the symptoms complained of. But if the catheter has passed easily, for nine or ten inches, and still no urine flows; and if, in addition, while following its course, the handle has become more than usually depressed, there will be little doubt in respect to the existence of prostatic enlargement. The ordinary catheter being inadequate to reach the bladder, a prostatic catheter, which measures from two to four inches longer, and possesses a larger curve than the ordinary catheter, may be employed. If it passes readily, the increased length of the urethra is easily ascertained, and the direction of the prostatic canal is calculated from the position of the shaft when the point enters the bladder. In some few cases, while the beak passes through the prostatic part of the urethra, the handle will be distinctly deflected to the right or left, from which fact, if verified by two or three trials, a greater degree of enlargement may be suspected to exist on the side towards which the handle turns.

For the purposes of accurate diagnosis, we shall employ with advantage an instrument of a different form, viz., an exploring rod with a very short curve at its extremity, or possessing a hook rather than a curve, which is shorter and more angular than that of the ordinary catheter. By its means not only can every part of the bladder be searched for calculus, but information respecting the form and degree of obstruction at its neck can also be acquired: see fig. 288. After the bladder has been traversed, the instrument should be gently withdrawn until the hook lies just within the urethro-vesical orifice, when, by turn-

ing it round to the right and left, the natural co-exist, of that part can be ascertained; or, on the the presence of tumour or of stone, the depth behind the prostate, and other relative points, c-mined. A light, delicate, and practised hand is its use.

This portion of the subject may be appropriately a few remarks on the diagnosis of prostatic enlargement, stricture of the urethra, vesical calculus, tumour of simple atony or inertia of the coats of the bladder.

In stricture of the urethra, the stream of urine small, in a confirmed case extremely so; in the position, though diminished in force, it is much less

FIG. 288.



The exploring sound, with short angular beak, for defining tumour, calculus: in this case represented in a bladder with enlarged

than in the previous case. The use of a full-sized ever, marks the distinction clearly. In stricture, is encountered almost invariably before six inches oment have disappeared, always before it arrives at the urethra. In enlarged prostate, obstruction is not until seven, eight, or nine inches have passed, and rarely then, for, provided that the instrument be long, it may pass into the bladder; but the handle depressed between the patient's legs in a manner in the normal state. Lastly, stricture almost invariably its appearance before middle life, prostatic hypertrophy until that period is passed.

In regard of calculus, while many of the symptoms

to both complaints, the occurrence of sudden cessation of stream of urine, of severe pain at the close of micturition, exacerbation of symptoms, especially of pain and the appearance of a little blood after exercise, may be looked upon as strongly indicating the presence of stone in the bladder. It may exist in the absence of most of these symptoms, the first-named especially, from the circumstance that the calculus is usually situated behind the enlarged prostate, and does not approach the more sensitive region of the internal urethra. The fact of small quantities of florid and unmixed urine being occasionally passed after exercise, more closely approaches in value to a pathognomonic sign than any other. In the frequency of making water is greater at night in the prostatic affection, but greater during the day when the disease is calculus. A persistent discharge of muco-pus streaked with blood should arouse suspicion. The use of the sound, however, can alone clear up the case satisfactorily.

The existence of non-prostatic tumour of the bladder is less fully affirmed. Compared with prostatic enlargement, there is much more pain in the introduction of instruments, the pain is frequently or generally mingled with sanious discharge and flocculi, to which sabulous matter is often seen adhering. Examination of these under the microscope may reveal the peculiar structure of villous growth; or may demonstrate that the flocculent matters consist of organised structures, not of organic materials, a fact possessing some significance.

Simple uncomplicated chronic cystitis, with catarrh, is by no means a common affection. Such symptoms are almost invariably due to the presence of a foreign body, to some form of obstruction, or to paralysis, or atony, depriving the patient of the power of expelling the contents of his bladder, a condition which is tantamount to obstruction. We may rely upon it, that in most of the obscurer cases there is a material cause, frequently calculus; the presence of which needs a more than ordinarily searching examination to verify.

Single or repeated acts of voluntary over-retention of urine are sometimes followed by atony or inertia of the muscular fibres of the bladder, and a state of chronic retention follows on their consequent inability to expel the vesical contents. The condition resulting resembles much the retention produced by an enlarged prostate, and requires frequent relief by the

catheter in the same way, at least for a time. Here the absence of physical signs, the suddenness of the attack, its connection with a cause generally recognised by the patient, and the diminished power of discharging the urine *after a catheter has been placed in the bladder*, especially when he is recumbent, are sufficient to distinguish this affection. Particular attention should be paid to the last-named point. In enlarged prostate, the urine often flows with considerable force when the influence of the obstruction is removed by the introduction of a catheter, and the current can be accelerated materially by the will of the patient, unless there be atony also, as there may be from undue distension; the atony, however, is not generally considerable, except in long-neglected cases. But when the cause of retained urine is not obstruction, but atony of the bladder, the urine barely flows from the catheter, and is not propelled, nor is it so much influenced by any efforts of the patient.

Lastly, there is paralysis of the bladder; a condition in which its nervous supply is either impaired or destroyed. It is almost always associated with a similar condition of the lower extremities, and this may result either from disease or injury of the encephalon or spinal cord. There can be, therefore, little doubt respecting its presence; and when it does exist, the indication which catheterism presents is characteristic. An instrument being introduced, the urine is propelled by the weight of the parts around, the will of the patient exerting no influence upon its flow unless the abdominal muscles should be in a normal condition, as in cases of injury (rare) occurring to the spinal cord between the sources of nervous supply to those muscles and to the bladder, in which case a slight influence is perceptible. Otherwise no impulse is noticeable, except through the agency of acts unassociated with micturition; such as deep inspiration, coughing, sneezing, and the like, by which a momentary pressure is communicated to the paralysed bladder, and the stream is temporarily accelerated.

Treatment may be regarded under three heads.

1. Treatment for the purpose of obviating the results of obstruction caused by hypertrophied prostate.

The fact being discovered that a certain quantity of urine is habitually retained in the bladder, however frequently or forcibly the efforts to evacuate it have been made, the first thing is to ensure the complete removal of the urine from the viscus at least once a day. It may be very desirable to do this twice, or

three or more times, daily, the necessity depending upon degree of obstruction, and the consequent amount of residual


And if the power of urinating is almost or quite lost, it is necessary to employ the catheter as often as a decided desire to micturate is felt. There are certain modifying circumstances which must be taken into account; such as the facilities which exist for passing instruments, and the condition of the urethra itself. If the patient possesses the ability to pass the catheter easily for himself,—and it is very rare that he cannot do so by tuition and practice,—he complies with the demands of the case. He should be made to understand that, having his sensations to guide him, he may soon attain considerable proficiency in the management of an instrument in his own ure-

But if this be in an extremely irritable state, which is aggravated by the frequent use of the catheter, however carefully it may be introduced, it is necessary to consider carefully the relative necessities of the urethra and the bladder, and some compromise must be made between them. In most cases, the removal of the urine night and morning suffices to maintain the reservoir in a tolerably sound and healthy condition, and it is extremely undesirable to resort to artificial aid with greater frequency than is absolutely necessary to accomplish this. For the use of the patient himself, and generally for the surgeon, the instruments are, in the majority of cases, the better and simpler kind. When silver instruments are necessary, the curves require to be larger, and more strongly marked than the ordinarily employed. Three good forms are shown in the accompanying woodcut. The fourth catheter is a flexible one, mounted on a well-curved stylet to preserve it for use in its original form.

When the necessity for a catheter is very frequent, some have advocated the plan of permitting a catheter to remain in the bladder for days together. As a rule, this is not desirable, for irritation and even ulceration may be soon produced by the instrument. Where, however, great frequency in making void exists, and ease and sleep are found to be promoted by the catheter, it should be adopted for twelve hours or so at a time; as during the night, for example. In this manner it may be extremely useful. But in circumstances of urinary retention, when the patient has been relieved with difficulty by the catheter, we may be justified in permitting the catheter to remain in the canal for a considerable period. The vulcanised india-rubber catheter

has for some years been recognised in this country as the easiest for the patient, and the least liable to be encrusted with phosphates when tied in for these cases. It may be introduced without a stylet sometimes, but owing to its extreme flexibility often requires to be mounted on a well-curved one, which is withdrawn and the tube left in place. Usually it is easy to retain it there by passing a piece of soft string or bobbin

FIG. 280.



the penis behind the glans, attached to the neck of catheter. If there is a tendency to slip out, it is usually overcome by applying a piece of soft adhesive plaster round whole of the penis, and so covering the retaining string. Holt appears to have experienced some difficulty in keeping the vulcanised catheter in its place, and has therefore suggested the attachment of 'wings' to its inner end to prevent ejection by the bladder. The circumstances are not frequent in which the vulcanised instrument is ever necessary still

ment is it to experience much difficulty in retaining it, in such circumstances, if they arise, the winged catheter may always be tried with advantage.*

The consequences of enlarged prostate already alluded to, the increasing retention of urine and habitual distension of the bladder, which accrue from not completely emptying it, form only a portion, although a very important one, of the evils which can be obviated by the habitual use of the catheter. It is impossible to overrate the benefits arising to the patient from its use; and the responsibility which is incurred by overlooking it should be ever present to the physician's mind in dealing with any signs of irritability of the bladder, or incompetence to retain perfectly the urine by patients in advanced years.

In relation to the treatment of chronic cystitis, catarrh, &c., bearing as they so frequently do in connection with hypertrophied prostate, see the sections thus headed, pp. 889, 890.

The general treatment of enlarged prostate.

It is of great importance to maintain all the functions of the system in healthy action, in treating patients who thus suffer. Any derangements in other parts of the system are very liable to augment the urinary symptoms. A simple catarrh, a cold, indigestion, or unrelieved constipation, are very apt to increase increased obstruction or greater irritability of the bladder.

Anything that tends to derange the stomach and bowels, to tax unduly the digestive powers, or to over-excite the circulation, should be avoided.

The clothing and the habits should be such as encourage and maintain a due action of the skin. Damp must be sedulously avoided, or removed after exposure, especially from the feet, without delay: the lower limbs should be kept habitually dry and warm; a habit of the first importance, as freedom of circulation and healthy vascular action here is one safeguard against the recurrence of congestion in the organ.

The question of exercise is one of importance. The subject of enlarged prostate must not be encouraged to believe himself much an invalid, but must exert his physical powers, as far as they exist, in daily exercise in the open air, of which walking

* *Lancet*, May 14, 1870.

3. Special treatment of the hyper

Although many agents have been proposed, all medical treatment has been of little efficiency, as to any power exhibited to reduce the size of the enlarged organ. The hypochlorate of ammonia have all been tried, but no utility has been claimed for them. It is said that there exists very little ground for the efficacy against senile hypertrophy. The time seemed to promise more success. It was employed by Mr. Stafford, who is said to have of greater value than others have said. His plan consisted in administering iodine to the rectum, occasionally by the mouth, and the urethral surface of the prostate with an ointment; commencing with one grain of iodine to the drachm of simple cerate, and increasing to twenty grains to the drachm, sometimes to a large quantity of iodine.

Regarding this as one of the most promising methods for the removal of glandular enlargement, to infer that the hypertrophied prostate is probably acted on either by iodine or by cerate, over simple enlargements of the ut

ances for effecting this purpose. Dilatation by water in leading india-rubber tubes has been employed by myself. I do not say, however, that any benefit is to be anticipated by any of these means.

Division of the obstructing portion at the neck of the bladder has been performed. Other operations have been also attempted for effecting a similar purpose, such as the excision or crushing of a protruding portion; and even the ligaturing of a polypoid outgrowth. Respecting the division of an obstruction, bar-like in its form, elevated from the posterior surface of the neck of the bladder, it is no doubt a proceeding that may be accomplished without much difficulty, with the exercise of ordinary care. In most cases, although not invariably, the obstruction is a prostatic development, and when well marked may be incised with advantage, and without serious injury to the patient. Such was the opinion of the late Mr. John Ericson. I cannot say, however, that I have yet seen a case in which it has appeared at all advisable to practise such an operation. So much can be done by management, by maintaining the bladder in a healthy condition by means of the catheter, that the case must be rare indeed in which such an operative procedure is indicated.

In estimating these proposals, most English surgeons will be content with awaiting further experiences in the hands of those who have hitherto seen fit to adopt them. We can cherish the hope that any benefit will be conferred on the patient by the methods of accomplishing the ends proposed, even grant that no doubt existed as to the possibility of carrying them into execution.

Atrophy of the prostate.—The prostate is sometimes atrophied with age, and occasionally it is so under certain circumstances of early adult life. It is but slightly developed in some malformations, and in constitutions in which the male sexual character is not strongly marked.

The normal weight being about four and a half drachms, it is occasionally found as low as two and a quarter, and not infrequently at three drachms, at ages between 60 and 80 years.

There are no special symptoms of atrophied prostate, nor is any special treatment indicated for patients who are the subjects

of some malignancy, the cancer is overlooked.

Judging from the data at present appears to be less commonly the seat of deposit; the latter only will be to the contrary is indicated.

Malignant disease of the prostate is rare. After a close examination of the prostate, it is difficult to adhere to the opinion expressed by the microscope, resulting at that time from an examination of the prostate, viz. that 'the evidence now possess, viz. that 'the evidence of true scirrhus of the prostate is defective.

Melanotic deposit is said to be associated with encephaloid of the prostate. In two cases; one at adult age, the other at childhood, it is to be forgotten that interstitially-developed growth may be mistaken for true cancer. In the naked eye it sometimes resembles a melanotic deposit.

Malignant disease has at present been recorded in childhood and at advancing age. No record between the ages of eight and fifteen of the disease, from the first appearance of the disease, appears to vary from one to three in adults, and from three to nine months in children.

The symptoms of the malignant disease of the prostate are as follows:

time or another, sometimes to an alarming extent. The urine is usually voided almost pure or unmixed, and frequently some attempt to urinate, which has been attended with greater exertion than usual. Less commonly is the hæmorrhage continuous for some time, as happens in villous tumour, for example.

The enlargement formed by the prostate itself, when examined by the rectum, is hard at first, and may or may not be irregular in outline or consistence. Softening may in the later stages be present, but the patient's powers do not always sustain him to so long a period as that in which the growth either softens or ulcerates. Consequently, on examination after death, the prostate may be simply enlarged; or there may be breach of surface and protrusion of soft granulations; or there may be a cavity of substance and a cavity, the last-named circumstance being to be rare. Frequently other organs are affected, but by no means invariably. But there are always diseased lymphatic glands adjacent, and sometimes the infection reaches to distant groups. The existence of such swellings in the course of the iliac vessels, and sometimes in the inguinal region, may frequently be verified by examination of the abdomen, and constitutes a valuable sign in relation to diagnosis.

The urine may be closely examined, in cases of a doubtful nature, for the presence of cells which may be regarded as malignant from inspection of their forms and constitution. A good deal of debris is seen in advanced cases, its presence indicating that the growth has fungated, and throws off portions of its tissue; and these will sometimes exhibit all the characters of a malignant growth elsewhere. It is not to be forgotten, however, that the urinary passages yield epithelium cells of all forms and sizes abundantly, and these, I suspect, are mistaken sometimes for the so-called 'cancer-cell.'

Regarding the general treatment of malignant disease of the prostate, nothing more can be suggested than applies to the complaint when occurring in any other part of the body. The treatment is palliative, and must be regulated according to the necessities which arise in the progress of the case.

Thus, accumulation of urine must be provided against at the smallest possible risk of irritating, much less of injuring, the rectum. If catheterism can be dispensed with altogether, so much the better. In no circumstances is it of more importance to be extremely gentle in the manipulation of instruments. The pain

must be relieved by anodynes administered both by mouth and rectum. The addition of conium to opium, by enema or suppository, is often useful; and by the mouth belladonna is sometimes a valuable auxiliary in mitigating pain, given in doses of from one-fourth to three-fourths of a grain twice or three times a day. Hæmorrhage must be treated on principles already illustrated (p. 878). The powers of life are to be supported by every means in our power. Nutritious food, both in the solid and fluid form, with a due proportion of alcoholic stimulant, must be supplied in accordance with the wants of the patient.

Tubercle of the prostate.—The prostate is very rarely the seat of tubercular deposit, and when it is so, appears generally to be somewhat increased in size, until the later stages of the complaint are reached, when, after suppuration and discharge, its volume may become smaller than natural.

At no period of the disease is the prostate affected alone, some other part of the genito-urinary tract being the primary seat of the affection. In most cases the deposit takes place first in the kidney; the organ next commonly affected, among the genito-urinary group, is the testicle. In fourteen cases collected by myself, tuberculosis of the kidney occurred in eleven, and of the testicle in six: in seven of these cases the lungs are stated to have been diseased, they were probably so in nearly all. It is difficult to isolate any special symptoms indicative of this affection. Undue frequency and pain in making water, occasionally blood in the urine, and at times signs of cystitis, are commonly experienced. The presence of pus in the urine, of occasional hæmaturia, of pains in the loins, perineum, and penis, give rise to suspicions of calculus, to be resolved sometimes only by a careful search; no foreign body being found, the nutrition of the patient, his history, and the condition of the lungs and other viscera, will probably lead to a correct diagnosis.

Nothing need be said of the constitutional treatment of tubercular disease, and little in relation to the local manifestation in the prostate. Mechanical interference is to be avoided, and every kind of irritating application. If suppuration takes the form of external abscess, it must be treated as other perineal or ischio-rectal abscesses. But more commonly the discharge of purulent and tubercular matter takes place into the urethra. The improvement of the health, by all those numerous

means which regulation of the diet, regimen, exercise, climate, medicine enable us commonly to achieve in tubercular affections, constitutes almost the whole of the treatment to be employed in the affection, when involving the urinary or genital organs. The diagnosis once established, it is of great importance that the patient should be kept free from all instrumental treatment, which, in such cases, provokes irritation, and aggravates the disease, without conferring upon him any benefit whatever.

Cysts of the prostate.—It is not at all uncommon, in making sections of an enlarged prostate, to find cavities, of an irregular shape, in its substance, not met with in the normal organ. These cavities have all the appearance of being dilated follicles of the glandular structure. Ducts are easily traced into them; and frequently numerous little dark concretions lie free within. I have seen from thirty to fifty of these minute bodies occupying a cavity about the size of a grain of wheat or of a small pea. But larger concretions, that is, of the size of pearl-barley, or prostatic calculi, may occupy each a separate recess of its cavity; and on removing the foreign body, a spherical, thin, and double-walled cavity is displayed. Sometimes hundreds of such cavities may be found in one prostate; but this is a very rare circumstance.

The formation of these cavities, or cysts as they have been called, depends a good deal on the prior formation of concretions. In all events, the two occurrences are closely associated. We know too little of either the one or the other to affirm anything confidently respecting the precise mode of their formation. But probably the cavities are nothing more than enlarged follicles, dilated caecal terminations of the glandular tubes. There are no isolated cysts in the prostate filled with fluids, giving no communication with the secreting structures around, as in the kidney; no formation, indeed, which can be regarded as analogous to that which may be considered as the type of simple fluid cysts. Although, in conformity with the practice of other authors, I have referred to 'cystic disease' of the prostate, the use of the term is scarcely warranted by the phenomena presented; and if retained, it must be held to signify a formation of a wholly different kind from that which is indicated by the same term in the breast or kidney.

The cavities referred to do not attain a sufficient size, nor, as

far as we know, do they give rise to any symptom whatever to render a knowledge of their presence possible during life. Generally speaking, they are capable of holding not more than a few minims of fluid. In relation to practice, the diagnosis is unimportant, as no indication for treatment would be presented by the fact of their existence, were it ascertained.

The prostate is, after long-continued suppuration, sometimes converted into a kind of cyst or membranous bag; this condition can in no respect be regarded as a form of cystic disease. The organ has, in fact, disappeared, and its capsule forms part of the sac of an abscess which has replaced the normal structure.

It is doubtful if hydatid cysts have ever been met with in the prostate. Cases are on record in which retention of urine and distension of the bladder occurred as a result of a hydatid cyst *between the bladder and rectum*, near to the neck of the former; but in which the prostate was not affected except by pressure. Prostatic enlargement was very closely simulated certainly in some of them, and in two the prostatic catheter was employed under the belief of its existence. Among these cases, one or two have at times been regarded as offering examples of hydatids formed in the prostate itself. But I think evidence is wanting to show that they were so.

THE MALE URETHRA.

Congenital malformations.

Injuries.

Inflammation.

<i>Stricture of the urethra</i>	{	organic.
		spasmodic.
		inflammatory.

Tumours of the urethra.

Urinary abscess, acute and chronic.

Urinary fistulæ.

Retention of urine.

Extravasation of urine.

Rupture of bladder (from retention of urine).

Congenital aberrations and malformations.—1. Absence of the urethra is occasionally met with, as in cases of extroversion of the bladder, in which there is neither any cavity nor reservoir

to contain the urine, nor any canal to carry it off; but a mucous membrane, corresponding with the posterior surface of the bladder and with the floor of the urethra, alone remains. For further description, see '*Malformations of the Bladder*,' p. 881.

2. The canal is sometimes occluded, and this produces retention of urine, and death during the early hours of life. The obstruction may consist merely of a membrane forming a diaphragm across the canal, or of an obliteration of some lines in length, and it may occupy any portion of the tract, but is usually found near the vesical orifice.

3. Deficiency of a portion of the urethra near the anterior orifice is the most common deformity. When a part of the upper covering is deficient, and the mucous membrane of the lower part, or floor, is exposed, the condition is termed *epispadias*; when the contrary condition is presented, the deficiency being that of the floor, the term *hypospadias* is applied to denote it. The degree of deficiency varies considerably; from a quarter of an inch to an inch and a half may be absent in either case. Occasionally, and this is only a variety of hypospadias, a second meatus exists, usually about an inch behind the normal one, opening externally through the floor of the urethra. But the opening may be much further back, and may even render impracticable the ejection of semen into the vagina.

4. Besides these, slight exaggerations in the size or form of the natural parts may be occasionally noted; as of the lacuna magna, the sinus of the bulb, and the sinus of the verumontanum. Not very unfrequently there is congenital narrowing of the external meatus, or of some portion of the passage situated within an inch of it.

Treatment.—The simple diaphragm may be perforated by a trocar, or point of a bistoury, when its existence has been clearly ascertained. The congenital narrowing of the external meatus may be dealt with as stricture in that situation. None of the other conditions, with one exception, are improved by the various procedures, by paring and suture, which have from time to time been practised, nor are they sufficiently important to be so treated. But hypospadias existing or extending far back, so as to produce a practical impotence, should not, under some circumstances, be refused the possibility of remedy by surgical operation. This has been accomplished by a combination, of perforating the anterior portion of the penis, paring the edges

of the opened urethra, and uniting by suture. Much care and skill, however, are necessary to give the patient any chance of a cure.

Injuries of the urethra.—(See vol. ii. pp. 726 et seq.)

Inflammation of the urethra.—(See the essay on GONORRHEA.)

Stricture of the urethra.—The term stricture implies an unnatural contraction of the urethral canal, congenital or acquired. Contraction may occur in two forms: it is the nature of the first to be permanent, and of the second to be transitory only, as regards its duration. A permanent contraction is due to organic deposit in or around the walls of the urethra, and is accordingly termed organic or permanent stricture. A transitory contraction is supposed to be due either to local vascular inflammation or congestion, causing temporary narrowing of some part of the urethra, hence the term inflammatory or congestive stricture; or to unwonted muscular action alone, of the voluntary or of the involuntary fibres, in which case it is designated spasmodic stricture. This last-named condition, that of spasm, may exist alone, but is usually found as a complication of the other kinds. The term spasmodic is understood to include only cases in which involuntary contraction of the muscular fibres constitutes the main source of obstruction. So also the term inflammatory stricture can only be employed when the diminished calibre of the urethra is mainly due to an attack of inflammation.

But the terms implying temporary stricture are liable to cause confusion and misapprehension. Obstructed micturition from inflammation has its origin in the prostate when swollen through an attack of that nature, and is improperly described as *stricture*; just as we have tonsillitis obstructing deglutition, but which is never spoken of as stricture of the throat. Obstructed micturition from mere spasm of the muscular parietes of the urethra must be excessively rare, although, no doubt, spasm complicates previously existing morbid changes there. Hence I prefer to restrict the term stricture of the urethra to those narrowed conditions of the canal which depend on organic changes in, and immediately surrounding, its walls; in other words, I shall use the word stricture as synonymous with the

is organic or permanent stricture, and as implying nothing else.

Anatomical classification.—Strictures present themselves in a variety of forms. For anatomical purposes all these may be arranged in four groups, by which means it is easy to designate a specimen, according to the physical characters of the stricture itself. It will then be only necessary, for the sake of further description, to speak of it as more or less narrow, and name the region in which it is situated.

1. *Linear stricture.*—The urethral canal may be obstructed by a thin membrane stretched across it, with an aperture in the centre, or on either side of it, and having an appearance, in relation to the rest of the passage, somewhat resembling that of the pyloric orifice of the stomach. It occasionally happens that a fold of the mucous membrane obstructs the passage at one of its sides only, forming a crescentic septum, and so constituting a segment of the calibre of the canal: these constitute what has been called ‘the bridle stricture.’

2. *Annular stricture.*—Those instances in which the contracted part is thicker and broader than the foregoing description would indicate, have been termed annular strictures, which present an appearance as if a piece of cord had been tied round the canal at one point, leaving the remainder free.

3. *Indurated annular stricture.*—This term may apply to those cases of confirmed stricture in which the induration is seen to extend into the tissues around the urethra, to the depth of half a line or a line: although it may be limited in extent from before backwards to a space occupying less than half an inch of the canal. The centre of the space is the point in which the contraction is most considerable, so that the affected portion presents a form somewhat resembling that of an hour-glass; and it is worthy of remark that the induration is generally thicker at the floor than the upper aspect of the urethra.

4. *Irregular or tortuous strictures.*—In a few instances, some of the natural rugæ of the urethra seem to be adherent, or even fused together, for the space of a few lines in length. In very rare cases a patch of indurated tissue is seen, resembling a matrix, around which the mucous membrane is puckered in radiating lines; the amount of contraction appearing to correspond with the extent of previous loss of substance. But

occasionally the contraction extends longitudinally for a considerable distance, and the canal is narrowed, and its thickened on all sides, for a length of one or more inches. In these cases the induration sometimes involves the entire substance of the corpus spongiosum, and gives rise to the obstinate and undilatable form of stricture.


Occasionally several separate strictures may be observed in the same subject. John Hunter records an instance in which he met with six strictures in one urethra. Lallemand and

FIG. 290.



Long and indurated stricture at the junction of the bulbous with the membranous urethra. Much dilatation of the urethra and its lacunæ behind.

FIG. 291.



An irregularly narrowed constricting the anterior part of the urethra, the narrowest part rather more than two inches from the orifice [drawing half nature]. Dilatation of the urethra and of lacunæ behind.

French writers describe seven or eight. I have never been able to find any such examples. Three, or at the most four distinct contractions is the highest number I have seen. The latter is rare.

Origin, nature, and site of stricture.—The first effect of inflammation upon the mucous membrane is a swelling or thickening of it, caused by engorgement of the vessels. Then exudation of an albuminous fluid takes place into the submucous tissue, which perhaps becomes absorbed under favourable circumstances. When the morbid action persists, more or less plastic mat-

is thrown out, the result of which is, the formation of fibrous tissue around the canal, causing adhesion between the mucous membrane and the submucous tissue, infiltrating the meshes of the latter, and even involving the substance of the corpus spongiosum itself.

A widely-differing condition to any of those above described has been referred to by some under the title of stricture. Sometimes, but by no means frequently, an exudation-deposit upon the surface of the urethral mucous membrane is said to cause occlusion of the canal. Few cases are on record, nor are many examples to be found in our museums; and it is now quite certain that this condition must be regarded as extremely rare.

Speaking in general terms, the degree of contraction is proportioned to the duration of the complaint, and to the extent of the inflammatory action which has existed; although the severity of the symptoms, the amount of distress, and the effects on the constitution, are by no means always commensurate with the amount of narrowing which exists. It is very rare indeed to find the urethra altogether impervious during life to the flow of urine. However contracted the canal may be, the urine still issues either in a very small stream or by drops. Retention does not depend on absolute organic impermeability. It is easy to conceive that when the canal is contracted to a mere pin-hole, the slightest cause may operate to occasion total obstruction; a little tumefaction of the part, a pellet of thick mucus, a flake of fibrinous deposit, or a very small calculus, are quite sufficient to block up the channel. Probably the sides of the urethra never adhere, and cause obliteration of the canal; unless indeed fistulæ have been established, when, although very rarely, this accident may happen in that part of the canal which is anterior to the stricture. But even then it occurs almost invariably in those strictures which are of traumatic origin. The urethra may be cut across by a wound in the perineum, and for want of proper attention the urine may pass entirely through the artificial opening, and adhesion seal up the proper passage. But such obliteration is a wholly different thing from stricture, and ought not to be confounded with it.

Locality of stricture.—After a laborious investigation of this subject, respecting which there has long been much difference of opinion, comprising the examination of 270 preparations in


body, which can by no means be admitted as evidence on subject.

Causes of organic stricture.—There are two principal and distinct species of lesion which give rise to the formation of stricture of the urethra. The first, and that which by far the most frequently operates, is inflammation in the canal; the second is injury by violence.

The first-named process, viz. the inflammatory, in the great majority of cases, fortunately terminates by resolution, and leaves the urethra as healthy as before. Whether the urethritis be gonorrhoeal, or of non-sexual origin, that is the ordinary result. In exceptional cases, occurring in some instances probably from indiscretion on the part of the patient, in others from violent treatment, and in others from a constitutional indisposition to maintain inflammatory action, this latter persists and produces the deposit which ends in stricture as before described. The last-named cause is not a speculative theory, for I am now fully satisfied that in certain families individuals inherit a propensity to the formation of stricture. It is not therefore acute gonorrhoea, but a persisting chronic or subacute affection, which seems to lay the foundation of organic stricture. Hence a long period often elapses between the acute attack and the appearance of symptoms of obstruction. These latter appear often very insidiously and slowly, and at first perhaps scarcely excite the attention of the patient. Thus it is that a period of many years is often passed before the existence of the organic change has become obvious.

The next principal cause of stricture is mechanical injury. Wounds on the perineum, bruising, lacerating, or partially destroying the urethra, are received in a variety of ways: by contact with some hard object, as across spars, scaffolding, ladders, gates, wheels, saddles, &c.; or on some sharp object which punctures the perineum, as palisading and other fences, earthenware vessels which break under the sitter, &c.; or, a laceration may occur from the bones in pelvic fracture; or, all violence in the use of surgical instruments must be admitted as another traumatic cause. In all these cases the injured urethra often unites irregularly, or loss of substance takes place, and the cicatrix resulting permanently contracts the part. The strictures so formed are among the most obstinate and difficult to treat which come under our care.

An occasional cause of stricture is cicatrization following



up to the period of adult life without convenience, indeed without being become the source of severe symptoms become in later life less extensive obstruction to the outflow of urine.


Symptoms.—Among the early symptoms charge is often noted, and its precise true cause. Pain is usually felt in stricture, at the time of micturition the bladder are emptied at shorter intervals. The stream is altered in form, becomes perhaps twisted, spirting, forked, and conditions are caused by the current of size and force to dilate and extend the externus; but it must not be existence of such a stream is a proof to many persons, from a tumid condition habitually pass such a one. As the stream grows smaller, the moment and in time the urine may issue. In speaking, the act of micturition to extent corresponding with the degree. Ultimately the acts of micturition are painful, some patients being compe

the perineum, the testicles, the loins; and the hypogastric are usually present in such cases.

In some few cases the most prominent symptom throughout retention. There may be but little irritability of bladder, the stream of urine, when passed, is not necessarily very feeble; a No. 6 or 7 catheter may pass through the urethra; but if a true organic stricture is present, and may perhaps be felt by the touch, as a ring of indurated material in the middle of the urethra. The patient is liable to frequent retention, and finds no relief but from catheterism, while attempts at micturition are followed by inability, more or less prolonged, to pass water, until an instrument of very large size has been introduced.

The urine also shows signs of change, which become more marked in proportion as the case advances. A portion being retained in the bladder, from inability to overcome the obstruction, the secretion becomes partially decomposed, and irritates the mucous membrane; and thus urine, cloudy, ammoniacal, depositing as it cools a quantity of pus and mucus, is not an infrequent accompaniment of stricture. It is usually turbid, deposits slimy tenacious mucus, and also a precipitate of prismatic crystals of the triple phosphate of ammonia and magnesia, of exudation or compound granular corpuscles, and sometimes pus; while on its surface an iridescent pellicle collects, commonly consisting of the triple phosphate, and sometimes of the phosphate of lime. Hæmaturia is also one of the occasional concomitants of stricture. Blood sometimes comes from the bladder, in small quantity, communicating a dark tint to the urine. It frequently follows the use of the catheter, but it appears also when no instrumental interference has taken place; or it may be due to the rupture of blood vessels during erection, the urethra being unduly confined by the presence of the stricture, and strained by the act (see hæmaturia, p. 876).

As the case advances, attacks of complete retention, depending on the causes above mentioned, become more frequent. At length the urine is at length habitually discharged by drops, so that a free stream cannot be said to exist. Sometimes the urine passes off involuntarily during the unconsciousness of sleep, or during muscular exertions; and at length the patient loses the power to retain it altogether. At this stage 'incontinence' is said to occur, which term does not here apply either to



organ remaining filled with the
unless it be frequently emptied b
state of retention, or of engorgemen
The extent of dulness on percussio
this condition, and also the size o
distended viscus. Under these c
bladder is increased, and disorgani
than before. Thus ulceration ma
brane, or in that of the urethra b
such circumstances, in a paroxysm
rupture of the urethra may take
urine occur into the cellular in
scrotum, and supra-pubic region (s

Besides the local symptoms of s
order of the digestive organs, follo
impaired nutritive function. Th
strength, looks anxious and care
the back and loins, and often bec
attacks of shivering. Others invar
the passage of a bougie, or if an i
larger than the accustomed size be
appears to be more common am
inhabited warm climates. The a
corrosive substance to the urethr

otherwise, to experience them, and who have suffered for a time from stricture. I have observed, on more than one occasion, suppression of urine, rapidly followed by death, to result from the introduction of an instrument larger than the patient has been accustomed to; or again, when the ordinary instrument has been less skilfully employed, and an abrasion, though only an exceedingly slight one, has been made in the mucous membrane of the urethra. The rapidity with which death may occur, under these circumstances, in patients who are the subjects of extensive chronic disease of the kidneys, and in whom an apparently exceedingly trifling lesion so caused, is remarkable. It seems to be due to uræmic poisoning; the post-mortem appearances, to the naked eye, do not necessarily exhibit traces of inflammation resulting from the particular lesion. In these cases it appears that the function which determines the elimination of urea suddenly ceases after slight injury to the urethra, as by the propagation of some shock to the excreting organ, in cases where the kidneys are largely diseased. And there are undoubtedly sometimes, although rarely, instances in which the fatal result ensues in this manner from the mere passing of a catheter, although it is absolutely certain that no mechanical injury whatever has been inflicted.


From the symptoms which present themselves three classes of uræmic stricture may be made, which will embrace all the varieties of the disease, and so serve as a means of describing in brief terms any particular example in the living patient.

1. *Simple stricture.*—Its chief sign is diminution in the size of the stream; there is increased frequency of micturition also, and some pain, although the amount of either varies greatly in different cases.

2. *Sensitive or irritable stricture.*—Proneness to disturbance of the nervous system, as evidenced by chilliness, irregular circulation, or even rigors on very slight irritation. Great pain is caused even by the gentle application of instruments, and it continues sometimes long afterwards. In a few cases also a disposition to hæmorrhage is manifested.

3. *Contractile or recurring stricture.*—There is constant tendency to become narrower in the absence of treatment; and contraction rapidly recurs after dilatation has been applied.


Diagnosis and treatment. Dilatation.—The symptoms just detailed are not sufficient alone to establish the presence of



the inflexible, and I advise them. They are of two kinds, each made of gum-catheter has a considerable but is very flexible if placed in water and retain when plunged into cold water when cated when warm. This quality in the treatment of enlarged prostate is required. Ordinarily 1 stylets, which maintain the desire is to be withdrawn from the catheter of the same material are susceptible. Formerly they were made of wax but have fallen into disuse, being replaced by


The French instruments are more than the English, and may be made not taking any other form than extreme flexibility enables them without pain or even discomfort even when solid instruments pass

FIG. 2



re of the instrument (see Fig. 292). The long and extremely flexible extremity readily insinuates itself through the passage; were the instrument to end in a fine point, this might easily be caught in some lacuna, but the terminal bulb prevents this, and the merest tyro can generally pass it at once into the bladder. In doing so, if the patient stands, the bougie, if oiled, is merely introduced and pushed horizontally forwards, without any attempt to direct its progress. The bulbous bougies and catheters, often but less appropriately termed 'syring bougies,' are often of extreme utility both to the surgeon and to the patient who requires them for his own use, but they are quite unsuitable to hot climates, where for the most part metallic instruments are necessary.

To introduce the silver catheter, and supposing the patient to be in the standing position, the following plan may be adopted. The handle or upper end of the instrument be lightly held between the thumb and the fore and middle fingers of the left hand, the concavity of the curve looking towards the left side of the patient, and the general direction of the instrument almost horizontal. The penis is to be raised with the right hand, the point of the instrument inserted into the urethra, and slowly carried onwards until four or five inches have disappeared; the handle is brought to the middle line at the same time, at first close to the patient's abdomen, after which it is to be gently and lightly depressed; and as the point is felt to surmount the sub-pubic curve, the handle is gradually brought forward towards the operator, until it sinks beneath the horizontal line, when the opposite extremity will be free in the bladder. If difficulty occurs, the direction of the instrument may be slightly altered, or it may be withdrawn an inch or so, and then re-introduced with the handle more upright than before, or the reverse, as the case may require. Notwithstanding all, the instrument is stopped point blank, at the same spot on each trial, there is reason to suppose a stricture is encountered. On the other hand, if the urethra is healthy, a solid sound of good size will pass almost of its own weight if lightly handled; at all events, a very slight pressure from the fore-finger upon its handle will be fully sufficient, if additional impetus be required. The main object is to allow the point to slide easily along the upper surface of the urethra. It will not then be caught in the depression of the bulb on the floor of the urethra, nor will it be so likely



course.) It is never to be forgotten that the handle is depressed on structures which may be injured by force being applied. Whatever the operator is forced to: temper, patience, and a light touch are all cases of difficulty. All attempts are wholly out of place, fraught only with pain and calculated to reflect discredit on the operator.

That mode of passing a catheter known as the 'tour de maître,' can be only attempted by a corpulent patient. It consists in placing the convexity of its curve upwards, with the concavity of its curve up in a perpendicular line beneath; the operator sweeps the perineal fascia in this direction, at the same time, in sweeping it round around the point, of which its point is the handle, gradually depressing the handle to the sub-pubic curve.

When much time is required, the patient is generally best. The head and shoulders are elevated by pillows, and the knees are drawn up from each other; the operator should sit on the couch, hold the catheter as be-

inferior wall of the canal, to avoid a lacuna on the roof, sometimes considerably developed; an accident which gives the patient pain, and sometimes disconcerts an operator. The fingers of the left hand gently draw the penis over the instrument as it glides easily on to the bulbous part, the handle still being horizontal, or nearly so; arrived at which, if some obstruction seems to offer, the instrument should be withdrawn an inch or so, and again passed, taking care not to elevate the handle so soon; after which, by gently raising it and using it to describe a curve along the middle line, the extremity will probably glide slowly upwards into the bladder as the handle sinks towards the interval between the patient's thighs. In passing an English gum-catheter, take care that it is well curved after the stylet is removed, as the warmth of the urethra tends to straighten the instrument, and so occasion it to catch on the floor on arriving at the membranous portion. The point should be well turned up, and the curve maintained in passing the first half of the catheter very close to the belly of the patient, after which a rather quick depression of the shaft carries the point easily into the bladder.

In exploring the urethra, especially if the symptoms are not well marked, we are not hastily to conclude, because a little obstruction presents itself in the passage, that stricture is of necessity present. The part is extremely sensitive, and resists only but gentle efforts to traverse it, the more so if it be the first time an instrument has been introduced. Sometimes

the neck of the bladder itself a little more than usual depression of the handle of the catheter is required, or it may be necessary to employ an instrument with a stronger and longer curve.

The size of an instrument to be first used is by no means a matter of indifference. As a rule, whatever statements the patient may make, we should *always* use a bougie or catheter, not smaller than No. 7 or 8, with a blunt, not conical or bulbous extremity, as the former affords a far more certain indication

of the situation of the obstruction than a small instrument, which might pass through a slight constriction altogether; or whose point may be entangled in a lacuna, or in a fold of mucous membrane, the first of which would have been closed, and the second obliterated, by the passage of an instrument sufficiently large to fill the urethra. By such entanglement, perhaps, injury may be inflicted on the canal, or the unfounded

belief in the existence of abnormal obstruction in it may be induced.

Suppose, however, that a stricture is encountered, its distance from the external meatus should be accurately noted and its situation remembered. Then a smaller instrument is to be passed, in order to determine the calibre of the stricture; if this enter the contracted part, it is at once obvious by its being 'held,' i.e. it has entered a narrower passage which fits closely to and retains it, so that on attempting to withdraw it, some force is necessary for the purpose. A trial of several instruments may be necessary, until at last one which is sufficiently small to enter the constriction is arrived at: but we should always bear in mind that the smaller the instrument the more careful and the more sparing of pressure must the operator be, since such an instrument will inflict a wound with ease directly proportioned to the smallness of the point.


The method just recommended I have found generally sufficient to determine the existence of stricture under ordinary circumstances. When it is desirable to have an exact knowledge of the length to which the canal is affected by narrowing, a bulbous-ended instrument, of which the shaft is slender, may be employed for the purpose. It is rarely necessary to employ them, although before performing urethrotomy it is as well to do so. For this purpose instruments must be used with bulbs of various sizes, from about No. 2 to No. 7, the stems of the smaller to be equal to that of a No. 1 catheter, of the larger about No. 2 or 3. One of these is found, the bulb of which will pass with slight pressure through the stricture, when it is felt to be free and easily movable beyond. On withdrawing, as it passes outwards through the constriction a little obstruction is encountered. In this manner, the extent of contracted urethra, generally comprised within a range of the fourth or half of an inch, is readily estimated. So also the existence of another stricture beyond the first may be demonstrated in the same way.

Of late years attempts have been made to examine the urethra and bladder, not merely through the sensations obtained by touch from an instrument, but by vision also. For this purpose a straight metal tube (the 'Endoscope,') which may vary in size between 9 or 12 of the catheter scale, is used for the urethra, and one in the form of a beaked sound with a piece of glass at the angle for the bladder. To the tube when introduced a lamp is adjusted, with

piece and reflector, so as to throw a pencil of powerful rays on every substance impinges directly on the bottom of the light tube. Whatever this is, being comprised within a circle about the $\frac{1}{4}$ or the $\frac{1}{2}$ of an inch in diameter, is visible to the observer looking through the eyepiece and the perforated reflector. The same tube may be employed also with an ordinary gas, or with sunlight. I have tested various Endoscopes on numerous occasions during many years past, and I am compelled to say that I do not find any advantage to be derived from their use in the treatment of stricture, or of any disease of the bladder. It is easy to see the colour of the mucous membrane of the urethra and of the bladder touched by the tube; to see that very small portion of a calculus against which the end of the tube rests. No illumination of the cavity of the bladder can be made; this being filled with urine, mostly in the more or less cloudy and opaque, less is visible there than in the urethra, and in the latter spot it is difficult, indeed rarely possible, to discover the orifice of a stricture, or to identify even characteristic structures proper to the canal. Differences of colour are those only which are readily appreciable. The employment of such an apparatus, moreover, is not to be regarded in any means as a painless proceeding, to say the least of it; on the contrary it is productive of much irritation. With a large endoscope suited for examining the rectum, the result is very much more satisfactory.

Hitherto, in speaking of the sizes of urethral instruments, the French scale has been always assumed, as being that best

FIG. 294.



in this country. The scale, however, of the French is π , as producing smaller intervals between each number, in making these intervals more regular, and the number itself to

express the measure of the calibre marked by each instrument. No. 1 is one millimetre in circumference, No. 2 is two millimetres, and so on up to 30. The Nos. 3 to 21 represent pretty generally the Nos. 1 to 12 of our scale, that is, about 18 numbers instead of 12. A fac-simile of the metrical scale is seen in the figure 294.

Treatment.—Its object is twofold :

First, to restore the natural calibre of the canal, so far as is consistent with the safety and comfort of the patient.

Secondly, to maintain the adequate patency of the canal afterwards.

Now, since strictures vary in amount of contraction, in dilatability, in disposition to return, in local sensibility, and in liability to manifest sympathy with other parts of the body, &c., various modes of treatment are necessary, appropriate to different cases. Hence innumerable inventions for the fulfilling of the above indications have been described, and modes of treatment proposed. All, however, may be resolved into three classes. The opposing tissue of the stricture is either dilated, gently or forcibly; or it is wholly or partially destroyed by chemical agents; or it is divided by some cutting instrument; and of course all these processes may be more or less combined with certain general or constitutional treatment.

Dilatation.—This is the mildest and the most desirable treatment to employ whenever the case admits of it; it is also the most generally applicable, and the best adapted to a very large proportion of all the cases presented to our notice. It is the method which almost all surgeons agree to use as the rule, availing themselves of other means when its action after fair trial is found to be inefficient. The history of surgery shows that it is still the most ancient mode, having been employed for the destruction of ‘carnosities’ in the time of Galen, and never having been superseded to the present day.

To apply it in ordinary cases, a flexible bougie, as large as the stricture will easily admit, should be passed fairly through it, and then be at once withdrawn with as much care and gentleness as was employed in introducing it: a note of the size should be recorded, and the patient desired to come again in two or three days: the same may then be passed, perhaps with greater ease than before; if so, it is to be withdrawn at once, and the next size larger introduced, provided it requires no undue force to do so. The visit is to be repeated as before, if neither pain, nor

ling, nor much smarting in micturition, follow and continue the operation. Sometimes a fit of shivering occurs, or the patient may be faint or sick, which are not unfrequent effects of passage of an instrument, more especially for the first time. Any of these phenomena take place, the interval should be lengthened a day or two, and the condition of the health examined. If the patient complains of soreness of the urethra, that micturition is painful, the urine is, perhaps, unduly, and it will be desirable to regulate his diet and habits so as to promote a healthy character in the secretions generally. At the same time, if it be so, he will generally derive benefit from a little bicarbonate or citrate of potash, combined or not, on occasion may require, with hyoscyamus, and taken three or four times a day; or, the alkali may be given in infusion of opium, if the mucous membrane of the bladder seems to be irritable, or disposed to secrete too freely. Irritability of the urethra, however, is much allayed by the gentle and careful use of instruments. Even when much suffering is produced at the attempt, it usually becomes notably less at every successful passage of the sound. Supposing none of these consequences happen, the same plan of gradual advance may be continued at each visit, so that No. 11 or 12 may be soon safely reached in such a case as that described, when a very fair amount of dilatation has been achieved: if the urethra is of the ordinary size, if the last step or two in the progress have been easily made, without pain or annoyance to the patient, it is well to go to No. 12 or 13. Finally, the maximum point of dilatation reached at should be maintained for a short time, the largest-sized instrument employed being used at gradually increasing intervals of time, in order to maintain the ground which has been won.

If, however, the case is more difficult, and after the use of a smaller instrument no penetration is effected, it is desirable to let the patient make water; if the stream is small, the size of the instrument should correspond with it; if it is not, the contraction cannot be very considerable, and some fold of membrane, or perhaps a false passage, has entangled the point; and in any case the bougie to be tried next should be as nearly as possible of the size of the stream. In introducing it, the floor of the urethra, as well as any lateral deviation in its course, is to be avoided, these being favourite situations for false passages. Failing of success in one direction, we should next

cautiously carry the point towards others, trying patiently for a short time to insinuate it either above, below, or on either side of the passage, if the slightest sensation of its being 'held' indicates that the orifice exists in any of these directions. And when the instrument has thus become a little grasped, we should endeavour to facilitate its progress by patient, continued, and moderate pressure, the precise amount of which should be proportioned to the degree which the patient will bear without much complaining. Some minutes should be devoted to the attempt, the success of which will much depend upon the steadiness and singleness of purpose with which it is pursued. Sometimes the introduction of the left fore-finger, previously oiled, into the rectum, will facilitate the progress of the instrument, either by permitting its point to be raised to some extent, or by enabling us to judge more precisely of its exact locality and relation to the parts around; above all, whether it is or not in a false passage.

The employment of continued pressure on the face or commencement of an indurated and not very sensitive stricture, is sometimes successful, either by inducing absorption, or by mechanical action upon the yielding materials of the obstruction. The operator, however, should be certain that he is really acting on the contraction, and not following or making a false passage. It is important to remember, as an invariable rule in relation to these attempts, that when the instrument is tightly grasped, the operator may infer that its point is safe within the strictured part, and that when the point feels free, movable, and capable of being withdrawn without appreciable effort, it is certainly not in the stricture; it may be, in such circumstances, in a false passage. If, after being grasped or 'held,' it advances suddenly for a short distance under pressure, and becomes movable, it is probable that a false passage has been made and the urethral walls perforated; after which unfortunate occurrence all further efforts must be given up, at least for several days, and the employment of instruments, when again resorted to, must be conducted with vigilant care, to avoid any re-opening of the lacerated part.


If after several applications of the instrument the stricture is still not passed, or if the case at its commencement appears to be more than usually difficult, we must take still further precautions to insure success. It is preferable to visit the patient when in bed, with the skin warm and moist from ample cover-

A hot hip-bath just before the visit is sometimes ad-
geous. When introducing the instrument, all unnecessary
ure should be carefully guarded against. Premising that
ituation of the stricture has been carefully verified before-
a silver catheter should be selected, the size of which
l be a little less than that of the stream of urine. Perhaps
ometimes desirable to apply the oil to the *urethra itself*,
r than to the instrument, injecting, by means of a common
yringe, four or five drachms of olive-oil. This proceed-
cilitates the progress of a small instrument to the stricture,
nds to open the latter by the penetration of the oil into
arrowed passage itself.

hen the stricture has been passed, considerable care is
ary in guiding onwards the point through the canal
ed, to prevent its becoming engaged in the enlarged
se which are commonly found here. These are usually on
loor; and not a little patience and management are some-
s required to carry the point of a slender instrument over
and lodge it safely in the bladder. Usually the point has
tilted upwards somewhat; and here it is that the finger
se rectum is most commonly of use.

Also passages are some of the most perplexing complications
narrow stricture which the operator has to deal with,
much as the difficulty of getting into the right opening is
ly increased by the readiness with which the instrument
as the wrong one. It is to be remembered that, as a rule,
passages most frequently commence on a level *below* that
se proper opening; and that the operator's finger when in
rectum, near to which the false passage is almost certain
an, will inform him as to the route which the catheter is
ng, whether it be too close to the gut or deviating to the
t or left of the median line, and will also assist him to
et the point in its true course.

he influence of chloroform has been sometimes found ex-
ely useful in facilitating the passage of a catheter or sound
ugh the urethra, especially when it is more than ordinarily
sitive, and the pain occasioned by instrumental interference
duces uncontrollable and involuntary efforts of resistance on
part of the patient. It is to be remembered that it is not
the purpose of permitting the instrument to be used with
ster force than before, but in order to produce relaxation of
muscular tissues, both of the voluntary and involuntary





or an instrument is followed by r
sometimes one of the most effic
adopted. In putting it into execu
finement to the house must be
having been introduced, it may be
the body; for this purpose one ma
hips, another is connected with
crest of the ilium and passed unde
to the crest, and then made fast.
by a piece of narrow tape to the b
tuber ischii. A peg of wood is
catheter, or, better still, a piece of
with the other end in a suitable
catheter, it may be attached to a
circling the penis.

In the management of this pro
attended to: first, the catheter wh
into the bladder, or at any rate b
distance is readily ascertained by
through it, and drawing the ins
stream just ceases; a very small
can then remain within it. Sec
passed has been a silver one, the s
be of gum-elastic, as causing less

amias or opium, if pain or irritation render it necessary. Matter may be given in the form of suppository; but this is only necessary if proper care is taken in the management of the catheter, which is the most important part of the treatment. However, blood appears in the urine, or fever, or swollen bladder, the catheter must be withdrawn at once. Usually, in about thirty-six hours, a purulent discharge is around the instrument, which soon becomes loose in the bladder, although when first introduced it may have been firmly fixed by the stricture. It may then be replaced by another one larger, which will probably enter easily. In a day or two a similar advance may be made; and if all go well, the treatment may be continued for five or six days; the patient is kept up daily for a few hours, if the gum-catheter is employed. If the patient is fatigued, a day or two's rest may be given, and the treatment resumed if necessary. When a full size has been reached, the catheter is to be withdrawn, and an instrument passed for a few days, gradually increasing the interval, but endeavouring to retain as much as possible of the calibre; a certain degree of recontraction at first, however, is to be met with.

Mechanical apparatus for effecting dilatation.—Among various kinds of special apparatus may be noticed a method adopted by Thomas Wakley, which consists in sliding tubes over a urethral guide first passed into the bladder; so that if the guide is at first correctly taken, all the other tubes will follow the same course. Different methods of using sliding tubes were first adopted by Dr. Buchanan of Glasgow (1831), Dr. Hutt of Dublin (1835); by Desault (1797), and Maisonneuve (1801). The attempt has been frequently made to dilate the urethra by means of a single instrument which expands in situ, in place of several instruments of increasing calibres which are to be successively pushed through the constricted part. For this object Mr. Luxmoor (1812) employed diverging rods; Leroy d'Etiolles the same method, some years later, and M. Perrève a somewhat similar plan (1847): Dr. Hutt commenced the employment of fluid expansion (1819). To a greater degree, the accomplishment of the same object has been attempted by the use of catgut bougies, and very recently bougies of Laminaria, a species of seaweed which expands in contact with moisture. A German surgeon formerly, and a Frenchman of Lyons more recently, have employed metallic



WHICH IS WELL KNOWN
the urethra. Moreover
the method of forcing
the instrument precisely
has operated with
greater freedom and
would generally ha-
bitually treat the stricture

In considering
means which have
to forbear remark
not rupture, com-
efficient or so desi-
simple bougie, wh-
with a light hand
affair of overcomin-
pedients of the eng-
themselves; but w-
prosecution of dil-
avoid producing l-
urethra and its in-
treated as tissues e-
relations of the hig-

On the other h-
which I have now

catheter-scale. If, however, forcible be applied to an old and rigid stricture the method in vogue thirty or forty (Mayor), that is by driving a large metal bougie through the obstruction, symptoms are almost infallibly incurred. employed, and not exerted from within as in the distending instruments re-pushes the stricture down the canal, cellular connections, and infallibly inflammation and fever.

He describes his mode of operating as follows: The instrument 'consists of two blades fixed in a divided handle, and between them a wire welded to their point. On this wire a tube (which when inserted between the blades corresponds to the calibre of the urethra) is quickly passed, ruptures or splits the obstruction.' He introduced it, and 'reached the bladder, it should be gently rotated, to prove it fairly within that viscus; and being removed, the surgeon is next to place the same tube he had previously selected between the blades, and thrust it forwards to the end. The stricture is fairly split, the dilator should be still further separate the sides of the stricture then be withdrawn; a catheter corresponding to the number of the tube being substituted for the purpose of removing the urine.'*


After the patient is then removed, the patient is given quinine and opium for the first twelve hours; and the same catheter inserted in forty-eight hours; and again on the next days for a week or two, gradually increasing the interval.

Under obligations to Mr. Holt for his highly pursuing his practice by rupture,

FIG. 296.



Distending instrument; the blades are opened to the full extent. The index is seen just below the handle.



...is with a cutting instrument cases which have been hitherto question offer advantages. To first sight a harsh and rough prejudiced examination of its personal practice of it, has character accorded to it here.

The plan which I have adopted consists of two long and narrow each other, but both forming whoment the size of No. 3, 4, or 5. end and throughout four inches c by a handle containing a screw, rods to diverge from each other a given spot, forming a spindle-sha in length, and the third or half c is an index near the handle which tension made, by pointing out th which is reached, as the operatio reason for carrying distension be of the urethra will admit, is the the great majority of strictures, t the bulbous portion of the ureth which naturally possesses a calib catheter-scale, and that ordin...

Before operating, the distance of the stricture from the external meatus is measured by passing a full-sized bougie down to the stricture; the slide is then placed upon the figure on the instrument which denotes that distance. The instrument is passed until the slide arrives at the meatus, and the blades are slowly separated: when the maximum distension is reached, the screw is turned back only about half-way, so as not to close the blades; the instrument is withdrawn; a full-sized gum catheter is then passed, and allowed to remain twenty-four hours. On the next day after the operation a large metallic sound is passed, and subsequently at longer intervals. The instrument may also be used rapidly to produce rupture if preferred.

We return to the subject of gradual dilatation, and particularly to the employment for old and irritable strictures: in such cases, especially when occurring in enfeebled constitutions, it is necessary to proceed with great caution. More progress will be made by employing those adjuvants, which a deranged state of the constitution requires, than by pertinaciously continuing the use of the bougie when the stricture is unusually firm; this is a maxim of the first importance. Further, when arrived at any given number in the scale, say 6 or 7, at which it is difficult to proceed, we may sometimes gain on by employing half or intermediate sizes, which I have been in the habit of thus using for many years. Or we may employ the French scale, which gives 18 numbers between our numbers 1 and 12, and so admits of more gradual steps in the treatment; the number 3 of the French scale equalling our number 1, and the 20 our number 12; these have the advantage of indicating the exact calibre of the urethra through which we have passed, the number of each indicating so many millimetres of circumference.

There are some cases unquestionably in which—however much pains and care, of adjuvants local and general, have been expended—the patient's existence is barely a tolerable one. The urethra and bladder are so sensitive, or the stricture is so constrictile, that exquisite suffering from instrumental interference, fever, or retention of urine constant or constantly impending, are the alternations presented. These facts, recognised by surgeons of experience, have led to proposals to treat the stricture by many other methods. The next to be considered is treatment by caustic agents.

Caustics.—During three centuries at least, escharotic sub-

stances have been employed by surgeons for the purpose of destroying the obstructions, in earlier times supposed to be 'callus' or 'caruncle,' which oppose the passage of the urine. With this view, savin, antimony, mercury, verdigris, quick-lime, vitriol, alum, and other active agents have been carried by some contrivance or another down to the urethra. John Hunter, in the latter part of last century, brought into note the nitrate of silver; and Mr. Whately, in the commencement of the present century, the caustic potash. These substances have been supposed by their advocates to act, first, by deadening sensibility, and thus allaying spasm; secondly, by causing to slough, and so destroying, the morbid tissue of the stricture itself; and lastly, by dissolving or softening that tissue, and thus permitting instruments to be passed. A very considerable experience has been obtained by some English and many French surgeons, the result of which is, that the practice of applying chemical agents has been for some considerable time steadily diminishing in the hands of the best surgeons. It has been very generally doubted if it be possible to limit the application of any soluble substance possessing the active qualities of those two agents, especially those of the potassa fusa, to any given portion of the urethra, particularly in those cases which are complicated with false passages, and which furnish by far the most difficult examples which the surgeon is called upon to treat. Their use also is by no means free from painful and even dangerous results, such as violent spasm, retention, and fever. Moreover, there is reason to believe that these caustics exert an action upon the urethra similar to that which follows their application on other parts of the human body, and liable to be followed by increased induration, and the formation of fresh contractile tissue. On the other hand, the advocates of potassa fusa especially claim, that after its use the catheter is more easily passed; and, that in cases in which they have been unable to pass any instrument at all, this agent has frequently been successful: while numerous results of this kind may, they say, be referred to. Now, it may fairly be granted as probable that a certain present advantage may be gained by the method in question, and that in this manner it may in some hands be undoubtedly useful; the remoter effects, however, already alluded to, should not be lost sight of; nor should the very important consideration that, by patience, care, and gentle management, there are very few cases indeed in which a catheter of some kind cannot be introduced fairly

high the stricture into the bladder; a result far more desirable than any which is only accomplished by means of caustic applications. The conviction gains ground, both here and abroad, that those cases which have been called 'impermeable,' in relation to catheterism, should very rarely be considered so. Even the opinions and practice of the profession abroad, in relation to caustics, cannot be more truly or concisely represented than in the recent words of Nélaton, who states that "irritation of any kind is rarely employed now, on account of uncertainty and the tumefaction of the urethra and retention of urine which it produces, and the new formation of contractile tissue which is likely to result."* In like manner, the widely held opinion in England is thus expressed by Mr. Erichsen: "The practice, stigmatised by Mr. Liston as "most atrocious," is now but few advocates; and, indeed, there appears to be no surgeon who effects but what can be accomplished much more safely and easily by a catheter or sound in an ordinarily skilful hand."† It may be added, that my own observation and experience lead me fully to coincide with these able surgeons.

Under such circumstances, little need be said here as to the mode of employing these agents. It suffices to state that the nitrate of silver may be applied by passing a small stilet, on which the salt has been fused, by means of a hollow canula down the stricture, with which the agent is permitted to remain in contact for a few seconds. The potassa fusa is ordinarily employed by carefully attaching a small fragment, such as the end or the half of a grain, to the end of a wax bougie, and passing it down and pressing it for a moment or two against that structure, whatever it may chance to be, which obstructs the instrument in its progress along the urethra.‡

Incisions.—Under this somewhat comprehensive term are included numerous modes of dividing the morbid structure which obstructs and narrows the urethra. All may be arranged in three classes: the first contains operations in which the incision is made altogether within the urethra; the second, those in which the incisions commence from without, usually in the peri-

Mémoires de Path. chir. Paris, 1858, tom. v. p. 409.

Science and Art of Surgery, 2nd edit. Lond. 1857, p. 925.

Further information on this subject, if required, may be found in the works of Phillips, who first advocated and subsequently repudiated the employment of caustics; of Dr. Jas. Arnott, Mr. Wade, and Mr. H. Smith.

neum, and are carried into the urethra and through the seat of stricture.


The first, or internal urethrotomy, will now be considered. This method of dealing with obstinate stricture has been practised for above a century. Long before that time, internal incisions had been combined with the use of escharotics, although it does not appear that they were employed alone much before the period named. Allies of France (1755), Physick of Philadelphia (1795), John Bell (1806), Charles Bell (1807), M'Ghie (1823), and Stafford (1827), have employed different instruments and different modes of performing internal incision. Modified forms of some of these have been occasionally adopted by modern surgeons in this country; but the method has been practised much more extensively in France, where very numerous forms of urethrotome have been designed.

All belong to one of two classes: in the first, section is made by pushing downwards a lancet-like blade, generally with a slender conductor in advance of it, into the obstruction to be divided—incision from before backwards. In the second, a portion of the instrument containing a small blade, sheathed, is first carried through the stricture, which is divided by protruding the blade and drawing it outwards through the whole of the contracted portion—incision from behind forwards.

Incision from before backwards.—Whenever this mode is practised, a guide should previously be passed through the stricture into the bladder, to insure an accurate course for the blade: formerly incisions were made without any guide—a method now wholly discountenanced, on account of its uncertainty and danger. This proceeding has usually been considered more applicable to obstinate strictures which are situated within three or four inches of the external meatus, provided they are not greatly indurated, than to those which are more deeply placed, although it is quite possible, especially with modern instruments, to employ it for these also. It has not been greatly used in this country, but has of late come more into vogue abroad, from the recent employment of a fine flexible guide, which precedes the urethrotome in its passage through the stricture and is then pushed onwards into the bladder. The practical value of this method cannot be endorsed here: a better one, with a grooved guide, has been employed by Mr. Wood of King's College Hospital; and more lately a 'catheter urethrotome' has been designed by the present writer, 'the peculiarity of which is that


of the process is mechanically certain and safe, after silver catheter has been passed.' The catheter having entered the urine, and thus proved that the instrument is

FIG. 297.



theter, size of No. 2, by which the withdrawn. B is a rod of the same size screws into the end of the catheter. urethrotome: *b* the blade sheathed; *c* the blade which unsheathes it. This instrument is passed down a groove in A, and, after incising the stricture, is withdrawn; when E, a silver catheter, is passed into the bladder with the catheter and rod, AB.

FIG. 298.



Civiale's urethrotome.

false passage, the blade slides along it through the stricture, is then withdrawn, and a large gum catheter is passed

over the small silver catheter, which, being also withdrawn, concludes the operation (see Fig. 297). This remains for twenty-four hours; and the subsequent treatment is the same as that described in the next section. More recently an instrument has been devised which bears the name of Trélat of Paris, more efficient, perhaps, than any of this kind which have preceded it, inasmuch as it may be used not only from before backwards, but if, after such an incision, a second is necessary, it can be made from behind forwards, and with augmented power (see Fig. 299).

FIG. 299.

Trélat's
urethrotome.

Incision from behind forwards.—Nevertheless, this mode of operating is for the most part more certain and effective than the preceding; and it is applicable equally to all parts of the canal. It requires generally, however, that the stricture should be dilated sufficiently beforehand to admit that part of the urethrotome which contains the blade to pass through and beyond, before the incision can be made; and this portion is necessarily equal in size to No. 3, 4, or 5. This preliminary dilatation is accomplished by the 'continuous' process. It has often been objected that if we can pass a No. 4 or 5 through a stricture, there can be no occasion to employ any cutting instrument at all. But such a remark is based on defective acquaintance with the subject. The indication for a cutting operation is not that the stricture is of very small calibre, but that it is non-dilatable; that is, that its calibre cannot be permanently increased by the most persevering and judicious application of instruments, or that so much fever and distress are produced by them as to counterbalance the benefit of any dilatation they may effect. Thus, a stricture may at the outset of treatment not admit even a No. 1 bougie, yet in a short time it may be easily dilated, and show only slight tendency to return, easily counteracted by occasionally passing an instrument; on the other hand, a stricture

which habitually admits No. 6 or 7 may be the cause of the greatest suffering, of almost daily retention, and be irritated instead of benefited by all attempts to dilate. It is in such cases, happily not frequent, that incision has proved of very great service. I have rarely experienced better results from any operation than those obtained from internal urethrotomy in cases of the kind described.

Before employing any form of internal urethrotomy, it is necessary to have an accurate idea of the site and extent of the stricture to be divided; this is obtained by employing on two or three occasions a solid sound with a bulbous extremity, just large enough to pass through the contracted part (p. 954).

I have employed by choice the urethrotome of Civiale,* and to the present day (1870) I prefer it to any other. In this instrument the blade lies concealed in the bulbous extremity, from which, by means of a simple contrivance in the upper end of the sheath or canula, it can be made to project one, two, three, or four degrees, according to the depth of the incision intended (see Fig. 298). The mode of proceeding is as follows: having sufficiently dilated the stricture, so that it will admit about a No. 4 or 5 bougie, the situation and extent of the obstruction are ascertained by means of the urethrotome itself, the bulbous end of which forms a useful sound. The bulb is then carried about one-third of an inch, or a little more, beyond the stricture, the blade projected, and the incision made by drawing it slowly but firmly outwards—that is, in a direction towards the external meatus—to the distance of an inch and a half or two inches, generally towards the floor of the urethra, so as to incise fully the stricture and a little of the sound urethra before and behind it. Experience shows that hazard is not increased by the length of the incision, which should always be fully adequate to the purpose; but it is quite otherwise in the important particular of depth: this should never exceed the amount produced by about half the power of projection possessed by the instrument in question. It is not an unwise proceeding to pass immediately after the incision the dilating or distending instrument to the fullest size, so as to ensure the perfect freedom of the canal at all points. After the operation, a full-sized elastic catheter is

* Various other methods and instruments have been devised, by Amussat, Leroy d'Etiolles, Sédillot, Maisonneuve, Ricord, Mercier, Reybard, Heurteloup, Bonnet, Boinet, and others; but it is deemed unnecessary to refer further to them here.

retained for twenty-four hours. During the first fortnight, a metallic sound is passed about every second day, taking care to press the convexity of the curve well downwards into the site of the wound, so as to keep the lips asunder, or at all events to extend the cicatrix. After this it is to be used every three days, every four, every week, and at last twice a month. Hæmorrhage to any considerable extent is rare; if it is free, a full-sized instrument in the urethra, and external cold, will stop it. Febrile symptoms occasionally show themselves after the operation, as in other modes of treatment, and disappear without remedial means. In relation to the accidents which may immediately follow the operation, the following is the result of perhaps a hundred cases of urethrotomy performed by myself in the manner described. In those practised for stricture situated in the urethra in front of the scrotum, I have seen no ill effects whatever. In those practised in the bulbous part of the urethra, I have twice seen free but never dangerous hæmorrhage; and sometimes a smart attack of fever, but never abscess, extravasation of urine, inflammation of the bladder or kidneys, or pyæmia. Such occurrences may occasionally be met with, doubtless, because any instrumental interference with the urethra—even the passing of a bougie—is apt in certain cases, happily very few in number, to give rise to these conditions. Indeed, no one who is experienced in the surgery of the urinary organs can question that such events must sometimes arise.

In relation to the positive benefits which are to be obtained, the immediate result is perfect. Properly incised, there is no longer the slightest grasp on the largest instrument introduced, and it is often difficult to recognise the seat of the former contraction.

In some cases these results are permanent; in others, the stricture shows a tendency to return; but as in external division, so in internal, the contractility and the nondilatability—those characters which have been before shown to be the indications for resort to operative procedure—are generally greatly diminished in intensity; and if subsequent treatment be necessary, dilatation usually suffices.

Such have been the general results of urethrotomy, when properly performed, and applied to suitable cases. I think it right, however, to add here, that my observation and experience of distension at a single sitting, or by rupture, have convinced me that we may thus produce excellent results at first, which how-

ever are often not durable; so that these methods will by no means supersede internal incisions. For all non-dilatable strictures situated from two to four inches from the external meatus, I decidedly prefer internal urethrotomy: the operation is perfectly safe and easy to perform, and its results more lasting and complete than any other.

Incision is also especially applicable to that not uncommon form of stricture which occurs at, or very near to, the orifice of the urethra. It may be congenital, the result of inflammation, of cicatrisation after chancre, or other lesion there. It may be the only obstruction in the canal, and yet give rise to the most painful and serious symptoms, and even to a fatal result. I have repeatedly given complete relief to distressing symptoms, the cause of which was not suspected, by dividing an external meatus, which admitted nevertheless a No. 5 or 6 catheter. The incision may be performed with a director, and a straight, narrow-bladed bistoury; but the most efficient instrument is a small *bistouri caché*, which is passed through the contracted part of the canal; pressure being made on the handle, the blade is opened to an extent previously determined and arranged by means of the screw; it is then drawn out, with the edge turned towards the frænum, and the section is made. A piece of dry lint, introduced by the meatus for an inch and left there, stops the bleeding. After a day or two, the patient introduces for himself a short conical metal bougie, about three inches long, and provided with a handle sufficiently large to prevent its slipping into the urethra.

The late Mr. Colles, of Dublin, adopted a method of treating some of these cases which deserves notice. It consists in drawing over the surfaces divided by the section described the two edges of urethral mucous membrane, and stitching them to the external margins of the wound, each lip of which is thus covered by membrane, and prevented from adhering to its fellow.

External urethrotomy.—The division of the tissue composing the stricture is sometimes effected by incisions commenced from the external surface. Various external operations, although designed mainly to relieve retention of urine, were made at a very early period, according to Rhazes (10th century), and Avicenna (11th century). The earliest operation for the cure of stricture apart from retention, but in which the surgeon could pass no catheter, is recorded by Wiseman (1652). A few years afterwards, Solingen, at Livourne, adopted a similar course.

François Tolet and Colet (1690) appear to have performed similar operations in France. J. A. Petit and Ledran (1740) had recourse to them also, and more frequently perhaps than their predecessors. In 1783 John Hunter performed the operation now known, and described hereafter, as the perineal section; but it was rarely employed until Mr. Grainger of Birmingham advocated it, in 1815. He was followed by Mr. Arnott: since which time it has been the proceeding ordinarily adopted for the treatment of certain extremely obstinate and complicated cases, in which the surgeon has found himself unable to pass any instrument whatever through the stricture. More recently (1844) Mr. Syme has advocated the division of the stricture for some exceptional cases, in which, *although a catheter can be passed into the bladder*, no other treatment has afforded sufficient or permanent relief. This operation he has called 'external division,' a term which should be reserved for his method; while it is equally desirable to limit the use of the term 'perineal section' to the proceeding just referred to in connection with strictures which have been found impervious to the catheter.

Perineal section will be first considered. This operation, or that without a guide, and employed only for so-called 'impermeable' stricture, is much less frequently performed now than it was some few years ago. There is a growing belief, which has greatly influenced practice, that few, if any, strictures exist through which, if the urine issues externally, a catheter of appropriate size may not be passed in the course of time, with care, gentleness, and skill. This opinion, after some years of experience of the treatment of stricture in its worst form, I entirely endorse; and have no hesitation in stating, as a general rule, admitting of very few exceptions, that no stricture permeable to urine is impermeable to an instrument, providing that that instrument is sufficiently small, and is used with the utmost gentleness and care. If, however, the surgeon meets with an exception to this rule, the following operation, which I may repeat I have had occasion to perform only twice, is considered admissible. The stricture being situated in the bulbous portion of the urethra, the patient is placed on a table, in a good light, is secured as for lithotomy, and the perineum is shaved. A large catheter is to be passed as far down the urethra as the obstruction will permit, taking care that the instrument rests on the face of the stricture, and not in a false passage; an assistant holds it firmly, and draws the scrotum upwards. An

an is made in the middle line of the perineum, along the
 from the base of the scrotum to near the margin of the
 and the point of the catheter is to be exposed by deeper
 ens. The sides of the opening are then to be held apart
 ely as possible with hooks, better still by two loops of
 one passed through each margin of the urethra at the
 of obstruction as soon as the canal has been laid open so
 give a clear view of the urethra and of the contracted open-
 This done, the operator, who should be provided with
 and directors of the smallest size, endeavours to carry one of
 through the contraction, and if he is successful, the division
 is made with ease and safety. He may not be able to pass
 rector more than two or three lines, until, having made a
 division so far, he may be enabled again to follow the track
 of contracted canal, and to divide another portion of it upon
 instrument. But if a director cannot be introduced, he must
 pour by dissection to follow the urethral canal as closely
 as possible. In either case, as soon as the continuity of the
 tube is restored, the catheter first employed is to be carried
 wards into the bladder, secured in the usual manner, and re-
 d for some days.

External division.—This proceeding consists in first passing
 under grooved steel sound through the stricture into the
 bladder, and in cutting down from the external surface to it, so
 divide entirely the stricture: then a large catheter is tied
 for forty-eight hours, and dilatation maintained at intervals
 wards.

The following are the details to be attended to in performing
 operation: First, a clear idea of the precise situation and
 extent of the stricture must be formed before commencing.
 The patient being placed as for lithotomy, a staff, the
 lower half of which is slender and grooved, and the upper of
 a size, is passed so that the shoulder or edge of the thick
 portion rests firmly against the face of the stricture, the slender
 part passing through it into the bladder. An assistant holds
 the staff in his right, and draws up the scrotum with the left,
 hand. The operator makes an incision in the line of the raphé
 above downwards, about two inches long, and steadily
 follows the direction of the median line, through the structures
 intervening between the skin and the staff, until he can dis-
 tinguish the shoulder of the staff, and by that is guided to the
 situation of the stricture, when, taking a sharp straight bistoury
 in his right hand, and the staff itself in his left, he engages the

point of the knife in the groove of the staff about an inch below the shoulder, and cuts upwards, dividing the tissues, to the upper end of the groove. If the stricture has been properly divided, the shoulder of the staff may be passed downwards through the incised part with ease, and if so, is withdrawn, and a catheter passed in its place. Since, however, the point of a catheter sometimes catches in the wound, and does not go readily into the bladder, perhaps from the urethra occasionally collapsing at the posterior limit of the incision, I prefer to introduce by the wound, before withdrawing the staff, a concave curved director along the convexity of the staff, on which the catheter glides securely and certainly into the bladder. If the catheter, however (which should not be smaller than No. 10), be obstructed at any point, and be obviously grasped after it has been passed, it is certain that some fibres still require division, and these should be at once incised.

At the end of forty-eight hours the catheter is to be withdrawn; and generally about three days after a full-sized sound should be passed. Mr. Syme likes to guard the wound from contact with urine as much as possible, by introducing a tube from the wound to the bladder, and leaving it there, as after lithotomy; and he employs, in that situation, one with a double curve, and about nine inches long, instead of the catheter, which he dispenses with altogether at first. Subsequently the catheter is passed every three or four days for a time, then once a week, then once a fortnight, and finally once a month; and the practice may be continued if any disposition to contract appears; but if otherwise, it is necessary only to make a trial of it from time to time.

Mr. Syme lays stress upon the importance of attending to the following particulars, which I add briefly here: 1. Maintain the median line in the incisions. 2. Make a direct opening down to the staff, not a tortuous one. 3. Divide the whole of the contracted part, rather more than less. 4. Do not cut so far back as to endanger the deep fascia of the perineum, and use the knife in the deep incisions with the cutting edge uppermost. 5. Do not close the end of the inlying catheter, lest urine be forced into or through the wound, for want of patency in the instrument. 6. Avoid escape or displacement of the instrument. 7. If incisions are made far back, introduce a curved tube through the wound when the catheter is withdrawn. 8. Do not neglect dilatation during the progress of recovery.

results of treatment by external incisions.—In reference to the operation without a guide, or the perineal section, the best that can be said of it is, that it is a hazardous proceeding, and no surgeon should entertain the idea of performing it except after thoroughly assuring himself that the stricture cannot be rendered amenable to instruments, and so made amenable to other modes of treatment. In a few cases the operation is performed successfully, but in most there can be little doubt that the course of the stricture itself is not followed, and that the knife creates a new channel through adjacent and often very unsound tissues, very inadequate to perform subsequently the function of the urethra. In many instances the attempt to make a channel into the bladder has wholly failed, and the patient has been removed from the operating-table unrelieved; and in not a few cases the result of the proceeding has been fatal. This termination, however, is to be anticipated in a large proportion of what must necessarily be the worst cases of their kind. Such views of the operation tend to impress us more strongly with the conclusion, that every chance of getting an instrument through a dilated urethra by protracted but gentle endeavours, combined with rest and constitutional treatment, should be exhausted, before we consent to resort to perineal section.

Respecting the operation of external division, or that performed on a grooved staff, it will be remembered that it is applicable only to cases which present an insuperable tendency to contract after dilatation, in which the existence of inextinguishable perineal fistulæ demands some such operation.

After a careful observation of a large number of cases so treated, there is ground for speaking with some degree of certainty respecting the immediate and remote results of this operation. Among 219 cases, collected with scrupulous care by myself, there was a mortality of between six and seven per cent. Of this percentage nearly two-thirds died of pyæmia, the remainder (one or two only excepted) from fever and suppression of urine. Such a mortality is by no means large when we consider what kind of subjects selected bad cases of stricture are, especially as they occur in hospital practice, by which most of this series was furnished.

Hæmorrhage, to a moderate but not dangerous extent, not infrequently occurs either at or after the operation. Very rarely is it serious; and when it is, there ought not to be much difficulty in stopping it. A full-sized catheter being in the

urethra, it is necessary to plug the wound with lint, and maintain a firm pad of the same material over it, by means of a T-bandage closely applied. In one case in which I operated, the perineum being greatly indurated with several fistulae, there was obstinate bleeding from vessels which the ligature would not hold, owing to the nature of the tissue in which their mouths were embedded; but I succeeded in stopping it by carrying a small curved needle beneath each bleeding point. In regard to the important question of remote results from this operation, there have been some few cases undoubtedly in which it has failed to be successful; in a much larger proportion, more or less of recontraction appears after the lapse of time. This has sometimes arisen from want of care to divide the entire contracted portion of the urethra; in other cases, from neglect of subsequent dilatation. On the other hand, in some cases, I can personally vouch for the fact that no narrowing has appeared seven years after the operation. In most, I believe, it will be always necessary, or at all events prudent, to maintain some dilatation afterwards; the condition usually obtained being a dilatable urethra, in the place of a non-dilatable and obstinate stricture.

SPASM OF THE URETHRA.

It is extremely rare that any considerable narrowing of the urethra takes place as the result of pure spasm of the muscles surrounding the passage. Granted, however, the presence of organic narrowing, or of inflammation in the canal, an undue action of the urethral muscles may be excited, so as still further to narrow it. There is no doubt that a slight degree of this action may be excited in any part of the passage; but it certainly becomes more obvious in the membranous part, where sphincteric muscles exist in addition to those unstriped fibres which surround the canal throughout the whole of its course. Nevertheless the existence of 'spasm' is very constantly and very frequently affirmed, often without any proof of its presence, and usually in order to account for unsuccessful catheterism. Organic difficulties are quite sufficient without it, in many bad cases, to baffle at times the most experienced and most dexterous operator.

As immediate causes of spasm may be named the following: the presence of organic stricture; of inflammation from repressed gonorrhœal discharge; irritation of the mucous membrane

from highly acid, and occasionally from acrid, urine; from foreign matters ingested, and expelled by the urine, as cantharides, turpentine, and perhaps some forms of condiments and of alcoholic drinks; and from voluntary retention of urine exerted during too long a period of time. Examples of more remote causes of spasm are rectal irritations, as hæmorrhoids, fissure, prolapsus, fistulæ, operations in the rectum, ascarides, prurigo about the anus, derangements of the digestive apparatus and of the cerebro-spinal system. But it is important to remember that the distinguishing feature which marks the phenomena thus ascribed to irregular muscular contraction, and by which they are contrasted with those of organic stricture, is their transitory character. The symptoms of a narrowed urethra may repeatedly occur, but the canal ordinarily possesses its natural degree of patency. This is never the case in the presence of organic stricture; the stream then varies, but it never attains the natural size.

In treating spasm, general principles must be kept in view, and applied according to the particular requirements of each case. Local treatment of the urethra itself is often of secondary importance, sometimes, indeed, prejudicial. The main thing, as in all spasmodic affections, is not to regard so much the sign or symptom itself as to investigate the cause, a correct appreciation of which is the only key to successful treatment. This must be carefully sought, not only in the urinary tract, but in adjacent and allied organs, and in the condition of the system at large. Speaking in general terms, it will be found that attention directed to the condition of the animal powers, to the improvement of the secretions, and to the regulation of the regimen and habits of the patient, will conduce more to the removal of the local symptoms than any measures apparently of more direct or special application.

If any degree of organic stricture coexists, it is essential to treat this in the usual way; but as far as the complication of spasm is concerned, the treatment consists in obviating all sources of derangement which can be discovered as likely to influence the condition of the local disease.

I have known a single instance of spasmodic retention to occur periodically, and to disappear before the use of quinine: Sir B. Brodie records one similar case. A very hot hip-bath, a cupping on the perineum, or the inhalation of chloroform may sometimes give instant relief. For the most part, however,

spasmodic action in the urethra is associated with inflammation affecting a part or the whole of the canal, and especially when inducing retention of urine. Any further notice of its treatment will therefore be referred to the management of that condition. (See Retention of Urine, p. 989.)

Inflammation.—Retention of urine is undoubtedly caused by inflammation of the urethra; but this is by no means inflammatory stricture in any sense in which the term can be logically employed; the pathological condition usually present in such cases is acute inflammation of the prostate, with swelling of that gland (see p. 940). The patient complains of heat, fulness, and tenderness in the perineum; the passing of the urine is exquisitely painful; the stream narrows rapidly during the act of passing it, and ceases before the bladder is emptied or the desire diminishes. Usually it is found on inquiry that a recent purulent discharge from the urethra has been suddenly checked by external cold or wet; the signs of inflammatory fever are usually present. As this condition almost invariably comes under the surgeon's notice for relief of retention of urine, the subject of treatment will be considered in the section devoted to that condition resulting from acute prostatitis.

Tumours in the urethra.—The cause of urinary obstruction was formerly supposed, in nearly all cases, to be the existence of 'carcinomata' or 'caruncles' growing from the mucous membrane of the urethra. Such bodies do exist, but they are excessively rare.

Pascal (1718) describes two cases of numerous small fungoid excrescences obstructing the urethra. Morgagni (1761), in his forty-second letter, speaks of one which he had seen among many examinations. Arnaud (1769) names three instances. Hunter met with two cases; one is in the Museum of the Royal College of Surgeons. Sir Charles Bell describes and figures two cases. Amussat, Civiale, Lallemand, Ricord, Chelius, Leroy, and Mercier, all describe occasional cases met with in their own individual experiences. Guthrie and others in this country have met with a case occasionally. A good specimen exists in the Museum of Guy's Hospital of polypoid growth springing from the mucous membrane of the membranous portion. I have myself met with a good example of small polypus springing from the verumontanum, in a man, aged fifty-four years, a patient in the St. Marylebone Infirmary. The only sign of its existence, which could be ascertained during life, was an increased frequency of micturition. Those which are found at the anterior part of the canal, and which appear almost confined, as regards situation, to the fossa navicularis, are usually soft, of a rose-red colour, bleed very readily, and are not very sensitive. Their close proximity to the erectile tissues beneath may be reasonably supposed to be the cause of their peculiar vascularity. That they are, but more rarely, to be found in the posterior parts of the canal, is proved by some of the prepa-

was referred to. On the other hand, almost all the specimens of the polypoid growths which I have seen are confined to the prostatic portion, and are sometimes accompanied by others at the neck of the bladder or within it, to which latter, indeed, they then have the appearance of being secondary formations, and they are more frequently found affecting only the lining of the bladder, and not that of the urethra at all. In such cases their structure amounts to more than hypertrophy of the mucous membrane.

The conclusions to be drawn from the facts known respecting growths into the urethral canal are:

First, that the existence of any excrescence so large as to attract observation as an independent growth in the urethra is extremely rare.

Secondly, that these bodies consist either of (a) *vascular excrescences*; (b) *polypoid formations* peculiar to the prostatic portion of the urethra; or (c) masses of *tubercular* and *cancerous* growth.

Lastly, that tubercular and cancerous deposits are rarely primary in the urethra, but are mostly secondary to disease of the kidney, bladder, or prostate, and rarely appear until the primary disease has been developed to a formidable extent elsewhere.

Urinary abscess—most frequently a result of obstruction in the urethral canal—may be regarded in reference to its situation as penile or ante-scrotal, scrotal, perineal, and intra-pelvic. Among these the perineal is the most common, next comes the scrotal; the intra-pelvic occurs usually in very advanced cases of stricture, &c., in which a fatal result is impending, and also as a consequence of operations; and the penile is rare, most frequently being caused by gonorrhoeal inflammation of the urethra.

Occasionally, but by no means commonly, urinary abscess in the perineum may be met with, when there is no obstruction in any part of the urethra. When this is the case, there is no ground for suspecting that disease exists at or near the neck of the bladder, particularly disease of the prostate. In chronic suppurative disease of that organ, perineal abscess sometimes appears.

Ordinary urinary abscess in the perineum may be acute or chronic: the former usually requires prompt interference on the part of the surgeon, and its presence is often indicated by constitutional symptoms before local evidence appears of a marked character. If these are severe, and there are also tension, swelling, and tenderness in the perineum, a free and deep

incision made in the middle line will often give instant relief; and if matter is not seen at once, such a proceeding will do no harm, although nothing else result but a little bleeding, and the relief of tension. No considerable hæmorrhage need be feared, although sometimes, when there has been much inflammation in the part, a smartish trickling may continue for some little time. When it has ceased, a poultice should be applied to the wound. Great improvement in the patient's condition often takes place almost immediately; the fever subsides, and complete recovery may follow in a very short time.

The importance of speedily evacuating such collections of matter, even at the commencement of their formation, cannot be overrated. Matter pent up behind the deep perineal fascia which forms a partition too dense to be penetrated by the action of absorption, will find its way into the cellular tissue of the pelvis, by the side of the bladder, between it and the rectum, and give rise to dangerous consequences, or in event of recovery, to urethro-rectal or vesico-rectal fistulæ. Otherwise it may burst into the urethra, and be discharged by the external meatus. The collection having been opened, pus in some quantity escapes, usually alone, sometimes mixed with urine, but not necessarily so. One of the main objects in making an early opening into a collection of matter in the perineum, is to prevent the occurrence of any lesion of the urethral walls; if its evacuation be soon and fully insured, we may hope to find the cavity gradually closing, and that no urine will penetrate it, when there will be less fear of its remaining long open, or of its becoming an abnormal passage for the urine.

In the case of chronic abscess, it is desirable that an incision should be made when the presence of matter is perfectly evident. These abscesses are usually situated in the neighbourhood of the bulbous portion of the urethra, and appear in the perineum and scrotum; sometimes as the result of a little ulceration behind the stricture, and escape of a drop or two of urine into the cellular tissue; oftener, perhaps, from the adjacent irritation, without having any direct communication at first with the urethra. This communication almost invariably takes place if the abscess breaks of itself, sometimes when it is opened by the lancet; but generally no urine appears when it is opened early, until some days after the operation, and occasionally not at all. An unnatural opening, however, having been established, the frequent passage of the urine

through it, which must occur at each act of micturition if the stricture be narrow, prevents its closure. This artificial canal, usually termed urinary fistula, is one of the common accompaniments of neglected stricture, and often forms a troublesome complication.

Urinary fistulæ.—The external openings of these passages are most commonly seen in the perineum and scrotum, which are traversed by them in various, and often circuitous, routes; less frequently they are observed in the groins, the upper part of the thighs, the adjacent part of the nates, or even above the pubic symphysis. In the last-named situation, the devious channel usually results from incisions made to relieve extravasation; but in the scrotum or perineum it is generally due to a previously-existing urinary abscess.

Under the term 'urinary fistulæ' all these conditions are commonly included; some of them simple, and easily amenable to treatment; others complicated, and requiring much care and time in order to attain a successful result. Some are merely narrow channels through nearly healthy parts; others pass through structures indurated and deformed by repeated deposits of plastic matter, and sometimes connected with cavities secreting pus, and detaining in their interior some quantity of the urinary secretion. The external orifices of the fistulous passage may be few or numerous; being in the latter case the outlets of sinuous channels springing from the original track, and giving exit to a number of small streams when the act of micturition is performed. And lastly, besides the foregoing, there are those openings into the urethra which have their origin in loss of substance by sloughing from extravasation, phagedænic ulceration, or violent injury to the parts; and these abnormal conditions are distinct in their nature from the two preceding classes. This mode of arranging the numerous and widely-differing lesions comprehended under the general term urinary fistulæ, indicates three forms of morbid condition, each requiring appropriate treatment.

1. Simple fistulæ. The first class embraces those cases where, in connection with stricture of the urethra, one or more fistulous passages exist, the surrounding parts being not much altered from their natural or healthy condition. These openings are the result of nature's mode of affording relief in cases of narrow stricture; they act as safety-valves to the pressure exerted upon important organs behind the obstruction. Thus

we often see patients enjoying fair health and comfort, notwithstanding large fistulous passages in the perineum, by which all their urine is passed. But the annoyance, sometimes the pain, besides the tendency to grow worse, which accompany urinary fistula, to say nothing of considerations arising in relation to the sexual function, demand the interference of the surgeon to bring about a natural state of things.

For these cases, as a rule, nothing else is required than to dilate fully the urethra. The urine will flow by the natural channel, and the fistulæ will heal of themselves, if we insure a free passage from the bladder. Those patients who form the exceptional instances to this rule are for the most part weak in constitution, have little reparative power, or are subjects of some chronic disease in addition to stricture of the urethra. The management of these is mostly that of the next class.

2. Cases in which the fistulæ pass through tissues indurated and deformed by inflammatory exudation.

In these cases also, the primary object is to dilate adequately the stricture, and to observe the effect induced. In some of them it is sufficient to enable the fistulous passages to heal slowly. Dilatation, however, having been maintained for some time, and little or no benefit having resulted, it will be desirable either to stimulate the walls of the fistulæ themselves, and so bring about adhesion of opposing surfaces; or to lay them open, in order to produce recent and healthy wounds, so that they may heal up soundly from the bottom. At the same time we must attend closely to the patient's general health, seeking to maintain the secretions in a natural condition. Various agents have been employed to stimulate the indolent fistulæ; one of the best is the concentrated tincture of cantharides, applied on a camel's-hair brush, or with a fine syringe. Solutions of the sulphate of zinc or copper, and of the nitrate of silver, injected by the syringe, sometimes give good results. An excellent mode is to introduce carefully, as far as possible, a small and flexible silver probe, coated with nitrate of silver; a plain probe having been introduced beforehand as a guide to the length and direction of the passage. It often happens that the external orifice of the sinus is smaller than any other part: in such a case, a little caustic potash should be applied for the purpose of enlarging it, and so permitting free removal of the discharge.

The application of compression to the fistulæ has been tried

several times, and success has been claimed for it in two or three cases; in one of which, all ordinary means having failed, a cure was obtained by making the patient apply firmly to the perineum an india-rubber ball, inflated with air, on every occasion before making water, and for some minutes afterwards. This plan was studiously followed during fifteen days, when the opening had soundly cicatrised. Four months after, the patient was perfectly well.

The cure of obstinate urinary fistula has also been attempted by introducing a catheter, and permitting it to remain in the urethra for days together, on the principle of insuring, as it has been supposed, the passage of the urine through the instrument, and thus preserving from irritation the fistulous passages. Little, however, is thus gained; for experience shows that, however large the instrument may be, and however closely it may fit the urethra at the present moment, before thirty-six hours have elapsed it will lie loosely in the canal, and urine will pass by its side. It is not possible, indeed, to remove urine from the bladder for any lengthened period by this means without inducing suppuration in the urethra, which is a bar to success.

It is better to withdraw all the patient's urine, by introducing a catheter three or four times a day, or more frequently if necessary. When the patient can do this cleverly for himself, and thus insure, at every want to pass urine during a period of several weeks, that it is carefully removed without contact with the urethra, a most successful result may be hoped for. I have had an instance of this under my own care, where a large portion of the floor of the urethra being lost, the opening closed after three or four months' attention to this practice.

Free incisions through the fistulæ, down to their origin in the urethra, or nearly so, have been found successful in inducing a healthy process of granulation from the bottom of the wound, and thus in ultimately closing the unnatural passages; provided always, however, that there is no obstruction to the free egress of the urine by the urethra, otherwise no such measures can be of any service. In some cases in which external division of the stricture on a grooved staff is indicated, this operation may be performed in such a manner as to include the fistulous opening in the incision, in which case a successful result may generally be reckoned on. Old chronic perineal fistulæ are sometimes better dealt with by the galvanic

cautery than by any other method. Another obstinate species of fistulous passage, communicating with the prostatic urethra, sometimes follows the operation of lithotomy; this also may be often successfully treated by the introduction of an iron wire heated to intensity; the best means of effecting which is undoubtedly the galvanic current, since it maintains as well as produces the required temperature.

A fistulous passage is sometimes prevented from closing by the presence of a small calculus in some part of its course. It may be a small concretion which has escaped from the bladder, or a deposit from unhealthy urine while passing through the sinus. Sometimes it occurs in fistulæ connected with diseased prostate, or with a prostate which is the subject of calculous formations. Fistula associated with the prostate may, however, exist in the absence of any of these causes; such as those which are sometimes connected with prostatic abscess, and which are generally exceedingly obstinate and irremediable.

Fistula is not necessarily a continuous passage from the urethra to the surface; opening from the urethra at one end, it may have a blind or cæcal extremity: hence 'blind urinary fistula' has been described. A small tumour, originally formed by a collection of matter, and having a communication with the urethra, constitutes the general form. Its origin has been variously accounted for; some believing it a result of stricture, others of inflammation of the mucous follicles of the urethra. Until the tumour is opened externally, it will not disappear, when it becomes a fistula of the ordinary kind, requiring treatment already indicated.

Urethro-rectal fistulæ sometimes occur as a consequence of stricture or abscess, and more rarely, perhaps, vesico-rectal. In either case their existence is usually first announced by the passage of liquid in an unusual manner and quantity by the anus; sometimes by the appearance of feculent matter in the urine or by the urethra.

I have recently treated a case of urethro-rectal fistula in a gentleman of about twenty years of age, in whom it resulted from acute prostatic abscess following gonorrhœa. The urine passed freely by the anus for some weeks: after trying various means, the use of the catheter, &c., without success, I discovered that by placing him flat on his face during the action of micturition, almost all the urine passed by the urethra. This plan was uniformly adopted for three months. Urine ceased to pass altogether by rectum in a month, and in two more he could pass it in the upright position without any escape. The fistula has remained soundly healed ever since.

If such a plan does not succeed, the actual cautery, and particularly that heated by the galvanic current, affords the best chance of success. It may be applied through the rectum; a speculum having been first introduced, and a full-sized sound carried into the bladder. On the day before the operation the bowels are to be freely purged, and cleared by an enema an hour or two before the application of the cautery, after which they must be prevented from acting for two or three days. The cautery is to be re-applied two or three times if necessary. If the opening is large, the edges should be pared and brought together by silver sutures, after the method of Dr. Marion Sims.

3. The third class of urinary fistulæ comprehends those unnatural openings into the urethra which depend upon actual destruction of substance from the walls of the urethra and superjacent parts. The common causes of these are, sloughing from extravasation of urine, simple and phagedænic ulceration, and mechanical injuries of various kinds. They are for the most part larger, although not invariably so, than any of those already referred to. Generally a portion of the floor of the urethra is destroyed, as well as the structures which have intervened between it and the external surface; so that in many cases more or less of the mucous membrane of the upper aspect of the canal is visible from the outer orifice. As a consequence the whole, or nearly the whole, of the urine passes by the artificial channel in a full stream. Such apertures may be found before the scrotum, and are called ante-scrotal or urethro-penal fistulæ; or in or behind the scrotum, known simply as scrotal and perineal fistulæ.

A broad distinction exists between these two classes in relation to their amenability to treatment, and to the nature of the operations necessary to their cure. Ante-scrotal urethral openings are the most difficult to close, since the coverings of the urethra here possess but little substance. And further, owing to the extreme mobility of the member, and its liability to erection, it is difficult to maintain that perfect steadiness of position so desirable in a part which is the subject of an autoplasmic operation. Nevertheless, with all these difficulties, in addition to that formidable one, the contact of urine, such openings, even when large, are not to be regarded as beyond the reach of surgical skill. The exercise of considerable patience, and of unremitting attention during a long period, is indispen-

sable both to the surgeon and to the patient in a case of urethro-penal fistula requiring a plastic procedure for its cure.

Openings in the perineum involving loss of substance, although by no means easy to close, are remediable with less difficulty, on account of the absence of the conditions just adverted to as the obstacles in ante-scrotal fistulæ.

Treatment.—Ante-scrotal fistulous openings which are of small size, but obviously depending upon loss of substance in some degree, have been closed by repeated applications of a caustic agent to their edges and to the surrounding parts. Of these the nitric acid, the nitrate of silver, or the strong tincture of cantharides have been successfully employed in very small openings.

Dieffenbach used the cantharides, and sometimes combined with it what he called 'the lace suture.' This he describes as applicable to small fistulæ in the anterior part of the canal. The margin of the fistula and the surrounding skin must be frequently touched during the day previous to the operation with the strong tincture of cantharides. Before using the suture, the loose epidermis is to be removed by scraping, and a sound introduced into the urethra beyond the opening. The operator is then to take 'a small curved needle, sharp at the point, but not at its edges, with a stout silk waxed thread, and by means of a needle-holder to introduce it beneath the skin at about three lines from the border of the fistula.' The point of the needle is to be carried deeply, but not into the urethra, and made to emerge at another point, about three lines from the margin of the fistulous opening. By three or four of these stitches, the thread is to be carried round the opening, until it finally emerges at the point at which the needle was originally entered. The thread is imbedded in the cellular tissue around the fistula, at about three or four lines' distance from it. Its ends are now to be drawn together slowly, so as gradually to approximate the borders of the fistulous orifice until it is obliterated, and then to be fastened by a knot. In three or four days the ligature may be divided and drawn away. Most commonly, however, some plastic operation is necessary for these openings. The proceedings of Cooper and Earle, the first on record, are the basis of many various proceedings since employed. Space does not admit of the necessary description here. Two of the modes employed for ante-scrotal opening are described in the Essay on PLASTIC SURGERY: I have given

in the second edition of my own work on Stricture and of *Fistulæ* in some detail.

Retention of urine.—By this term is generally understood a condition in which the patient is unable to pass any urine whatever, or only an extremely small quantity; and is in danger of serious consequences if not relieved at no distant period. Urine, however, and often is, habitually retained in the bladder, while the patient nevertheless passes as much in the twenty-four hours as his kidneys furnish, the reservoir remaining more or less full. This is engorgement of the bladder, and is not retention, the term is held to indicate an acute and not a chronic condition.

Retention of urine may be caused by simple prostatitis, usually occurring in young and middle-aged subjects; by stricture of the urethra, usually in middle life; and by hypertrophied prostate in elderly persons only.

In the consideration of the first and second forms will come in the proper order here.

Complete retention.—When any degree of organic stricture exists, however slight, retention of urine is always possible if the influence of certain exciting causes takes place. Obstruction of the passage may be caused by inflammatory enlargement, or unwonted muscular contraction; more frequently, however, by a combination of both. Sometimes it may be caused by a foreign body, as a small calculus, a portion of a stone, or the like, obstructing the stricture; and this is probably the rarest form.

In treating retention, the catheter is the first means required. In nine cases out of ten, and often the only means. The first point to be ascertained is, is it a case of merely temporary obstruction by inflammation following gonorrhœa, or does it depend on permanent stricture of some standing? The next points are the duration of the attack and the condition of the bladder, bearing in mind that while some patients may exhibit distension of the bladder almost to the umbilicus, the subject of an old stricture may be in a state of greater danger, although no distension is perceptible above the pubes, owing to the contracted condition of the bladder, which has become natural to him. The treatment is somewhat different in the two cases just briefly described. In the case of acute, or that of temporary inflammatory obstruction, it has been recommended to employ baths, opium, and depletion before

using the catheter; and relief may certainly be afforded in this manner, although at the expense of prolonged suffering to the patient. The object of this plan is to avoid injury to the urethra in its inflamed condition; and if the surgeon is not gentle, or is not skilled in the use of the catheter, it is probably the safest course to follow. But if he employs a light and careful hand, a catheter of rather small size, say No. 4 or 5, of gum-elastic or silver, the former producing less pain, should be at once passed into the bladder; and in these cases this is generally easily accomplished, if the flexible instrument retains a good curve after the stilet is withdrawn. If the surgeon does not succeed in passing an instrument, the patient should be put into a hot bath, 100° to 103° or 104° , to relax spasm and relieve internal congestion; a full dose of opium should be given by the mouth; and if there is much straining, an enema or suppository containing opium may be also administered with advantage.

If there is much inflammation about the parts, as shown by a swollen penis, urethral discharge, and tender perineum, cupping or a dozen leeches on the latter region are decidedly useful, in the event of retention persisting; generally, however, some relief follows the use of the bath and the opium, which latter may be repeated two or three times, to insure its full action on the system. There are very few cases among those here regarded as inflammatory or spasmodic that will not yield; sometimes slowly without further catheterism, sometimes by means of an instrument, which may often be passed with comparative ease after the treatment described.

There are some instances where the difficulty seems more spasmodic than inflammatory, in which repeated doses of the tinct. ferri sesquichl., that is to say 15 or 20 minims every quarter of an hour, administered four or six times, relieve the retention. When, on the other hand, we have to deal with retention resulting from confirmed organic stricture, it is better at once to examine the urethra in the usual way, that is, with a full-sized instrument; and, having ascertained the locality of the obstruction, patiently to devote a long time to the careful use of the smallest catheters, silver and gum. In relation to its management, see pp. 957, 958, on Catheterism in difficult cases. If this proves unsuccessful, the hot bath must be resorted to; and after faintness has been induced, the catheter may be again employed while the patient is still in the bath.

Further treatment, if necessary, must depend upon the condition of the patient. Generally, opium should be given by enema and by mouth, after which the catheter is again to be used. The time which is to be devoted to the employment of all these means must be regulated by the judgment of the surgeon. The condition of the patient, and a knowledge of the time during which absolute retention has existed, will enable him to decide the question of affording relief by some other measures. Chloroform is an agent of great value in obstinate retention. It relaxes the muscles; by causing insensibility it overcomes involuntary resistance and straining, and thus enables the surgeon to use the catheter with greater advantage, although not with more force than without its influence.

But supposing none of the treatment hitherto employed has been successful, that the patient's condition is urgent, the next proceeding is to afford relief by making an artificial opening either into the bladder or the urethra.

To effect this, the following operations are employed :

The stricture may be 'forced;' or, the urethra may be opened at or behind the stricture; and the bladder may be punctured, either by the rectum, above the pubes, or through the pubic symphysis.

1. The first method, or 'forcing' a stricture, is rarely performed now with deliberate intention. It consists in driving a catheter of moderate size forcibly through the obstacles which oppose its course, in the direction of the bladder, which when reached is evacuated, and the instrument is tied in. I have seen it successful in old, hard strictures. But it is not a proceeding which can be recommended in competition with those less harsh and dangerous proceedings which are generally practised by modern surgeons.

2. The urethra may be opened, first, by a dissection from the surface to the stricture, and through it, according to the method already described as perineal section (p. 974); secondly, the urethra may be opened behind the stricture, the bladder relieved, and the obstruction divided subsequently. Both these are difficult operations. Especially when the perineum is thickened and indurated, it is by no means always certainly possible to dissect through the stricture from the front, or to hit the urethra behind the obstruction. The perineal section, under such circumstances, often fails to trace the narrowed channel, and sometimes even to discover the urethra at all; and

a prolonged operation, with much loss of blood, not unfrequently results in failure to relieve the patient.

The method of opening the urethra by an incision made altogether behind the stricture was recommended in this country by Guthrie and Liston, but is very rarely performed, the puncture of the bladder per rectum being now generally preferred. The operation is performed by introducing the left forefinger into the rectum, to distinguish the apex of the prostate, and entering a straight sharp-pointed bistoury with the right hand just in front of the anus, so as to hit upon the urethra at the point indicated by the finger in the rectum. The knife is carried upwards exactly in the median line, so as to make an incision of one or two inches in length. In this manner the urethra is to be sought, and the stricture afterwards divided forwards or not, as the operator determines to be most desirable.

3. The bladder may be punctured in two principal ways, viz. by the rectum, and above the pubes; to which may be added a third, through the pubic symphysis. Formerly it was also punctured by a direct stab from the perineum—a dangerous proceeding, now wholly obsolete.

Puncture by the rectum is the operation most commonly adopted at the present day. Its results have been carefully studied of late years, and have been brought before the profession more prominently by Mr. Cock, who has done much to test its value. The result of his conclusions is, that it is fraught with less danger, and is more easy of performance, than any other which is adopted for the relief of retention. The objections made to it are: the liability to abscess between the rectum and the bladder; a persistent fistulous opening there; injury to the seminal vesicles, leading to suppuration of the testicle; and the danger of perforating the peritonæum with the trocar, and thus inducing inflammation of that membrane and a fatal result. The last-named, and the most serious, of these accidents is happily that which may most easily be avoided. After an examination of several cases by dissection, it is my opinion that the peritonæum is out of reach of the trocar when employed with a moderate degree of care. The puncture usually falls short of the peritoneal pouch by an inch at least. The other results do occasionally, but only rarely, occur.

The operation is thus performed:—The rectum being emptied by an enema, the patient is placed in the position for lithotomy,

firmly held by two assistants. The operator then introduces the left forefinger, oiled, into the rectum, and defines the posterior margin of the prostate, beyond which the finger should be extended if possible. Fluctuation should be obtained there, the contents of the bladder, by a tap made on the hypogastric region, unless the viscus be very contracted indeed, in which case the propriety of operating is doubtful, since the finger may enter the opposite coat of the bladder, from absence of the requisite amount of distension. Having determined the point at which fluctuation is most distinct, and directed an assistant to place a hand on each side of the supra-pubic region to support the bladder, a well-curved trocar, seven or eight inches long, is carried along the finger in the middle line to the point indicated, the handle well depressed, and the point carried through the coats of the rectum and bladder, until it is felt free in the cavity of the latter. The stilet is withdrawn, and the canula, or a flexible tube passed through it, retained by a bandage and tapes. The length of time it should remain will depend on the amenability of the stricture to treatment. If this fails, the urine will pass through the natural channel, and the opening in the rectum will readily close. Little fear need be entertained of the continuance of a fistulous opening; for on several occasions on which the canula has escaped by accident, it has been impossible to replace it, and a fresh puncture has been necessary. When it is impossible to find fluctuation by the rectum, or when the prostate is enormously enlarged, this operation, as a rule, is not to be employed.

Puncture of the bladder above the pubes is performed as follows: the patient being placed in a half-reclining position, and the pubes shaved, a vertical incision of the integument is made directly above the symphysis pubis, about an inch and a half or two inches in length at the surface; this is to be carried downwards through the linea alba, so as just to admit the tip of the trocar to reach the distended bladder. Meantime an assistant, standing behind the patient, places one of his hands on either side against the abdominal walls, so as to steady the bladder. A straight or a slightly-curved trocar (if the latter, the convexity of the curve should be upwards) is then to be carried with a little inclination downwards into the bladder. After the operation the canula should be exchanged for a silver tube, specially adapted to slide through it, secured by tapes and a bandage, which may remain a day or two until lymph has

been effused upon the edges of the wound, when it may be withdrawn, and a gum-elastic catheter worn in its place, an instrument which is generally better tolerated by the bladder than one made of metal. For most cases of retention in which this operation is required, this mode of performing it will probably be still the most desirable to adopt.

But a very ingenious proposal, originating in Paris last year with Dr. Dieulafoy, and since twice realised on the living patient, must not be left without notice. Dr. Dieulafoy has applied his 'aspiration' apparatus to the distended bladder successfully in the following manner. He inserts a very fine trocar, no larger than a small exploring needle, close above the pubes into the tumour; if it reaches the bladder a drop or so of urine may issue; he now applies the vacuum instrument to it, and a strong jet of urine instantly appears and soon fills the instrument; this is discharged by a side tap, a new vacuum is created by a stroke of the piston, and the process is repeated. In this manner three or four pints of urine may be withdrawn by this minute channel in fifteen or twenty minutes, and the trocar withdrawn with a scarcely perceptible puncture. I have never yet applied this apparatus to the bladder, although I have proved its exceedingly perfect action twice in paracentesis thoracis. If one evacuation of the bladder were sufficient, this very simple operation would suffice, but if a route for succeeding acts of micturition is required, as is almost always the case, a larger opening is necessary. Still the large trocar generally employed is by no means necessary: thus my own trocar for puncture above the pubes is only equal to a No. 6 catheter in size, and it is sufficiently large for all purposes. By means of a stilet lengthened by screwing to its end an additional length, forming a guide, a gum-elastic tube may be slipped into the bladder in place of the trocar and canula when these are withdrawn.

The puncture through the symphysis pubis was first proposed by Dr. J. M. Brander of Jersey, in 1825, and several successful cases have since occurred in the practice of himself and others. He has usually employed a hydrocele trocar of medium size for the operation, which is thus performed. The patient should recline, and the trocar is introduced—whether after a small preliminary division of the integuments or without it, appears to be immaterial—about the centre of the symphysis, reckoning from above downwards, and in a direction at about right angles to the vertical axis of the body. Dr. Brander says, 'somewhat ob-

liquely downwards and backwards towards the sacrum, varying the direction according to circumstances; a piece of flexible catheter is then to be introduced through the canula,' and retained by a tape. I attempted to perform this operation some time ago, and could not penetrate the symphysis, although the trocar was strong and not large in calibre. With more improved methods of supra-pubic puncture, as above referred to, and by rectum, I should think this mode should not be in future considered, certainly in the cases of elderly persons where the cartilage is very firm or ossified.

In considering these methods of affording relief to the distended bladder in reference to any case which requires an operation, the question to be first solved is the following:—

Are the patient's powers and condition such as to compel us to prefer the simplest method of affording immediate relief, without regard to ulterior results? If so, unless the urethra can be felt in the perineum distended with urine, which may sometimes be the case, especially if that region be not thickened or deformed, the rectal puncture of the bladder, supposing the prostate not to interfere (putting aside the puncture by the symphysis, as not yet sufficiently tested by experience to be considered on equal terms with the older operations), is generally the simplest method, as it affords instantaneous relief at the smallest possible expense to the patient's powers. But if it is probable that the canula may be required for a longer period of time following the operation, I then prefer the supra-pubic method, especially if the bladder is distended, as it is inconvenient and distressing to a patient to maintain a canula in the rectum for more than a few days; whereas I have seen patients wearing with perfect comfort a tube for many years above the symphysis pubis. Again—but this is rare—if distension of the urethra is perceived in the perineum, a lancet or sharp-pointed bistoury may be carried into it, and a female catheter introduced by its side before it is withdrawn; or if the powers of the patient are good, and the condition of the parts is natural, perineal section may be employed. Having relieved the bladder, we can make a careful attempt to divide the stricture, by introducing a fine probe from below upwards through the stricture, and passing a catheter into the bladder. If, however, induration and deformity of the perineum exist, as above indicated, the crisis of retention is not generally a favourable time for adopting this proceeding.

The treatment of the prostate.—When the disease is due to local causes, it is not to be treated only after middle age, but at an earlier period, if much obstruction exists, or if there has been much excess in wine and spirits, or if there has been much intemperance. When necessary; for the removal of the stone, measures already mentioned, such as the use of the hot bath and of the catheter, are at once to be resorted to. A gum catheter, or a catheter with a stilet, will often be sufficient to have been kept in the urethra in order to improve the passage of the stilet, which is often scarcely possible with the ordinary stilet. But in cases where the stone is large, it is necessary to employ a silver catheter of a size larger than No. 9. The catheter is to be from the rings of the catheter, and the curved catheter is to be of the third of a circle, and of a diameter; the most useful size.

In passing the catheter, the catheter with is usually sufficient. When it arrives at the neck of the bladder, it requires to be turned back, and to count a beaked catheter, will be sufficient. Occasionally a silver catheter of essential service. The use of the instrument in practice a matter of course, having been or not, which may be the case. The stilet, having been passed, is then withdrawn.

sing the curve and elevating the point of the catheter, so
 en to carry it over the enlarged portion in a manner less
 accomplished in any other way.

e question occasionally arises, is it desirable at once to
 ate the entire contents of the bladder when retention has
 d for a considerable period of time? In very rare instances
 removal of a large quantity of urine, amounting to several
 , especially when an upright position of the patient has
 incautiously permitted, has been followed by fainting and
 session, from which he has never rallied. When the extent
 esical dulness is very considerable, it is therefore prudent
 ford relief in a gradual manner; and, supposing that the
 eter is retained, this may easily be accomplished. The re-
 al of some thirty or forty ounces will probably afford com-
 ease, and after the lapse of half an hour or an hour another
 ion may be withdrawn; in this manner the bladder may be
 dually brought to adapt itself to the normal condition of
 traction, which subsequently, as a rule, must be insured at
 t once or twice a day. It is not long since my evidence was
 ired in a court of law, in a case in which death had occurred
 antly after six pints of urine had been suddenly withdrawn
 he upright position.

point of importance remains. A catheter having been in-
 uced with difficulty for the relief of retention, should it be
 mitted to remain? The answer must depend on circum-
 ces. In support of the negative, if the parts are already in
 ate of considerable irritation, it is undesirable to permit any
 uces of adding to it, of which the presence of a catheter may
 ne. On the other hand, the bladder, after long retention,
 very soon fill again, and less hazard may be incurred by the
 ence of the instrument than by a repetition of the efforts to
 e it there, especially if great difficulty was experienced in
 ing it in the first instance. Greater weight usually attaches
 e latter consideration, as a rule, than to the former. I
 e often seen great danger incurred by the too early re-
 al of a catheter which had relieved an urgent attack of
 ation.

, however, which very rarely happens, catheterism proves
 ecessful, and relief is necessary, the bladder may be punc-
 d or the prostate perforated. For the methods of per-
 ing the former operation, see page 992. If the prostate
 ery large, the rectum is not so convenient a situation for

the proceeding as the supra-pubic region, which is therefore generally adopted. I have punctured by rectum with the best results for retention from enlarged prostate, but then a fluctuating point was within reach of the finger, in which case it would be safe; this condition, however, is to be regarded as exceptional.

Perforation of the obstructing portion of the prostate may be performed with a strong silver catheter, about No. 9 or 10 in size, and rather longer than the ordinary catheter, but not possessing the large curve of the full prostatic instrument. The operator introduces this to the seat of obstruction, and satisfies himself by means of a finger in the rectum that it lies fairly in the urethra, and is engaged in the prostate; he then steadily carries it onward towards the cavity of the bladder, by pressing the point firmly forwards, and at the same time slowly depressing the handle; and he desists on feeling that the point is free in a cavity, and on finding that the urine flows through the instrument. This was done by Home and Brodie. Liston employed a cutting stilet 'carried through a slightly-curved and long canula,' and 'practised the operation a few times successfully.' Whatever be the instrument employed, the surgeon is careful to maintain it in the middle line, and also to aim at making the point emerge just behind the neck of the bladder, neither too near the pubes on the one extreme, nor the posterior wall of the bladder on the other. The catheter or canula, as the case may be, should be retained not less than forty-eight hours afterwards in the bladder, that the tissues around may consolidate, and no difficulty be experienced in replacing it by a catheter when withdrawn.

Extravasation of urine from rupture of the urethra may take place during unrelieved retention of urine by the giving way of the urethra at some point; it is very rare indeed that the bladder itself is ruptured. In either case, however, mechanical distension is not the direct nor the only cause. Ulceration of the mucous membrane behind the stricture, increased by the influence of urine in almost constant contact with it, causes a breach of the urethral walls. The bladder contracting drives the urine with great force into the cellular interspaces of the superficial fascia of the scrotum and abdomen, when the rupture occurs anterior to the membranous portion, as is almost invariably the case. The situation of the swelling and inflammation

is characteristic in these cases, because the course of the extravasated urine is defined and limited by the union of the superficial with the deep layers of fascia at certain points. Thus the scrotum is first gently distended; a bag, as it were, being formed by the union of the superficial and deep fasciæ across the perineum, and by their connection with the pubic rami. As the fluid increases, it rises into the coverings of the penis and over the pubes and abdomen, but does not descend to the thighs, because in the line of Poupart's ligament these fasciæ are similarly blended, and prevent the passage of urine beyond their junction. The consequences of this extravasation are extremely disastrous; an unhealthy inflammation results from the irritating fluid wherever it goes, and the areolar connections of the skin and subjacent tissues are destroyed. If unrelieved, the distension becomes enormous, the scrotum sometimes attaining the size of a human head; the skin assumes a dusky red hue, or becomes purplish and livid; and gangrenous spots are sometimes apparent; not unfrequently such a one is seen on the dorsum of the penis, and is generally regarded as a very unfavourable sign, although by no means one that is necessarily fatal. The general condition of the system is one of great depression; and unless complete relief be afforded, the patient must inevitably soon succumb. He is frequently delirious, with brown tongue and extremely feeble pulse, when the extravasation is considerable.

In these circumstances it is necessary at once to make free and depending incisions into the distended parts. The scrotum especially demands a deep incision on either side the middle lines, frequently the pubes, and the penis itself also. These incisions will not only free the parts from the extravasated urine, but provide for its passage from the bladder. By these openings the purulent matters, débris, and sloughs will subsequently come away, often in very large quantity. No immediate measures for the cure of the stricture are necessary, as in the present state of the patient they are neither practicable nor advisable; although it is probable that when the retention is relieved a catheter may be passed into the bladder by the urethra; however, there is no occasion to do this until the system has rallied, which it often does to a marvellous extent. In a few hours the sufferer may emerge from a state of utter prostration to one of comparative comfort and promise; the symptoms of depression and exhaustion sometimes disappearing



with the scrotum to a greater or sometimes completely stripped bare in the wound, and even has process the removal of the pro cleanliness of the parts must applications, frequently change tices, linseed-meal or charcoal chlorides of lime or soda well cations. The use of the disinfectant or of powdered charcoal, and for be adopted.

If the extravasation have taken of the deep perineal fascia, swelling may sometimes, but not perineum. This is to be at once behind the fascia altogether—finds its way upwards around the fatal result is inevitable.

Rupture of the bladder.—This except as the result of direct violence from a blow on the abdomen or distended with urine. It may be bone when the pelvis is fractured

The symptoms of vesical rupture take place after a prolonged but not necessarily absolute retention, for some surplus of urine may have been previously escaping by the urethra. The patient usually states that he has felt something give way. Acute abdominal pain then sets in; the belly becomes exceedingly tender and distended; the features are pinched and anxious; the breathing hurried; obstinate hiccough occurs, sometimes vomiting; the pulse is sharp, quick, and irregular; the flow of urine ceases altogether, as also does the straining to void it. General fluctuation may be sometimes found in the abdomen, and inordinate distension of the bladder, before felt in the rectum beyond the prostate, has now disappeared. Sometimes the patient is delirious, and even maniacal. And after a period varying from thirty-six hours to four or five days from the time of the accident, during which the patient's agonies are extreme, death takes place.

The indications of treatment which, in the absence of experience, we should endeavour to fulfil, would be as follows: To provide for the free exit of the urine from the bladder by puncture; to alleviate suffering by large doses of opium, with hot fomentations and rubefacients to the abdomen; and to support the powers of life. Whether an attempt to remove the urine in case of extravasation into the abdominal cavity, by puncture of its walls, should ever be entertained, is an unsolved question. Such a proceeding affords the only chance (exceedingly slender as it is) of recovery which surgical aid could afford.

HENRY THOMPSON.

URINARY CALCULI AND LITHOTOMY.

OF all the ailments to which the human frame is liable, that of the formation of a stone or calculus in the kidney, bladder, or urinary passages is undoubtedly one of the most formidable and important. And on account of the tendency of this formation to become lodged in certain dilated portions of the urinary apparatus behind natural necessary narrowings, it is thereby rendered liable to increase in size, and consequently soon to become impassable and incapable of expulsion. The stone now acts as a foreign body, and irritates the delicate structure of the part in which it is retained, causing at first local disturbance and mischief, afterwards seriously involving the constitution, and ultimately leading to a miserable existence, which most frequently terminates in a lingering death.

In order to comprehend more fully the subject in all its bearings, it behoves us to take into consideration the causes and formation of stone; and in doing so we must necessarily review cursorily the circumstances which may give rise to this affection, by passing in survey the more simple and slight deviations of the urinary fluid, then its more aggravated changes from the condition of health, until we arrive at those still more complex and dangerous forms which cause the amalgamation of the deposits into a mass, constituting a stone.

But previous to these inquiries, it is the essential duty of the surgeon to be well acquainted with the urine in a state of health, and in its varied conditions under every circumstance. He should know its chemical constitution, its specific gravity, its amount of acidity, its variety of colour, its microscopical appearances, &c.; and not only this, but he must also bear in mind the changes which the urine undergoes during certain conditions of the body, and during certain periods of the day;

thus there is the urine of the blood, passed early in the morning before any repast; the urine after the use of drinks; and the urine after the partaking of solids—all modifying the condition of the constituents. All these points are carefully detailed in our standard chemical and physiological works, and in the numerous special monographs on the urine, to which the reader is referred.

In making a careful analysis of the constituents of the urine, we find that there are two sources from which an abnormal condition may arise; and that an excess or deficiency of the elements from either of these sources may disturb the balance, and thus dispose to a precipitation of one or other of the constituents, causing deposits. The one source is from the organic elements, and of these urea and uric acid are the most frequent. Urea bears the proportion of 12 or 15 to 30 or 40 parts in 1000 parts of urine; it is exceedingly soluble and liable to decomposition with a development of ammonia, which is accelerated by mucus or other animal substances becoming putrid; this may take place in the urine in the bladder under certain diseased actions, and prove an abundant source for calculus. Uric acid, on the contrary, exists in very small proportions,—half a gr. to one gr. in 1000 grs. of urine; it is very insoluble in water, so that any excess of living, such as increased animal diet, will cause a ready deposition of this substance; hence the frequency of deposits and calculi of uric acid; and from this being the chief constituent of most stones, the deposit has derived the name of lithic acid. The other source is from the inorganic constituents, the salts of the urine; these comprise saline and mineral ingredients. The saline are soluble in water, and consist of sulphuric, phosphoric, and hydrochloric acids, in combination with their bases, potash and soda. The mineral salts are insoluble in water, and consist of phosphoric and occasionally carbonic acids, in combination with their bases, lime, magnesia, and sometimes alumina; traces of silica are occasionally met with. Of these salts the phosphates are the most important, as being more readily prone to be deposited.

*Urinary deposits.**—These may be advantageously arranged

* In giving a description of urinary deposits and urinary calculi, the author has availed himself of the standard works: for more minute detail reference must be made to the works of Bird, Beale, Thudichum, Bence Jones, the Catalogue of Calculi of R.C.S. England, Hassall, &c.

products, such as blood, pus, mucus, milk, fatty matter, &c.; and the highly doubtful origin, viz. cyanourine, melanic blue. Two rare forms of deposit, the first will be alluded to under the head of calculi.

1. The *uric acid deposits* may be simply a lateritious sediment, composed of various

FIG. 300.



Uric acid.

these crystals, however, may collect in masses, producing a kind of sand, of about one-fourth of an inch in diameter, and mostly globular. Such masses may concrete and form a

The crystal is much affected by the strength of the acid which is added; thus, where the acid is small in quantity, the crystals are regular, mostly tables, squares and lozenges; where the acid is in larger quantity, added to a strong solution of urate of ammonia, they consist of large and long tables, with very elongated lozenges; and where the acid is strong, and ammoniacal urate itself used, the crystals most frequent are acicular prisms. The causes which give rise to this deposit are described by Dr. G. Bird* to be: 1, the waste of tissues being more than the supply, as in fever, rheumatism, &c.; 2, the supply of nitrogen in the food being greater than is required for the reparation of the tissues, as in over-indulgence, especially in the use of animal food; 3, the process of digestion being inefficient to assimilate an ordinary and normal supply of food, as in dyspepsia; 4, obstruction to the cutaneous outlet for nitrogenised secretions, as met with in diseases of the skin, invariability of the climate, &c.; 5, congestion of the kidneys, following injury or disease of the organ. Imperfect respiration is also said to be a cause of an undue proportion of uric acid in the urine, owing to an excess of carbon and oxygen uniting to form urea. The treatment to be adopted will of course be to control and remedy the cause by abstinence from over-indulgence in animal food, regular diet, exercise, free action on the skin by baths and moderate diaphoretics, mild diuretics, diluents, and liberal use of good water, and so on.


The *urates*, generally known as the urates of ammonia, form the most frequent urinary deposits. Heintz† regarded these deposits as consisting of urate of soda and urate of ammonia, with traces of urates of lime and magnesia. Their colour is a pale fawn, but it may vary from snow-white, through every shade, to brick-red, pink, or purple. Dr. Bird remarks, that in the snow-white, fawn, or brick-red deposit there is a deficient state of the cutaneous functions; whilst in the pink, crimson, or purple variety, there is more or less evidence of functional or organic derangement of the liver, spleen, or other organs influenced by the portal circulation, which is attributed by some to the presence of purpurine, and by others to the obstructed excretion of carbon, as in rheumatism, gout, and diseases of the liver or spleen. The alkaline urates generally form an amorphous deposit of minute spheres, having protruding there-

On Urinary Deposits, 5th edit. p. 159.

† *Lehrbuch der Zoochemie*.

from acicular spiculæ of sharp crystals of uric acid. Some describe two forms of fawn-coloured deposit of urate of ammonia: the one, in which the precipitate mixes readily with the supernatant fluid by slight agitation, appears to form a very homogeneous mixture, and subsides exceedingly slowly by repose; the other, with frequent admixture of crystals of pure

FIG. 301.



Urate of ammonia.

uric acid forming a pale urine, with diffusion of the deposit through the fluid in detached bran-like particles, which readily subside, and occasionally form dense clouds in the fluid, like puriform mucus, disappearing on heating; the urine is of low specific gravity. Sometimes the deposit becomes quite white, and often assumes a mucilaginous or jelly-like character.

The causes and treatment are, for the most part, those of the uric acid deposit. Open air, healthy residence, and tonics, either mineral or vegetable, are essential.

3. The *uric oxide* deposits, otherwise called *xanthic oxide* or *xanthine*, closely resemble uric acid, but the conditions of the urine depositing them are unknown. It is supposed that these depend upon an imperfect oxidation of the materials from which uric acid is eliminated by the ordinary processes of the kidney.

4. The *oxalates* are usually observed in the form of oxalate of lime, the crystals consisting of two varieties. One of these is the quadratic octahedron class, which generally have one axis much shorter than the other two. Their varieties in shape are due to the position in which they are viewed, some even being

square, and resembling the back of a letter e other variety is the dumb-bell-shaped crystal: erer by many to be formed by oxalurate of others to consist of uric acid, both of which explained in Dr. Beale's work, pp. 300 et seq. crystals often unite into a mass, and form the concretion called the hemp-seed calculus. Dr. erer these deposits to be due to an increased he uric acid formation, and believed that they healthy urine, although their constituents are one form or another. Lime is always present in bined with phosphoric and perhaps other acids; the insolubility of the oxalate of lime, and so

FIG. 302.



Oxalate of lime.

tual affinity of its constituents, that the addition as a soluble oxalate instantly determines the prehe salt. The particular manner in which the he oxalic acid found in the urine occurs can garded as at all satisfactorily explained; most, to regard it as a secondary product of the oxida-charine matter occasionally present in the secre- appear inclined to draw a close analogy between n which sugar and that in which oxalic acid is r. Prout maintained that oxalic acid is one of imperfect digestion, and that it is derived from , more especially such as contains sugar. Liebig arises from the imperfect oxidation of uric acid amorphosis of the living tissues; if the supply of o a certain extent, the products of combustion

in the lungs are urea, oxalic acid, and the substance called allantoin. Dr. Owen Rees* considers that the oxalate of lime concretion is formed by the decomposition of uric acid, or of an urate in the kidneys or bladder; these are converted into oxalic acid, which unites with the lime, so that the oxalitic diathesis is an accidental and unimportant modification of the uric acid diathesis.

The causes of this condition have been attributed to great nervous depression, as from grief, loss of blood, malaria; and of course the treatment will be to counteract these causes, and to pay great attention to diet and the action of the skin; as also to avoid all articles containing or liable to produce sugar, as fruits, fermenting liquors, and substances into which oxalic acid enters as a constituent; mineral acids are highly beneficial.

5. The *cystic oxide* deposits, or cystine, are considered to be modifications in which the urea and uric acid are replaced by an excessive elimination of sulphur. The deposit is described to

FIG. 303.



Cystic oxide.

be bulky and easily diffusible, resembling in appearance the white or pale lithates, and to consist of more or less distinct six-sided plates, variously super-imposed one upon the other, the centre portion of the crystals being somewhat opaque. Dr. G. Bird believed that it does not exist in healthy urine, and rarely in morbid; but is probably derived from the sulphur extractive matter of the urine; and he inferred that it proceeds

* *Croonian Lectures*, London, 1856.

the waste of tissues and of the hæmato-globuline. In the majority of the sufferers the general health and nutrition have been bad. The indications for treatment are acids and tonics; as nitro-hydrochloric acid, chalybeates, &c.

We now come to the deposits, for the most part of inorganic kind:—

The phosphatic deposits. The phosphatic salts in healthy urine are of two kinds, the alkaline and the earthy phosphates. The alkaline includes the phosphate of soda, the acid phosphate of soda, and the phosphates of soda and ammonia, which are easily soluble, and do not give rise to deposits or concretions. The earthy phosphates, viz. phosphate of lime and magnesia, exist in very small quantities, and are eliminated in a perfect state of solution, but under certain circumstances and certain morbid actions are readily deposited and form concretions. The earthy phosphates are insoluble in water, but are dissolved by mineral acids; they are soluble in organic acids; mucinous substances dissolve them, but the salts are again easily precipitated by ammonia. Thus the phosphatic deposits derived from the earthy phosphates of the urine, and comprise three varieties:—

1st. The phosphate of lime. The deposits of this salt, Dr. Williams observes, have led to more errors than any other class.


FIG. 304.



Phosphate of lime.


When it exists in the urine in solution, or is precipitable by heat, it has repeatedly been mistaken for albumen; and where it is formed the great mass of a deposit, has as frequently been mistaken for pus or purulent mucus. The deposits consist generally of a white amorphous matter, not in crystals, but as minute granules.

PL. IV. S T



an oblique, six-sided, prismatic
ammonio-phosphate of magnesia (1

FIG.



Triple pho

and ammonia) generally assume the
 prisms with truncated extremities
 are sometimes bevelled off; or of
 six-sided plates; or again, of stars
 has been added to the urine. They
 of a natural colour, of rather high
 times neutral, often acid, and

alkaline state of the urine, from a diminution of the re-acidity; this alkaline condition being due to local injury of the urinary organs, the presence of foreign matter, or to general morbid conditions, such as over-exertion, depression, spinal disease, insufficient food, or abuse of food and medicines. The source of this alkali may be an excess of carbonate of ammonia, probably effected through the action of a ferment converting the urea into a volatile alkali. The mucus altered by an unhealthy condition of the mucous membrane may effect the conversion of urea into an alkali; so also the presence of a fixed alkali, such as the phosphate of soda, produced during digestion. Dr. G. O. Rees maintains that the alkali which causes the precipitation of the phosphate is secreted by the mucous membrane of the bladder; the earthy phosphates are precipitated from the urine, and not secreted from the mucous membrane, as was formerly supposed. The treatment of such deposits will be the removal of all recognisable causes, such as imperfect digestion and assimilation, improving the patient's powers, and correcting the morbid conditions of the system and mucous membrane of the bladder.

The *carbonate of lime* deposits are of rare existence, generally of secondary formation, and usually occurring in an amorphous form. Under the microscope they appear as small and white crystalline spherules. These often congregate together, and assume the shape of a drum-stick. The deposit is stated to be the result of such a state of enervation, that the bond of affinity, which in a healthy condition of the nervous system firmly united the elements of urea, becomes loosened; a rearrangement of atoms occurs, the urea becomes carbonate of ammonia, and, under the influence of ordinary chemical affinity, carbonate of lime is generated at the expense of the calcareous matter of the urine.*

Having thus taken a cursory survey of the formation of deposits due to changes in the urinary fluid, we now pass on to more aggravated conditions, viz. the amalgamation or conglomeration of such deposits into a mass, forming a stone or calculus. Formation of a calculus, in a practical point of view, may be attributed to one of three causes: 1st, it may take place in the secreting cells of the kidney, by precipitation of the salts, as occurs naturally in the kidneys of reptiles and birds;

* *Guy's Hosp. Reports*, series i. vol. vii. p. 224.

deposition of urate of ammonium and amorphous ingredients, according to the state of irritability or innervation of the bladder.

The simplest mode of witnessing the formation of a stone is in cases where a gum-eucalypti stone remains in the bladder for some time. Urinary crystals and deposits of uric acid, Foreign bodies, no matter of what kind in the bladder, cause decomposition of the mucous membrane of the bladder, and the solution of its salts upon it. Solid foreign bodies, which comes in contact with the mucous membrane, causes the latter to secrete a mucus which comes in contact with the foreign body, and precipitated around the foreign body. In the contrary, such as mucus, fibrin, and other organic matter, coat, but cause a precipitation of uric acid around them, viz. the uric acid, and form stone.

Another, and by far the most common mode of formation of a vesical calculus is the escape of uric acid which cannot pass with the urine, and it is the uric acid which acts in fact as a foreign body, and precipitates on its surface. Renal calculi +

ing in bulk as the hardening process continues. Others maintain that the calculus is formed by direct precipitation of salt. If an excess of uric acid be separated by the kidneys, salts of uric acid are thrown down; the latter may also be freed from decomposition of the urate of ammonia—the more ordinary source.

The oxalate of lime calculus of the kidney, according to Dr. Prout's researches, is not generated in a perfectly healthy kidney, and two conditions are necessary to its formation: the first, that the oxalic acid should exist in the system, and be mixed with the urine; the second, that lime, in some shape or other (that is, the phosphate or carbonate), shall be furnished by the mucous membrane of the infundibulum. The phosphate of ammonia and magnesia is rarely deposited in the kidney, but it may form a coating for the other calculi, and linger for a considerable time in the organ. Its formation, according to Dr. Prout, takes place in the following manner: Urine, under ordinary circumstances, contains the phosphate of magnesia, which is held in solution, being a highly soluble salt. But in some cases of disease the urea of the urine becomes decomposed in the kidneys, and ammonia is evolved, which combines with the phosphate of magnesia so as to make the triple salt.

Dr. Beale* observes: 'It is not uncommon to meet with microscopic uric-acid calculi; aggregations consisting of uric acid crystals, which if retained might receive deposits of fresh material on the outside, until small calculi, varying in size from mustard-seed to that of a pea, or larger, are formed. Microscopic calculi of phosphate of lime are by no means uncommon, and are often found in the kidney; but until a few years ago I never had an opportunity of watching the formation of calculi composed of oxalate of lime. The nucleus of these calculi does not consist of mucus or epithelium, as in the phosphatic, but is of the same composition as the exterior.'

The formation of a calculus in the bladder without any previous nucleus of renal origin, or the presence of a foreign body, is due either to an excess of insoluble material and the immediate precipitation of the salts of the urine,—as in the urinary forms of uric acid, urate, and oxalate concretions,—or stagnation of the urine in the bladder, in consequence of

* *On Urinary Deposits*, 2nd edit. p. 392.

formation to fermentation.

The whole of the chemical could be held together by a kind of matter derived from animal matter; and some affirm that it is or even pus. Mr. Coulson (in *Bladder*, p. 343) thus sums up : secretion of the bladder ; Fourcroy and sometimes to gelatine with a however, could not determine fibrin, albumen, gaseous matter, it to consist of a mixture of gelatin ' that the smaller calculi are a mucus, albumen, or some other which entangle, and ultimately the more insoluble ingredients of as quoted by Gross, extends this of the concretion ; he found the animal matter enveloping the part of salts, bore a great resemblance frequently detected in the central proportion of epithelial scales from fibrinous casts from the uriniferous

Urinary calculi may be arranged in distinct classes, to which a third name is given. The first will include

lements of calculi, yet there are other chemical ingredients present in more or less quantities; such as organic matters, carbonate of magnesia, silica, oxide of iron, benzoate of ammonia, oxalate of ammonia, phosphate of iron, urea, clay- &c.

The number of urinary calculi present varies; the renal are usually two or more, the vesical usually only one, and the prostatic very numerous: the vesical, however, may amount to 242, 307, 678, and even to 1000, as stated by Dr. Physick.* The size also is extremely variable, viz. from that of a pea, or almond, to a cricket-ball; the phosphatic calculi usually of the largest size.

The weight depends not so much upon the size as the composition of the stone, for the phosphatic are very light, and the uric very heavy; it may vary from a few grains to many ounces. The majority are under an ounce; the heaviest one I saw was 6lbs. 3ozs.† Recent calculi are heavier than old, as they contain moisture.

The shape is not uniform, depending greatly upon the situation and composition of the stone; thus renal calculi are irregular and often moulded to the form of the calyces, pelvis, and infundibulum of the kidney; those in the ureter are generally cylindrical; the vesical are more regular and of an ovoid form; the prostatic numerous, small and faceted, or pea-shaped. The uric acid and urate of ammonia calculi are generally smooth and regular; the oxalates tuberculated like a mulberry, and the latter appellation; the phosphates irregular and conical. But there are an infinite variety of forms, such as conical, pyramidal, triangular, cubical, square, rhomboidal, pyriform, pisiform, cardiform, cuneiform, semilunar, &c. Some are as if they had been actually divided by a firm cutting instrument; and in one case, in the Guy's Museum (No. 2136³⁰), apparently divided portions seem as if they had again become cemented and framed in by a subsequent deposit. In the same collection (No. 2145³⁵) there is a cystic oxide calculus in the shape of an ear-drop. Some remarkable forms of calculi have been detailed in the Guy's Hospital Reports, series iii. vol. i. p. 351, where the appearance resembles that of two calculi joined by a transverse bar, from which circumstance they are

* Gibson's *Institutes*, 5th ed. vol. ii. p. 220.

† Coulson, op. cit. 5th ed. p. 337.

termed barshot or dumb-bell shaped calculi; these forms are generally met with in calculi which have originated or become lodged in the prostate, and grown backwards into the bladder. Such shapes have also been found in encysted calculi of the bladder.

The colour of a calculus depends for the most part on the chemical nature of the crust or surface, but this by no means indicates the actual nature of the interior of a calculus: a white surface shows a phosphatic crust; a cinder-grey, that of urate of ammonia; a yellow pale-brown, or brown, the uric acid; cinnamon-brown, the uric or xanthic oxide; a brown, brownish-green, or even blackish-green, the oxalate of lime; a grey-greenish hue, the cystic oxide. These colours, however, are liable to modification, and give but presumptive evidence.

The odour of recently extracted calculi is sometimes peculiar. The phosphatic calculi are very ammoniacal, fetid, and acid; oxalate of lime calculi on section are said to emit a faint odour of semen. Some are affirmed to possess an aromatic smell, like that of castor or musk.

The consistence depends in a measure on the chemical composition, on the amount of cohesion of the particles, and on the presence of moisture or organic matter: it varies from that of soft mortar or sand to that of hard marble. Phosphatic calculi are generally soft, the uric acid hard, and the oxalate very hard. The crust is usually the softest part of a stone, and the nucleus the most dense. A stone of rapid formation is less dense than one of long existence and of small size.

The section of a calculus shows its internal construction, and ordinarily presents a centre or nucleus, a body or intermediate part, and a crust or outside coating. When all these consist of one and the same chemical material, the calculus is called uniform or simple; when it is made up of one or more elements, the stone is termed mixed, heterogeneous, or compound. Again, these elements may be so disposed as to form a solid mass, without any visible distinction of arrangement; but generally they are deposited in concentric layers or laminæ, and in some instances there are seen lines radiating from the centre to the circumference, as if the laminæ were composed of perpendicular crystalline fibres. Dr. Prout* thus explains the laminated condition: 'Between the different intervals at which the different

* Prout, *On Gravel, Calculus, &c.*, p. 361.

laminæ have been formed, periods have intervened during which no deposition has taken place. This remark not only applies to the different laminæ of a heterogeneous calculus, but to the different laminæ of calculi composed of the same substance; as, for instance, to the different laminæ of which lithic acid concretions usually consist. This explanation is in perfect accordance with the circumstances attending the formation of calculi, which often, as is well known, remain in the bladder for a great number of years without attaining any remarkable size. Moreover, the constant state of change alone, to which the urine in all individuals is liable, almost precludes the notion of homogeneity in a calculus. We may suppose, therefore, that certain changes take place in the urine during which the law of continuity of deposition is suspended, and the surface of the concretion becomes, as it were, *water-eaten*, and less apt for future accretion; in short, assumes all the properties of a heterogeneous substance. Under these circumstances, when a tendency to deposition occurs, it will have to commence *de novo*, and, as it were, on the surface of a foreign body. The consequence will be, that the adhesion between the new and the old coats or laminæ will be less firm than in the intermediate parts, and that a calculus thus formed will be disposed, when broken, to separate into concentric laminæ.'

Dr. Hodgkin* gives the following account of the *Structure and Character of Calculi*:—

A. Calculi of a Crystalline Character.—In many instances, the particles separated or precipitated from the urine to form calculi, assume a crystalline character; and the mode in which this crystallisation takes place around a small nucleus materially influences the structural character and the form of a calculus. The structure is radiated, and the form more or less rounded. Such specimens of calculi bear considerable analogy to certain zoolites, and to some specimens of radiated quartz. The cystic oxide affords one of the best specimens of decidedly crystalline calculi. Some of the phosphates appear to come next; and calculi consisting of oxalate of lime frequently present evident traces of crystallisation.

B. Amorphous or Massive Calculi.—The molecules which unite in the composition of these calculi do not possess a perceptible crystalline form, and the resulting substance is what the mineralogists would call massive, with no other indication of structural arrangement than more or less distinct traces of superposition in concentric layers. There is considerable variety in calculi of this description; some possessing granular texture, which might be compared to a coarse sort of sandstone, whilst in others it is soft and earthy, like some of the softest specimens of chalk or tufa. The lamellar arrangement is very dif-

* *Guy's Hospital Reports*, series i. vol. ii. pp. 268 et seq.

... it is strongly marked, and ... in small flakes; whilst others, ... pick up almost indifferently in

4. *Amphioxus*.—These are seen in the form of a section of a single fibre, or a bundle of short fibres, arising from an assemblage of adaxial cells in the specimens which

... with a model calculus, viz. the
... as urea is the chief con-
... composed of oxalate of
... as uric acid or urates
... the crust formed of phos-
... When a calculus consists of
... another in a regular
... Fig. 9.)

... the composition and relative proportion
 ... as regards the nucleus.
 ... the reader to the standard works
 ... especially to the Catalogue of
 ... the Royal College of Surgeons of
 ... there are excellent coloured plates
 ... In general, calculi are classified
 ... of the nucleus; and for the
 ... although Dr. Beale
 ... it is important to bear
 ... the calculus which is visible to
 ... the nucleus, while the real
 ... of a different composition to
 ... is it. The nucleus of
 ... consists of uric acid, is really
 ... which the uric acid has

Dr. G. Bird's observations, is that the nucleus of the calculus, or nearly so, is usually remarkably eccentric, as in some cases several distinct nuclei, or centres, may be seen within. In some rare instances, the nucleus is found loose within

• • • • •, 1961, p. 354.

— *Journal of the History of Ideas*, Vol. VII, p. 182.

body of the entire calculus; a circumstance, in all probability, arising from a layer of blood or mucus having concreted round the nucleus, and the matter forming the body of the calculus having been deposited on this layer. In this case, on the whole becoming dry, the mucus or blood would be diminished to a very thin film, and the calculus would appear to contain a nucleus. In a few instances calculi appear to possess no nucleus, the centre being occupied by a cavity full of stalactites or mammillated projections, giving the idea of the external having been first formed, and the mammillated portions subsequently in the interior. In the Museum of Guy's Hospital this state occurs only in the uric-acid calculi. In one specimen in the collection (No. 2154) the central cavity is lined with fine scales of triple-phosphate, resembling the crystals of quartz often found lining cavities in flints. The explanation of the formation of such cavities is, that the original nucleus was of blood or mucus, which has disappeared by desiccation.

Though in an essay of this kind we cannot enter fully into the chemical details of urinary calculi, yet we have deemed it advisable to introduce the following oft-quoted table of Dr. Bence Jones,* somewhat modified:

Urinary Calculi.

Destroyed by heat; combustible; leaving only a small residue:
Become red on the addition of nitric acid, and form a murexide:

- | | |
|---|--------------|
| a. Soluble in carbonate of potash, evolving <i>no</i> ammonia; soluble in caustic ammonia, or potash; on the addition of an excess of acid crystallises in angular crystals; not soluble in water | } Uric acid. |
| b. Soluble in carbonate of potash, evolving ammonia; soluble in water when boiled; solution in water with a few drops of ammonia, when evaporated, crystallises in needles | |

} Urate of ammonia.

Do not become red on the addition of nitric acid.

- | | |
|--|--------------------------|
| a. Soluble in ammonia, <i>not</i> crystallising when evaporated; insoluble in carbonate of potash; dissolves without effervescing in nitric acid, leaving a lemon-coloured residue; soluble in strong sulphuric acid, not precipitated by dilution | } Uric or xanthic oxide. |
| b. Soluble in ammonia, crystallising in six-sided plates when evaporated; soluble in strong caustic potash; the solution, when boiled for a few moments, on the addition of a drop of dilute acetate of lead, gives sulphuret of lead | |

} Cystic oxide, or cystine.

* *Medical Times*, October 1851.

bonated alkalies and oxalate
dilute acetic acid, with efferves

2. Soluble with hydrochloric acid ;
soluble in mineral acids with
in acid when neutralised gives
carbonated alkalies and oxala
in acetic acid ; decomposed
yielding carbonic acid and c
boiled with carbonate of soda
in the solution and precipitate
3. Soluble with hydrochloric acid ;
before or after heating :
 - a. Solution in acid with excess o
crystalline precipitate ; wi
phate of lime (bone-earth
blow-pipe, and gives off a
solves in acetic acid with
 - b. Solution in acid with excess
phous precipitate ; with
phate of ammonia and
before the blow-pipe . .
 - c. Solution in acid with excess
partly crystalline, partly
without addition easily fu
4. Not acted upon by acids or alkali
bulk of carbonate of soda forms

We will now give a general

that of lime, giving a white appearance; and the nucleus, though consisting of pure uric acid according to Dr. G. O. may be quite white, owing to the absence of colouring matter. The size varies greatly, and it often attains considerable weight and specific gravity is intermediate between oxalate and the phosphate; the form is generally ovoid, slightly compressed and smooth, but sometimes beset with tubercles, which may be increased to such an extent as that the stone resemble the oxalate of lime;* the calculus also moulds itself into irregular shapes in the locality in which it may be placed. The appearance on section varies greatly; Dr. Thudichum† describes it to consist of concentric layers of variable thickness, every layer preserving its own whiteness pretty regularly all round the calculus; the texture is seen on the surface; in the hard and pure varieties it is crystalline and fibrous, the fibres of each layer converging like spokes towards the centre of the stone, and their fracture being also parallel with those fibres; the less dense and pure calculi are of a more earthy character and amorphous in fracture, and have no laminae. Some few are so hard as to impart a ringing sound on percussion, a sharp sound like a pebble. Here the fracture is conchoidal, and such stones when broken, as in the majority of cases, form sharp angular fragments, which are dangerous on account of their tendency to wound the bladder and urethra.

The *urate-of-ammonia calculus* is not common; it was discovered by Fourcroy and Vauquelin in 1798 (see Plate, Fig. 2). In the Museum of the Royal College of Surgeons, Edinburgh, there are only 14 in 649 calculi, being 1 in 46; at Guy's Hospital Museum there are 7 in 394: but this substance forms an important constituent of compound calculi, varying from 1 in 3½ to 1 in 100 of all calculi. The calculi are seldom large, rarely exceeding half an inch in size, are generally ovoid and smooth, or sometimes slightly tuberculated on the surface. Their colour is very generally characteristic, being of a slate, pale greyish-fawn, or clay colour; not unfrequently they more nearly resemble pipe-clay than any other; sometimes there is an admixture of red or brown, and in some instances pink layers are observed near the centre. In the Guy's Hospital Museum (No. 2213) there is a remarkable collection of 142 calculi removed from the bladder of one patient

* See preparation No. 2125, Guy's Hospital Museum.

† Thudichum, *On Urinary Deposits*, &c. p. 111.

by Sir Astley Cooper, all of the same figure, being cubes rounded at the edges and angles; these have the colour of pipe-clay, and are chiefly composed of urate of ammonia: it is curious that the patient had afterwards another calculus, which on examination proved to be of a different kind from these. Their consistence is usually compact, earthy, and very brittle, and they have a fine earthy fracture; this is well shown in a calculus (Guy's, No. 2136⁵⁰), where there is a remarkable fissure running through it, as if it had been fractured and subsequently reframed by a deposit of urate of ammonia. They seldom present the distinct concentric arrangement of uric-acid calculi, generally appearing homogeneous, but at times an indistinct thin laminated condition may be traced; this circumstance is readily explained by the frequent amorphous condition in which the urate is deposited in sediments. These calculi are generally observed in children, and are said to give rise to little constitutional disturbance.

Calculi of urates of soda or lime are not met with; the salts are generally mixed with those of urate of ammonia or uric acid in the several layers of compound calculi. In one instance, however (Guy's Hospital Museum, No. 2154³⁶), there is a fusible calculus with a nucleus of urate of lime.

3. *Uric oxide, or xanthic oxide, or xanthine*, discovered by Dr. Marcet in 1815, is a very rare calculus, of which only four specimens are recorded (see Plate, Fig. 5).

The calculus in which Dr. Marcet first detected this substance weighed but eight grains, and nothing is known regarding the portion not consumed in the analysis: it does not exist in the collection at Guy's Hospital, and neither Dr. Prout, Dr. Yelloly, nor Dr. Babington can give any information on this point. It was rediscovered by Dr. Stromeyer: * the calculus was removed by Langenbeck from a peasant's child, eight years of age, at Hanover. In shape it resembled a flattened pullet's egg: it broke into three pieces during the operation. The whole calculus weighed nearly 330 English grains; its section was partly of a lustrous bright brown colour, and partly earthy and pale brownish flesh colour. It was composed of concentric separable layers, without any appearance of a crystalline or fibrous texture: it possessed a distinct nucleus, which did not, however, differ in chemical characters from the body of the calculus. It was as hard as uric-acid calculi generally are, and on slight friction it assumed a wax-like lustre. Professor Merx of Göttingen sent two fragments of the calculus to Dr. Willis, one of which is in the Museum of the Royal College of Surgeons, Edinburgh, and the other is in the Guy's Hospital Museum (No. 2145⁹⁰). In the Catalogue of Calculi at the College of Surgeons of England there is a plate of this calculus (see Plate xii.).

* *Annalen der Physik*, Band 41, p. 393.

calculus bears a close relation to uric acid, and seems to depend on an imperfect oxidation of the material from which, in the ordinary processes of the kidney, uric acid is eliminated.

The *oxalate-of-lime calculus*, discovered by Wollaston in 1784, is the next most frequent after the uric-acid calculus, and the ratio of 1 in 14½ to 1 in 20 of other calculi. These do not generally attain a very large size, but are very hard, dense, and proportionately heavy. As a nucleus in uric calculi, its proportion varies from 1 in 4½ to 1 in 7½ (Plate, Fig. 4).

G. Bird* thus describes these calculi: Their form or local configuration is almost invariably that of a tubercular, lobular, or spinous character, and rarely perfectly smooth, but, together with their colour, varying from grey to a rich brown or almost black, has gained for them the specific name *mulberry*. In some calculi the surface is studded with spines so acute and slender as to resemble thorns; in others there is a coating of acute octahedrons of transparent oxalate of lime, giving an extremely beautiful appearance. Sometimes the crystals are opaque, and the octahedron is remarkably modified: the calculus then looks as if studded with pearl-spar.† J. Bird observes: 'It has been supposed that the spinous or irregular form of these calculi depends upon their being formed in the kidney, and that they were casts of the pelvis and vessels of the organ. This, however, is by no means sufficient account for their figure; for no calculus so closely becomes a copy of the interior of the kidney as the fusible variety, and yet it ever presents a tolerably smooth surface, however contorted in figure. It is, therefore, more likely that the constantly crystalline state in which the deposit of oxalate occurs has more to do with it; and without calling to our aid any very fertile imagination, a tendency to a cubic or octahedral outline can easily be traced to the entire calculus.' Sometimes these interstices between the projections or spines are filled up by some amorphous substance, as urates, or phosphates, thus giving the calculus an ovoid character. The appearance on section is generally that of an imperfectly lamellated structure, the consecutive layers forming waving lines, and often resembling knotted

* *Guy's Hospital Reports*, series i. vol. vii. p. 213.

† See Preparation 2139²⁵ in the Guy's Hospital Museum.

heart of oak; occasionally there is seen a layer of oxalate of lime arranged round the interior one with great regularity, having a remarkably radiated appearance, like a series of infinitely minute needles placed side by side, and presenting a perfectly porcellaneous structure. In compound calculi the oxalate of lime deposition gives to the character of a stone a remarkably beautiful appearance resembling that of fortification agate.

The rarer varieties of oxalate-of-lime calculus are the following:—

1st. The small, smooth, globular ‘hemp-seed calculus’ described by Dr. Wollaston, and thus alluded to: ‘It is of a much lighter colour, so as to resemble in hue as well as smoothness the surface of a hemp-seed.’ There may be one or many, and they are generally found in the kidney.

2nd. Where the character of the calculus is crystalline throughout, and it has a white or pale brown colour. These calculi consist of nearly pure oxalate of lime in a crystalline form. Dr. Yelloly, on analysing the calculi in the Norwich Hospital Museum, met with not less than twenty examples of such crystallisation. The author has removed by lithotomy a calculus consisting of pure oxalate of lime, and of a pale brown colour, which crumbled to pieces immediately after extraction, a circumstance which was due to the absence of any binding material, there being no trace of animal matter.

The third variety is the pure white oxalate-of-lime calculus. This is of a milk-white colour, possesses a highly polished surface, is of extreme rarity, and is generally, if not always, found in the kidney; its external surface presents no crystals, but is perfectly smooth, though it may be spinous.* This variety seems to have escaped the attention of writers on Stone.

In the Museum of the Norfolk and Norwich Hospital are three specimens: the first, a moist preparation representing a renal calculus of cauliflower shape and of great magnitude; it occurs in the form of a single mass, filling a greatly distended pelvis. From this mass there branch out other smaller masses, which fill up the infundibula. The whole kidney is greatly enlarged. The stone is extremely white, quite smooth, very hard and compact, and polished: it resembles Parian marble. The only circumstance known of its history is, that the preparation was removed from a man after death.

The second example is a small calculus, very smooth and shining, weighing

* For the detailed account of this variety we are indebted to Mr. Charles Williams, formerly House-Surgeon to the Norwich Hospital.

was, about the size of a horse-bean, said to have been passed by an adult. On analysis it yielded oxalate of lime.

Third specimen is nearly spheroid in shape, and about the size of an egg-marble; it is of a white colour, though not so polished as the above; its external surface is covered with spines and tubercles peculiar to 'cherry calculus,' which it most closely resembles in everything but its tubercles are smooth, i.e. they are devoid of crystals; its weight is 1.5 grains. The catalogue informs us it was passed by a female, aged 40, who was pregnant at the time. On scraping its surface, and placing it under the microscope, very minute octahedra of oxalate of lime were detected; they were insoluble in potash and acetic acid, and thus distinguished from other substances which they resemble.

In reference to this Mr. Williams remarks:—'I am inclined to believe the pure white colour of the third variety of oxalate-of-lime calculus is due to the circumstance that it is met with only in the kidney, and that while in the organ it is not permitted to lie continuously in urine, and therefore does not acquire the dark brown or almost black hue which the oxalate-of-lime generally assumes when situated in the urinary bladder. The tuberculate form of this calculus, in consequence of its roughness, causes the membrane of that viscus (the bladder) to pour out blood far more copiously than any other variety of calculus; and there can be but little doubt that the colour is derived from the hæmotosin, which has been absorbed by the urine from the blood-corpuscles, and remains suspended in

the urine in the kidney, the constant trickling of urine over this variety of stone gives rise to its smooth white surface.

The *cystic oxide*, or *cystine calculus*, discovered by Wollaston in 1800, is a rare calculus (see Plate, Fig. 3). It differs from all others in containing a considerable proportion of sulphur, no more than two atoms being present in every equivalent of the acid.

The calculus is not a product of the bladder, as it would seem to imply, but of the kidneys; and there is a remarkable hereditary disposition to its formation, for out of 10 collected cases, 10 occurred in four families, and in 3 cases the subjects of the complaint were brothers. Ten specimens in the collection of Guy's Hospital were passed by the same patient, whose age was 30; 3 were passed in the year 1814 (No. 1144), 6 small ones subsequently (No. 2145), and in 1828 a very remarkable one, of the shape of an ear-drop (No. 2146).

These calculi are usually small in size; the largest is in the University College Museum, and weighs 850 grs.; at St. Bar-

tholomew's Hospital, hæmorrhagic urine, filtered so frequently as to lose all the corpuscles, will not possess the dark blood-stained colour notwithstanding.

tholomew's there is one weighing 740 grs. The one at Guy's Hospital (No. 2143), the second specimen analysed by Dr. Wollaston (the discoverer), is described by him in the *Phil. Trans.* for 1810; it is an inch and a-fifth long and one inch broad.

They are described to be generally rounded and smooth, but may be covered either with smooth tubercles or sharp projections externally; they have a wax-like lustre, appear semi-transparent and glistening, and resemble very much the ammoniaco-magnesian phosphate calculus. When recent, their colour nearly approaches that of the uric-acid calculus, being of a pale yellowish brown; they undergo, however, a remarkable change by long keeping, turning slowly from brown to grey or green. Thus the calculus in Guy's Hospital Museum (No. 2143) was brown in 1817, now it possesses a rich bluish-green colour. Some consider that the greenish-blue or dirty-green colour is due to the presence of earthy phosphates. Dr. Bird, however, remarks, 'It has been suggested to me by both Dr. Prout and Dr. Willis, that this alteration in tint may in some way depend upon changes produced by the sulphur.' Their consistence is soft, and on section they present a very imperfectly radiated structure, and exhibit no tendency to a development of concentric layers; when scraped, they yield easily to the knife, and form a perfectly white powder, whether the calculus be brown or green. The fracture is crystalline.

6. The *phosphate-of-lime calculus*, discovered by Wollaston in 1797 (see Plate, Fig. 6), is likewise rarely found in the bladder uncombined with other salts; and it seldom forms the nucleus of other calculi. Calculi chiefly or entirely composed of it are tolerably hard and smooth externally, not presenting the angular asperities of the oxalate of lime, or the contorted and irregular figure frequently assumed by the fusible calculus: sometimes they consist of two or three portions fitting into each other in a variety of ways.

Of three specimens in the Guy's Hospital Museum, the first (No. 2148) presents a porcellaneous appearance externally, admitting of considerable polish by friction; is conchoidal in its fracture, and is of a greyish-white colour; the second (No. 2149) is more regular in figure, and is made up of a series of concentric layers of the phosphate, readily distinguished from each other by their tint, being alternately white and fawn-coloured; the third (No. 2150) is light-coloured, and has a spongy, cancellated structure, and looks more like a bony concretion than a deposition.

There are, however, two varieties of this form of calculus:

as, as described by Wollaston, of renal origin, and consisting of stral phosphate of lime; and these are usually pale brown, with a smooth polished surface regularly laminated, and the pieces so slightly adherent as to be easily separable into lamellar crusts; in some, radiating lines are seen in a direction perpendicular to the laminæ; these calculi contain a considerable proportion of animal matter. The other form is of vesical origin, and composed of phosphate of lime, similar to that of the former, and hence often called 'bone-earth' calculi; they are more common than the former, and constitute irregular masses resembling mortar, or a granular semi-crystalline powder, embedded in a tenacious mucus. Dr. Taylor, in the Catalogue of the Royal College of Surgeons, England, has described several specimens as consisting entirely of this salt.

Phosphates are scarcely ever succeeded by any other form of calculus, but there are three instances in which a phosphatic calculus was succeeded by uric acid or urate of ammonia, and in one instance it was followed by oxalate of lime; the general law in which they succeed other deposits is 1 to $4\frac{1}{2}$. Foreign bodies as a rule have earthy phosphates deposited upon them; there is one remarkable exception (Cat. of the Royal College of Surgeons of England, Plate iv. Fig. 6) where a slender piece of shell formed the nucleus of a large oval calculus, consisting entirely of uric acid.

Respecting the earthy phosphates in calculi Dr. Thudichum remarks:—'Not quite 10 per cent. of all calculi have a nucleus of mixed phosphates; but these substances enter into the composition of about 34 per cent. of all calculi, forming either the body, one or more layers, or the crust. This shows that the presence of deposits of mixed phosphates in the urine rarely ever gives rise to the formation of calculus, but that the presence in the bladder of other calculi frequently cause a deposit of phosphates to be formed around them, in which case every calculus is nothing else than a foreign body. For a calculus to form, it requires the presence of some foreign material, such asropy mucus, or a clot of blood or of blood, in hæmaturia and chronic disease of the mucous membrane of the bladder.'

The *triple-phosphate calculus*, or ammoniaco-magnesian phosphate, also discovered by Wollaston in 1797 (see Plate, Fig. 1) is rare.

There are three specimens of this calculus in the Museum of the Royal

College of Surgeons of England, and one or two only at Guy's Hospital: in the former collection they appear as a white crystalline mass, radiating from the centre, and having the surface studded with shining crystals, which, when recent, are nearly transparent, but by exposure become opaque; their texture is earthy, friable, and imperfectly lamellar; some are hard and compact, semi-transparent, and crystalline in fracture like alabaster. In the Guy's collection (No. 2154) a calculus of this kind is to be seen which has no nucleus, but a central cavity lined with delicate crystals of triple phosphate, resembling the crystals of quartz in the cavities of flints; and No. 2152 is a section of a large calculus of the kind on a nucleus of a tobacco-pipe. Dr. Thompson mentions a calculus of this sort weighing nearly 2 lbs.

8. The *fusible calculus*, or mixed phosphate, or the phosphate-of-lime and phosphate-of-magnesia and ammonia calculus, discovered by Wollaston in 1797 (see Plate, Fig. 8), is the most frequent of the phosphatic calculi, and forms rather more than $\frac{1}{12}$ of all calculi, being rarely succeeded by any other species of urinary deposit; their relative proportions in the several collections are various, from 1 in $8\frac{1}{2}$ to 1 in $13\frac{1}{2}$.

These calculi grow to a considerable extent, and are comparatively light and of low specific gravity; they are generally very irregular, and mould themselves to the situation in which they are placed, often filling the interior of the bladder, and becoming impressed with its rugæ; sometimes they are globular and ovoid, at other times taking on the most extraordinary shapes; where there are two or more, they assume a cubic or tetrahedral form. Their colour is white, grey, or dull yellow, and their consistence is more friable and earthy than any other variety, and is sometimes so soft as to resemble moist chalk.

Their appearance on section is thus described by Dr. Taylor (op. cit.): 'They are frequently composed of concentric laminae, which in general adhere but slightly to each other; between the laminae shining crystals of the triple-phosphate are often observed; in some the laminae are entirely wanting, and these form a white friable mass like chalk; in others they appear semi-crystalline, as if made of numerous small crystals confusedly aggregated together. The relative proportion of the constituents of this calculus is exceedingly various, and the predominance of one or the other salt gives peculiar characters; in those which have a crystalline and glistening texture the triple-phosphate is most abundant, while the calcareous phosphate is in excess in those which have an amorphous earthy appearance.'

It generally constitutes the calculous masses deposited on foreign bodies accidentally or intentionally introduced into the bladder.

9. The *carbonate-of-lime* calculus, discovered by Brugnatelli in 1819, is a very rare form indeed.

He describes* forty-eight small concretions extracted from the bladder of a young man: these were of the size of a pea, lamellar in structure, and broke with a shining fracture. He also mentions several ash-coloured calculi, sixteen in number, and of the size of a nut, which were composed of carbonate of lime with a trace of carbonate of iron: these were removed after death from the bladder of a woman. Dr. Prout has also seen small calculi of this salt, and these were perfectly white and very friable.

In Guy's Museum, No. 2182⁷, there are some calculi of carbonate of lime, with mere traces of carbonate of magnesia and alumina; in one specimen, No. 2187⁵⁰, it constitutes the body of a calculus, being deposited in a snow-white colour, on a nucleus of uric acid. There are several in the Oxford Museum, among the collection of Mr. Hitchins, and there is in the possession of Mr. R. Smith of Bristol a remarkable collection of these calculi, some of which are figured in the Catalogue of the Royal College of Surgeons of England, Plate xii. Figs. 3, 4, 5, 6, and 7. Of these calculi, five were extracted by the lateral operation from the bladder of a boy aged sixteen, and the other fifteen were passed by the urethra previous to the operation: these latter were of the size of peas, of a rounded figure, and had flattened surfaces; they were compact and lamellar in structure, and their external surface was of a light brown colour. Those extracted by operation were exceedingly irregular, and their external surface dusted over with white powder: the largest was of the size and figure of a large almond; when sawn through, the concretion showed no concentric laminæ, but irregular waved lines similar to those of the mulberry calculus. The largest was so hard as to require the assistance of the lapidary's wheel to divide it.

Prostatic calculi sometimes consist almost entirely of this substance, but, as Dr. Thudichum observes, 'it is always questionable whether the lime or the carbonic acid were in any case derived from the urine.'

10. The *fibrinous calculus* of Dr. Marcet and of Dr. Prout is by some not ranked among calculi, in consequence of its differing so totally from those other concretions. Dr. Bird† says:— 'It must be regarded as a portion of dried inspissated albuminous matter exuded from an irritated kidney, rather than as a calculus produced under circumstances at all analagous to those of other concretions. Several specimens exist in the Guy's Museum of the pelvis of the kidneys and ureters being obstructed by clots of fibrin; but none of them present the hard

* *Litologia Umana*, Pavia, 1819.

† *Guy's Hospital Reports*, series i. vol. vii. p. 180.

concrete condition of the calculus described by Dr. Marcet. I am not aware of this variety having been mentioned by any other author except Brugnattelli, who, in his *Litologia Umana*, describes some calculi as consisting of crystallised albumen; they were passed by one individual, and each was about the size of a nut. These pseudo-calculi appeared to consist of dried coagulated albumen, which not unfrequently presents considerable lustre and a vitreous fracture, although scarcely sufficient to justify its being regarded as crystallised.* Some have described it to resemble yellow wax in appearance, and others have regarded it as an elastic organic substance closely allied to fibrine.

11. The *uro-stealith calculus* is another of the pseudo-forms: it is not decided whether it is a resin or fat, and its elementary condition is entirely unknown.

It has been described in Heller's *Archives* (1844, p. 97, and 1845, p. 1), and it consisted of several small bodies composed of a peculiar form of fatty matter, which were passed by a man aged twenty-four, who suffered from symptoms of stone in the bladder. Dr. Moore of Dublin has confirmed the existence of this substance, and has examined several specimens sent to him by Dr. Robert Adams of Dublin.*

12. *Blood calculi* have been but recently discovered, and are likewise pseudo-calculi.

Those described by Dr. Alison† were found in the infundibula and pelvis of the kidney of a painter, aged 52, who died of consumption; black calculi occupied the pelvis, while the infundibula were tenanted by a few calculi of a whitish-grey colour, with one exception small in size, about the magnitude of pear-seeds, and wanting the ordinary physical characters of phosphate of lime. One calculus which occupied the infundibulum was the size of a horse-bean, looked somewhat worn and disintegrated, and at one point resembled a piece of decayed wood, and was black on one side from the presence of altered blood; it was very light in weight, and was composed of blood and phosphate of lime. The black calculi, which formed the chief point of interest, were about six in number, and ranged from the size of a coriander-seed to that of a small horse-bean; when found, these black calculi were tolerably hard, but being friable they broke asunder on handling; the fractured surface varied a little in colour, in some parts presenting a dark rusty appearance. Liquor potassæ dissolved them, and they were capable of partial combustion; the microscope revealed only amorphous particles, but Dr. G. O. Rees, with the assistance of a neutral saline solution, discovered forms which he considered to be the remains of blood-corpuscles.

Silicious concretions are not met with, although silica has been

* *Dublin Quarterly Journal of Med. Science*, May 1854, p. 473.

† Dr. Scott Alison, *Archives of Medicine*, vol. i. p. 245.

found in other calculi, by Berzelius, Vauquelin, Fourcroy, and others. Those which have been extracted from the bladder have been pebbles or small stones introduced from without.

There are certain predisposing causes and conditions which are supposed by some to influence the formation of stone, or tend to its production; to these we can only cursorily allude. Thus, respecting climate and locality, it is stated that calculus is more frequent in temperate than in warm and cold regions, but more especially in humid and moist countries of moderate and changeable temperature, such as Holland, France, England, and Germany. It is, however, met with very frequently in Egypt, Isle of France, Bagdad, Russia, and both the Indies. Even in some parts of the same country it is more prevalent than in others, as in Norfolk in England.

With regard to race, it is asserted that the disease is much more frequent among the white than the dark; and Liebig attempts to explain this by suggesting that the exemption of the people of China from gout and calculus is due to the urine containing large quantities of tartar. However, stone is excessively common among the natives of India.

In some cases it may be traced to hereditary transmission, especially where there is gouty diathesis.

The sex modifies the relative proportion, stone being less frequent in females than males, their proportion being 1 to 20 or 1 to 23. This is due to the peculiar anatomical disposition of the urethra in the female, viz. its shortness, and its capability of dilatation allowing the easy passage of a small stone; as also to the comparatively temperate and regular habits of life of females.

The age at which calculi are most frequently met with is said to be in the young and in old people. This can only be ascertained by statistics, which are for the most part imperfect. The table furnished by M. Civiale gives the following results of 5,376 cases: Under 20 years of age, 55·56 per cent.; between 20 and 40, 14·69 per cent.; between 40 and 60, 14·95 per cent.; and above 60 years, 14·76 per cent. That of Mr. Coulson,* of 3,264 cases: Under 20 years of age, 71·20 per cent.; between 21 and 40 years, 12·10 per cent.; between 41 and 60, 10·87 per cent.; and between 61 and 80, 5·72 per cent. That of Sir Henry Thompson,† of 1,827 cases: Under 20 years of age, 60·42 per

* Coulson, *On the Bladder*, &c. p. 402.

† Thompson, *On practical Lithotomy and Lithotrity*, p. 269.

cent. ; between
17·56 per cent. ;

Sir H. Thomp-
rience, during g
Guy's, St. Tho
Oxford, Birmin
a foot-note,* a
from this table
life. Thus, und
940 cases, or 51
may subdivide i
1 to 5 years, 47
6 to 13 years, 4
during adolesce
years, 228 cases
ceeding years, 1
At adult life, fr
vancing life, fro
in old age, from

Mr. Coulson
ber of calculous
liability of ind

* *Table of preva-
lence by ages :*

Age.	No. of cases
1 . . .	7
2 . . .	74
3 . . .	116
4 . . .	153
5 . . .	123
6 . . .	90
7 . . .	86
8 . . .	49
9 . . .	57
10 . . .	60
11 . . .	35
12 . . .	58
13 . . .	32
14 . . .	35
15 . . .	26
16 . . .	27
17 . . .	18
18 . . .	26
19 . . .	19
20 . . .	13
21 . . .	12

Here, as in many other statistical tables, an error has crept in from not taking care to distinguish between absolute and relative numbers of persons living at the several periods of life enumerated. Corrected in this manner, the tables would show that children and young persons are less liable to calculous disorders than has been commonly supposed, and that from twenty years and upwards the tendency goes on increasing in a very remarkable manner to the end of life; or, as Sir H. Thompson puts it, 'that the proportion of elderly calculous patients to the existing population at their own ages is larger than the proportion of children afflicted is to the number of existing children.'

The condition and habits of life have some effect as a cause. Thus sedentary habits impair the functions of the skin, and give increased work for the kidneys; so also luxurious feeding and use of malt-liquors and nitrogenised food in excess furnish a source from which concretions are derived; again, indigestion, mal-assimilation and fermentation, impairment of the vital powers and acidity generated in the stomach, are frequent causes.

Respecting the exemption of sailors and soldiers from this disease, as alleged by Mr. Copland Hutchinson and others, we have not sufficient data to go upon.

Morbid states of the urinary passages may predispose to calculous formations, and more especially the presence of a foreign body, which is sure to cause a precipitation and deposit on its surface.

With regard to the influence of certain kinds of waters and spas, as a beverage, in inducing stone, it has been proved to be entirely imaginary; none of the forms of calculi correspond with the salts which exist in natural waters.

We shall now proceed to the description of the symptoms, effects, and treatment of urinary calculi as met with in the bladder, prostate, and urethra. Those of calculi in the kidney and ureter are described in the essay on DISEASES OF THE URINARY ORGANS, p. 187.

Stone in the bladder.—A vesical calculus may consist of every possible variety of the urinary concretions already described. Whatever may be the nature of the calculus, when once it exists in the bladder, it acts as a foreign body, and will give rise to sufficient evidence of its presence by inducing certain

symptoms, so as to lead the surgeon to its detection. Of course all these symptoms are not uniformly met with; they may vary in degree and intensity, depending upon a variety of general and accidental circumstances, so that it will be prudent to have some standard as indicating the symptoms which are generally met with in patients suffering from a stone in the bladder. The symptoms are premonitory and actual. The premonitory symptoms can exist only in those cases where the stone is of renal origin, so that in many adults and old people the vesical troubles are preceded by paroxysms of nephritic colic; but in general these antecedent complaints, attending the formation of stone in the kidney and its passage along the ureter, are so severe, that directly the calculus has reached the bladder a more or less long interval of repose and quietude allays all further suspicion, and nothing more is thought of the complaint until the calculus has reached a sufficient size to be incapable of being expelled through the neck of the bladder.

The actual symptoms or signs of a stone in the bladder consist of two kinds; first, those evinced by the patient's sensations, &c., constituting the subjective or rational symptoms, which offer but presumptive evidence of calculus; and secondly, those signs detected through the aid of mechanical means, and termed physical signs, the only real method of determining and diagnosing the presence of a stone.

The subjective or rational symptoms consist of the following complaints:—There is pain and sensibility, of a more or less acute character, referred to the region of the bladder or its neck; this may amount to a mere uneasiness or sensation of weight in the bladder, perineum, or hypogastrium, which may be so slight as to be unnoticed by the patient; or, on the other hand, the pain may be very severe and intense, occurring in paroxysms of great agony.* At first the pain is generally only intermittent, occurring during and after micturition, and more especially with the last drops of urine; and in children this is so frequent, and is attended with such irritation at the extremity of the penis, that they endeavour to allay it by drawing the penis and prepuce forwards; and this gives rise to an elongated condition of the prepuce, which is generally sodden

* The pain was so severe, continuous, and intense, that a blacksmith at Amsterdam, and a cooper at Königsberg, performed self-lithotomy.

with urine, as are also the fingers, causing them to emit a urinous smell. The pain is also aggravated in the acts of walking, riding, jumping, sudden exertions, jolting of the body, &c., and its intensity generally increases with the advance of the disease. Oftentimes the pain is reflected along certain nerves, inducing morbid sensations at their peripheral extremity; this pain is experienced in the extremity of the glans penis, in the scrotum and testicle, inducing contraction of the scrotum and swelling of the testicle, in the perineum, kidneys, down the thighs, &c.

John Hunter* quotes the remarkable case of the father of Lord Cavendish, who had pain in the left arm associated with calculus in the bladder, and this was the only indication of a want to make water. Some have had tormenting violent pains in the soles of the feet, with numbness and tingling.

The pain may last a considerable time, and then suddenly cease altogether, giving rise to the supposition of a cure.

Such was the case with the patient of M. Morand, mentioned in the *Mémoires de l'Académie des Sciences*, 1740, where a catheter detected a vesical calculus, but after that all symptoms gradually subsided, so that M. Morand was considered to be in error, and the patient bequeathed him his body to teach him a lesson; but on the post-mortem examination three calculi of the size of apricot-stones were found on one side of the bladder.

On the contrary, a stone may exist and grow to a considerable size without any evidence of pain, and then pain may all at once occur after some act of exertion.

Thus it happened in the case of a priest, who was in the act of taking some books off a shelf, when sudden symptoms occurred, after which the presence of a large stone was detected: here in all probability the calculus had been lodged and then suddenly ejected from its hiding-place.

Pain may be altogether absent; and such is the case when there is paralysis of the bladder, since then there is no desire to urinate, nor any efforts to expel the urine. After all, pain is no positive sign, for it may occur in other diseases; hence it merely directs the attention of the patient and surgeon.

In gouty patients a calculus in the bladder is attended with much pain and irritability, which is sometimes so severe that relief is anxiously implored.

Associated with pain is an alteration in the act of micturition: ordinarily there is a frequent desire, from irritation of the neck

* Hunter's Works, edited by Palmer, vol. i. p. 321.

of the bladder, and there are futile attempts to evacuate the last drop; again, during the flow of urine there may be one or more sudden stoppages, but for a moment only, the stream reappearing in the ordinary manner. At times there is experienced considerable difficulty in urinating, and in some rare instances this has amounted to retention. In cases where the stone has attained a very large size, filling up and occupying the whole cavity of the bladder, incontinence of urine takes place, so that the urine flows in drops, the bladder having no room for receiving the fluid. All these varieties are not peculiar or essential to the presence of stone, for they are met with in other diseases of the bladder and urethra.

The character of the urine is of little assistance in the diagnosis, for urinary deposits may exist without calculus, and calculus may be present without any deposit whatever. The urine may remain perfectly clear and natural for a long period; it may also contain deposits, which may assist in ascertaining the nature of the calculus; but it may become at any moment turbid and mixed with mucus, which will be deposited in the shape of a tenacious matter remaining at the bottom and sides of the utensil, as seen in catarrh of the bladder. This secretion may become retained in the bladder, and may undergo decomposition, giving rise to a foetid dirty-grey discharge; or it may even assume a puriform character; or it may subside and entirely disappear for a short time. Albuminous urine is now and then met with, and may be due either to the presence of pus or blood in the urine, or it may be owing to diseased kidneys. Some practical deductions might be derived by investigating the relative importance of the different states and quantities of albumen in the urine in cases of stone; but hitherto this has not been done. Lithotomy has been successfully performed on many occasions in cases in which there has been a good quantity of albumen, even without pus or blood.

Hæmaturia is not always present: it generally takes place on expelling the last drop of urine, and occurs from the violent contraction of the bladder upon the stone; it may also result from sudden exertions—the jolting in a cart or carriage, the act of jumping, &c.; but by rest and the recumbent position it will speedily disappear.

Sometimes an erection of the penis is an attendant sym-

om, and in adults now and then involuntary emissions of semen cur.


In some patients, especially adults, there is tenesmus and a sense of bearing down of the rectum, with frequent and fruitless attempts to pass a stool; in others, especially children, there is frequent defecation, and often fæces are passed at every act of micturition; prolapsus of the rectum occurs very frequently in children, and sometimes in old men.

The physical signs of a stone in the bladder are to be derived from artificial or mechanical aid from without, giving to the senses of touch and hearing on the part of the surgeon distinct evidence of its presence. The means of attaining this is by the exploration of the bladder with a solid metallic instrument, which on coming in contact with the stone, enables one to feel the foreign body, and on striking it produces a ringing noise or sound; hence the term 'sounding' given to the operation, and 'sound' given to the instrument used.

The sound is somewhat similar to a catheter both in size and form, but is made of solid steel, which is well polished on its exterior, its length and size varying according to the age of the patient undergoing examination. The handle of this instrument is expanded and flattened, and also polished, in order to give a larger extent of surface for the fingers and thumb of the operator to receive tactile vibrations. The curve of a sound is generally that of the ordinary catheter; some prefer it shorter, curving at an angle of 45° , and others almost straight, except at its lower extremity, where it is bent or curved, and made somewhat bulbous (see p. 926). The reason why catheters are not ordinarily used is, that they are bad conductors of sound, and the presence of the stilet in the instrument may give rise to a rattling noise, a source of fallacy. A sound with a large curve may glide over a calculus, when lodged in a depression behind the prostate: hence in such cases, one with a short beak and at right angles should be employed.

The position of the patient in sounding should be the recumbent one; but in obscure cases it is necessary to vary this, and even to explore in the standing posture. The rectum should be empty, and the bladder moderately distended with urine. L. Civiale generally injects water previous to sounding.

The sound should be introduced carefully and slowly, and allowed to glide along the canal by its own weight; no force whatever is required, and in general no pain need be given.



right side; large calculi are general in the bladder, small and medium sized calculi on the right or left side.

Sounding not only enables the surgeon to detect the presence of calculi, but it assists him in ascertaining their number, and sometimes density. When first traversing the surface of the stone, the sound from side to side, and then from before backward. The presence of calculi may be sometimes detected by careful exploration. The results may prove deceptive where a stone is very small, or where a stone may likewise be made out by the sound, free, or fixed and encysted; whether on the right or left side, at the base or in the walls, of the bladder.

The noise emitted on the striking of a stone has led to opinions that the size of a stone may be thus detected; for some writers state that very movable calculi give but a feeble sound, while large calculi, as lithic acid and oxalate of lime, give a sonorous character; the light, friable calculi, as phosphates, emit a dull and scarce sound. The smooth surface of a lithic-acid stone, or the exterior of the oxalate of lime, may in some cases be detected by the instrument.

Difficulties and sources of fallacy

evacuation of the water in the bladder by means of a catheter may cause the stone to be felt.

Sir A. Cooper (*Lectures on Surgery*, vol. i. p. 348) says:—‘I have myself sounded and not detected a stone at one time, which I have afterwards felt. I have sounded and not discovered a stone which another surgeon has afterwards perceived. I cut a patient and extracted thirty-seven stones from his bladder, who had been sounded and declared not to have a stone.’

S. Cooper (*First Lines of Surgery*, p. 688) remarks:—‘In sounding, how possible it is to mistake a thickened indurated bladder for a stone in that organ may well be conceived, when it is considered that Cheseldon, with all his judgment and experience, actually cut no less than three patients, none of whom had any stone in the bladder at the time of the operation.’

The principal difficulties and sources of fallacy are, however, a large and deformed bladder,—a contracted bladder, unable to contain urine,—a sacculated or encysted bladder,—an irregular hour-glass contraction of the bladder,—calcareous matter adhering to the walls of the bladder,—an enlargement and roughening of the prostate gland,—calculi in the prostate and urethra, and tumours at the neck of the bladder. Again, the calculus may be coated with blood or mucus. Some of these difficulties may be overcome by varying the position of the patient, by the use of sounds of different sizes and curves, and by moderately distending the bladder with two or three ounces of tepid water. The presence of a stricture in the urethra may offer difficulties, which must be overcome in the ordinary manner by dilatation.


Sounding may produce cystitis and peritonitis; hence caution and gentleness are necessary in its use.

In some instances exploration with a lithotrite has cleared up a difficulty; but care must be taken not to mistake the opening and closing of the blades for the sensation of a stone. When using the lithotrite and engaging the stone, we can take a very fair measurement of the axis and size of a stone, a point which will be referred to in the essay on LITHOTRITY.

Sir Henry Thompson has suggested and employed a modification of the lithotrite,* which may act as a sound, catheter, and measurer.

Exploration of the base of the bladder by the introduction of the finger into the rectum may lift the calculus out of the

* Weiss's catheter-scoop with stop-cock. See Thompson, *On Pract. Lithot.* p. 219. This instrument in its original form was constructed for injecting and removing the debris of calculi from the bladder; by the addition of the stop-cock, it is also useful for measuring the size of the stone and for sounding. See the essay on SURGICAL INSTRUMENTS.



many stones the patients in the operation themselves, and matter.

The prognosis of this complaint character; there are no means of cure, unless by removal, through the urethra, or by the removal of the offending foreign body; otherwise, at a more or less remote period, the stone will pass through the urethra, now small, and in some rare instances causing the stone to be extruded from the rectum, vagina, or perineum. In exceptional cases. The surgeon always finds that although a stone has been successfully removed, in some isolated cases a second or even a third may develop.

The progress and termination of the disease is affected with calculi in the bladder, and is productive of much anxiety and suffering according to the condition of the patient, and the age and temperature of the body. In some rare instances there are no symptoms, and the stone is only discovered after the post-mortem examination, when a stone has been ascertained. I

irritated. The skin becomes dry and the excretion of urine frequent and distressing; and hypertrophy of the bladder takes place, so that there is no escape of the urine except in the reclining posture: in consequence, becomes in part retained and under fermentation, assuming an ammoniacal and offensive character. Now suffering now ensues; exhaustion, loss of appetite, and low fever set in; next, colic, delirium, and speedy death.

Ordinary results; but other changes may take place in the progress of the case: thus, acute cystitis may supervene in the patient, as evinced by the intensity of the pain, frequency of micturition, and muco-purulent condition of the urine; or the mucous membrane of the bladder may ensue; or chronic inflammation of the kidney, or even acute inflammation of the kidney, may set in, and carry the patient off by uræmia or pyæmia.

There may be also hyperæmia, hypertrophy of the coats of the bladder, irritable bladder, catarrh of the bladder, dysuria, hæmaturia, ulceration and sloughing, with perforation allowing of the calculus into the peritoneum, rectum, vagina, or uterus; or pelvic cellulitis may be induced, followed by suppuration.

Stones may become encysted either by means of the formation of a hernial protrusion of the mucous membrane through the muscular coats of the bladder, or by the presence of a parietal pouch of the bladder opening internally and receiving the stone in its sac. Sometimes a large and heavy stone becomes lodged behind the neck of the bladder, and here forming a nidulum or bed for itself, becomes more or less fixed.

The post-mortem appearances of those who have died with a calculus in the bladder will of course vary, according to the mode of death and the complications which have previously existed. Ordinarily they will consist of chronic cystitis and chronic inflammation of the ureters and pelvis of the kidney, with slow changes in the glandular structure of the organ.

The treatment will consist in palliating and relieving the immediate distress, and then resorting to such means as may produce a permanent cure. When a stone has arrived in the bladder we should acquaint the patient of the fact.

The palliative measures will comprise such treatment as may improve the general health and the secretions and excretions,

as well as relieve any complications which may exist, such as catarrh, cystitis, ischuria, enuresis, nephritis, prostatitis, &c. by the methods usually adopted in such diseases. Great attention must be paid to the diathesis of the patient suffering from calculus, as evinced by the nature of the sediment passed, whether it be lithic acid, phosphatic, or oxalate, and this must be treated as laid down in the early part of this essay,—by careful attention to diet and regimen, due regard to the action of the skin and bowels, and the use of sedatives and anodynes to relieve pain; not forgetting to correct the nature of the morbid urine by appropriate medicines, such as mucilaginous drinks, diluents, &c. Oftentimes the paroxysms of pain are relieved by the recumbent position, with the thighs and buttocks raised, and the body bent forwards to relax the abdominal muscles. A decoction of the *triticum repens* is highly recommended by Sir H. Thompson as giving great relief.

The curative means will comprise the removal of the stone from the bladder; and three methods have been advised and adopted: first, the attempted dissolution of the stone by chemical and other agents, such as the far-famed ‘lithontriptics,’ and the solution by electrolysis; secondly, the removal of the stone by mechanical means without a cutting operation, such as dilatation of the urethra, and lithrotrity; and thirdly, the extraction of the calculus by operative measures, commonly known by the term lithotomy.

The cure by lithontriptics has engaged the attention of surgeons, physicians, physiologists, and chemists, and has been had recourse to by quacks and empirics of all descriptions. Dr. Prout has fully pointed out the importance of the administration of increased quantities of fluid in certain calculous and other affections, showing that many comparatively insoluble substances are slowly dissolved away by the frequent renewals of the fluid in contact with them.

Lithontriptics have been employed internally, through the general system, by the administration of certain medicines and waters, as well as locally by the injection of solvents into the bladder and the use of electricity.

The constitutional remedies generally made use of consist in the prolonged use of alkalies, but often to the detriment of the general health; these are inadmissible in oxalate-of-lime calculi, as the latter resist their influence; so also in phosphatic calculi they are prejudicial, inasmuch as acids are of more

beneficial effect in arresting the growth of such stones ; but in the lithic acid variety much relief has accrued, and in some instances the growth of the calculus has been decidedly cut short by the use of such alkalies. The alkaline waters are supposed by some to act not merely by a process of solution, but by causing a kind of disintegration of the component parts of a calculus. Liquor potassæ and the carbonates and bicarbonates of soda and potash are the usual medicines employed. The Vichy waters have become renowned as a favourite and useful remedy in calculous disorders; and in some instances persons having undeniable stone, as ascertained by sounding, have had their symptoms relieved, and even removed, whilst partaking of the waters. The Académie des Sciences, in reporting upon the use of alkalies in cases of calculi, thus sums up this subject: 'Although we admit that alkaline bicarbonates are not without influence on calculi, yet these present sufficiently great difficulties in their mode of application to cause them to be regarded with caution.'

Some prefer the use of the borate of soda, as being more energetic ; and, again, others have suggested the use of gastric juice, malic acid, and the nitro-saccharate of lead as having dissolvent powers. The celebrated nostrum of Mrs. Stevens was made of calcined eggshells and soap.

However, up to the present time no satisfactory remedy has been discovered. The local use of lithontriptics, viz. the injection of solvents into the bladder, has also been sedulously tried, and many experiments have been performed on calculi to ascertain their solvents, which have then been employed by means of injection into the bladder through a catheter. Unfortunately the solution, if strong enough to be of any use, endangers the coats of the bladder, and when diluted its action is uncertain in the extreme. Diluted alkaline solutions are recommended for uric-acid calculi ; diluted acids, such as the hydrochloric acid, for phosphatic ; * and dilute nitric acid for oxalate of lime. Salts of lead, carbonate of lithia, nitro-saccharate of lead have been suggested ; the latter as being less injurious to the

* Sir B. Brodie has shown that phosphatic calculi might be greatly reduced in size, or entirely dissolved, by injecting a weak solution of nitric acid, two to two and a half minims of strong acid to the ounce of distilled water.

Dr. Hoskins employed a weak solution of acetate of lead (gr. j. ad ʒj.) with a mere trace of free acid. With a phosphatic stone double decomposition occurs : phosphate of lead, in the form of a fine granular precipitate, and an acetate of lime and magnesia are formed.

bladder. The free use of water, by introducing it through a double catheter and keeping up a continued stream for half an hour every two or three days, has been employed: Dr. Willis has recommended that the fluid should be placed in a reservoir at a sufficient height above the patient, and connected with the catheter by a tube provided with a stop-cock, by which means the flow of the solvent may be carefully regulated.

M. Pelouse, in the 14th vol. of the *Comptes Rendus* of the Académie des Sciences, thus sums up the conclusions that have been arrived at: 1st, the effects of different agents upon urinary calculi proceed very slowly, and are chiefly evinced upon the animal matter which enters into their composition; 2ndly, drinks and baths never effect a cure; 3rdly, the results of injection are problematical, and the danger of inflammation is not counterbalanced, as in lithotrity, by a rapid destruction of the calculus; 4thly, although the combination of lithotrity with injections increases the probability of success, yet it is most desirable to proceed with lithotrity.

The dissolution of calculi by electricity (electrolysis) has had its advocates. Sir W. B. O'Shaugnessy has recommended the transmission of an electric current by means of a properly constructed instrument; but it is difficult in its application, and is attended with risks to the bladder. The latest and most successful efforts are those of Dr. Bence Jones, who employed a solution of nitrate of potash, and decomposed this by the aid of a powerful galvanic battery. The nitric acid set free at the positive electrode would decompose the uric acid exposed to its influence, and the potassa evolved at the negative electrode would dissolve it; so that a uric-acid calculus placed between them would be disintegrated at both points. The battery employed was from five to twenty pairs of Grove's plates. From two to nine grains of uric-acid calculus were dissolved per hour at the temperature of the body; of oxalate of lime, half a grain to two grains per hour only were dissolved—the action was four times as slow as upon uric-acid calculi; of oxalate of lime and uric acid in alternating layers, four and a half to five grains were dissolved per hour; of phosphatic calculi, upwards of twenty-five grains were dissolved per hour. Dr. L. Melicher (*Oester. Medic. Jahrbuch*, 1848, vol. i. p. 154) tried to dissolve a calculus by the aid of electricity; it is stated that two experiments on the living body were successful.

Gruithuisen, in 1823, suggested the dissolution of stone by means of the galvanic pile, but no experiment has been made on man.

The removal of a calculus by mechanical means without the use of the knife is confined to a certain class of cases. In the early stage, when the calculus is small, we may assist nature in its expulsion from the bladder by administering alkaline purges, by dilating the urethra with a large bougie, and giving diuretics; the bladder should be allowed to retain as much urine as possible, and then a free and copious evacuation may carry the stone along with it. Should the calculus become lodged in the urethra, it may be extracted by means of forceps, such as described and used by Hunter, Civiale, Sir A. Cooper, and others.

In females this plan is very practicable, and we shall refer further on to it.

The removal of calculi by lithotrity, together with the arguments in favour of this proceeding, will be the subject of special consideration in the next essay.

We now come to the extraction of a calculus by means of an artificial external opening, a necessary operation when lithotrity is deemed inadvisable. This operation is termed lithotomy. There are three situations in which it may be performed, viz. in the perineum, in the hypogastric or supra-pubic region, and through the rectum; each of which will have to be considered.

1. *Perineal lithotomy.*—There are several methods by which we may remove a stone from the bladder through the perineum; but we shall at present only refer to the ordinary lateral operation, leaving the other modes to be described as attempted improvements thereon. The necessary requirements for the operation consist in having a firm, solid table, of sufficient height to bring the perineum in a line with the operator's face, he being seated upon a stool sufficiently low to suit the height of the table; the table should be narrow, so as to enable the assistants to have more complete command of the patient. Two rollers or bandages should be at hand for securing the proper position of the extremities.

The instruments employed are the following: a staff, which is to act as a guide or conductor for the knife to enter the bladder; it varies in length and size, according to the age of the patient, and consists of a grooved steel sound, curved like the ordinary

catheter, the groove being deeply made betwixt the lateral and convex aspects, and terminating within a quarter of an inch of the extremity, so as to prevent any further onward movement of the knife. The staff has been modified in shape, with the hope of facilitating the operation; thus Aston Key suggested and used a straight staff, which has been and is still employed by all the surgeons of Guy's Hospital; it is nothing more than a very long director, slightly bent at its extremity, having the groove placed in the median line. The knife also varies in shape and size, according to the operator's fancy; some use a common scalpel for the first part of the operation, and then it is exchanged for a button-pointed bistoury, to complete the section. Liston performed the whole operation with a common sharp-pointed, scalpel-shaped bistoury; and Key did the same, but his knife was slightly convex on the back near the point; it was longer in the blade than an ordinary scalpel, but about the same breadth. The knife used for children is to be of smaller and narrower dimensions, but not to any great extent. A probe-pointed bistoury is likewise necessary, should any subsequent enlargement of the section be requisite.

Again, some experienced lithotomists, and especially those of early days, prefer completing the operation with an instrument called a gorget; this is a broad wedge-shaped instrument, either *blunted* at the point and sides, or sharp and *cutting*; although in occasional use, it is seldom had recourse to nowadays—excepting, it may be, the blunt one, in cases of very deep perineum and of enlarged prostate—the surgeon preferring to make his requisite incisions to certain, instead of uncertain measurements. The forceps are of different sizes and shapes, to suit every variety of difficulty; ordinarily they are straight, and the blades so constructed as to allow of an oval space between them when closed, in order that the calculus, when seized, may not be crushed in the efforts at extraction; some are curved, so as to seize the stone should it be engaged in the upper part of the bladder, or fallen into a pouch behind an enlarged prostate, or encysted elsewhere. In children, a pair of common dressing forceps, somewhat lengthened in the handle, will answer all purposes. A scoop is likewise necessary, and proves at times an invaluable instrument; it is chiefly employed in levering out fragments when a stone, from its friability, has crumbled to pieces; but it is also of great assistance when the stone is small and eludes the forceps, and on its introduction,

together with the aid of the finger, the stone may be readily drawn out by a kind of leverage of the scoop and finger in opposite directions. In Russia a kind of spoon is used in almost all cases, and with great success and ease. A syringe and tube should likewise be within reach, in case it is necessary to wash out the bladder. A female staff and sound is sometimes needed; and a strong lithotrite or crushing forceps, for facilitating the extraction of a large stone.

In all cases having to undergo lithotomy, certain preliminary precautions have to be attended to, which may materially influence a successful issue. Key paid great attention to this point, but Liston was not an advocate of too much preparation. The secretions and excretions, if at fault, should be corrected by appropriate remedies, and the nervous system quieted by anodynes; the urine, especially, should be rendered as healthy as possible, by acids or alkalies, as suits the individual case; sometimes, however, the suffering is so intense that immediate operation is indispensable. It is highly necessary to have an empty rectum; hence it is usual to prescribe a mild purgative powder overnight, to be followed in the morning by an enema of castor-oil, taking care that a free evacuation takes place before operating. Some, as Coulson, prefer giving two enemata, the one to be administered three hours previous to the operation, and to consist of gruel, olive-oil, and salt; the second to be employed about one hour before the time, and to be composed of gruel, with the addition of twenty drops of tincture of opium. The perineum should be shaved if necessary.

The patient is placed upon the table, and chloroform administered; notwithstanding its depreciation by some lithotomists, it materially facilitates the operation, and its effects have not been found in any way to interfere with a successful result.

Four assistants are required besides the administrator of chloroform, viz. one for each side, so as to hold the patient in position, a third to take charge of the staff, and a fourth to assist the operator, in handing the instruments, &c. When under the influence of chloroform, the patient is placed and fixed in the proper position so as to expose the perineum to its fullest extent, and in an entirely perpendicular direction. This is done by flexing the knee and hip joints to such a degree that the patient's hands may be made to grasp the foot, when they are to be firmly secured by a well-adapted figure-of-8 bandage

over hand, foot, and ankle; the knees are now widely separated, and each held by an assistant, encircling it with his proximate forearm and hand, and embracing and drawing the foot outwards with the other hand, and thus steadily maintaining the proper position until the completion of the operation. The shoulders and chest have to be raised to the requisite extent, and the patient then to be slid along to the edge of the table. Some use a neck ligature, which passes under both knees and around the neck, but it is at present generally dispensed with. Some surgeons do not tie up their patients, but use a kind of table with upright props on either side of the lower end, which are connected by a transverse bar; over this the patient's legs are placed, and there maintained in position. This table is so made as to suit all cases. Lately, padded leather bracelets, fixed by hooks to firm leather anklets, have been recommended and used. Children need not be tied up.

The patient being now in an accurate position, the surgeon introduces the staff into the bladder: this is to be done in the same way as an ordinary catheter, whatever shape the instrument may be. Some introduce it before the patient is tied up, as being more easy; but this plan has the drawback of causing the escape of urine during the subsequent process. Again, the orifice of the urethra may be narrow, which is to be rectified by enlargement with a bistoury. The staff should always be of as large a size as the urethra will permit, but not of too great a size, as it may tear the urethra. It should always be made to strike the stone, and left in contact with it. When stricture exists with stone, the former should be rigorously attended to before proceeding to operate. Care must be taken not to mistake the promontory of the sacrum for a stone in sounding. We shall now give an account of the ordinary lateral operation with the curved staff.

A curved staff, of sufficient calibre and grooved deeply between the lateral and convex surfaces, is introduced into the bladder, and the end made to rest upon the stone; this is intrusted to an assistant to maintain *in situ* as delivered to him by the surgeon from first to last, with instructions not to depress the handle as the knife enters the bladder. Some surgeons prefer making the convexity of the staff to bulge in the perineum, so as to be more readily felt; others, as Liston, require it to be hooked against the symphysis, as a more steady security. The surgeon's left hand is thus left at liberty

throughout the whole operation to guide the knife, guard important parts, &c. &c.

The operator proceeds with the first step of the operation, viz. to lay bare the groove of the staff by an oblique perineal incision; and this of course requires careful previous anatomical knowledge of the neighbouring parts, and the course the knife ought to take to escape important structures, such as the bulb and its artery above, the rectum in the median line, and the internal pudic vessels to the outside. There is the raphé in the median line, succeeded by the anal aperture, and on either side the ramus and tuberosity of the ischium, leaving an intermediate triangular space almost free from danger. The point of commencement is a little to the left of the median line, at a variable distance in front of the anus, and the incision is to be carried obliquely downwards and outwards, midway between the anus and tuber ischii, to the extent of 3 to 3½ inches. Some discrepancy exists as to the exact spot of commencing: thus Cheselden, Keith, and Crichton, all of whom have been remarkably successful, recommend 1 inch in front of the anus; Brodie, Stanley, and Skey, about 1¼ inches in front; Fergusson and others 1½ inches. All these measurements refer to the adult perineum. The best practical direction is to commence about midway between the anus and scrotum.

The knife is plunged freely into the perineum and carried downwards and outwards, dividing the skin and superficial fascia, as also the transverse perineal muscle and its artery, and some fibres of the levator ani muscle, laying open the perineal triangle and ischio-rectal space. The forefinger of the left hand is then introduced into the wound through the loose cellular tissue to gain the groove of the staff behind the bulb, and just anterior to the resisting deep fascia, when the point of the knife, being slipped along the nail of the finger and thus guided by it, is made to enter the groove at this spot, viz. the membranous portion of the urethra. The left forefinger can also be used to depress and protect the rectum.

The second step is the deeper section, which is effected by carrying the knife along the groove of the staff through the deep fascia, the membranous portion of the urethra, with its muscles, the left lobe of the prostate gland, and a small portion of the neck of the bladder. This is done by maintaining the left forefinger still in the groove, and making perfectly sure that the knife is securely within it; then, by a slight movement

of the right hand, the knife is lateralised so as to correspond to the direction of the external wound, and steadily pushed on-wards, until a feeling of loss of resistance or a gush of urine takes place. This is one of the most important parts of the operation, namely, the mode and extent of dividing the prostate gland; and no surgeon should undertake the performance of lateral lithotomy in the adult without a due knowledge of the size, shape, and condition of the organ, and its relation with such vital surrounding structures as the recto-vesical fascia and peritoneum, the great object being to have a sufficient opening to allow of the passage of the finger, without bruising or tearing the parts. Some surgeons recommend small incisions and subsequent dilatation; others make a free section corresponding to the size of the stone, without regard of exceeding the limits of the gland. Liston is very minute and particular on this point.*

The third step is the introduction of the forceps and the removal of the stone. Now, although the second proceeding is one of the utmost delicacy and moment, the present far excels it in difficulty of execution, and is often the cause of the delay and trouble witnessed in lithotomy. The knife having been withdrawn, the left forefinger is carried on along the staff into

* 'This internal incision must be very limited indeed; it should certainly not extend beyond six or seven lines from the urethra outwards and downwards; for the less that is cut the greater will be the patient's safety.' At p. 510 (op. cit.) he introduces a sketch 'to show what is indispensable to cut—it is a section of the dense, unyielding fibrous tissue at the base and in front of the gland into which the muscular fibres are inserted; it appears as a white triangular band, effectually preventing dilatation or enlargement of the orifice of the bladder beyond a certain and very limited extent, without laceration, dreadful suffering, and imminent danger. The incision required to divide this need not pass very far from the groove of the staff; and if that is effected, the dilatation can be carried to any required extent. The object in following this method is to avoid all interference with the reflexion of the ilio-vesical fascia, from the sides of the pelvic cavity over the base of the gland and side of the bladder. If this natural boundary betwixt the external and internal cellular tissue is broken up, there is scarcely a possibility of preventing infiltration of urine, which must almost certainly prove fatal. The prostate and other parts around the neck of the bladder are very elastic and yielding, so that, without much solution of their continuity and without the least laceration, the opening can be so dilated as to admit the forefinger readily; through the same wound the forceps can be introduced upon this as a guide: and they can also be removed, along with a stone of considerable dimensions, say from three to nearly five inches in circumference in one direction, and from four to six in the largest.'

bladder; and on feeling the stone, the sound may be moved; the surgeon now ascertains its position and size, and proceeds with a suitable pair of forceps to seize and the calculus. The forceps are introduced shut, and by the finger into the bladder, the handles being gently so as to gain the fundus of the bladder, when the blades separated, one of which is to be made to slip under the calculus, and the instrument then closed upon the stone. The is then gently raised, by depressing the handles, so as to it towards the external incision, when extraction is to be in the direction of the axis of the pelvis slowly and gradually; the instrument is moved forwards and backwards so as to dilate the parts, and these are pushed back, as upon the stone, by the finger of the left hand, whilst the acting force is kept up.

surgeon may in this stage meet with perplexities. He introduces his forceps through the section he has made and finds evidence of stone, although the calculus was clearly discovered previous to the operation by its ringing sound and by evidence of other surgeons. This casualty must be borne in mind and the sources of fallacy considered: viz. 1st, the stone may have been a small one and expelled with the first of urine on section of the prostate; the stone may fall out to the ground, or become driven into the anterior part of the urethra, or have escaped into the rectum, should the latter have been wounded; 2ndly, the stone may be encysted, and hidden in one of the diverticula of the bladder, perhaps entering into the orifice of the ureter; 3rdly, the presence of a tumour at the neck of the bladder, behind which the stone has become lodged; 4thly, the section through the prostate not having been completed, so that the forceps, instead of entering the bladder, has passed between it and the rectum.

After the removal of the calculus, the finger must again be introduced into the bladder, so as to ascertain the existence or absence of any other stone; and if the perineum be very deep, the finger or sound may be passed.

Lateral lithotomy with the straight staff was first introduced by Mr. Aston Key at Guy's Hospital, where it is still used by the surgeons. It has its advantages and disadvantages; but upon the whole the straight is as satisfactory an instrument for a lateral incision into the bladder as the curved staff. Of course some modification in the mode of conducting the operation is

required, although the steps are almost precisely the same as in the foregoing operation, viz. to open the perineum and urethra, to open the bladder, and to extract the stone.

The operator with the straight staff must bear in mind, in the first place, that it is more liable to be tilted out of the bladder than the curved one; hence the surgeon is always very careful to ascertain that the end of the staff rests on the calculus previous to his commencing the incision. Again, it is also not so easily introduced along the urethra into the bladder when the patient is placed in the position for operation; but this difficulty is readily overcome by the practised operator. During the first step of the operation with a straight staff, there is generally a twofold or subdivided proceeding, viz. laying open the perineum first of all, and then laying open the urethra. This is necessary with the straight staff, as the membranous portion of the urethra is placed at a very great depth, and any attempt to enter the urethra early would involve the bulb and its artery. This is the chief objection to the use of this instrument. Thus, then, in the first incision the integuments and superficial fascia are cut through, and the knife made to traverse the transversus perinei muscle and fibres of the levator ani, giving a good external opening; the next point is to pass the forefinger of the left hand through the loose cellular tissue in the triangular space until it impinges upon the staff beyond the bulb, necessitating a much longer track than with the curved staff. The bulb is now protected, and the knife made to enter the membranous portion of the urethra; this must be thoroughly and unquestionably exposed: a point on which the success of the operation entirely depends.

The surgeon, while maintaining the knife in the groove of the staff, withdraws the left forefinger, and now takes charge of the staff from the assistant with his left hand; then depressing the handle and bringing it parallel with the axis of the pelvis, thus presenting a direct passage to the bladder, he lateralises both staff and knife simultaneously to the left side of the patient, so as to correspond to the external incision, and carries the knife onwards to the bladder, regulating the section of the prostate at the same time, or on its withdrawal. Key's knife has a double convex extremity; hence great caution is required to keep the point upwards in the groove of the staff at an angle of 45° : for if the staff and knife be held too parallel to each other the upper convex border of the latter will be tilted out of the

is just anterior to the prostate, and instead of entering the urethra, it will pass into the pelvic cavity. The third step is in laying the knife aside, and transferring the staff to the right hand, so that the left forefinger may be introduced into this wound, and guided by the staff into the bladder, so as to feel the stone, and when satisfied of its presence, maintaining it *in situ*, while withdrawing the staff. The forceps is then introduced along the finger, and the operation of extraction is completed with as already detailed.

After the removal of the stone, provided no casualties or difficulties have occurred, some experienced operators recommend the introduction of a gum-elastic tube through the wound into the bladder, where it is to be secured properly by tapes: this is done in children, for twenty-four hours; and in adults, for eight hours. Liston says, 'It is especially adapted for the purpose,' and he attributes much of his success to its employment, 'on occasions no inconvenience, and affords great facility for preventing any oozing that may arise from the vessels.' In the female, however, it is dispensed with.

The patient is unbound, carried to bed, and laid on his back, his body raised and his knees bent over a pillow, with the feet resting upon a draw-sheet; so that the latter may be changed away at intervals, as the urine saturates the sheet.

Some surgeons, as Crichton, recommend immediate closure of the wound, in order to gain union by the first intention. In some cases this has succeeded well, but others have been followed by severe inflammatory symptoms: hence, as a rule, it is not advisable or justifiable to adopt this measure.

Little or no medical treatment is required, except, perhaps, the exhibition of opium when there is great restlessness; and if the bowels are not relieved within six days a castor-oil emulsion, carefully given, will be sufficient.

Before adverting to the several varieties of operative measures substituted in lieu of lateral lithotomy, for removing and remedying the defects and difficulties which are encountered in that operation, and for correcting its faults, if there be any, let us briefly survey the several accidents and difficulties, as well as the causes of death, which are generally met with.

With regard to the passing of the staff, great attention and care must be paid to its use in children, as a false passage is easily made, so that the instrument enters the pelvis below the bladder instead of passing into it; and in some rare

instances unusual force has separated the urethra from the prostate. Hence we are never to operate unless the staff is left in contact with the stone. In children prolapse of the rectum is also very frequent; the gut must be returned, and the sphincter allowed to contract, before operating. The presence of the cicatrix of a former operation need not interfere with a second or even a third incision in the same situation. On several occasions at Guy's Hospital lithotomy has been performed on the same patient two or three times along the original seat of incision, and without difficulty.* Some recommend fresh incisions to be made on the right side of the perineum, should there be much induration and rigidity, with fistulous abscesses and sinuous tracks; or the surgeon may dilate, if possible, the original opening, cutting along the finger among the altered structures, guided of course by a staff in the urethra. If a perineal hernia exists, it is liable to be cut into; but these cases are exceedingly rare, and the accident is scarcely to be avoided.

Enlarged superficial vessels, or the transverse perineal artery, may be wounded and bleed, but are readily tied, if necessary: with care and due attention the bulb and its artery ought not to be injured (except in cases of very deep perineum), and the accident is mainly due to too great haste and anxiety to enter the groove of the staff early: when the bulb or its artery has been wounded, it is best to proceed with the operation, and endeavour to secure the bleeding vessel afterwards, by means of the tenaculum, or compressing forceps: of course if the hæmorrhage be violent, the vessel must be secured without delay.

During the second step, several serious accidents may arise; for if the knife be carried too much outwards, under the ramus of the pubes, the pudic artery may be wounded; or if carried too much inwards, with a full rectum and prolapsus of the organ, that tube may be involved; and lastly, which is by far the most important point, there may be too great or too little section of the prostate gland, and wound of the prostatic plexus of veins. In wounds of the internal pudic artery, it is best to request an assistant to compress the artery against the pubes, and to complete the operation of extraction, when the artery may be attempted to be secured by ligature; should this fail, a short gum catheter, around which some oil-silk is fixed, like to

* On this subject see *infra* the table of cases of relapse of stone occurring at the Norfolk and Norwich Hospital, with remarks by Mr. C. Williams, pp. 1065, 1068.

umbrella, is to be introduced into the wound, and the space between it and the catheter plugged with lint, cotton, or charcoal; this is the *canule à chemise* of Dupuytren. Sometimes there is an unusual distribution of vessels: but the case must be treated in the same manner. Where there is general oozing of venous hæmorrhage, cold water and ice to the perineum above the pubes are useful.

If wounding of the rectum has occurred to some of our operators, during the struggling of the patient, when the rectum was not in vogue; sloughing of the rectum from pressure and injury during the extraction of a large stone has also taken place. The urine will flow in part by the rectum, and a cure cannot readily take place. The earlier the sphincter is divided, as for the cure of complete fistula in ano, the sooner will be the patient's chance of being freed from annoyance.

As regards the section of the prostate gland, much weight is attached to the amount of its division in adults; whereas in children and young children there is no such fear, as the gland is rudimentary. Sir H. Thompson (*op. cit.*) remarks, that the prostate is so exceedingly small as to be almost always, if variably, cut wholly through in lateral lithotomy, yet without the occurrence of any accident. Indeed, it is not possible that either forceps or finger can pass into the bladder, as the incision exceeds the thickness of the organ.' Again, in children a difficulty may arise from the staff having escaped from the urethra by a false passage, so that the knife cannot be introduced into the bladder; or the urethra may be separated from the prostate; or again, the bladder may recede before the point of the knife: in the two former cases the staff must be partially withdrawn and manipulated, so as to reach the bladder by the natural passage, and a grooved director introduced through the wound may sometimes be passed. When the bladder recedes before the knife, so that the finger cannot be made to enter the urethra, Liston suggests the use of a searcher, whereby he would feel the opening of the bladder, as it were, down upon the finger; should, however, it be possible to introduce the bent first joint of the finger, the bladder can be pulled down and the stone felt. In the adult, the section should never exceed the external limits of the prostate; this has already been alluded to. Sometimes the patient is remarkably corpulent, and has a very fatty perineum, and probably also an enlarged and rigid

prostate and a receding bladder, making a very long space to be traversed, so that there is difficulty in making the internal opening freely, and the finger will scarcely reach the prostate; in such cases, a probe-pointed knife may be introduced to complete the section,—a lithotome may be used; but by far the most judicious proceeding is the introduction of a blunt gorget, carefully directed onwards to the bladder, and not allowed to escape from the staff: in fact, the gorget used formerly to be, and is sometimes now, employed in completing the ordinary lateral operation. A hypertrophied prostate, or one deformed by tumour, may give rise to difficulties; so may also a deformed pelvis. Martineau used the blunt gorget with success. His object in this part of the operation was threefold: to make a small incision in the gland with the knife, to enlarge the incision with the blade of the blunt gorget, and to take time in drawing the stone from the bladder. In so conducting the operation, the gland is not bruised, but torn; the slow introduction of the gorget separates the fibres of the gland and enlarges the incision as far as the deep fascia, without dividing it; for the fascia yields and stretches as the gorget enters the bladder. Sudden violence in the act of introducing the gorget is nearly as much to be deprecated as the violent extraction of the stone. The kind of violence which the blunt gorget, properly employed, inflicts, is well borne by the gland, and symptoms of high inflammation are rare after such an operation.

Respecting the difficulties encountered in the extraction of stone, they are almost innumerable, and must be met according to circumstances: a few of these we shall just allude to. In cases of large calculi, Liston (*op. cit.*) maintains that ‘the single lateral incision affords sufficient room for removal of the stone in nineteen cases out of twenty; and there can be no purpose served, therefore, in always making a cut in both sides of the gland, and thus endangering the emasculation of the individual.’ He again, however, observes: ‘Should it be impossible to bring it through an opening on one side of the prostate such as described, without bruising and laceration of the parts, it will then be proper to provide more room for its exit; and this is gained by making the incision on the opposite side of the neck of the bladder. This is done by passing a narrow-bladed and blunt-pointed bistoury along the finger, and directing its edge towards the right tuberosity of the ischium. If the external incision has been well planned, properly made, and of

lent extent and low in the perineum, there is no occasion for further division of the skin and superficial parts. No complicated machine is required to make this bilateral division, it is quite time enough to have recourse to the proceeding if the necessity has been fully ascertained to exist.'

Stieaneau* remarks, that 'should the stone be large, or there be difficulty in the extraction, rather than use much force the forceps have firm hold of the stone, I give the handle to the assistant . . . while the part forming the stricture is which is easily done, as the broad part of the blade becomes anterior to the knife; and rather than lacerate, I have often avoided this enlargement of the inner wound two or three times.'

Some surgeons, more especially recently, have recommended the crushing and breaking up of a large calculus by a mallet, or strong crushing forceps, or cutting it in two by appropriate instruments, and then extracting. This has been sometimes effected successfully; but great danger and risk is necessarily incurred, for the bladder is generally firmly contracted on the stone, so that the coats and mucous membrane can hardly escape injury. Where it is so large, and the above measures fail, resort must be had to the recto-vesical or supra-pubic operation. If the stone is soft and friable, or brittle, crumbling to pieces under the application of the forceps, much of it may be removed therewith, and the rest evacuated by the scoop. Washing out of the bladder with warm water, by means of a syringe and tube, is highly useful, and attended with excellent results, in detaching any débris or concretion from the walls of the bladder without injury.

When the calculus assumes an elongated form, or any unusual shape, careful manipulation with the finger and forceps may bring it into such a favourable position as to allow of its extraction.

Where there is much spasm of the bladder, and a kind of ur-glass contraction, it is best to wait a few moments until the spasm subsided, and then carefully coax the forceps between the calculus and the walls, gentleness and patience being the chief requisites. Sometimes a stone is caught in the upper part of the bladder, and can almost be felt above the pelvis; by gentle pressure in the hypogastric region, it may be made to descend so as to be seized with the forceps. Encysted calculi often occasion an infinite deal of trouble, and sometimes will defy

* *Med.-Chir. Trans.* vol. ii. p. 411.

all efforts at extraction. In extracting phosphatic calculi, the operator must be on his guard, and recollect that sometimes calcareous matter is firmly deposited on the mucous membrane, and any attempt to remove it by the forceps will only tear away the tissue with it. Another point, which is alluded to by Sir H. Thompson, is, that sometimes in elderly patients portions of the prostate gland have been removed by laceration, such as the small rounded tumours in the substance of the organ, and the pedunculated growths met with in this situation.

The extraction of the stone is attended with more hazard to the gland. In this step of the operation, time must be given for the parts that embrace the stone to dilate; the muscular structure at the neck of the bladder, the firm substance of the prostate gland, and the deep fascia, will each stretch and yield to the sustained efforts of a firm but gentle hand. The gland is sometimes injured in withdrawing the stone, by the forceps embracing a portion of it, and tearing it away from the body. This may be avoided by passing the finger below the forceps, and disengaging the gland from the forceps after the stone is seized.

A nervous or a violent operator, feeling some resistance to his efforts at extraction, redoubles them, until he finds the stone obeying the force that he employs; and the operator, in ignorance of what he is doing, drags the gland before the stone, separates it from its attachment to the deep fascia, and brings it nearly to the external aperture before his efforts succeed. The consequence of such an injurious proceeding is to bruise the gland, to cause it to slough, and to render infiltration of urine almost inevitable, by the laceration of the deep prostatic connections.

Before proceeding to the question of mortality after lateral lithotomy, we will just allude to the complications which may ensue, although the patient has made a good recovery; thus, the wound may not entirely close, and leave a urinary fistula, which must be treated as other urinary fistulæ; incontinence of urine may result, either temporary or permanent. This is not very common: it is mostly met with in children, and may be cured by attention to the passing of the urine and the general health; as age advances, it generally passes off. Key* remarks, that 'in the young subject partial incontinence will sometimes occur, if the patient is allowed to leave his bed too soon after the operation, before the neck of the bladder is firmly healed and the

* *Guy's Hospital Reports*, vol. ii. p. 25.

ster has recovered its tone.' Impotency, from injury done to the ducts and vesiculæ seminales, is very rare, and is irremediable. Fæcal fistula sometimes follows, either from the use of the knife, or from the sloughing consequent upon laceration of the rectum during the extraction of the stone.

The causes of death after lateral lithotomy are the following : Death from the shock of the operation, without any disabling lesion, may occur, as in any of the ordinary operations ; it is very unfrequent, generally occurring in old people, and sometimes in children ; it seems to depend upon some peculiar idiosyncrasy of constitution and nervous system.

Death from chloroform may likewise occur, and the same remarks as just made about shock apply to these cases. Lithotomy in no wise militates against the use of anæsthetics.

Death from hæmorrhage and exhaustion may ensue, either primarily immediately after the operation, or secondarily some time after, from separation of the plug of coagulum in the lacerated vessel, or separation of a slough from a bruised vessel, or extension of a sloughing wound involving the vessels. Death from primary hæmorrhage rarely occurs, and although men can little stand any great amount of loss of blood, yet we sometimes see them rally after having been thoroughly blanched in an apparently hopeless condition. Liston states that the average is about 1 death in 100 cases of death ; M. Begin affirms that hæmorrhage is the cause of death in 1 out of every 4 deaths ; this is not so in this country. Although primary and secondary hæmorrhage may not *per se* cause death, yet it will so act as to cause great constitutional depression and debility, and frequently prevent the reparative process, and lay the foundation for general decline, or place the patient in a condition favorable for pyæmia.

Death from pelvic cellulitis is by far the most frequent in lithotomy. It consists in an acute inflammation of the tissues around the outside of the bladder, more especially the neck, and sides ; and is generally produced by urinary infiltration into the cellular tissue of the pelvis. Some consider it to be due to too great a section of the prostate, involving the deep perineal fascia, and allowing the urine to pass directly into the cellular tissue, and thus cause the inflammation ; others maintain that the infiltration of the urine is not a mechanical process, but generally follows inflammatory action, and that this inflammation is due to mechanical violence inflicted in removal of the

stone from insufficient section of the gland. In some instances death has taken place from cellulitis, quite independent of urinary extravasation. This inflammation may extend by contiguity to the peritoneum, inducing peritonitis; but it generally leads to purulent infiltration, deposits of pus, and pyæmia. The symptoms are those of severe constitutional irritation, attended with rigors, great prostration and anxiety, and soon followed by symptoms of low typhoid fever; the wound assuming an unhealthy aspect, and giving exit to a foetid discharge.

5. Death from acute peritonitis (*per se*) is rare, seldom occurring in adults, but now and then in children where the anatomical position of the bladder is closer to the peritoneum, so that too large a section of the neck and base of the bladder may involve it. Peritonitis used formerly to be considered as the most frequent cause of death after lithotomy, but subsequent investigations have disproved this.

6. Death from acute inflammation of the mucous membrane of the bladder (cystitis), extending to the kidney or peritoneum, is also rare, although Boyer has asserted it to be the cause of three-fourths of the deaths; this is not the case.

7. Death from inflammation of the prostatic veins (phlebitis) and its effects is not unfrequent, and generally occurs in persons of advanced life; the disease does not often set in until after a week or a fortnight, when recovery is hopefully anticipated.

8. Death from pyelitis, or inflammation of the kidney, or suppuration of the kidney, has been also occasionally met with; and at the post-mortem examination the advanced state of the disease has given evidence of its having existed long prior to the operation.

9. Death may also occur from absorption of the products from the urine, thus poisoning the blood and causing rapid death.

Erysipelas, diarrhœa, fever, or any other accidental attack, may render the operation fatal.

Many statistical inquiries have been made respecting the mortality after the lateral operation of lithotomy. These have been recorded by Crosse, Civiale, South, Coulson, and others, to whose works we must refer those of our readers anxious to investigate the subject. For the most part, these statistics cannot be relied upon as offering any direct assistance, except so far as a general survey, in estimating the value of the operation: they comprise compilations, selections, and returns, in which are many faults of omission and commission. This has been alluded to by Sir Henry Thompson (*op. cit.*), who has attempted to lay before the profession a more correct table, and this table may be used as a tolerably complete epitome of the mortality of lithotomy under British surgeons:

ved from reliable sources, mentioned in the text of his work, and 1,827 cases.

rates at all ages in 1,827 cases of lateral operation, in metropolitan and provincial hospitals.

	Cases.	Deaths.		Cases.
sh (Crosse)	669	91	about 1 in	7½
e that time	124	15	„	1 in 8½
.	110	14	„	1 in 8
er	90	8	„	1 in 11
.	29	4	„	1 in 7½
gham	102	10	„	1 in 10
Hospital	230	33	„	1 in 7
omas's Hospital	200	29	„	1 in 7
ity College Hospital	90	12	„	1 in 7½
dge	183	13	„	1 in 14
	<hr/> 1827	<hr/> 229		nearly 1 in 8

ave based the rate of mortality on the weight of the stone (see Crosse), drawn a conclusion that the mortality increases in nearly the same weight.

and perhaps with more reason and practical tendency, have considered the mortality in proportion to the different ages of the patients. This has been done by Sir H. Thompson in the following table. Still even this is open to objection, inasmuch as a stone may have been formed in the bladder and have remained there a long time previous to the operation.

Mortality rates at different ages.

the years inclusive	Cases.	Deaths.		Cases.
.	473	33	or about 1 in	14½
„	377	16	„	1 in 23½
„	178	19	„	1 in 9½
„	76	11	„	1 in 7
„	86	11	„	1 in 8
„	75	7	„	1 in 10½
„	100	17	„	1 in 6
„	191	40	„	1 in 4¾
„	233	63	„	1 in 3¾
„	38	12	„	1 in 3½
	<hr/> 1827	<hr/> 220		

we have taken the liberty to make a slight modification of the above, so as to show the mortality at the different epochs of life, which may be taken in some measure; but it is somewhat arbitrary:

Mortality rates at different epochs of life.

1 to 5, infancy and childhood, 473 cases, 33 deaths, about 1 in 14½ cases	Cases. 850 . .	Deaths. 49 . .	Cases. about 1 in 17½
6 to 11, boyhood, 377 cases, 16 deaths, about 1 in 23½			
12 to 16, boyhood, with increasing de- velopment and setting-in of puberty	178 . .	19 . .	„ 1 in 9½
17 to 29, adolescence and manhood	162 . .	22 . .	„ 1 in 7½
30 to 48, adults	175 . .	24 . .	„ 1 in 7½
49 to 70, advancing life	424 . .	103 . .	„ 1 in 4½
71 to 81, advanced life	38 . .	12 . .	„ 1 in 3½
	1827	229	avr. nearly 1 in 8

We have here inserted a table drawn up by Mr. C. Williams, comprising the cases operated on at the Norfolk and Norwich Hospital, during a period of ninety-eight years, from January 1772 to December 1869.

[See Table on page 1063.]

970 were male patients, of whom 871 were operated on by lateral lithotomy, with a mortality of 116; 64 by the median operation, of whom 13 died. Lithotripsy was resorted to in 34 cases, with a loss of 3; and 45 were female patients, of whom 2 died. The total number of patients on whom some form of operation was performed for the removal of the foreign body amounts to 1,015, which gives an average proportion of 10·46 cases per year. The total number does not represent the actual number of patients suffering from stone admitted into the Institution during the 97 years, but simply the number operated upon. During that period a much larger number of cases presented themselves, some of whom refused to submit to any operation; others were found to be suffering from disease of important organs—a condition which precluded the hope of a recovery, had any operative proceeding been adopted. The Hospital Reports for the last twelve years show that 205 cases of stone were admitted, and 170 only were operated on.

The lateral method having been performed on so large a number of patients as 871, renders it desirable to abstract that number from the above table, and give the result of that operation at the different epochs of life, in a separate and convenient form.

[See Table on page 1064.]

TABLE CONTAINING AN ANALYSIS OF ALL CASES OF STONE IN THE BLADDER OPERATED ON AT THE NORFOLK AND NORWICH HOSPITAL DURING A PERIOD OF NINETEEN YEARS, ENDING DECEMBER 1869.

Compiled by Mr. CHARLES WILLIAMS, Assistant Surgeon to the Hospital.

Age of cases operated upon.	No. of operations.	Recoveries.	Deaths.	Per centage fatal.	Proportion fatal.	No. of operations.	Recoveries.	Deaths.	Per centage fatal.	Proportion fatal.	No. of operations.	Recoveries.	Deaths.	Per centage fatal.	Proportion fatal.				
1 to 5	206	189	17	8.25	1 in 12.11	364	342	22	6.04	1 in 16.54	623	578	45	7.22	1 in 13.84				
5 „ 10	158	153	5	3.16	1 „ 31.6														
10 „ 15	81	74	7	8.64	1 „ 11.57	127	114	13	10.23	1 „ 9.76									
15 „ 20	46	40	6	13.04	1 „ 7.6														
20 „ 30	64	59	5	7.81	1 „ 12.8	132	122	10	7.57	1 „ 13.2									
30 „ 40	68	63	5	7.35	1 „ 13.6														
40 „ 50	65	51	14	21.53	1 „ 4.64	206	167	39	18.93	1 „ 5.28				392	305	87	22.19	1 „ 4.5	
50 „ 60	141	116	25	17.73	1 „ 5.64														
60 „ 70	154	116	38	24.67	1 „ 4.05	186	138	48	25.80	1 „ 3.83									
70 „ 80	32	22	10	31.25	1 „ 3.2*														
Total	1015	883	132	13	1 in 7.68	1015	883	132	13	1 in 7.68	1015	883	132	13	1 in 7.68				

TABLE OF CASES ON WHOM LATERAL LITHOTOMY WAS PERFORMED AT THE NORFOLK AND NORWICH HOSPITAL FROM JANUARY, 1772, TO DECEMBER, 1869 (A PERIOD OF 97 YEARS).

Compiled by Mr. CHARLES WILLIAMS, Assistant Surgeon to the Hospital.

[illegible]

TABLE OF CASES ON WHOM LITHOTOMY WAS PERFORMED A SECOND TIME AT THE NORWICH HOSPITAL. THE TABLE IS INTENDED TO SHOW THE AGE IN YEARS OF EACH PATIENT AT THE TIME OF HIS BEING FIRST OPERATED ON; THE INTERVAL IN MONTHS BETWEEN THE FIRST AND SECOND OPERATION; THE WEIGHT OF THE CALCULI REMOVED; THE RESULT; AND SOME OBSERVATIONS ON THE GENERAL CHARACTER OF THE CALCULI.

By Mr. CHARLES WILLIAMS, Assistant Surgeon, Norfolk and Norwich Hospital.

No.	Age at 1st operation.	Interval between 1st and 2nd operation.	Weight of calculi.		Result.	OBSERVATIONS.
			1st operation.	2nd operation.		
1	2½	15 months	3jss	3j	Cured	The first calculus phosphatic, and broken in the extraction; the second likewise phosphatic, and broken.
2	3	14 "	6 gra.	3ijss	"	One small oxalate of lime removed entire at first operation, and a mixed one at the second.
3	3½	24 "	3ijss	3ij 3ij	"	The first calculus was entire and phosphatic; the second likewise phosphatic and unbroken.
4	7	24 "	3jss	3ij	"	A perfect oxalate of lime stone taken away on first occasion; a phosphatic one on second.
5	8	17 "	3vj	3ivss	"	One entire lithic-acid calculus removed at the first operation, which left a recto-vesical fistula; the second was an entire phosphatic calculus.
6	15	16 "	ijj	3j	"	Three broken phosphatic calculi extracted at first; on the second occasion the calculus was of the same description.
7	18	8 "	3vj	3ivss	"	One phosphatic calculus, broken into four fragments, taken away at first; and one of the same composition at the second operation.
8	24	24 "	3xvj	3vj	"	A broken phosphatic stone removed on the first, and the same on the second occasion.
9	31	22 "	3j	3vj	"	First calculus (a mixed one) perfect; an encysted stone left undiscovered, which was removed at the second operation (median), together with one lying loose in the bladder.
10	33	104 "	3xvj	3ix	"	A single calculus of considerable size, consisting of alternate layers of dried blood and calculeous material which broke down on extraction, was removed at the first operation; a broken phosphatic one at the second. Lateral lithotomy performed on first and median on second occasion.
11	37	46 "	3v	3ijss	"	One perfect phosphatic calculus at first operation; also one of the same composition at the second.
12	48	12 "	5ij	3v	"	One flat lithic-acid calculus first removed entire; the stone removed at the second operation was of the same consistence.
13	60	34 "	3vij	3iv	"	Four entire uric-acid calculi removed on the first occasion, and one of the same material on the second.

(Table continued from preceding page.)

No.	Age at 1st operation.	Interval be- tween 1st and 2nd operation.		Weight of calculi.		Result.	OBSERVATIONS.
		1st operation.	2nd operation.	1st operation.	2nd operation.		
14	61	30 months		3iv	3v	Cured	This man had undergone lithotripsy twice previously to the first lithotomy, at an interval of twenty-eight months; an interval of thirteen months existed between the last lithotripsy and the first lithotomy. The debris after lithotripsy consisted principally of uric acid and phosphates; several phosphatic calculi, many of them broken, were removed at the first lithotomy, as well as at the second; a perineal fistula remained after the former operation; median lithotomy adopted on latter occasion.
15	63	32 "		5iv	5ijj	"	One entire lithic-acid stone at the first, and the same at the second operation; had stone a <i>third time</i> , but was deemed unfit for operation.
16	66	4 "		3j	3viij	"	A broken lithic-acid stone at the first operation, and five capped with phosphates at the second; median lithotomy resorted to in both instances; a perineal fistula resulted from the first, which was cured by the second operation.
17	68	7 "		3viij	3vj	"	Five entire and two broken uric-acid calculi came away at the first operation, and two perfect uric-acid at the second, on which occasion median lithotomy was performed.
18	71	28 "		5ixss	3xij		Three unbroken uric-acid calculi were removed on the first occasion, and one large phosphatic one on the second.
19	38	74 "		3viiij	5iv	Died	One entire lithic-acid calculus at first operation; a phosphatic one at the second.
20	46	12 "		3xss	3iv	"	One large entire lithic-acid calculus removed at the first operation, during which the rectum was wounded; and a recto-vesical fistula was present when the second was performed, on which occasion a phosphatic stone was taken away.
21	55	142 "		3viijss	3xvj	"	The first was a large oval unbroken uric-acid stone; second was of the same form and composition.
22	60	111 "		5iv ij	3xxv 3ij	"	The first calculus was a perfect lithic acid; the second, a large <i>urethro-vesical</i> lithic-acid stone; symptoms of recurrence of stone showed themselves three years after the first operation, lithotripsy was resorted to, and subsequently lithotomy.
23	62	70 "		3iv	6 grs.	"	One entire and two broken lithic-acid calculi removed at the first operation, and a small angular one at the second, on which occasion median lithotomy was performed.

CASES ON WHOM LITHOTOMY WAS PERFORMED A THIRD TIME.

No.	Age at 1st operation.	Interval between		Weight of Calculi.			Result.	OBSERVATIONS.
		1st and 2nd operation.	2nd and 3rd operation.	1st operation.	2nd operation.	3rd operation.		
24	7	16 months	11 months	3ij	3x	3iv	Cured	Two entire lithic-acid calculi were removed at the first operation; a phosphatic one at the second, and a perfect oblong phosphatic one at the third. The lateral operation adopted on each occasion.
25	57	78 "	11 "	3xijss	3ij	3ij	"	Five unbroken uric-acid stones at first operation; one phosphatic and broken at the second, and a single calculus of phosphates and urates removed entire at the third operation, which was lateral; the two former were median.
26	59	58 "	15½ "	3ij	3v	3ij gr. xv	"	One entire lithic-acid stone at first operation, and one of the same composition at each subsequent operation. The lateral resorted to on each occasion.
27	69	9 "	34 "	3viij	3ijss	3i	"	At the first operation several small phosphatic stones were taken away; a perineal fistula present at the second and third operations, which were performed according to the median plan; soft phosphatic calculi removed on both occasions. This patient had been lithotomised ten years before his first operation at the hospital, by Mr. C. M. Gibson, of Norwich; he had therefore been operated on <i>four</i> times, twice by the lateral method and twice by the median. He lived four years after the last operation.

Remarks by Mr. C. Williams.

This table presents 27 cases of recurrence of stone. Lithotomy was performed a second time in 23 instances, and a third time in four. The proportion that such cases bear to the whole number—1015—is one to 37·58.

Twenty-two were cured; five died; five had stone a third time, four were cut and recovered, the fifth was deemed unfit for operation; all the patients were males, no instance of recurrence having shown itself in the female.

Six of the cases were below 10 years of age; two between 10 and 20; one between 20 and 30; four between 30 and 40; two between 40 and 50; three between 50 and 60; eight between 60 and 70; and seven between 70 and 80. One death occurred below 40 years of age, and four above that period of life; the former (38 years of age), died on the 17th day, the cause being unrecorded; of the latter, one aged 47 died on the 20th day, cause unknown; the second, aged 65, died on the fifth day, cause unknown; the third, aged 70, died of exhaustion on the 74th day, and the fourth case also died of exhaustion on the 14th day.

The second operation took place, in six cases, within one year from the date of dismissal from the Hospital; in nine within two years; in four within three years; in three within seven years; in two within nine years; and in one within twelve years.

In four cases lithotomy was performed three times; in two of these patients, at the expiration of eleven months from dismissal; in one, fifteen months; and in the fourth, thirty-four months.

The lateral operation was performed twice in 17 cases, and three times in two cases. The median preceded the lateral once in one patient, and twice in another patient; and in five instances the lateral preceded the median. One patient was cut twice by the median plan, and another patient once by the lateral, and twice by the median.* The operation was performed in all the cases along the original seat of the incision, and without difficulty.

In 18, the calculi were removed in an entire condition;

* One patient had been operated on *four* times—twice by the lateral and twice by the median.


in seven they were broken previously to extraction; in two instances, where the stones were small and numerous, some were removed in a broken state, and others were perfect; in one case, a sacculated calculus was left undetected in the bladder (Case 9).

It does not appear that the calculus removed at the second operation is necessarily of the same character, in all cases, as that removed at the first; in 18 the second formation was of the same composition as the first, nine being composed of the phosphates, and nine of lithic acid and lithates; the phosphates succeeded the lithates in seven, and the oxalates in two.

Such are the practical points in connection with ordinary lateral lithotomy; we shall now cursorily allude to the several other modes of proceeding in the perineal section, as adopted in this country and elsewhere.

1. Lateral lithotomy modified as regards the section of the prostate:—

Fig. 306.



Vertical section of prostate, showing the various incisions recommended.

Boyer made a transverse incision of the prostate with the lithotome caché, dividing the gland in its greatest diameter (see Fig. 306, BB'); but the objection to the operation was the division of the perineal vessels, and consequent hæmorrhage. Senn combined the two, by making the ordinary oblique incision through the left lobe, and the transverse one through the right (see Fig. 306, AB').

2. The perineal incision, made in a transverse direction and of a semilunar curve in front of the anus. This was advocated by Dupuytren in his well-known bilateral section of the prostate gland in cases of large stone. He divided the tissues towards the symphysis, so as to avoid the rectum; the urethra was opened from above downwards (median) with a double-edged bistoury, and then his curved double lithotome was introduced into the bladder; the blades were then opened and

made to cut their way out, dividing the prostate in its oblique diameters (see Fig. 306, DD). The results of this operation gave 19 deaths in 85 cases, or 1 in $4\frac{1}{2}$ cases.

3. *Median lithotomy*, where the incision is made in the central line of the perineum, formerly known as the Marian operation. This comprises two modes of proceeding, viz. with section of the prostate gland, or without section of the gland, by merely opening the urethra, with perhaps partial division and incision of the apex of the organ. Vacca was the first to revive the median operation, and he practised a vertical section of the prostate; but, although there was no fear of hæmorrhage, the rectum was endangered, so that the operation could only be performed when the stone was small and in adult prostates (see Fig. 306, c).

Civiale in 1836 modified this procedure by his medio-bilateral operation. Having made his perineal incision in the median line, he avoided the bulb, and opened the membranous portion of the urethra, when he introduced a straight double-bladed lithotome into the bladder, and by withdrawing it opened, made a transverse section of the gland, as in Boyer's operation.

Buchanan* has further simplified this method by the following mode of proceeding: he uses a rectangular staff, which is one bent at right angles three inches from the point, and having a deep lateral groove, with a posterior opening. The instrument is introduced into the bladder, assisted by the left forefinger in the rectum, and thus the angle made to correspond in situation with the apex of the prostate gland, the lower or grooved branch lying parallel to the rectum. The angle rests on the farthest extremity of the membranous portion of the urethra, so that when the knife is plunged into the groove of the staff the membranous portion escapes all injury. The staff is intrusted to an assistant, and the operator, keeping the left forefinger in the rectum, takes a long straight bistoury, holding it in his right hand with the palm upwards, the blade horizontal and the edge directed to the left; he then enters the perineum opposite the angle of the staff, and passes the knife straight into and along the groove, as far as the stop at its extremity, into the bladder. Next he withdraws the bistoury slowly, making a lateral section of the prostate; but as he does so cuts outwards and downwards, a distance rather more than equal to

* *Edin. Monthly Journal*, Feb. 1848.

at one-fourth of a circle round the upper and left side of the anus, in which organ his finger still remains: an external and surrounding the corresponding part of the anus about $1\frac{1}{2}$ inches in length results from the operation. His arguments in favor of this procedure are: 1. the more easy and rapid method of reaching the prostate gland, being only two lines from the surface of the skin, the rectum pressed out of the way and the knife passed straight forwards into the bladder; 2. the membranous portion of the urethra is avoided and less incision required; 3. all blood-vessels are out of the way, and consequently there is no hæmorrhage; 4. the rectum, notwithstanding its proximity to the incision, is less likely to be injured; and 5. there is less risk of deep-seated infiltration of urine.

Millar's* operation is also performed in the median line, he does not make any section of the prostate gland, except a but partial, just incising the apex. The ordinary staff is introduced into the bladder, and then the left forefinger passed into the rectum, so as to feel the staff at the prostate: he then fixes with a straight bistoury the perineum in the median line about half an inch above the anus, carrying the knife on it strikes the groove of the staff at the membranous portion in front of the prostate; the urethra is pierced at this spot, and on pushing the knife towards the bladder for a few lines, it is withdrawn, at the same time cutting upwards, dividing the urethra a little, and finishing so as to have an external incision of $\frac{1}{2}$ to $1\frac{1}{2}$ inches. A long ball-pointed probe is then passed into the bladder along the groove of the staff, when the latter is withdrawn, and the left index-finger slid along the probe into the bladder, dilating the prostate and neck of the bladder to the requisite extent, and serving as a guide to the forceps. Where the stone is large he uses Weiss's three-bladed female dilator, Arnott's hydraulic dilator.

In cases of large stone Vidal de Cassis has advised a quadrilateral section of the prostate gland; he says, 'no matter what kind of external incision, as long as it is not too small, whether it be transverse, oblique, vertical, or curved: the point in view is to have one large external incision and many small internal ones. The two first sections of the prostate are to be made along the two inferior oblique diameters of the gland, which will be sufficient when the stone is of moderate size; but when

* *On Median Lithotomy.*

large, the two superior oblique divisions are to be made, first one and then the other' (see Fig. 306, AEEE).

We shall now pass on to *lithotomy through the rectum*,—the *recto-vesical operation*. This has been recommended in consequence of its supposed readiness of performance, and the easy passage of instruments, besides being free from hæmorrhage. The operation consists in placing the patient in the same position as for ordinary lithotomy; a grooved curved staff is then introduced into the bladder, and held by an assistant firmly and perpendicularly, so that the groove is in the median line. The operator takes a sharp-pointed straight bistoury in the right hand, resting the blade flatways on the palm or surface of the left forefinger, which latter is then introduced into the rectum to the extent of 10 or 12 lines. With the right hand the edge of the knife is turned upwards, and its point thrust through the anterior wall of the rectum, so as to gain the groove of the staff, and as the knife is withdrawn, it cuts through the rectum, the external sphincter and cellular tissue covering the urethra and integument in the median line, to the extent of one inch. The left forefinger is next carried into the wound of the sphincter, and the nail inserted into the groove of the staff, when the bistoury, with its edge downwards, is guided along it through the wall of the urethra into the groove, and pushed in the direction corresponding to the raphé, dividing the neck of the bladder and the prostate to a greater or less extent according to the presumed size of the stone; the staff having been removed, the finger is introduced into the bladder, and the calculus extracted by the forceps.

The drawbacks to the operation are the gliding of the mucous membrane of the rectum before the knife, the great risk of wounding the peritoneum and vesiculæ seminales, the subsequent occurrence of urinary infiltration, the passage of fæces into the bladder, fistulous sinuses, &c.

Recto-Vesical Lithotomy was in former years greatly in vogue with the Italian surgeons Vacca Berlinghieri, Giorgi, Cavarra and others; it was also adopted by the French surgeons Sanson, Dupuytren, &c. By some English surgeons it has been occasionally had recourse to, but is at the present time an operation generally discarded; so much so, that Malgaigne has erased the operation and mode of its performance from his recent edition of *Operative Surgery*. The best summary and

of this operation is that given by König,* which we modified and appended.

Age.	Cases.	Cures.	Recoveries with fistula.	Deaths.	Proportion of deaths.
Years	12	8	1	3	} 1 in 5
Under 10	13	9	2	2	
" 20	16	13	1	2	1 " 8
" 30	5	3	1	1	1 " 5
" 40	5	4	—	1	1 " 5
" 50	3	2	—	1	1 " 3
" 60	9	4	2	3	1 " 3
" 70	6	6	—	—	—
And over	9	4	2	3	1 " 3
And	5	3	2	—	—
	83	56	11	16	1 " $5\frac{2}{18}$

Hypogastric or high operation.—Epicystotomia is considered by some surgeons the most direct, short, and least dangerous operation. Its advocates argue that there are no major anatomical difficulties, with the exception of the incision, which is easily avoided. It suits all sizes of stone, and the surgeon to have a free opening into the bladder, and not so much danger of infiltration of urine as has been the case. The wounding of the peritoneum must be regarded as a minor anatomical manipulation. Pitha remarks, that in children and young persons below 20, as well as in females, the bladder stands high above the pubes, and presents an unobstructed portion freely to surgical manipulation; under 8 years of age the reflection from the bladder does not generally extend more than $1\frac{1}{2}$ to 2 inches from the navel. In old persons, especially when emaciated, the bladder lies deep in the pelvis, behind the symphysis, and is difficult to reach; hence the improbability of operating by this method in such cases.

Some surgeons maintain that there is great risk of peritonitis, extravasation of urine, and wound of the peritoneum, and that the operation is only applicable in cases of deformed pelvis; and where there is much fat it is rendered very difficult.

One old method was that of Franco, who introduced a finger into the rectum and pushed the stone up towards the hypogastrium, and its prominence formed the guide for the incision. A more cautious proceeding was afterwards adopted. The preparation for the operation consists in emptying the rectum

* *Journ. der Chirurg. von Graefe u. Walker*, Bd. viii. p. 529, 1825.

and distending the bladder, either by the patient retaining his urine or by injecting a sufficient quantity of tepid water to produce distension without over-excess, which latter is best performed under chloroform at the time.

The patient is placed on the back, with the pelvis raised, and the operation is performed in three steps. The first consists in laying bare the anterior wall of the bladder, by a vertical incision in the median line immediately above the pubes, cutting through the linea alba and fascia transversalis, and exposing the loose subfascial cellular tissue, in which some fat is often found.

The second step is that of opening the bladder, and is the most important part of the operation, requiring the greatest precision and care; the opening ought to be made close to the symphysis, and quite in the median line. Some surgeons have suggested and used an instrument called the sonde-à-darde, which consists of a trocar concealed in a catheter. This is introduced through the urethra into the bladder, and then the trocar is extruded and pushed through the fundus of the bladder at the external opening. This is considered both imprudent and mischievous. It is best to transfix the bladder through the wound, and thus secure it. The bladder is then to be opened by a median incision from below upwards. A few authors recommend a transverse incision. The external incision in the meanwhile must be kept open by retractors, in consequence of the tendency of the parietes to contraction.

The third step consists in the extraction of the stone, which may be readily done by a pair of straight forceps.

Civiale uses the sonde-à-darde, and has otherwise modified the operation.*

The after-treatment is now confined to simple measures; formerly a gum catheter was left in the bladder, others made a counter-opening in the perineum to drain off the water, and some used sutures in the bladder; but it is now considered more judicious to place the patient in bed on the back or sides with the legs drawn up, to employ water-dressing, and nothing else. No catheter, no canula, no other measure, is required.

Vidal recommends the operation to be performed 'en deux temps.' An incision is to be made in the median line, dividing all the tissues and exposing the bladder; then the operation is to be set aside, and the wound filled with charpie, which is to be

* See Thompson, op. cit. p. 72.

placed daily, so as to prevent union. If at the end of six, seven, eight days there be abundant suppuration, the operation is to be completed by opening the bladder with a straight bistoury, extracting the calculus.

Supra-pubic lithotomy had at one time many advocates, and it was strongly urged in those cases where there is a very deep abscess, with enlarged prostate, and a large-sized calculus. The most comprehensive work on the subject is that by M. Belmas,* whose carefully collected table of 100 cases will give the general idea as to the success of the operation.

Ages.	Cases.	Recoveries.	Deaths.	Proportion of deaths.
Under 5 years	9	8	1	1 in 25 cases 1 " 7½ " 1 " 9 " 1 " 1½ " 1 " 7 " 1 " 2½ " 1 " 1½ " 1 " 3½ "
From 5 to under 10	16	16	0	
" 10 " 20	15	13	2	
" 20 " 30	9	8	1	
" 30 " 40	7	4	3	
" 40 " 50	7	6	1	
" 50 " 60	7	4	3	
" 60 " 70	16	6	10	
" 70 and upwards	14	10	4	
	100	75	25	1 " 4 "

CIVIALE† gives a very unfavourable account of this operation performed in Paris, for of 75 cases, 35 died, and 35 recovered, 1 of these latter 5 had a fistula remaining. In 5 cases the result is not given. Gross‡ has constructed a table of 180 cases, but it is probable that many of the cases of one collector included in that of another. However, we have thought fit to offer the table as it stands in his work.

TABLE SHOWING RESULTS OF 180 CASES OF THE SUPRA-PUBIC OPERATION.

Operative collectors.	Cases.	Cures.	Deaths.	Proportions.
Mère Côme	100	81	19	1 in 5½
Longlas	4	3	1	1 " 4
Beeselden	9	8	1	1 " 9
Amal	4	4	0	
Dejeu	4	1	3	3 " 4
Calgill	4	3	1	1 " 4
Hornhill	16	13	3	1 " 5½
Seuberbeille	39	28	11	1 " 3¾
	180	141	39	1 " 4¾

* Belmas: *Traité de la Cystotomie sus-pubienne*. Paris, 1827.

† Civiale: *Parallèle entre la Cystotomie, etc.* Paris, 1836.

‡ Gross: *Diseases of the Urinary Bladder*, 1855, p. 636.

Dr. Humphry* of Cambridge operated successfully on a boy æt. 14, where the stone was of the size and shape of a small hen's egg, and weighed $1\frac{1}{2}$ ounce. He gives a table of 104 cases, mostly derived from the foregoing ones, and with the following result.

'In the 104 cases tabulated 31 were fatal, or somewhat less than 1 in 3. The age of the patient makes no great difference in the result, for—

In 38 cases	under 20 years	9 were fatal.
" 8 "	between 20 to 50 years	3 "
" 19 "	from 50 to 70 years	5 "
" 18 "	above 70 years	6 "
<hr/>							
83							23

'The cause of death appears to have been generally peritonitis, or infiltration of urine into the loose cellular tissues about the bladder.

'Maund is said to have lost only 5 in 40 operations; and of 100 patients operated on by *Frère Côme*, no more than 19 are reported to have died. These results are so favourable that I have been unwilling to admit them in the general table.'

In taking a general survey or making any critical comparison of the various principal operations and modes of removing a calculus from the bladder, much will depend upon the age and condition of the patient, as well as the size and nature of the stone. In the first place, the great point will be to determine whether the case be fit for the operation of lithotritry. This subject, however, is alluded to in the essay on that subject. When lithotomy is determined upon, there can be no doubt for a moment that in children under 12 years of age the lateral operation is the safest and most successful, the mortality being 1 in $17\frac{1}{2}$ cases only. Of course this does not preclude the performance of the median operation, where the stone is small or even of moderate size. At puberty and adolescence, and in manhood and adults, from 12 to 48 years, the mortality of the lateral operation rises to nearly 1 in 8, and from 49 to 81 years it amounts to 1 in 4 cases. It is in these periods of life, then, that some other modes of operating have been suggested to lessen the mortality. Statistics have not yet shown so much improvement as might be anticipated. Thus in Allarton's median operation on 139 cases at all ages, 13 deaths ensued, being 1 in 11; and when restricting this to adults, the mortality was 1 in 7; and

* *Trans. of Prov. Med. and Surg. Association*, new series, vol. v. p. 102, 1851.

Mr. Williams's report the mortality over 50 years of age is only 1 in 2 $\frac{1}{2}$. Of 60 cases operated upon by Buchanan's method, the result seems to be the same as in Allarton's operation, except that in adults the mortality is 1 in 8.

The bilateral operation of Dupuytren bears a death-ratio of 1 in 4 $\frac{1}{2}$, the recto-vesical about 1 in 5, and the supra-pubic 1 in 8.

These latter proceedings are only had recourse to in cases of very large stone and where complications exist preventing ineal lithotomy.

As far as our present information extends, the median operation may be performed with safety in cases where the stone is small; but the lateral operation is to be preferred in all other cases where lithotripsy is inexpedient.

The main issue, as regards the success of any one form of operation, seems, after all, to be chiefly dependent upon the mode of enlarging the opening of the urethra through the prostate and the neck of the bladder. This resolves itself into one of two proceedings; either dilatation of the prostate and neck of the bladder, as in the median operation, or section of these structures, as in lateral lithotomy. Is it safer to dilate, bruise, or lacerate the prostate gland and neck of the bladder, or to make a clean, careful incision with the knife? Small stones are treated whatever as to the value of either mode of operation, but are calculi in children; for in the latter, whichever mode is adopted, a recovery is expected.* It is in adults that we are always trying to lessen the mortality after the lateral operation, but none of the recent improvements seem to have supplied the deficiency. Even in Mr. Allarton's recent table, p. 499, 2d edit., the mortality is 1 in 7.

We should have gladly quoted the table inserted in that work, which consists of 170 cases of median lithotomy, but these do not in the least represent the actual state of affairs; the cases are for the most part taken from all sorts of reports of a successful issue; many known cases of death after the median operation are unrecorded; in two instances the author himself has been obliged, in consequence of the size of the stone, to abandon the operation, and terminate it by section of the prostate gland;

* Thus, in the last two years, at Guy's Hospital, fifteen children under four years of age underwent lateral lithotomy, without any unfavourable result; it is not improbable that median lithotomy might have been similarly successful.

which latter proceeding undoubtedly led to the successful termination. Median lithotomy, therefore, necessitates the bruising and laceration of the prostate gland, in cases of stone above a certain size, and in consequence will almost inevitably lead to a fatal result. The lateral section still maintains its position as the safest; it was so with Cheselden, Crosse, and numerous other surgeons.

The most perfect and unique collection of cases of the median operation, performed at a hospital of well-established reputation for lateral lithotomy, will give more insight into the comparative merits of the two operations; and we gladly avail ourselves of the opportunity of presenting to our readers one that has been so formed. By permission of Mr. Williams we are enabled to insert his tables, and thus compare the results of 64 cases of median lithotomy with the last 64 current cases of lateral lithotomy performed at the Norfolk and Norwich Hospital up to November 1869.

TABLE OF THE LAST SIXTY-FOUR CASES OF LATERAL LITHOTOMY
PERFORMED AT THE NORWICH HOSPITAL.

Ages of the patients operated upon.	Cured.	Died.	Total.	
Under 5 years	16		16	
5 to 10	11		11	
10 „ 15	1	1	2	
15 „ 20	1		1	
20 „ 30	1		1	
30 „ 40	1		1	
40 „ 50	2	1	3	
50 „ 60	6	1	7	
60 „ 70	15	4	19	
70 „ 80	2	1	3	
Total	56	8	64	

ABSTRACT OF THE CASES OF MEDIAN LITHOTOMY.

Ages of the patients operated upon.	Cured.	Died.	Total.	
Under 5 years	11	1	12	
5 to 10	4		4	
10 „ 15	4		4	
15 „ 20	1		1	
20 „ 30	1		1	
30 „ 40	5	1	6	
40 „ 50	1		1	
50 „ 60	7	2	9	
60 „ 70	14	9	23	
70 „ 80	3		3	
Total	51	13	64	

TABLE OF CASES ON WHOM MEDIAN LITHOTOMY WAS PERFORMED AT THE NORFOLK AND NORWICH HOSPITAL. THE TABLE SHOWS THE AGE OF EACH PATIENT—THE RESULT—THE NUMBER OF DAYS UNDER CARE AFTER THE PERFORMANCE OF THE OPERATION—THE NUMBER OF CALCULI REMOVED FROM EACH CASE—THE WEIGHT AND DIMENSIONS OF EACH CALCULUS—WHEN MORE THAN ONE STONE WAS PRESENT, THE DIMENSIONS OF THE LARGEST ARE GIVEN.

By CHARLES WILLIAMS, F.R.C.S., Assistant-Surgeon to the Hospital.

Age.	Result.	Days under care.	No. of calculi.	Weight.	Dimensions in inches.			Remarks.
					Length.	Breadth.	Depth.	
2	cured	22	1	14 grains.	$\frac{5}{16}$	$\frac{4}{16}$	$\frac{3}{16}$	
2	"	35	1	34 grs.	$\frac{5}{16}$	$\frac{4}{16}$	$\frac{3}{16}$	
3	"	23	1	14 grs.				
3	"	29	1	3ij	$1\frac{1}{16}$	$\frac{7}{16}$	$\frac{5}{16}$	
4	"	29	1	35 grs.				Broken in extraction.
4	"	15	1	16 grs.				Broken.
4	"	29	3	30 grs.				Small calculi.
4	"	36	1	47 grs.	1	$\frac{5}{8}$	$\frac{3}{8}$	
5	"	50	1	15 grs.				
5	"	22	1	9ij	$\frac{6}{16}$	$\frac{4}{16}$	$\frac{4}{16}$	
5	"	29	1	8 grs.	$\frac{6}{16}$	$\frac{4}{16}$	$\frac{3}{16}$	
6	"	22	1	3jss				Broken.
8	"	35	1	9v	1	$\frac{6}{16}$	$\frac{4}{16}$	
10	"	16	1	3 grs.		$\frac{5}{16}$	$\frac{3}{16}$	
10	"	29	1	9v	$\frac{7}{8}$	$\frac{5}{8}$	$\frac{3}{8}$	Removed entire. Lithates.
11	"	21	3	3iij 9ij				All broken.
12	"	28	1	6 grs.				
13	"	29	1	9j				Broken.
13	"	36	1	3ij gr. x	$\frac{3}{16}$	$\frac{3}{16}$	$\frac{3}{16}$	Oxalate of lime. Removed entire.
19	"	56	1	3iij gr. x	$1\frac{1}{16}$	$1\frac{1}{16}$	$\frac{7}{16}$	A perineal fistula remained.
21	"	29	1	5vjss	$1\frac{1}{16}$	1	1	Oxalate; broken.
33	"	22	1	3ij gr. x	$1\frac{1}{16}$	$1\frac{1}{16}$	$\frac{3}{16}$	
33	"	15	2	9xiv	$1\frac{1}{16}$	1	$\frac{3}{16}$	One weighed 9viij, the other 9vj. The dimensions of the larger only are given.
38	"	36	1	3j	$\frac{5}{16}$	$\frac{4}{16}$	$\frac{3}{16}$	Broken into 3 pieces; phosphates and lithates.
38	"	22	1	3j	$\frac{7}{16}$	$\frac{7}{16}$	$\frac{3}{16}$	Lithates; slightly fractured.
40	"	78		3ix 9j				Phosphatic; broken into fragments; a fistula remained.
47	"	28	2	3viij	$1\frac{1}{16}$	$\frac{5}{16}$	$\frac{3}{16}$	Lithic acid.
52	"	21	3	5v 9ij	1	$\frac{5}{16}$	$\frac{3}{16}$	Lithic acid. One removed perfect; the rest broken.
57	"	29	5	3xiij 9ij	1	1	$\frac{4}{16}$	Each weighed about 3ij.
57	"	29	1	3iij	$1\frac{1}{16}$	$1\frac{1}{16}$	1	
59	"	16	1	5i gr. x	$1\frac{1}{16}$	1	$\frac{3}{16}$	
59	"	36	2	3iv	1	$\frac{5}{16}$	$\frac{3}{16}$	Lithic acid; both removed entire.
59	"	63	1	3vss	$1\frac{1}{16}$	$1\frac{1}{16}$	$\frac{3}{16}$	Lithic acid; entire.
60	"	15	4	3viij	$1\frac{1}{16}$	1	$\frac{3}{16}$	Each weighed about 3ij.

Table continued from page 1079.

No.	Age.	Result.	Days under care.	No. of calculi.	Weight.	Dimensions in inches.			Remarks.
						Length.	Breadth.	Depth.	
35	61	cured	37	1	3v				Broken during extraction.
36	61	"	31	10	3ij				Several were broken.
37	61	"	28	1	3j	$\frac{7}{8}$	$\frac{2}{3}$	$\frac{3}{16}$	Lithates and phosphates; unbroken.
38	62	"	58	several	3viiij				Some entire; others broken. Lithates and phosphates.
39	62	"	36	1	3iv	$\frac{7}{8}$	$\frac{6}{8}$	$\frac{3}{16}$	Lithic acid; removed entire.
40	63	"	69	1	3iv 3v	$2\frac{6}{8}$	2	$1\frac{1}{4}$	A recto-vesical fistula remained.
41	64	"	50	1	3iv 3ij	$1\frac{3}{8}$	$1\frac{1}{8}$	$\frac{3}{16}$	Lithic; entire.
42	64	"	29	1	46 grs.				Phosphates; broken.
43	66	"	42	1	22 grs.				Broken during extraction.
44	66	"	36	5	3viiij				Median operation performed a second time; a fistula remained after the first, which was cured by the second. Same case as No. 43.
45	67	"	30	1	3ij 3ij	$1\frac{3}{8}$	$1\frac{1}{8}$	$\frac{3}{16}$	Phosphatic and broken; a perineal fistula present from a former lateral operation.
46	68	"	29	2	3ij	$1\frac{3}{8}$	$\frac{1}{8}$	$\frac{3}{16}$	
47	70	"	22	1	3ij				
48	70	"	37	1	3ij	$1\frac{3}{8}$	1	$\frac{3}{16}$	Lithic; entire.
49	72	"	50	1	3j				Broken; same patient as No. 47; fistula remained unhealed.
50	73	"	42	several	3ijss				Phosphatic; all broken.
51	73	"	29	1	3ix	$1\frac{6}{8}$	$1\frac{1}{8}$	1	Entire; coated with phosphates.
52	$1\frac{1}{3}$	died	1	2	5ss				Both broken.
53	37	"	13	1	3vij				
54	55	"	3	2	3vij	$1\frac{1}{16}$	$1\frac{1}{16}$	$\frac{1}{16}$	
55	57	"	11	1	3j 3ij	$1\frac{1}{16}$	$1\frac{1}{16}$	$\frac{1}{16}$	Each weighed about 3v.
56	61	"	9	2	3x 3ij	$1\frac{1}{16}$	$1\frac{1}{16}$	$\frac{1}{16}$	
57	61	"	9	1	3ijss	$1\frac{1}{16}$	$1\frac{1}{16}$	$\frac{1}{16}$	Lithic; entire.
58	62	"	18	1	3vj	$1\frac{1}{16}$	$1\frac{1}{16}$	$\frac{1}{16}$	
59	64	"	86	3	3x				
60	65	"	11	several	3ij gr. x				Broken previously to extraction; several found in the bladder after death.
61	66	"	4	1	3vss	$1\frac{1}{16}$	$1\frac{1}{16}$	$\frac{1}{16}$	Lithic, all broken: 9 found in the bladder after death.
62	66	"	2	1	3j	2	$1\frac{1}{16}$	$\frac{1}{16}$	
63	66	"	4	1	3xij	2	$1\frac{1}{16}$	$\frac{1}{16}$	
64	68	"	14	1	6 grs.				

The average number of days that each of the 51 cases of cure was under care, after the performance of the median operation, was 34.

The average number of days that each of the 56 cases of cure was under care, after lateral lithotomy, was 42.

In 64 lateral lithotomies 8 died, or one death to seven recoveries. In 64 median lithotomies 13 died, or one death to four recoveries. The average number of days each case survived the former operation was ten, the latter fourteen.

The Number and Weight of the Calculi removed from those who died after Median Lithotomy.

Under 5 years (æst. 1½) . . .	1 calculus, weighing	3ss.
From 30 to 40 years . . .	1 " "	3vij.
From 50 to 60 " . . .	{ 2 calculi " 3vij.	
	{ 1 calculus " 3j 3ij.	
	{ 1 " " grs. vj.	
	{ 1 " " 3vss.	
	{ 1 " " 3j.	
	{ 1 " " 3xij.	
From 60 to 70 " . . .	{ 1 " " 3vj.	
	{ 1 " " 3iiss.	
	{ 2 calculi " 3x.	
	{ 3 " " 3x 3ij.	
	{ several " 3ij gr. x.	

The Number and Weight of the Calculi removed from those who died after Lateral Lithotomy.

From 10 to 20 years . . .	1 calculus, which weighed	3ij 3j.
" 40 " 50 " . . .	1 " " "	3ix.
" 50 " 60 " . . .	2 calculi, weighing	3ij.
	{ 1 calculus " 3ivss.	
" 60 " 70 " . . .	{ 1 " " 3j.	
	{ 1 " " 3iv 3j.	
	{ 1 " " 3x.	
" 70 " 80 " . . .	1 " " "	3iv.

Remarks by Mr. Williams on the cases of median lithotomy.

Two cases were cut twice by the median plan, at intervals of nine and four months respectively.

In no case did recovery result when the calculus exceeded 3 drachms 2 scruples; except in one case, in which the stone weighed upwards of 4½ oz., but a portion of the rectum and perineum sloughed, and a perineo-recto-vesical fistula was established.

In no case did a cure result when the long diameter of the calculus exceeded $1\frac{1}{2}$ inches, and the short $1\frac{1}{8}$ inch, except in the case in which the stone weighed upwards of $4\frac{1}{2}$ oz.

Calculi in the prostate gland.

The prostate gland, like other glands, is liable to an inspissation of its secretion, producing small yellow, sometimes red, pale or colourless bodies scattered throughout the follicular structure. These at first are said to consist entirely of organic matter, which Virchow believes to be derived from a peculiar insoluble protein substance mixed with the semen; but sooner or later these formations are believed to irritate the mucous membrane, causing phosphatic depositions, which become encrusted upon the organic matter, and thus the genuine prostatic calculi are formed.

Prostatic calculi, therefore, have their development and origin in the gland-tissue. Their number varies from 1, 2, 3, to 100 or more, like a bag of marbles. Their size may be that of grains of sand, or may increase to that of a poppy-seed or cherry-stone. As they gradually enlarge, they come in contact with each other, and undergo a degree of attrition, whereby they form facets or articulations, and the surfaces are often rendered as smooth as porcelain.

Their form, then, generally is very irregular, sometimes triangular, square, &c.

Their colour is generally that of brown chestnut, sometimes whitish, black, or red; the brown coating is supposed to be derived from the secretion of the gland; sometimes they are transparent, pearly, like pearl-barley.

Their structure and consistence also vary; they are sometimes of not very great consistence, and amorphous; but generally in concentric layers around a central nucleus: occasionally with radiating striæ; and sometimes they are compact, firm, and brittle.

Their chemical composition, according to Dr. Wollaston, is:

Phosphate of lime	.	.	84.5
Carbonate of lime	.	.	0.5
Animal matter.	.	.	15.0
			<hr/>
			100.0

Although prostatic calculi in their early formation are seated

the gland-substance, becoming enveloped on all sides by its capsules, and forming for themselves a bed in the lateral lobes, when enlarged encroaching upon the rectum, so as to be felt by digital examination; yet more frequently they distend prostatic ducts, converting them into pouches, and appear as open mouths, so as to be felt by the sound in its passage through the prostatic portion of the urethra. These calculi may extend into the urethra, and thus have a free urethral surface and an adherent prostatic portion. Either of these surfaces become enlarged, the prostatic portion attaining the size of an ordinary urinary calculus, and its urethral portion may extend backwards into the bladder, forming a prostatovesical calculus.*

Sometimes one or more prostatic calculi may be detached from their recess and lie free in an open duct.

The other forms of calculi in the prostate are derived from a different source; thus, a urinary calculus may escape from the bladder when small, and become lodged in the prostate: it may either obstruct the canal, or may be expelled along the urethra; or it may form a bed for itself in the prostate and either by ulceration or pressure. These calculi are readily recognised on section as presenting all the characters of the ordinary urinary calculus, and as being totally different from the true prostatic calculus. But they so far resemble each other, that where a urinary calculus becomes lodged in the prostate, the successive deposits hereafter ensuing consist of phosphate and carbonate of lime; so that in external appearance they are somewhat like that of the true prostatic calculus. After lithotrity, and even after lithotomy, fragments of stone may become arrested or left in the prostate; so that after one, two, three, or more years, a prostatic calculus may be developed, the nucleus being a fragment of a urinary calculus.

There is one other rare variety (the vesico-prostatic); it is that in which a vesical calculus is situated within the bladder, and its neck, which on increasing in size sends a prolongation through the vesical opening of the urethra into the prostate gland. Here again it enlarges, both vesical and prostatic portions increase, and their bar of union remaining stationary produces the same condition as is observed in the prostatovesical.

*See collection of cases in the *Guy's Hospital Reports*, 3rd series, vol. iii. 51.

vesical variety, viz. a dumb-bell or bar-shot-shaped appearance. It is singular that in the majority of instances the urethra above was perfectly pervious, allowing a free flow of urine and the passage of a sound over the stone.

The symptoms depend, of course, upon the size, number, and situation of the calculi, and vary exceedingly; being sometimes very slight. Indeed in some instances there have been no symptoms whatever, and the presence of the calculi was only ascertained after death: when the calculi are large, and encroach upon the urethra, there is a sense of pain and weight in the perineum, and frequent micturition, irritation and uneasiness at the neck of the bladder; the prostate may become inflamed, and abscesses follow. Sometimes there are continual semi-erectations of the penis, and difficulty in the emission of semen.

The diagnosis is arrived at by exploration of the urethra by means of a sound or lithotrite, and the introduction of the finger into the rectum. Soft bougies of a large size have been recommended, which are to receive the impressions of the stone; but these impressions may become obliterated on withdrawal of the bougie.

The treatment will consist in removing them by means of a properly constructed forceps,* when the calculi are small and free; but, generally speaking, an external incision is requisite. The operation is conducted in the same way as for lithotomy, with these exceptions: the incision is to be made in the median line, and should include a sufficient section of the prostate for the escape of the calculus. If there are no symptoms, some recommend that the case should be left alone; others advise that when the symptoms are urgent the calculus should be pushed back into the bladder: this latter suggestion only applies to lodged urinary calculi or fragments of calculi, where the urethral forceps have failed. Where the stone has attained a large size, lateral lithotomy must be performed.

Calculi in the urethra are for the most part derived from the bladder and kidney, or from fragments of calculi becoming arrested at the narrower parts of the canal, more especially the membranous portion, there lodging and setting up various

* Such as Weiss's long urethral forceps, $8\frac{1}{2}$ inches long; the ends of the blades are not pointed, should not meet each other, and should be slightly scooped; or Weiss's canula forceps.

ptoms. However, in some very rare instances the calculus is formed in the urethra, in this wise: urine collects and a stricture or constriction, and a pouch or diverticulum ensues, wherein urinary deposit may take place and form nucleus of a calculus, generally of the lithic-acid variety; it is more probable that a particle of calculous matter or all particles of gravel have become arrested in this pouch and rise to stone.

The size of the stone varies; where there is only one, it is usually of a larger calibre than the urethra in its transverse diameter, and assumes an elongated form; they may acquire a large size; where there are several, they are often small faceted, generally plane and smooth.

The symptoms are very variable—in some cases there are none at all; in others there is acute pain at the site of the calculus, and if this be rough, the pain is very excruciating. Urine may escape freely, or it may be passed frequently, or there may be a frequent desire with inability to micturate, or there may be actual retention; this latter symptom is more recently present where the stone is lodged in the fossa navicularis. When left alone, dilatation of the urethra ensues, evasions and pouches are induced, ulceration of the urethra follows, and sometimes rupture, causing infiltration and extravasation of urine, involving the scrotum; this is attended with irritation, fever, and constitutional disturbance. Suppuration may follow, and the abscess burst externally, allowing exit of the stone, which is succeeded by urinary fistula. Stricture and enuresis may likewise be produced; and in some instances suppression of urine and uræmia may follow.

The diagnosis is generally easy by careful exploration of the canal with a small-sized sound or a long probe, the finger at the same time being passed along the outside of the canal, so that when the instrument impinges upon the stone, the finger may be placed on its vesical side, so as to prevent the pushing back of the stone and its slipping into the bladder. Rectal examination is oftentimes of service.

The treatment by endeavouring to facilitate the escape fords by means of suction and insufflation, as practised in former years, is obsolete, and inefficient for its purpose. There is always a gradual tendency of the urine to push the calculus forwards; hence the surgeon often directs compression of the urethra to be made in front at the moment of micturition: when

the urine has been retained in the bladder and urethra as long as possible, the compression is to be suddenly withdrawn, and the calculus may often follow in the gush. This failing, a warm bath should be ordered, and the urethra dilated with a bougie, which may prove successful. The case, however, generally resolves itself into the attempt at the removal of the calculus by instruments passed into the urethra; and here numerous instruments have been devised both in this and in other countries, and with some satisfactory results. In some instances these have succeeded in their object; in others a failure has been experienced. We must bear in mind that too much endeavour at extraction by means of forceps is bad, and detrimental to the canal and the patient. The quickest, easiest, and safest remedy, after failure of the forceps, is the incision in the median line over the seat of the stone. Make a good long clean cut at once down upon the stone, and turn it out, taking care it does not slip back by planting the finger or thumb of the left hand firmly upon the urethra behind the stone. There is no risk from a clean incision either of extravasation or fistula; for the incision generally heals quickly without trouble, and without any subsequent stricture or fistula. When the stone is opposite the scrotum, provided it cannot be extracted per urethram, it is best to push it back into the perineum, and thence extract it by incision. Incisions into the urethra in front of the scrotum are very often difficult to heal, and leave a fistulous opening. Should the calculus slip back into the bladder during the median operation, a long pair of slender forceps may be introduced, and the stone removed.

Calculus in the female, although not very frequent, is by no means so rare as is usually supposed. The peculiar anatomical arrangements of the urethral canal from the bladder to the exterior render any obstruction to the passage of a small calculus comparatively unfrequent; there is no prostate gland, and there are no natural contractions of the canal; the urethra is short and readily dilatable; there is scarcely any curve. So that although females may be prone to calculous depositions, as well as males, yet they escape the dire consequences of its retention in the bladder by the above arrangement of the parts. We cannot judge of the frequency of the disease in females as compared with males by the proportion of cases operated upon; for in the one instance a large proportion of the calculi escape,

re removed by dilatation without operation, while, on the other hand, nearly all the males undergo lithotomy or lithotripsy; hence we have no data to rest upon.

The symptoms, when a stone has become so large as not to pass from the bladder, will consist of many of those already detailed in the male; but the special symptoms in the female consist in,—1st, bearing-down pains and pains along the urethra; 2nd, incontinence of urine. These are sometimes very distressing, and call for urgent attention.

The symptoms often simulate those of uterine disease, for which they are not unfrequently mistaken; so that the surgeon should be on his guard, and on failing to detect uterine mischief, he is bound to pay special attention to the bladder and urethra. Sounding and vaginal examination seldom fail to detect the stone.

The natural progress of calculus in the bladder which is not controlled by nature, and where no remedial measures have been resorted to, will be the ulceration of the stone through the bladder into the vagina, by which a cure is sometimes effected, at the expense of a vesico-vaginal fistula; a circumstance which is always to be avoided if possible.

The methods for extracting calculi from the female bladder are the following:

1st. *Urethral dilatation*.—This mode is often performed by natural means, and even calculi of large size have been expelled through the urethra. Thus, there is a case on record where a young girl, æt. 18, passed spontaneously a very large calculus, 1½ inches long, 1⅓ inch broad, 1⅓ inch thick, and weighing 10 grains. She made a good recovery and escaped any after consequences.*

The dilatation may be artificially performed by the surgeon, either by the slow or rapid process. The slow, gradual, and tedious method is very tedious, and consists in the introduction of hard or flexible bougies, gum-elastic catheters, prepared sponges, or the use of dilators, the speculum or blunt gorget,—increasing the dilatation to the extent that the calculus may be removed with the forceps. The second method is that by rapid dilatation under chloroform by means of Weiss's dilators, so that the stone is removed by one sitting:—this proceeding is the one usually adopted now-a-days, and has been successfully

* *Guy's Hospital Reports*, 1st series, vol. iii. p. 167, with plate.

dilatation.

2nd. Dilatation with partial dilatation has been carried on t stone is too large for extractio necessary. This is seldom neces

3rd. Lithotritry—alluded to operation under chloroform is r ments removed at once: the ar cessful cases of the kind; the l be adapted to the age of the ind

4th. Lithotomy. The patien the same position as for males duced into the urethra. A bist that are required. The follow performing the operation: (1) common and ordinary method, l sible direction, generally implic The incision into the urethra been analogous to the lateral op viz. by introducing the knife in cutting downwards and outwa: degrees. This is called the obli tion; the objection to it is the r nence of urine. Others have r

vided vertically, by some in the direction upward towards
es, but the space was found too small, even with subse-
lilatation; and by others downwards, as suggested by

Attempts have been made to extract the calculus by
; the bladder directly, as suggested by Celsus. This
m consists in making a transverse incision between the
sis and urethra, and thus attacking the anterior parietes
bladder; and M. Lisfranc modified this into a curved
, under the name of the vestibular operation: these
ms have found no recent advocates.


he calculus has been removed by incision through the
a very easy and expeditious operation, but which is
l almost necessarily by vesico-vaginal fistula and incon-
Some make the incision longitudinally, and others in
verse direction. M. Vidal's experience is derived from
operations without any death: he introduces a staff
the urethra into the bladder which is held in the median
l its convexity made prominent in the vagina; a blunt
is next passed into the vagina deeply, so as to depress
rior wall of the vagina (a speculum with one valve will
l with the left index-finger the groove in the staff is felt,
bistoury plunged into it, taking care to avoid wounding
thra; the stone is extracted by the forceps; a female
r is introduced through the urethra into the bladder;
edges of the wound in the vagina brought together by
M. Vidal remarks that there is generally no hæmorrhage
onitis: but that fistula often follows.

nal lithotomy has recently had some warm advocates, and
dered by them as the only justifiable operation for stone
emale bladder. The whole subject has been thoroughly
ated by Dr. Aveling:* he enters into the literature of
ject, and describes the operation minutely. The wound
the operation is to be treated on the same principles
he operation for vesico-vaginal fistula. The mortality
l to be 1 in 35.

t has been recommended to remove the calculus by the
ubic operation, and the same remarks as were made
ng the male will hold good for females.

stics respecting stone in the female, the operation and

* *Obstetric Transactions*, vol. v. pp. 1 et seq. 1864.



there is a report of a case by v
was obstructed by a stone in th
after death, was found to weigh
surement being $3\frac{1}{8}$ inches long
nature of the case was not p
catheter was passed to ascertain
the child's life sacrificed by per

Velpeau (*Art des Accouchements*)
in which a calculus weighing
bladder of a woman four months
who afterwards was delivered w

An obstructing calculus is usually
pain; but it sometimes happens
wedged between the head of the
pubes.

If the calculus, during the delivery,
upper part of the vagina, it will
head descending into the pelvis,
obstruction; but if the calculus
and be placed below the head,
difficult.

Treatment.—If the calculus is large,
it should be removed before labour
be practised with safety during

ed back into the cavity of the bladder above the brim of pelvis. This *reposition* is recommended by Smellie, Den-Baudelocque, Dubois, &c.

When the calculus is fixed and does not admit of reposition, it must be extracted in one of three ways. 1st. *Dilatation of urethra*. 2nd. *Lithotomy*. 3rd. The high operation, or *supra-lithotomy*.

ALFRED POLAND.

FIG. 1. *Uric Acid* external layers are cor consists of the crystal which gives it a reddi Catalogue of Calculi, modified.

FIG. 2. *Urate of A* earthy phosphates dis brittle and homogene boy et. 5, by B. B. C

FIG. 3. *Cystic Oxid* structure, of a radiatin transparent and glist bluish-green colour. cit).

FIG. 4. *Oxalate of* oxalate of lime, arran agate; its colour is of is tuberculated and 1 moved from the bladd ptoms of stone until tr

FIG. 5. *Uric Oxide* Fig. 2, Catalogue of C flesh-colour, and its str


FIG. 6. *Phosphate of* laminated variety, and mortar. It is of a g Museum.

FIG. 7. *Triple Pho* nucleus. It is confus laminated, and is of a

FIG. 8. *Phosho Calc* challs, white, more fri composed of concentric from Prop. 2158, Guy's

FIG. 9. *Alternating* (Calculi, Royal College monia, mixed with oxa secondly, by uric acid; the earthy phosphates.

URINARY CALCULI.



Gosse del. lith. et ad nat.

M. & N. Harcourt, imp.



ON LITHOTRITY.

THE amount of mortality attending the operation of lithotomy in the adult has always made it a question of paramount importance with surgeons to discover some more successful method of removing a calculus from the bladder.

It is not necessary here to allude to the attempts made from time to time to get rid of the calculi by injecting solvents into the bladder.

The removal of calculi from the bladder through the urethra was effected by Sir Astley Cooper and Sir Benjamin Brodie before the introduction of lithotripsy into practice. These surgeons were in the habit of removing small calculi with an instrument called an urethra-forceps. Sir William Blizard is also said to have performed this operation. And in this way patients were successfully relieved of small calculi; in one case, as many as a hundred being thus removed.

To give a detailed account of the early history of lithotripsy would now be of little interest. Whether the credit of originating the operation is due to the Spanish monk or to the Italian officer—who both performed an operation on themselves for the purpose of breaking a stone in the bladder—is now of much importance. It is admitted that to M. Civiale we are indebted for the operation as practised at present. Among those who have written on the subject, and whose works may be consulted for a history of the operation in all earlier stages, are Gruithuisen, Elderton, Civiale, Amussat, Roy d'Étiolles, Heurteloup, King, Costello, Bellinaye, Coulson, and others.

Baron Heurteloup* was among the earliest who performed the operation in this country. At first, several instruments

* See *Principles of Lithotripsy*, by Baron Heurteloup. London, Whittaker and Sons, 1831.

were used by him, such as virgule, évideur or forceps, &c the stone, and break it up by made use of an instrument in sliding forceps, to which he applied broken by percussion. All the operating have been, for many now in general use, viz., crural lithotrite.

Sir Benjamin Brodie, who published on its first introduction, and simplicity of the present method. 'Lectures on Diseases of the Urinary Organs,' 1846. In 1847, Mr. Weiss made an instrument which I have called the sliding forceps. It was intended for the purpose of dividing the stone into fragments; but it was of no use, as it then was, was certainly not successful. 'Baron Heurteloup attempted it, but finding it impracticable, he adopted the principle of the instrument proposed by Mr. Weiss, at the same time, to render it more convenient for use, and for seizing and retaining the stone. This he made another change, by adding a screw, and substituting for the handle a ring, which enabled him to crush the stone. Now the first of these alterations, I believe to have been of essential use, the instrument would have been of no useful purpose. But as to the second, that anything that I have seen of it, or that of others, would lead me to recommend it, I cannot say.

* *Brodie's Works*, edited by Charles Brodie, London, 1865.

† The application of the screw was first made by Mr. Brodie, who was formerly surgeon to the Birmingham General Hospital. He had been a successful operator, many times, with only four unsuccessful cases. Sir Benjamin Brodie in his work on Lithotomy recommended, in lithotomy in the female, to divide the symphysis of the pubes.

Sir Benjamin Brodie in his early operations was accustomed to use an instrument—a scoop lithotrite—so constructed that a portion of the crushed calculus always remained within the urethra; by this means a considerable quantity was removed at each operation by the repeated introduction of the instrument. But this method was attended with great objections; the withdrawing the instrument loaded with fragments stretched the urethra beyond its natural size, giving much pain at the time, as well as afterwards. When the urine was passed, it was occasionally attended with some bleeding, and in some instances the urethra was torn, fragments lodging in the torn portion; infiltration of urine, followed by perineal abscess, were the consequences, and in two such cases death resulted. In referring to these cases, in his *Notes on Lithotomy* in the 38th volume of the *Medico-Chirurgical Transactions*, he says: 'The experience of these cases led me some years since to discontinue the use of the forceps already referred to, or at least to have recourse to it very rarely, and only under some special circumstances, and to substitute for it a forceps made by Mr. Weiss, in which there is a longitudinal opening in the curved part of the fixed blade, with a corresponding projection in the opposite or sliding blade. The effect of this instrument is to crush a calculus very completely, and in such a way that a part of it remains between the blades, the whole being intended to be passed with the urine afterwards. The ultimate recovery of the patient may in some instances be thus a little (but not greatly) protracted; but this inconvenience is more than compensated by the smaller amount of pain which the patient suffers, the smaller liability to rigors, and the complete absence of danger from the infiltration of urine and perineal abscess.'

This instrument, now called the 'screw lithotrite,' is the one in very general use. Alterations have been made in it. The rack and pinion, applied instead of the screw, was originated by Sir William Ferguson, and is used by that surgeon. Dr Henry Thompson uses an instrument that he strongly recommends. These are delineated in the article on SURGICAL INSTRUMENTS. Mr. Coxeter, and more recently Mr. Weiss, have also made alterations, by which it is intended to allow of more facility in operating, and to gain lightness in the make of the instrument; but I must confess that, notwithstanding the high authorities in favour of these different instruments, the

instrument just described has, I think, still advantages over any other; it is simple in construction, quite as light as it is safe to have it made, and it is not so liable to get out of order as those of a more complicated construction.

It may be as well first to describe the methods of performing the operation, and then to consider the cases to which it may be applied.

When the symptoms are such as to lead one to suspect that a patient has a calculus in the bladder, should there be any considerable amount of irritation of that viscus, and instruments have not previously been used, it is well to keep the patient confined to the sofa for a day or two before any examination is made; and should the general health need it, to give such medicines as may be necessary.

In examining the bladder, instead of using a sound, the better mode is, having placed the patient on a sofa, with the pelvis raised by means of a pillow, to inject into the bladder through a silver catheter about four or five ounces of water; to the catheter should be attached a stop-cock, so that the water may be retained in the bladder during the examination. If the stone be of any size, it is usually detected by means of the catheter; but should this not be the case, the lithotrite may be introduced, and an examination made in the same manner as will be described when it is used for crushing the calculus.

Having ascertained the presence of a stone, and decided that lithotritry is the operation to be performed, it is necessary to place the patient in as favourable a state as possible for what he has to undergo. One of the first requisites is, that the bladder should be able to contain a sufficient quantity of water to render the necessary use of instruments safe; and it is not unfrequently required, on account of the intolerance of the bladder to a sufficient quantity, to inject the bladder, and keep the water in for a short time on one or two occasions previous to the introduction of the lithotrite. After such treatment the bladder usually retains the necessary quantity. If an extreme irritability should continue, so that at least four ounces of water is not borne, the administration of an injection, containing twenty or thirty drops of laudanum, per rectum most frequently accomplishes the desired object. It is also necessary, in some cases where instruments have not been previously used, or where there is any difficulty in introducing them, to pass an

trument a few times on different occasions before proceeding to crush the stone.

The bladder and urethra being thus prepared, the following is the mode of proceeding to be adopted. The patient should take a dose of aperient medicine the day before, and be confined to the house for a couple of days, so that the urinary organs should be in as quiet a state as possible. The patient being placed on a sofa, with the pelvis raised as before described, from four to six ounces of warm water should be injected into the bladder; it is as well to have the piston of the syringe graduated, so as to know exactly the quantity of water injected. Should there be much spasm, so that the bladder resists the introduction of the water, this part of the operation must be conducted slowly, only a small quantity being introduced at a time; indeed, if it is found that the bladder will not retain the requisite quantity, the further performance of the operation must be postponed to a future occasion. In some cases, where the intolerance continues, the following plan may be adopted: directing the patient to retain the urine for as long a time as possible previous to the operation being attempted, and the surgeon being satisfied that a certain quantity of urine is in the bladder, he may introduce the lithotrite, and crush the stone *once*; when a stone has been once crushed, this irritation of the bladder not unfrequently ceases. It is the practice of some surgeons not to inject any water into the bladder before the introduction of the lithotrite. I have found that the bladder more readily retains the water injected than the quantity that may be in the bladder when the lithotrite is introduced.

The principal requirement in the operation of lithotrity is, that the calculus should be broken into pieces of such size as may be readily passed; and it is to be considered what is the best method of accomplishing this with as little disturbance to the bladder as possible. The size of the instrument must depend in some measure on the supposed size of the stone, which there is no mode of ascertaining with any very great accuracy; if the stone is not large, a medium size is all that will be required; but should there be any doubt as to the size of the stone, a large instrument should always be used on the first occasion of operating.

The introduction of a lithotrite, from its shape, is more difficult than a catheter; and the passage of a lithotrite through

the neck of the bladder requires caution and dexterity, and some amount of pressure is occasionally necessary. Where there exists much enlargement of the prostate gland, considerable difficulty is occasionally met with; but with proper manipulation this is overcome. It is of the greatest moment, in this step of the operation, that no attempt should be made to open the blades of the instrument until it is completely in the bladder; much mischief is sure to ensue if such an attempt is made. The next step is the manipulation necessary for the seizing of the stone; on this great authorities differ. Sir Benjamin Brodie, in his *Notes on Lithotrity*, says: 'The rule should be to *move the forceps in the bladder as little as possible*, never using it as a sound for the purpose of exploring the bladder, or ascertaining the position of the calculus. Such an examination does not assist the surgeon in seizing the calculus afterwards; it gives pain to the patient, excites the bladder to contract and expel the water which had been previously injected; and I know that instances have occurred, though not in my own practice, in which a rough handling of the forceps has caused great injury to the bladder, ending in the death of the patient. The rule for seizing the calculus (which I must acknowledge to have first learned from witnessing the very dexterous operations of M. Heurteloup) is as simple as possible. The patient lying on his back, the handle of the forceps is elevated, which of course brings the convex part of the curved extremity of it in contact with the posterior surface of the bladder, where it is contiguous to the rectum. The forceps is then to be opened by withdrawing the sliding blade to a greater or less extent, according to the probable size of the calculus, the fixed blade being at the same time pressed gently downwards in the direction of the rectum. The object of this manipulation is, that the forceps, being below the level of the other parts of the bladder, the calculus may fall into it by its own weight; and it is generally successful. If it should not do so, the forceps without being moved from its situation, may be gently struck with the hand on one side, or on its anterior part, and the slight concussion thus communicated to the bladder will probably be sufficient to dislodge the calculus, and bring it within the grasp of the instrument. If it should be otherwise, the forceps, being closed, may be very gently and cautiously turned to one side or the other, so that the curved extremity of it may make an angle of 25° or even 30° with the vertical line of the

ly, then opened, and pressed in the direction of the rectum the manner already described.

'When the prostate gland is much enlarged, there is sometimes a difficulty in seizing the calculus, arising either from it lying under that part of the gland which projects into the bladder, from the impediment which it offers to the elevation of the handle of the instrument. For such cases the operating-table invented by M. Heurteloup, which enables the patient's shoulders to be suddenly lowered, is very convenient; or the same purpose will be answered sufficiently well if the patient is placed on a light sofa, the end of which may be raised by an assistant. The calculus is then seized, not in that part of the bladder which adjoins the rectum, but in the fundus, it being rendered the lowest point by the elevation of the vis.'

FIG. 307.




Diagram of the ordinary method (Brodie's) of seizing the stone.

M. Civiale recommends another method, thus described by Sir Henry Thompson in his work *On Lithotomy and Lithotritry* :* 'The other mode is that of Civiale. Its principle is the reverse of the preceding. By position of the patient, the centre of the bladder and space beneath it are selected as the area of operation ; no depression is made ; contact between the walls of the bladder and the instrument is, as much as possible, avoided. The instrument is applied to the stone in the situation which this naturally takes, and the operator carefully avoids moving it, or any movements of concussion whatever, however slight. It is only due to the distinguished operators first named to say that this, the modern, and it is believed the improved, method, is in part due to the mechanical improvements which have been made in the lithotrite of late years. The method was scarcely possible until the present instruments existed.

'We shall now consider it in detail. The blades having entered the cavity of the bladder, the instrument slides easily and smoothly down the trigone, which in the living and healthy organ is an inclined plane, although quite otherwise in the atonied and in the dead bladder.

'In many cases the stone is grazed by the instrument as it passes, and the slightest lateral movement of the blades right or left will determine on which side it lies. If so, the operator is careful not to disturb it, but he inclines the blades *slightly away from the side on which it lies*, carrying the instrument gently in towards the posterior wall of the bladder, while the male blade is slowly withdrawn. It is important always to bear in mind, that as long as the blades are near the neck of the bladder, the male blade cannot be withdrawn, since it would impinge on that sensitive part, and cause pain or injury. Having done so, he now inclines the well-opened lithotrite towards the stone, slowly closes, and almost certainly seizes it.

'But suppose no stone was felt on entering, he is then directed simply to withdraw the male blade an inch or more in the middle line, to incline the blades to the right side about 45° , and then to close them, without altering the axis of the shaft, or otherwise disturbing the central position of the instrument. Thus in almost all positions the stone is seized sideways

* London, Churchill, 1863.

the blades of the lithotrite, and very rarely by their extremities. If no stone is felt, he turns them, opened to the left in a similar manner, and then closes them. Observe, that the blades are always to be opened before they are turned, for this reason: if the turn is first made and the blades are subsequently opened, the chance is that the male blade as it is withdrawn will move the stone away; whereas if the blades are inclined while open, the stone, if there, is almost certainly seized. This is one of the many apparently minute but extremely important points of which systematic lithotripsy is made up. To return: it is very rare that the stone will elude the search thus far; but if it does, depress the handle of the lithotrite half an inch or so, which raises the blades very slightly from the floor of the bladder, and turn them another 45° to the left, bringing, in fact, the blades horizontal to the left; close: if unsuccessful, turn them entirely to horizontal on the right, and close. These five positions (vertical, right and left incline, right and left horizontal) explore the bladder fully, middle, right, and left, and will almost certainly find any stone of a moderate size in a healthy bladder. The object is at the same time strictly to avoid communicating any jerk to the instrument or to the bladder.

In all these movements, if properly executed, there has been merely contact of the lithotrite with the vesical walls; at all events, no pressure, nothing to provoke undue pain, or cause contractions of the bladder. If, however, there is an enlarged prostate, causing an eminence at the neck of the bladder, a compression behind it, or the stone is very small, or we are exploring for some fragment, at the close of the case, which is suspected to have eluded previous search, the blades may be reversed so as to point downwards to the floor, and the object sought may then often be secured with ease. If seeking for a small stone or for fragments, we shall employ a lithotrite with short blades, which can therefore be reversed with much greater ease than one with long blades.

‘In order to do this properly in the normal bladder, the handle of the lithotrite is depressed another inch or so, between the patient’s thighs, so that the line of the instrument, instead of being directed obliquely a little upwards, is level with, or even points a little below, the horizon; the blades, supposed to have been already brought to the horizontal as before described, are cautiously turned, about 45° say, to the right (right reversed incline), so as to point obliquely to the floor, which should be

barely felt, or very lightly touched by them. No pressure should be made on this part of the bladder by any part of the instrument, and it is easily avoided by depressing sufficiently the handle of the lithotrite. Then close the blades; next turn them back, that is upwards, over to the left (left reversed incline), and close. Lastly, they may be brought round, to the reversed vertical position, and the floor lightly swept; this requires the maximum depression of the handle, and is only necessary to pick up small fragments with a short-bladed instrument. But when the prostate is considerably enlarged, and a stone or fragments have to be sought behind it, the lithotrite is reversed without depressing the handle.

‘All these movements are to be executed at or beyond the centre of the vesical cavity, the proper area for operating, without hurry, rapid movement, or any other which partakes of the nature of a jerk or concussion, and, if in a fairly healthy bladder, without causing more than a very slight degree of pain to the patient. The operator’s eye is also to be so familiar with the scale marked on the sliding-rod, that he knows at a glance the exact interval which it indicates as existing between the blades in the bladder.

‘It is essential to good practice, while manipulating the lithotrite, to maintain the axis of the instrument, as far as possible, always in the same direction. The blades only are to be moved: the shaft should occupy the same inclination, unless when this is intentionally altered. In screwing home the male blade, the operator is very apt to move the lithotrite also, at each turn of the screw, unless he is conscious of the care necessary to avoid this evil. All lateral movements, all vibration and concussion, necessarily tell on the neck of the bladder and prostatic urethra, where the instrument is most closely embraced, and its mobility is most limited. To that part of the lithotrite which occupies the anterior portion of the urethra much freedom of lateral movement is permitted, and in the bladder the instrument is free, although in a less degree; but the axis, or fixed point, as regards lateral movement, is at the part indicated, which is also the most sensitive spot of the entire passage. Hence the aim of the operator should be to produce in this situation no motion of the lithotrite, except that on its own axis. Few of the details of the operation require more practice to master than this.

‘There is one important rule with reference to the situation

the calculus in the bladder. The larger it is, the more certain it is to be found lying near to the neck of the bladder in the ordinary recumbent position, while a small one is usually ejected at the back of the trigone. This position of the large one requires a different method, and it will be found almost invariably successful. The moment the lithotrite enters the bladder, it is not to be pushed onwards to the bottom of the cavity: first, let the blades be inclined away from the side on which the stone is felt, then push on the female portion of the instrument only, by itself as far as it will go, maintaining the

FIG. 308.




Diagram of Civiale's method of seizing the stone. From Sir H. Thompson *On Lithotomy and Lithotripsy*, p. 177. 1863.

the blade at the neck of the bladder; it is now only necessary to incline towards the stone, and it will be seized almost certainly at once. But if the operator commences by pushing the whole instrument, and then withdraws the male blade according to the ordinary custom, this blade is infallibly drawn against the large stone, which it therefore fails to catch, and presses it back against the neck of the bladder, producing

pain, irritation, and perhaps bleeding: this is a practical rule of importance. As already intimated, for a small stone, the instrument glides down to the posterior wall of the bladder, the male blade is withdrawn, and the stone caught in the usual manner.

‘Such then, in the main, is the method of Civiale for finding the stone; the other, or English method, is without doubt an efficient one, but I believe it to be more irritating to the bladder, and less certain for removing every minute fragment towards the end of the operation, while it fails to deal efficiently with a stone lying behind an enlarged prostate. Hence the crushing operation has been often said to be inapplicable when such disease exists, a conclusion wholly unwarranted by the practice and results of modern lithotrity. Having tried both methods myself, I have no hesitation in preferring the former.’

Notwithstanding the opinion thus strongly expressed by one so competent to come to a right conclusion on this matter as Sir Henry Thompson, I am still inclined to prefer and to recommend the method practised with so much success by Sir Benjamin Brodie, as the one less likely to cause irritation of the bladder; and I have not found it attended with the disadvantages described by Sir Henry Thompson.

The calculus being seized by either method, the question then arises—how often should the process of crushing be repeated? In the first operation, the surgeon should often be satisfied with crushing the stone but once, especially if the bladder is irritable. One crushing frequently relieves the patient of much pain and irritation of the bladder, particularly in the case of large stones, and it is well to give the patient as little inconvenience as possible at the first operation, so that he may look to a repetition without dread. On the occasion of first seizing the stone, I usually move the instrument gently to either side, to ascertain if there be more than one stone. The lithotrite being carefully withdrawn—care being taken that the blades are quite closed—the patient should at once go to bed, taking a glass of warm wine-and-water or brandy-and-water, and be kept warm, in order to prevent his having a rigor; he should for the next twelve hours make water in the recumbent posture, and be confined to the house until the next operation.

The period that should elapse between the operations must vary; if at the first operation the calculus has been crushed once only, unless it be a small one, in all probability no frag-

its will be passed; in which case the operation may be repeated usually in three or four days. During the intervals between the succeeding operations, the patient may be allowed to take a certain amount of exercise.

It is of considerable importance to determine how much should be done at each operation; on this question surgeons differ. Some do a great deal, whilst others are satisfied with seizing the stone only once or twice, and repeating the operation at short intervals. Some place the patient under the influence of chloroform, and continue seizing and crushing the stone until the whole is broken up, at one operation. This latter plan is accompanied with this disadvantage: where a large stone is thus crushed into pieces capable of being passed, the passage is liable to be blocked up with the fragments, which leads to a difficulty in getting rid of them, and much consequent irritation of the parts. The wiser plan is to adopt an immediate method, and to do at each operation so much only as the patient can bear without great inconvenience, so as to require as few operations as possible necessary; whilst on the other hand, the patient is not distressed by the manipulations being continued too long at one operation. From five to ten minutes is the average time the lithotrite should be employed in the bladder.

The interval there should be between the performance of the operations, is the next question to be considered. As a rule, I never repeat the operation whilst any fragments are being passed, and I usually allow one or two days to elapse: the fragments have ceased to come away; this usually requires about a week between each operation.

With regard to the removal of the fragments by means of an instrument, I have found no reason to alter an opinion I formerly expressed, and which was published in the *Transactions of the Pathological Society* for the year 1850:—

In relation to the occurrence of infiltration of urine, and especially after lithotrity, Mr. Charles Hawkins thus expresses himself: He believed this accident might be entirely avoided, if the operation of crushing be performed with the lithotrite in such a manner as to satisfy the operator with crushing the stone, without attempting to withdraw the fragments between the jaws of the instrument. He believed this accident never occurred except where the urethra had been lacerated with a fragment of stone during the withdrawal of the instrument; then

impaction of a fragment led to the results which the preparations exhibit; but where the urethra has not been lacerated, impaction is of little importance, and no dangerous symptoms are likely to result from such an occurrence. . . . Since no attempts have been made to remove fragments from the bladder in the instrument, he had met with no such accident. . . . As far as his experience went, it was not necessary to attempt the removal of stone in the scoop-lithotrite. If the stone is *well crushed*, it may be left to nature for the bladder to be evacuated of the fragments: or where there was a difficulty in passing them, he had removed much by means of washing out the bladder.*

It is not necessary to introduce the lithotrite more than once at each operation, if the plan described be adopted; nor is it requisite to draw off the water when the lithotrite is withdrawn. It is desirable that some water should be left in the bladder; it renders it better able to bear the presence of the broken stone. In performing lithotritry, this rule should be invariably followed: *to introduce an instrument into the bladder as few times as possible.*

When there is much enlargement of the prostate gland, and the patient on this account is unable to pass very small fragments, and their removal has not been effected by means of the catheter, the best mode of proceeding is to make use of the scoop-lithotrite; and one end of the sofa on which the patient lies being raised by an assistant, the fragments fall into the instrument at the fundus of the bladder, that part by this plan being now the lowest portion. Mr. Clover has invented a very good apparatus for removing the fragments in such cases.

Among the difficulties that may arise during the treatment of a case are the following: Sometimes after the first operation retention of urine occurs, but not very frequently when the bladder is in a healthy state, and where too much has not been attempted: this must be met in the usual way, by the use of the catheter. A fragment of stone may be retained in the passage, after it has left the bladder, in cases where there has been no laceration of the parts: as before stated, this state of things, although troublesome, is not dangerous; and when the fragment lodges in the membranous or prostatic portion of the

* *Path. Trans.* part i. of vol. iii. 1850, pp. 123, 124.

thra, a full-sized gum catheter should be used, and the fragment gently dislodged and pushed back into the bladder. Many ingenious instruments have been devised for the purpose of seizing and removing fragments thus arrested; but of all, the gum catheter will be found the most simple and most useful. Should the fragment be retained in the anterior portion of the urethra, it usually makes its way out in course of time; to facilitate this, the patient should retain the urine for as long a time as possible, and pressing the penis at the other side of the urethra for a short time when the bladder is about to act, the fragment is washed out. Fragments sometimes lodge just behind the urethral meatus; with a common dressing-forceps they may be removed from this position; in a few cases it may become necessary to make a slight incision in the lining membrane of the urethra. The passing of fragments sometimes gives rise to inflammation of one or both testicles; this may delay the operation. The chief cause of mischief to be apprehended is when there is considerable irritation, followed by inflammation of the bladder, the urine depositing a quantity of ropy mucus, which may give rise to great pain in the passage of water, and a constant desire to pass it. These cases require very careful treatment; depletion is rarely necessary: with the use of the warm bath and the administration of opium and hyoscyamus, the patient being kept to his bed or the sofa, the attack usually subsides off. But these are signs that the bladder will not bear lengthened operations. This state of bladder is sometimes caused by a fragment of stone, which is both too large and too small—small enough to enter the neck of the bladder, but too large to pass further; in this case, notwithstanding the irritation that may be present, the lithotrite should be introduced to crush the fragment, when generally the symptoms subside. The bladder will in many cases continue to secrete a quantity of ropy mucus as long as any stone remains, so that there is no reason for not proceeding with the operation; the patient may take daily half a pint of the decoction of the pira brava; but the treatment really requisite is the removal of the stone. Nevertheless these cases are occasionally not so easily managed, and sometimes it may be necessary to cease all operative proceedings for a time, until the bladder becomes in a more healthy state; it may, indeed, happen that the attempt to remove the stone by lithotritry must be abandoned: this, however, occurs but very rarely; usually with proper treatment,

these unfavourable symptoms are overcome, but not always. Sir Benjamin Brodie, in his *Notes on Lithotrity*, relates the following case :—

In a fourth case, a very small calculus was crushed with great ease by a single operation. The operation was succeeded by a rigor, which terminated, as is usual, in a perspiration. A disturbed state of the system followed, marked by a frequent pulse, a furred tongue, and much prostration of strength; and attended with a deposit of adhesive mucus, but not in large quantity, from the urine. These symptoms continued, and after some time an abscess presented itself in one groin. The abscess having been opened, a considerable discharge of matter took place, and was followed by great relief as to the general symptoms. The patient seemed to be in an improving state, when between four and five weeks after the operation, he suddenly expired.

On examining the body, no remains of the calculus were discovered in the bladder. The mucous membrane of the bladder bore marks of inflammation. There was an abscess of the pelvis, occupying the space between the bladder and rectum, and extending in the direction of the abdomen as high as the groin, in which the puncture had been made. The parts were carefully dissected and examined by the late Mr. Vincent (who had attended the patient with me in consultation), Mr. Charles Hawkins, the Curator of the Museum of St. George's Hospital, and myself; but no lesion could be detected of the bladder, nor any kind of communication between the bladder or urethra and the abscess. Still I cannot doubt that the abscess was somehow the result of the operation. Probably a very small splinter of the calculus might have penetrated the coats of the bladder, allowing the escape of a very minute quantity of urine into the cellular membrane. This would be quite sufficient to account for an extensive suppuration, at the same time that it is easy to suppose that so small a puncture might at once have closed, so as to be invisible afterwards.

In this case the cause of death I believe to have been fatty degeneration of the heart; and had this disease not existed, I have little doubt but that the patient would have got well of the operation.

In those cases where a fragment becomes impacted in the urethra, usually in the membranous portion, and where the parts have been lacerated, in all probability such an accident will be followed by infiltration of urine and perineal abscess; the surgeon must lay open the abscess freely. In two cases where this treatment was followed, the patient ultimately recovered.

The number of times it may be requisite to repeat the operation must necessarily depend upon the size of the stone—the amount of crushing that is performed at each operation—and the facility with which the fragments are passed. Some patients pass very considerable-sized fragments; others require the pieces to be very small, necessitating more frequent

rations. In the cases operated on by Sir Benjamin Brodie, and those that have occurred in my own practice, the number of operations averaged between six and seven; one or two operations sufficing in many cases of small stones. Mr. Prescott Hewett has operated in one case as many as nineteen times with ultimate success.

It is occasionally a wise plan, in cases in which there have been many operations, and the patient begins to feel the constraint and the recurrence of the operations, to cease altogether for a time, so as to allow him to recover his usual health; and, of course before dismissing the patient as cured, the surgeon must be satisfied, by examining the bladder, that no stone remains. In some cases where the fragments are too small to be crushed by the lithotrite, or rather where, from their size, they cannot pass through the opening in the blades of the instrument, and do not pass, the introduction of a full-sized catheter (which could be retained in the bladder for a short time) will enable the patient to void them.

The placing the patient under the influence of chloroform is by some surgeons much recommended. As a rule, I do not think it advisable to make use of it, but only to have recourse to it in those cases where circumstances may render it necessary. In ordinary cases there is not only no occasion for it (lithotomy, if properly performed, is not a painful operation), but, on the contrary, it is as well to do without it on account of the operator being thus better able to ascertain to what extent the patient can bear the operation at each time, so that he should not proceed so far as to cause the bladder to resent the operative proceedings. The bladder does not retain the water well when chloroform is employed. When patients are peculiarly sensitive with regard to the operation, chloroform must be administered; for without it such patients could not be induced to submit to the operation, however favourable their cases might be in other respects.

I operated on a patient twenty-four years of age, who had symptoms of calculus when a child. They had for many years remained dormant, but had returned about four years before he had consulted me, at which time he was passing water incessantly and with intolerable pain, the urine depositing a large quantity of ropy mucus mixed with blood; in fact, the bladder being in as unamiable a state as possible for any operation. Having seized a very large stone with the lithotrite, I was unable to proceed with the operation on account of the patient being incapable of exercising the necessary control over himself; but, after chloroform, I was enabled to effect a complete cure in five operations. It

would have been impossible to have done this without the aid of chloroform. In the 41st volume of the *Medico-Chirurgical Transactions*, I have related a case in which I operated successfully on a patient in whom there was a communication between the bladder and intestine. The calculus, the nucleus of which was some vegetable matter, was of such a size that I could scarcely seize it with a very large lithotrite. The patient was enduring such dreadful sufferings that he was anxious to be placed under the influence of chloroform; but on the third occasion of operating, it was done, at his request, without chloroform: the operation was borne as well as when chloroform was used, and the bladder held more water.

In such cases chloroform is very useful; but in the majority of cases, it is not requisite, and not being requisite, it is as well not to have recourse to it.

In considering in what cases lithotritry is to be preferred to lithotomy, it may be observed that the early writers on the operation were of opinion that only where the calculus was very small, requiring but one or two operations—the bladder and kidneys perfectly healthy—was lithotritry available. Further experience, together with the great improvement in the instruments used and in the mode of operating, have tended in a great measure to alter this opinion. No doubt the frequent introduction of instruments when the fragments were removed within the blades, causing unnecessary disturbance to the bladder, and the stretching and not unfrequently the laceration of the urethra, rendering it in a very unfit state for the passage of fragments, led to most dangerous accidents; but since the practice of removing the fragments within the blades of the instrument has been laid aside, many cases which formerly would have been unfavourable for lithotritry are now successfully treated. The inability of the bladder to expel all the fragments was also considered a state in which lithotritry was not applicable. Yet now, after the stone has been well crushed, these cases can be managed by washing out the débris through a silver catheter. It may be well to mention that the eye of the catheter used for this purpose should always be on the concave side of the curve—for the reason, that if a fragment should chance to lodge in the aperture and not be removable, although on the withdrawal of the instrument the fragment might lacerate the urethra, the consequences that follow such an accident are of much less importance than if the lower part of the urethra were injured. A catheter has lately been invented with the opening in the convex part—for the injecting the bladder—under the idea that the stream of water coming

are immediately in contact with the fragments, their removal is more readily effected; which is rather a theoretical than a practical improvement, and the catheter has the disadvantage of having the aperture in the lower curve. There is no doubt that a healthy state of the bladder is very desirable when lithotrity is to be employed; but I have performed it with success when the bladder has been in a very considerable state of irritation and secreting muchropy mucus. This irritation and secretion of mucus diminish as the operations are performed, and nearly cease before the whole of the calculus has been removed. These cases certainly require much caution in their management, still they are by no means cases in which lithotrity is to be rejected. The inability of the bladder to hold the requisite quantity of water might at first deter a surgeon from selecting such cases for the operation; but, on the other hand, a little management, by injecting the bladder for some time previous to operating, generally overcomes this obstacle, and not unfrequently, where a bladder has been very intolerant of water, this inconvenience has subsided after the stone has been once crushed. Where there is serious disease of the kidneys, lithotrity is no doubt as likely to be attended with ill consequences as lithotomy. But cases do occur where the presence of kidney disease would inevitably render lithotomy an unsuccessful operation, in which lithotrity may be employed, if not with entire success, with at least the alleviation of much suffering, and prolongation of life. The worst form of kidney disease, albuminuria, is fortunately not frequently found in combination with calculus in the bladder at the period of life at which it is desirable to perform lithotrity. In cases of very bad stricture of the urethra, rendering the introduction of the lithotrite impossible, lithotomy must be had recourse to; but in cases in which a considerable amount of stricture existed, I have been enabled, by using a small lithotrite, and crushing the stone very fine, to relieve the bladder of considerable-sized stone. But now that stricture is so readily removed by Mr. Barnard Holt's 'immediate treatment,' it ceases to be so serious an objection. Malignant disease of the bladder, in combination with calculus, of course precludes lithotrity as well as lithotomy.

The removal of foreign bodies that have been introduced into the bladder, such as bougies, may be effected with a scoop-lithotrite.

cases of lithotritry and lithotomy. *Notes on Lithotritry*, makes

expect that any method of lithotomy is so certainly fatal if left to its own misery, as calculus of the bladder is, and uniformly successful. If lithotritry has its dangers also; and the practical surgeon to consider is, which of the two. Of the nine cases of lithotritry it may well be doubted as to one of which was the immediate cause of the death connected with the operation; while the result was to be attributed to a mode of operation which my later experience has proved fatal. But, even if we admit the whole nine cases to be the average failure of the operation, the ratio of deaths to recoveries is somewhat less than 1

I might compare this with the proportion of deaths after lithotomy, I have referred to a paper by the late Mr. Smith of Bristol, published in the eleventh volume of the *Transactions* of this Society, and entitled *An Inquiry into the Causes of Stone in the Bladder*; and I there find it stated, that at Bristol Hospital, during a series of many years, the proportion of deaths after lithotomy was 1 in $4\frac{1}{2}$; in the hospitals, 1 in 5; and in the hospital at Norwich 1 in $7\frac{1}{4}$. Twenty-five years have elapsed since the publication of these statistics, but there is no reason to believe that the success of lithotomy is greater now than it was when Mr. Smith collected his observations. The editor of a weekly journal (the *Medical Times and Gazette*) has for some time past published the statistics of the various operations performed in the London hospitals, including lithotomy. The facts seem to have been collected with some care, and are probably a near approximation to the truth. Mr. Charles Hawkins has been at the pains to collect from the various numbers of that journal, published during the year 1854, the facts relating to lithotomy; and it appears that of 59 patients who underwent that operation, as many as 10 died, being in the proportion of rather more than 1 in 6.

‘But here two other facts must not be overlooked, without

which no just comparison can be made of the results of operations. First, while cases of vesical calculus in and under the age of puberty, in private practice, and among more affluent classes of society, are of rare occurrence, they form the very great majority of those which are admitted to hospitals; and, secondly, the proportion of deaths after lithotomy among children is very much less than it is among adults. Both these facts are sufficiently obvious to those who have had the opportunity of witnessing the practice of lithotomy in hospitals. From the data furnished by the medical journals, which I have already referred to, it appears that in the hospitals during the last year children formed a small minority, more than three-fourths of the whole number of those who underwent lithotomy; that among them the deaths were in the proportion of one to fourteen recoveries; while among adults the deaths and recoveries were equal. That in this respect the large proportion of deaths among adults was beyond the average, and depending on accidental circumstances, will well be a matter of doubt, and indeed it is plain that no general rule can be drawn from the limited number of cases which have occurred in the space of a single year. Still, as even in the best hospital, where there is reason to believe that lithotomy has been on the whole more successful than in any other institution, the proportion of deaths among adults is so much greater than to have been four times as large as that among children, it is evident that the difference in the degree of danger at different periods of life is very great; and it must always be borne in mind that, in estimating the comparative value of the two operations, it is only the results met with in adults that should be introduced into our calculation.*

There have not as yet been any very extensive statistics of lithotritry published. Among the latest are those of M. Civiale, of which Sir Henry Thompson, in his work before referred to, says:—

‘Take, for example, the practice of my esteemed friend, M. Civiale, during the last year. It is a fair sample of his usual experience, which he is good enough to publish annually, and has of late presented to the Académie des Sciences. He treated, during the year 1862, 69 cases of patients—66 men, 2 women, and a child; 45 in private practice

* *Medico-Chirurgical Transactions*, vol. xxi. p. 32.

at Hôpital Neckar. Fifty-eight of these were operated on : were submitted to lithotrity ; of these 8 were partially cured, it was successful in all the remainder but 1.

Ten were treated by lithotomy ; 3 were cured, 2 relieved, 5 died. Three were treated by a combination of lithotomy & lithotrity ; 2 were cured, the other had incontinence of urine.

In *eleven cases* operative means have been adjourned or considered impossible.

Now, although the lithotrity here recorded is extremely successful, every English surgeon will feel surprised to find one every six adult cases placed in the category last named, and half of the lithotomy cases fatal. It must be obvious to all who are familiar with the practice of this country, that nothing like this proportion of cases is adjudged unfit for operation.

It will be seen by the tables so carefully prepared by Sir Henry Thompson, that in 1,827 cases of lithotomy, 1,116 occurred in patients between the ages of one year and twenty years, leaving but 717 from that period up to the age of thirty-one, and of these 462 cases occurred between fifty and thirty-one years, the period of life (at least in private practice) are most called upon to operate for stone ; and in these cases there were 115 deaths—about one in 4 cases. Now if we compare these statistics of lithotomy with those as yet given of lithotrity, it will, I think, leave no doubt in most minds which operation the surgeon should prefer, at least in the large majority of adult patients.

One of the great advantages of lithotrity is, that patients are more ready to submit to the operation when the stone is small, and when its presence has not given rise to any serious mischief in the bladder or kidneys ; when, on the other hand, the irritation so serious and so dreaded as lithotomy is driven off, as a last resource.

In reviewing all the bearings of the two operations, I think there is every reason to concur in the opinion expressed by Sir Benjamin Brodie : ' My own experience has certainly led me to the conclusion that lithotrity, *if prudently and carefully performed, with due attention to minute circumstances*, is liable to all objection than almost any other of the capital operations of surgery. The cases, indeed, to which it is not applicable are very few indeed, and they are chiefly those in which, from the calculus having attained an unusual size, the danger and

great that no surgeon would as a matter of duty, undertake it.' I will to consider how far it may be possible to draw a distinction as to which of the operations may be selected. There is some distinction between private and hospital practice. The greatest success attending lithotomy is met with in the latter class. These patients are not so likely to submit to an operation at an earlier period than those in private practice, are consequently in a more advanced stage of disease for lithotomy; and not being so amenable to treatment between the operations of lithotritry and lithotomy, the latter classes (for lithotritry, although a simple operation, requires, for a successful issue, great care and attention to the most minute points during the whole time of treatment), when the treatment becomes more advanced, are liable to become unfavourably influenced by the disease. To hospital patients *time* is of much more value than it is to the more affluent; so it may be more advisable to have recourse to lithotomy, on admission into hospitals, when in private practice lithotritry would doubt be the operation selected. Such are the reasons why in hospitals lithotomy is as frequently resorted to in adult patients than in private practice.

In order to have accurate information as to the progress of lithotritry in making in hospital practice in London, while this was originally going through the press, I applied to the authorities on this point to all the London hospitals, and was generally and readily supplied with what I required from all, with the exception of the Hospital for Stone; the authorities of that hospital made no reply to my request. I must also state that the return from University College Hospital is not quite complete, as I failed to obtain any information from one of the wards of that hospital. If the numbers reported from University College Hospital be compared with other hospitals of the same size, it will be seen that the number of cases unrequiring operation cannot be large. The absence of information from one hospital cannot in any great degree influence the result obtained from the foregoing table: from which it will be seen that out of 1000 patients admitted, in two years, into the London hospitals with stone in the bladder, only 32 were treated by lithotritry, and 968 underwent no operation.

IENTS WITH STONE IN THE BLADDER ADMITTED INTO THE LONDON
HOSPITALS IN THE YEARS 1862-63.

Hospitals.	Total.	Children.	Adults.	Lithotrity.	Lithotomy.
y's	31	15	16	7	24
Bartholomew's	25	16	9	3	22
St. George's College	22	7	15	9	13
St. Andrew's	17	5	12	2	15
University College	16	6	10	5	11
St. George's	13	2	11	3	8
Mary's	12	9	3	1	11
St. Thomas's	11	7	4	1	8
St. Mary's Free	7	7	—	—	7
St. Andrew's	6	2	4	1	3
St. Martin's	5	3	2	—	5
St. Martin's Free	4	1	3	—	4
St. Andrew's Cross	3	2	1	—	3
St. Andrew's Children	3	3	—	—	3
St. Andrew's Northern	2	1	1	—	2
	177	86	91	32	139

since the above was originally printed I received a communication from Mr. Armstrong Todd, stating that my application to the Hospital for Stone had inadvertently remained unanswered. He was good enough to forward me the following return of patients admitted into the hospital during two years, with stone in the bladder:—The total number of patients admitted were 17, viz. children, 5; adults, 12. Lithotrity performed on 2; lithotomy on 7; not operated on 8. This increases the number of adult patients with stone to 103, of which only 34 were lithotritised. From all I can learn it does appear that lithotrity has since made much advance as an operation in London hospitals.

Sir William Fergusson in his lectures delivered at the Royal College of Surgeons in 1864 and 1865 (since published), has given his enlarged experience in the practice of lithotrity. To that work I must refer those who wish to be well informed on that refers to this operation; but what he wishes greatly to press on operators is the advantage of the removal of fragments by the aid of instruments. He says: 'No doubt in most cases fragments come away spontaneously in a satisfactory manner, but even in many cases these would be rapidly expelled were such means taken.' But he says the practice cannot be followed without instruments properly adapted for the purpose.

An important communication has been made relative to the statistics of lithotritry since this essay first appeared, in a paper read before the Royal Medical and Chirurgical Society, by Sir Henry Thompson, on May 10, 1870, of which the following is an abstract.*

The author presented a series of 184 consecutive cases of lithotritry in the adult, operated upon within a recent period; all treated by the same method and with the same instruments. He furnished all the most important details relating to each case, and presented the stone itself in almost every instance, preserved for inspection. His object was to make an impartial estimate of the crushing operation, to ascertain its real value, and its place amongst surgical operations. Although this had never yet been fully done, he regarded Sir B. Brodie's last communication to the Royal Medical and Chirurgical Society as perhaps the most trustworthy and valuable record, so far as it goes, which exists on the subject. In order to accomplish this object, he had made carefully written records of every case; and he cited the following circumstances as necessary to be taken into consideration: that the 184 cases had been treated by a uniform method, within a comparatively brief period of time; that all were adults, and embraced much variety of constitution; that all the important facts relative to each were noted in a history of each one, which was attached to the paper as an appendix; and that a large proportion of the calculi were of considerable size. And the author believed he was correct in saying that so complete an opportunity for studying the results of lithotritry had not been offered hitherto, since, as far as he was aware, the data necessary for the formation of a judgment had not been presented to the profession, either in this country or elsewhere.

The results of the operation were discussed under the following heads:—
1st. The rate per cent. of recovery after the operation, and the causes of death when it occurred. 2nd. The general condition of the patient after the operation. 3rd. The frequency of recurrence of stone after lithotritry.

The chief facts relative to the 184 cases were as follows: that they were consecutive in point of time, no case being omitted; that all were adults, and mostly of advanced age; that they included many individuals of very feeble health and constitution; that they were chiefly British, although several were from other nations. The mean age of the 184 cases was no less than sixty-one years. The youngest was twenty-two years old. Only three were below thirty years. The oldest was eighty-four years. There were forty-six cases of seventy years and upward. With very few exceptions, all stones of an ounce and upward were reserved for lithotomy. All obviously below that were crushed. Not one case was refused operation, not one was left unfinished, and in no instance was an operation of lithotritry completed by lithotomy. The recoveries, reckoning every kind of casualty following the operation, were 93 per cent.; but omitting five cases of death not by any means due to it, the mortality amounted to only 4 per cent. A second operation for recurrence of the stone was performed for 13 of the 184 cases; 122 were uric acid and the urates; 16 were mixed; 40 were phosphatic; 4 oxalate of lime; 1 cystic oxide; and 1 pure phosphate of lime.

* The paper is published *in extenso* in the *Med.-Chir. Trans.* vol. liii.


the important logical conclusion to be derived from the mass of facts considered was, that lithotrity is an eminently successful operation. For a certain number of cases, its success may be regarded as a certainty—absolutely without fear of any contingency, except such as attends the minor operations of surgery—for example, the opening of a small abscess, or the passing of a catheter. For the author stated that he had never lost a patient in the whole course of his experience after crushing a stone which was no larger than a walnut; and this he considered was a size at which, with few exceptions, any stone ought to be discovered. But this very fact led the author to remark that the success of lithotrity cannot therefore be considered apart from knowledge of the extent, in regard of the magnitude of the stone and the constitution of the patient, to which the capabilities of the operation have been limited. When it is employed for stones as large as a date, or a small chestnut—and it is impossible to deny the excellent chance of success which this method offers to the subjects of such stones—a certain, but still only small, portion of deaths must be expected. And the rate of mortality will correspond with augmentation in the size of the stone, and with the amount of pre-existing disease and age on the part of the patient. Given a small stone in a healthy person, and success is certain; the possibility of a fatal result in a case depending only on the presence of those remote and excessively unfavourable conditions which will make for an individual here and there the mere use of a catheter a cause of death.


The rule observed had been, for the most part, to apply lithotrity to all stones obviously less than an ounce in weight, easily discovered by sounding, and to operate on all larger ones by lithotomy.


Since this communication appeared, the number of Sir Henry Thompson's cases have reached 204, with one more death, making 13 deaths in the whole number.

CHARLES HAWKINS.

END OF THE FOURTH VOLUME.







M31 Holmes, T., ed. v. 4, 7788

H74 A system of surgery

v. 4

1870

NAME

DATE DUE

ATTIC

ATTIC